

||Jai Sri Gurudev||

S. J. C. INSTITUTE OF TECHNOLOGY, CHICKBALLAPUR

Department of Mathematics

Lecture Notes

Calculus & Linear Algebra (18MAT11)

Prepared By:

Purushotham.P

Assistant Professor

Email id: ppurushothamp48@gmail.com

SJC Institute of Technology

Phone: 8197481658


CALCULUS AND LINEAR ALGEBRA

MODULE - 01

POLAR CURVES AND EVALUATES

Introduction :-

Let \overrightarrow{Ox} be the initial line or initial ray, there is any point on the plane $P(r, \theta)$ with the radius of Vector $OP = r$ and $\angle xOP = \theta$ and $r = f(\theta)$ be a polar curve at the point




$P(r, \theta)$. Here, the coordinates of P is called as the Polar coordinates.

A. ANGLE BETWEEN RADIUS VECTOR AND TANGENT TO THE POLAR CURVE

Let \overrightarrow{Ox} be the initial line $P(r, \theta)$ be a point in the plane and $\overrightarrow{OP} = r$ be the

radius of vector and let $r = f(\theta)$ be a polar curve at the point 'P', 'T' be the tangent to the curve $r = f(\theta)$ at P and making angle with the initial line ψ



let ϕ be the angle between radius of vector and tangent to the curve $r = f(\theta)$ at P

W.K.T

$$\psi = \phi + \theta$$

$$\Rightarrow \tan \psi = \tan(\phi + \theta)$$

$$\tan \psi = \frac{\tan \phi + \tan \theta}{1 - \tan \phi \tan \theta} \rightarrow ①$$

also W.K.T The slope of the tangent T

$$\text{is } m = \tan \psi = \frac{dy}{dx} \rightarrow ②$$

let $x = r \cos \theta$ and $y = r \sin \theta$

differentiate x and y w.r.t θ

$$\therefore \frac{dx}{d\theta} = \frac{dr}{d\theta} \cos \theta - r \sin \theta$$

$$\frac{dy}{d\theta} = \frac{dr}{d\theta} \sin \theta + r \cos \theta$$

$$② \Rightarrow \tan \psi = \frac{dy}{dx} = \frac{dy/d\theta}{dx/d\theta}$$

$$\Rightarrow \tan \psi = \frac{\frac{dr}{d\theta} \sin \theta + r \cos \theta}{\frac{dr}{d\theta} \cos \theta - r \sin \theta}$$

$$= \frac{\frac{\frac{dr}{d\theta} \sin \theta}{\frac{dr}{d\theta} \cos \theta} + \frac{r \cos \theta}{\frac{dr}{d\theta} \cos \theta}}{1 - \frac{r \sin \theta}{\frac{dr}{d\theta} \cos \theta}}$$

$$\Rightarrow \tan \psi = \tan \theta + r \frac{d\theta}{dr} / 1 - r \frac{d\theta}{dr} \tan \theta$$

$$\frac{\tan \theta + \tan \phi}{1 - \tan \phi \tan \theta} = \frac{\tan \theta + r \frac{d\theta}{dr}}{1 - r \frac{d\theta}{dr} \tan \theta}$$


$$\boxed{\tan \phi = r \frac{d\theta}{dr}}$$

$$\phi = \tan^{-1} r \frac{d\theta}{dr}$$

In terms of, $\cot \phi = \frac{1}{r} \frac{dr}{d\theta}$

B. ANGLE BETWEEN TWO POLAR CURVES

Let $r = f_1(\theta)$, $r = f_2(\theta)$ be the given two polar curves having the tangents T_1 and T_2 and making an angles of radius of vector ϕ_1 and ϕ_2 then the angle between the




given two polar curves is the angle
between there two tangent = $|\phi_2 - \phi_1|$

WITH USUAL NOTATION, PROVE THAT

$$\frac{1}{P^2} = \frac{1}{r^2} + \frac{1}{r^4} \left(\frac{dr}{d\theta} \right)^2$$

Let \overrightarrow{Ox} be the initial line $r = f(\theta)$ be a polar curve at $P(r, \theta)$ and $r = OP = r$ be the radius of vector.



Let T be the tangent to the curve $r = f(\theta)$. At P and M is M the foot of the \perp of the pole $O(0,0)$ having the perpendicular distance $OM = p$ from the ΔOPM we have $\angle OMP = 90^\circ$.

$$\therefore \sin \phi = \frac{OM}{OP}$$

$$\Rightarrow \sin \phi = \frac{p}{r}$$

$$\Rightarrow P = r \sin \phi \quad \rightarrow ①$$

Square on both Sides

$$\Rightarrow P^2 = r^2 \sin^2 \phi$$

$$\Rightarrow \frac{1}{P^2} = \frac{1}{r^2 \sin^2 \phi}$$

$$\Rightarrow \frac{1}{P^2} = \frac{1}{r^2} \csc^2 \phi$$

$$\Rightarrow \frac{1}{P^2} = \frac{1}{r^2} (1 + \cot^2 \phi)$$

$$\Rightarrow \frac{1}{P^2} = \frac{1}{r^2} \left[1 + \left(\frac{1}{r} \left(\frac{dr}{d\theta} \right)^2 \right) \right]$$

$$\Rightarrow \frac{1}{P^2} = \frac{1}{r^2} \left[1 + \frac{1}{r^2} \left(\frac{dr}{d\theta} \right)^2 \right]$$

$$\Rightarrow \frac{1}{P^2} = \frac{1}{r^2} + \frac{1}{r^4} \left(\frac{dr}{d\theta} \right)^2$$

I. Find the angle between the radius vector and tangent for the following polar curves.

$$① r = a(1 - \cos \theta)$$

Given,

$$r = a(1 - \cos \theta) \rightarrow ①$$

differentiate ① w.r.t. θ

$$\frac{dr}{d\theta} = a(0 + \sin \theta)$$

$$\frac{dr}{d\theta} = a \sin \theta$$

$$\frac{1}{r} \frac{dr}{d\theta} = \frac{a \sin \theta}{r}$$

$$\frac{1}{r} \frac{dr}{d\theta} = \frac{a \sin \theta}{a(1 - \cos \theta)}$$

$$\Rightarrow \cot \phi = \frac{\sin \theta}{(1 - \cos \theta)}$$

$$\Rightarrow \cot \phi = \frac{2 \sin \theta/2 \cos \theta/2}{2 \sin^2 \theta/2}$$

$$\Rightarrow \cot \phi = \cot \theta/2$$

$$\boxed{\phi = \theta/2}$$

When, $\theta = \frac{\pi}{3}$

$$\phi = \frac{\pi/3}{2}$$

$$\phi = \frac{\pi}{6} = 30^\circ //$$

② $r = a(1 + \cos \theta)$

Given

$$r = a(1 + \cos \theta) \longrightarrow ①$$

Differentiate ① w.r.t. θ

$$\frac{dr}{d\theta} = a(-\sin \theta)$$

$$\frac{dr}{d\theta} = a(-\sin \theta)$$

$$\frac{1}{r} \frac{dr}{d\theta} = \frac{-a \sin \theta}{r}$$

$$\frac{1}{r} \frac{dr}{d\theta} = \frac{-a \sin \theta}{a(1 + \cos \theta)}$$

$$\frac{1}{r} \frac{dr}{d\theta} = \frac{-\sin \theta}{(1 + \cos \theta)}$$

$$\frac{1}{r} \frac{dr}{d\theta} = -\frac{2 \sin \theta/2 \cos \theta/2}{2 \cos^2 \theta/2}$$

$$\Rightarrow \cot \phi = -\tan \theta/2$$

$$\Rightarrow \cot \phi = \cot (\pi/2 + \theta/2)$$

$$\phi = \frac{\pi}{2} + \frac{\theta}{2} //$$

$$③ r = a(1 + \sin \theta)$$

Given

$$r = a(1 + \sin \theta) \rightarrow ①$$

Differentiate ① w.r.t. θ

$$① \Rightarrow \frac{dr}{d\theta} = a(0 + \cos \theta)$$

$$\Rightarrow \frac{dr}{d\theta} = a \cos \theta$$

$$\Rightarrow \frac{1}{r} \frac{dr}{d\theta} = \frac{a \cos \theta}{r}$$

$$\Rightarrow \frac{1}{r} \frac{dr}{d\theta} = \frac{a \cos \theta}{a(1 + \sin \theta)}$$

$$\Rightarrow \cot \phi = \frac{\cos \theta}{1 + \sin \theta}$$

$$\Rightarrow \cot \phi = \frac{\cos^2 \theta/2 - \sin^2 \theta/2}{\cos^2 \theta/2 + \sin^2 \theta/2 + 2 \sin \theta/2 \cos \theta/2}$$

$$= \frac{(\cos \theta/2 + \sin \theta/2)(\cos \theta/2 - \sin \theta/2)}{(\cos \theta/2 + \sin \theta/2)^2}$$

$$= \frac{\cos \theta/2 - \sin \theta/2}{\cos \theta/2 + \sin \theta/2}$$

$$\begin{aligned}
 &= \frac{1 - \tan \theta/2}{1 + \sin \theta/2 / \cos \theta/2} \\
 &= \frac{1 - \tan \theta/2}{1 + \tan \theta/2} \\
 &= \frac{\tan(\pi/4) - \tan(\theta/2)}{1 + \tan(\pi/4) \tan(\theta/2)}
 \end{aligned}$$

$$\Rightarrow \cot \phi = \tan\left(\frac{\pi}{4} - \frac{\theta}{2}\right)$$

$$\Rightarrow \cot \phi = \cot\left[\frac{\pi}{2} - \left(\frac{\pi}{4} - \frac{\theta}{2}\right)\right]$$

$$\Rightarrow \phi = \frac{\pi}{2} - \frac{\pi}{4} + \frac{\theta}{2}$$

$$\Rightarrow \phi = \frac{\pi}{4} + \frac{\theta}{2} //$$

II. Show that the following pairs of curves intersect each other orthogonally.

$$①. r = a(1 + \cos \theta), r = b(1 - \cos \theta)$$

Given

$$r = a(1 + \cos \theta) \rightarrow ①$$

Differentiate ① w.r.t. θ

$$\frac{dr}{d\theta} = -a \sin \theta$$

$$\frac{1}{r} \frac{dr}{d\theta} = \frac{-a \sin \theta}{r}$$

$$\Rightarrow \cot \phi_1 = \frac{-a \sin \theta}{a(1 + \cos \theta)}$$

$$\Rightarrow \cot \phi_1 = \frac{-\sin \theta}{1 + \cos \theta}$$

$$\Rightarrow \cot \phi_1 = \frac{-2 \sin \theta/2 \cos \theta/2}{2 \cos^2 \theta/2}$$

$$\Rightarrow \cot \phi_1 = -\tan \theta/2$$

$$\cot \phi = \cot \left(\frac{\pi}{2} + \frac{\theta}{2} \right)$$

$$\therefore \phi_1 = \frac{\pi}{2} + \frac{\theta}{2}$$

Similarly differentiate ② w.r.t. θ

$$\frac{dr}{d\theta} = b \sin \theta$$

$$\Rightarrow \frac{1}{r} \frac{dr}{d\theta} = \frac{b \sin \theta}{r}$$

$$\Rightarrow \cot \phi_2 = \frac{b \sin \theta}{b(1 - \cos \theta)}$$

$$\Rightarrow \cot \phi_2 = \frac{\sin \theta}{1 - \cos \theta}$$

$$\Rightarrow \cot \phi_2 = \frac{2 \sin \theta/2 \cos \theta/2}{2 \sin^2 \theta/2}$$

$$\therefore \cot \phi_2 = \theta/2$$

$$\therefore |\phi_1 - \phi_2| = \left| \frac{\pi}{2} + \frac{\theta}{2} - \frac{\theta}{2} \right| = \frac{\pi}{2} //$$

\therefore the Given Curves are intersecting orthogonally.

$$\textcircled{2} \quad r = a(1 + \sin \theta), \quad r = a(1 - \sin \theta)$$

Given

$$r = a(1 + \sin \theta) \longrightarrow \textcircled{1}$$

Differentiate \textcircled{1} w.r.t. \theta

$$\frac{dr}{d\theta} = a \cos \theta$$

$$\frac{1}{r} \frac{dr}{d\theta} = \frac{a \cos \theta}{r}$$

$$\Rightarrow \cot \phi_1 = \frac{a \cos \theta}{a(1 + \sin \theta)}$$

$$\Rightarrow \cot \phi_1 = \frac{\cos \theta}{(1 + \sin \theta)}$$

$$\Rightarrow \cot \phi_1 = \frac{\cos^2 \theta/2 - \sin^2 \theta/2}{\cos^2 \theta/2 + \sin^2 \theta/2 + 2 \sin \theta/2 \cos \theta/2}$$

$$= \frac{\cos \theta/2 - \sin \theta/2}{\cos \theta/2 + \sin \theta/2}$$

$$= \frac{1 - \tan \theta/2}{1 + \tan \theta/2}$$

$$\therefore \cot \phi_1 = \tan \left(\frac{\pi}{4} - \frac{\theta}{2} \right)$$

$$\Rightarrow \cot \phi_1 = \cot \left[\frac{\pi}{2} - \left(\frac{\pi}{4} - \frac{\theta}{2} \right) \right]$$

$$\phi_1 = \frac{\pi}{4} + \frac{\theta}{2}$$

Similarly differentiate \textcircled{2} w.r.t. \theta

$$\textcircled{2} \Rightarrow \frac{dr}{d\theta} = -a \cos \theta$$

$$\Rightarrow \frac{1}{r} \frac{dr}{d\theta} = \frac{-a \cos \theta}{a(1 - \sin \theta)}$$

$$\Rightarrow \cot \phi_2 = - \left[\frac{\cos^2 \theta/2 - \sin^2 \theta/2}{(\cos \theta/2 - \sin \theta/2)^2} \right]$$

$$\Rightarrow \cot \phi_2 = - \left[\frac{\cos \theta/2 + \sin \theta/2}{\cos \theta/2 - \sin \theta/2} \right]$$

$$= - \left[\frac{1 + \tan \theta/2}{1 - \tan \theta/2} \right]$$

$$= - \tan \left[\frac{\pi}{4} + \theta/2 \right]$$

$$\cot \phi_2 = \cot \left[\frac{\pi}{2} + \left(\frac{\pi}{4} + \frac{\theta}{2} \right) \right]$$

$$\therefore \phi_2 = \frac{\pi}{2} + \frac{\pi}{4} + \frac{\theta}{2}$$

$$\therefore |\phi_1 - \phi_2| = \left| \frac{\pi}{4} + \frac{\theta}{2} - \frac{\pi}{2} - \frac{\pi}{4} - \frac{\theta}{2} \right| = \frac{\pi}{2} //$$

\therefore The Given curves are intersecting orthogonally

$$\textcircled{3} \quad r^n = a^n (\cos n\theta), \quad r^n = b^n (\sin n\theta)$$

Given

$$r^n = a^n (\cos n\theta) \rightarrow \textcircled{1}$$

Differentiate $\textcircled{1}$ w.r.t θ

$$\Rightarrow n r^{n-1} \frac{dr}{d\theta} = -a^n (\sin n\theta)$$

$$\Rightarrow \frac{r^n}{r} \frac{dr}{d\theta} = -a^n \sin n\theta$$

$$\Rightarrow \frac{1}{r} \frac{dr}{d\theta} = \frac{-a^n \sin n\theta}{r^n}$$

$$\cot \phi_1 = \frac{-a^n \sin n\theta}{a^n \cos n\theta}$$

$$\cot \phi_1 = -\tan \theta$$

$$\cot \phi_1 = \cot \left(\frac{\pi}{2} + n\theta \right)$$

$$\phi_1 = \frac{\pi}{2} + n\theta$$

Similarly differentiate ② w.r.t θ

$$② \Rightarrow n r^{n-1} \frac{dr}{d\theta} = b^n n \cos \theta$$

$$\Rightarrow \frac{r^n}{r} \frac{dr}{d\theta} = b^n \cos n\theta$$

$$\frac{1}{r} \frac{dr}{d\theta} = \frac{b^n \cos n\theta}{r^n}$$

$$\Rightarrow \cot \phi_2 = \frac{b^n \cos n\theta}{r^n}$$

$$\cot \phi_2 = \cot n\theta$$

$$\phi_2 = n\theta$$

$$\therefore |\phi_1 - \phi_2| = \left| \frac{\pi}{2} + n\theta - n\theta \right| = \pi/2 //$$

\therefore the Given curves are intersecting orthogonally

III. Find the angle between the Given pairs of curves.

$$① r = \sin \theta + \cos \theta \text{ and } \gamma = 2 \sin \theta$$

Given :-

$$r = \sin \theta + \cos \theta \longrightarrow ①$$

$$r = 2 \sin \theta \rightarrow ②$$

differentiate equation ① w.r.t. θ

$$\text{①} \Rightarrow \frac{dr}{d\theta} = \cos \theta - \sin \theta$$

$$\Rightarrow \frac{1}{r} \frac{dr}{d\theta} = \frac{\cos \theta - \sin \theta}{r}$$

$$\Rightarrow \frac{1}{r} \frac{dr}{d\theta} = \frac{\cos \theta - \sin \theta}{\sin \theta + \cos \theta}$$

$$= \frac{1 - \tan \theta}{1 + \tan \theta}$$

$$= \frac{\tan(\pi/4) - \tan \theta}{\tan(\pi/4) + \tan \theta}$$

$$\Rightarrow \cot \phi_1 = \tan(\pi/4 - \theta)$$

$$\Rightarrow \cot \phi_1 = \cot \left[\frac{\pi}{2} - [\frac{\pi}{4} - \theta] \right]$$

$$\Rightarrow \phi_1 = \frac{\pi}{2} - \frac{\pi}{4} + \theta$$

$$\text{III}^{\text{q}} ② \Rightarrow \frac{dr}{d\theta} = 2 \cos \theta$$

$$\Rightarrow \frac{1}{r} \frac{dr}{d\theta} = \frac{2 \cos \theta}{r}$$

$$\Rightarrow \frac{1}{r} \frac{dr}{d\theta} = \frac{2 \cos \theta}{2 \sin \theta}$$

$$\Rightarrow \cot \phi_2 = \cot \theta$$

$$\Rightarrow \phi_2 = \theta$$

$$\therefore |\phi_1 - \phi_2| = \left| \frac{\pi}{2} - \frac{\pi}{4} + \theta - \theta \right|$$

$$\Rightarrow |\phi_1 - \phi_2| = \pi/4 //$$

\therefore The Given pairs of Curves are intersecting at
 $\frac{\pi}{4}$ [45°]

② $r = a \log \theta$ and $r = \frac{a}{\log \theta}$

Given

$$r = a \log \theta \rightarrow ①$$

$$r = \frac{a}{\log \theta} \rightarrow ②$$

differentiate equation ① and ② w.r.t. θ

$$① \Rightarrow \frac{dr}{d\theta} = a \cdot \left(\frac{1}{\theta}\right)$$

$$\Rightarrow \frac{dr}{d\theta} = \frac{a}{\theta}$$

$$\Rightarrow \frac{d\theta}{dr} = \frac{\theta}{a}$$

$$\Rightarrow r \frac{d\theta}{dr} = \frac{\theta}{a}$$

$$\Rightarrow \tan \phi_1 = \frac{(a \log \theta) \theta}{a}$$

$$\Rightarrow \tan \phi_1 = \theta \log \theta \rightarrow ③$$

IIIrd differentiate equation ② w.r.t θ

$$② \Rightarrow \frac{dr}{d\theta} \log \theta + \frac{r}{\theta} = 0$$

$$\Rightarrow \log \theta \frac{dr}{d\theta} = -\frac{r}{\theta} = 0$$

$$\Rightarrow \frac{1}{r} \frac{dr}{d\theta} = -\frac{1}{\theta \log \theta}$$

$$\Rightarrow r \frac{d\theta}{dr} = -\theta \log \theta$$

$$\Rightarrow \tan \phi_2 = -\theta \log \theta \rightarrow ④$$

W. K. T

$$\tan(\phi_1 - \phi_2) = \frac{\tan \phi_1 - \tan \phi_2}{1 + \tan \phi_1 \tan \phi_2}$$

$$\Rightarrow \tan(\phi_1 - \phi_2) = \frac{\theta \log \theta + \theta \log \theta}{1 + (\theta \log \theta)(-\theta \log \theta)}$$

$$\Rightarrow \tan(\phi_1 - \phi_2) = \frac{2 \theta \log \theta}{1 - \theta^2 (\log \theta)^2} \rightarrow ⑤$$

∴ from equation ① and ②

$$a \log \theta = \frac{a}{\log \theta}$$

$$\Rightarrow (\log \theta)^2 = 1$$

$$\Rightarrow \log e^\theta = 1$$

$$\Rightarrow \theta = e^1 = e$$

$$\therefore ⑤ \Rightarrow \tan(\phi_1 - \phi_2) = \frac{2e}{1-e^2}$$

$$\therefore |\phi_1 - \phi_2| = \tan^{-1} \left(\frac{2e}{1-e^2} \right) = 2 \tan^{-1} e //$$

∴ The given two curves or pair of curves
are intersecting at $a \tan^{-1} e$.

③ $r^2 \sin 2\theta = 4$ and $r^2 = 16 \sin 2\theta$

Given

$$r^2 \sin 2\theta = 4 \quad \rightarrow \textcircled{1} \quad \leftarrow \quad r^2 = \frac{4}{\sin 2\theta}$$
$$r^2 = 16 \sin 2\theta \quad \rightarrow \textcircled{2}$$

∴ from equation ① and ②

$$\frac{4}{\sin 2\theta} = 16 \sin 2\theta$$

$$\Rightarrow 4 \sin^2 2\theta = 1$$

$$\Rightarrow \sin^2 2\theta = \frac{1}{4}$$

$$\Rightarrow \sin 2\theta = \frac{1}{2}$$

$$\Rightarrow \sin 2\theta = \frac{1}{2}$$

$$\Rightarrow 2\theta = \sin^{-1}(1/2)$$

$$\Rightarrow 2\theta = \frac{\pi}{6}$$

$$\Rightarrow \theta = \frac{\pi}{12}$$

Differentiate equation ① and ② w.r.t. θ

$$\textcircled{1} \Rightarrow 2r \frac{dr}{d\theta} \sin 2\theta + r^2 \cos(2\theta) \cdot 2 = 0$$

$$\Rightarrow r \frac{dr}{d\theta} \cdot \sin 2\theta + r^2 \cos 2\theta = 0$$

$$\Rightarrow r \frac{dr}{d\theta} \cdot \sin 2\theta = -r^2 \cos 2\theta$$

$$\Rightarrow r \frac{dr}{d\theta} = -\frac{r^2 \cos 2\theta}{\sin \theta}$$

$$\Rightarrow \frac{1}{r} \frac{dr}{d\theta} = -\cot 2\theta$$

$$\Rightarrow \cot \phi_1 = \cot (-2\theta)$$

$$\Rightarrow \phi_1 = -2\theta$$

$$\textcircled{2} \Rightarrow 2r \cdot \frac{dr}{d\theta} = 16 (\cos 2\theta) (2)$$

$$\Rightarrow r \frac{dr}{d\theta} = 16 \cos 2\theta$$

$$\Rightarrow \frac{1}{r} \frac{dr}{d\theta} = \frac{16 \cos 2\theta}{r^2}$$

$$\Rightarrow \cot \phi_2 = \frac{16 \cos 2\theta}{16 \sin 2\theta}$$

$$\Rightarrow \cot \phi_2 = \cot 2\theta$$

$$\Rightarrow \phi_2 = 2\theta$$

$$\therefore |\phi_1 - \phi_2| = |-2\theta - 2\theta| = 4\theta$$

$$\therefore |\phi_1 - \phi_2| = 4\left(\frac{\pi}{12}\right) = \frac{\pi}{3} //$$

∴ The given pair of curves are intersecting

at $\frac{\pi}{3}$

$$\textcircled{4} \quad r = a(1 - \cos \theta) \quad \text{and} \quad r = 2a \cos \theta$$

Given,

$$r = a(1 - \cos \theta) \rightarrow \textcircled{1}$$

$$r = 2a \cos \theta \rightarrow ②$$

differentiate equation ① and ② w.r.t. θ

$$① \Rightarrow \frac{dr}{d\theta} = a(-\sin \theta)$$

$$\Rightarrow \frac{dr}{d\theta} = a \sin \theta$$

$$\Rightarrow \frac{1}{r} \frac{dr}{d\theta} = \frac{a \sin \theta}{r}$$

$$\Rightarrow \frac{1}{r} \frac{dr}{d\theta} = \frac{a \sin \theta}{a(1 - \cos \theta)}$$

$$\Rightarrow \frac{1}{r} \frac{dr}{d\theta} = \frac{\sin \theta}{(1 - \cos \theta)}$$

$$\Rightarrow \cot \phi_1 = \frac{2 \sin \theta/2 \cos \theta/2}{2 \sin^2 \theta/2}$$

$$\Rightarrow \cot \phi_1 = \frac{\cos \theta/2}{\sin \theta/2}$$

$$\Rightarrow \cot \phi_1 = \cot \theta/2 \quad \therefore \phi_1 = \theta/2$$

III^{ly} ② d. w. r. t. θ

$$\frac{dr}{d\theta} = -2a \sin \theta$$

$$\Rightarrow \frac{1}{r} \frac{dr}{d\theta} = -\frac{2a \sin \theta}{2a \cos \theta}$$

$$\Rightarrow \frac{1}{r} \frac{dr}{d\theta} = -\tan \theta$$

$$\Rightarrow \cot \phi_2 = -\tan \theta$$

$$\Rightarrow \cot \phi_2 = \cot(\pi/2 + \theta)$$

$$\phi_2 = \frac{\pi}{2} + \theta$$

$$\therefore |\phi_1 - \phi_2| = \left| \frac{\theta}{2} - \frac{\pi}{2} - \theta \right|$$

$$|\phi_1 - \phi_2| = \left| \frac{\pi}{2} + \frac{\theta}{2} \right|$$

From ① and ②

$$a(1-\cos\theta) = 2a\cos\theta$$

$$\Rightarrow (1-\cos\theta) = 2\cos\theta$$

$$\Rightarrow (1-\cos\theta) = 2\cos\theta$$

$$\Rightarrow 3\cos\theta = 1$$

$$\Rightarrow \cos\theta = 1/3$$

$$\Rightarrow \theta = \cos^{-1}(1/3)$$

$$\therefore |\phi_1 - \phi_2| = \left| \pi/2 + 1/2 \cos^{-1}(1/3) \right| //$$

∴ The Given two curves or pair of curves
are intersecting at $\left| \pi/2 + 1/2 \cos^{-1}(1/3) \right|$

⑤ $r = \frac{a}{1+\cos\theta}$ and $r = \frac{b}{1-\cos\theta}$

Given,

$$r = \frac{a}{1+\cos\theta} \text{ or } r(1+\cos\theta) = a \rightarrow ①$$

① d w. r t θ

$$(1 + \cos\theta) \frac{dr}{d\theta} - r \sin\theta = 0$$

$$\Rightarrow \frac{1}{r} \frac{dr}{d\theta} = \frac{\sin\theta}{1 + \cos\theta}$$

$$\Rightarrow \cot\phi_1 = \frac{2\sin\theta/2\cos\theta/2}{2\cos^2\theta/2}$$

$$\Rightarrow \cot\phi_1 = \tan\theta/2$$

$$\Rightarrow \cot\phi_1 = \cot(\pi/2 - \theta/2)$$

$$\therefore \phi_1 = (\pi/2 - \theta/2)$$

Similarly differentiate ② w. r t. θ

$$r = \frac{b}{1 - \cos\theta} \text{ or } r(1 - \cos\theta) = b \rightarrow ②$$

$$\Rightarrow \frac{1}{r} \frac{dr}{d\theta} = \frac{-\sin\theta}{1 - \cos\theta} = -\cot\theta/2$$

$$\Rightarrow \cot\phi_2 = -\frac{\theta}{2}$$

$$\therefore |\phi_2 - \phi_1| = \left| -\frac{\theta}{2} - \frac{\pi}{2} + \frac{\theta}{2} \right| = \frac{\pi}{2}$$

\therefore The Given pair of curves are intersecting at the angle of $\frac{\pi}{2}$ or 90° .

IV. Find the pedal equation for the following polar curves.

$$\textcircled{1} \quad r = a e^{\theta \cot \alpha}$$

Let,

$$r = a e^{\theta \cot \alpha} \longrightarrow \textcircled{1}$$

\textcircled{1} equation differentiate w.r.t. \theta

$$\textcircled{1} \Rightarrow \frac{dr}{d\theta} = a \cdot e^{\theta \cot \alpha} \cdot \cot \alpha$$

$$\Rightarrow \frac{dr}{d\theta} = (a \cot \alpha) e^{\theta \cot \alpha}$$

$$\Rightarrow \frac{1}{r} \frac{dr}{d\theta} = \frac{(a \cot \alpha) e^{\theta \cot \alpha}}{r}$$

$$\Rightarrow \cot \phi = \frac{(a \cot \alpha) e^{\theta \cot \alpha}}{a e^{\theta \cot \alpha}}$$

$$\Rightarrow \cot \phi = \cot \alpha$$

$$\Rightarrow \phi = \alpha$$

w. k. t. the pedal equation

$$P = r \sin \phi$$

$$\Rightarrow P = r \sin \alpha$$

$$\textcircled{2} \quad r^n = a^n \cos n\theta$$

Given.

$$r^n = a^n \cos n\theta \longrightarrow \textcircled{1}$$

\textcircled{1} equation differentiate w.r.t. \theta

$$\textcircled{1} \Rightarrow n r^{n-1} \frac{dr}{d\theta} = a^n (-\sin n\theta)^n$$

$$\Rightarrow \frac{r^n}{r} \frac{dr}{d\theta} = -a^n \sin n\theta$$

$$\Rightarrow \frac{1}{r} \frac{dr}{d\theta} = -\frac{a^n \sin n\theta}{r^n}$$

$$\Rightarrow \cot \phi = \frac{-a^n \sin n\theta}{a^n \cos n\theta}$$

$$\Rightarrow \cot \phi = -\tan n\theta$$

W.K.T The pedal equation

$$\frac{1}{P^2} = \frac{1}{r^2} (1 + \cot^2 \phi)$$

$$\Rightarrow \frac{1}{P^2} = \frac{1}{r^2} [1 + \tan^2 n\theta]$$

$$\Rightarrow \frac{1}{P^2} = \frac{1}{r^2} \sec^2 n\theta$$

$$\Rightarrow \frac{1}{P^2} = \frac{1}{r^2 \cos^2 n\theta}$$

$$\Rightarrow P^2 = r^2 \cos^2 n\theta$$

$$\Rightarrow P = r \cos n\theta$$

$$\Rightarrow P = r \left(\frac{r^n}{a^n} \right) (\because ①)$$

$$\Rightarrow P = \frac{r^n + 1}{a^n}$$

$$\Rightarrow a^n P = r^{n+1}$$

$$③ \frac{2a}{r} = 1 + \cos \theta$$

Given.

$$\frac{\alpha}{r} = 1 + \cos\theta \quad \text{or} \quad \alpha = r(1 + \cos\theta) \rightarrow ①$$

② equation differentiate w.r.t. θ

$$① \Rightarrow (1 + \cos\theta) \frac{dr}{d\theta} + r(-\sin\theta) = 0$$

$$\Rightarrow (1 + \cos\theta) \frac{dr}{d\theta} - r \sin\theta = 0$$

$$\Rightarrow \frac{1}{r} \frac{dr}{d\theta} = \frac{\sin\theta}{1 + \cos\theta}$$

$$\Rightarrow \cot\phi = \frac{2 \sin\theta/2 \cos\theta/2}{2 \cos^2\theta/2}$$

$$\Rightarrow \cot\phi = \tan\theta/2$$

∴ W.K.T The pedal equation

$$\frac{1}{P^2} = \frac{1}{r^2} (1 + \cot^2\phi)$$

$$\Rightarrow \frac{1}{P^2} = \frac{1}{r^2} (1 + \tan^2\theta/2)$$

$$\Rightarrow \frac{1}{P^2} = \frac{1}{r^2} (\sec^2\theta/2)$$

$$\Rightarrow \frac{1}{P^2} = \frac{1}{r^2 \cos^2\theta/2}$$

$$\Rightarrow P^2 = r^2 \cos^2\theta/2$$

$$\Rightarrow P^2 = r^2 \left(\frac{1 + \cos\theta}{2} \right)$$

$$\Rightarrow 2P^2 = r^2 (1 + \cos\theta)$$

$$\Rightarrow \dot{r}^2 = r^2 \left(\frac{\dot{\theta}}{\theta} \right)$$

$$\Rightarrow \dot{r}^2 = \alpha r$$

$$④ r^m = a^m (\cos m\theta + \sin m\theta)$$

Given

$$r^m = a^m [\cos m\theta + \sin m\theta] \rightarrow ①$$

① Differentiate w.r.t. θ

$$① \Rightarrow m r^{m-1} \frac{dr}{d\theta} = a^m [-\sin m\theta(m) + \cos m\theta(m)]$$

$$\Rightarrow \frac{r^m}{r} \frac{dr}{d\theta} = \frac{a^m (\cos m\theta - \sin m\theta)}{r^m}$$

$$\Rightarrow \cot \phi = \frac{a^m (\cos m\theta - \sin m\theta)}{a^m (\cos m\theta + \sin m\theta)}$$

$$\Rightarrow \cot \phi = \frac{\cos m\theta - \sin m\theta}{\cos m\theta + \sin m\theta}$$

$$\therefore 1 + \cot^2 \phi = \frac{1 + (\cos m\theta - \sin m\theta)^2}{(\cos m\theta + \sin m\theta)^2}$$

$$\Rightarrow 1 + \cot^2 \phi = \frac{(\cos m\theta + \sin m\theta)^2 + (\cos m\theta - \sin m\theta)^2}{(\cos m\theta + \sin m\theta)^2}$$

$$\Rightarrow 1 + \cot^2 \phi = \frac{2(\cos^2 m\theta + \sin^2 m\theta)}{\left(\frac{r^m}{a^m}\right)^2}$$

$$\Rightarrow 1 + \cot^2 \phi = \frac{2}{r^{2m}/a^{2m}}$$

$$\Rightarrow 1 + \cot^2 \phi = \frac{2a^{2m}}{r^{2m}}$$

∴ the pedal equation

$$\frac{1}{p^2} = \frac{1}{r^2} (1 + \cot^2 \phi)$$

$$\Rightarrow \frac{1}{p^2} = \frac{1}{r^2} \cdot \frac{2a^{2m}}{r^{2m}}$$

$$\Rightarrow \frac{1}{p^2} = \frac{2a^{2m}}{r^{2m+2}}$$

$$\Rightarrow 2a^{2m} p^2 = r^{2m+2}$$

$$⑤ \frac{l}{r} = 1 + e \cos \theta$$

Given,

$$\frac{l}{r} = 1 + e \cos \theta \rightarrow ①$$

differentiate w.r.t. θ

$$① \Rightarrow (1 + e \cos \theta) \frac{dr}{d\theta} + r(0 - e \sin \theta) = 0$$

$$\Rightarrow 1 + e \cos \theta \frac{dr}{d\theta} = r e \sin \theta$$

$$\Rightarrow \frac{1}{r} \frac{dr}{d\theta} = \frac{e \sin \theta}{1 + e \cos \theta} = \cot \phi$$

$$\therefore 1 + \cot^2 \phi = \frac{1 + e^2 \sin^2 \theta}{(1 + e \cos \theta)^2}$$

$$= \frac{(1 + e \cos \theta)^2 + e^2 \sin^2 \theta}{(1 + e \cos \theta)^2}$$

$$= \frac{1 + 2e \cos \theta + e^2 \cos^2 \theta + e^2 \sin^2 \theta}{(1 + e \cos \theta)^2}$$

$$= \frac{1 + 2e \cos \theta + e^2(1)}{(1 + e \cos \theta)^2}$$

$$= \frac{1 + 2e \cos \theta + e^2}{(1 + e \cos \theta)^2}$$

$$\Rightarrow 1 + \cot^2 \phi = \frac{1 + e^2 + 2\left(\frac{l}{r} - 1\right)}{\left(\frac{l}{r}\right)^2}$$

$$= \frac{1 + e^2 + \frac{2l}{r} - 2}{\frac{l^2}{r^2}}$$

$$= \frac{e^2 + \frac{2l}{r} - 1}{\frac{l^2}{r^2}}$$

$$1 + \cot^2 \phi = \frac{r^2(e^2 + 2l/r - 1)}{l^2}$$

\therefore the pedal equation

$$\frac{1}{p^2} = \frac{1}{r^2} (1 + \cot^2 \phi)$$

$$\Rightarrow \frac{1}{p^2} = \frac{1}{r^2} \cdot r^2 \left(\frac{e^2 + 2l/r - 1}{l^2} \right)$$

$$\boxed{\frac{1}{p^2} = \frac{e^2 + 2l/r - 1}{l^2}} //$$

Radius of Curvature

Generally the Curvature of any curve can be denoted by 'k' and the reciprocal of the curvature will be called as the radius of curvature and it will be denoted

as,

$$\rho = \frac{1}{K}$$

Note:-

- the radius of curvature can be evaluated in the Cartesian form by the following formula.

$$\rho = \frac{(1+y_1^2)^{3/2}}{y_2}, y_2 \neq 0$$

where $y_1 = \frac{dy}{dx}$, $y_2 = \frac{d^2y}{dx^2}$ at any point P

In other way $\rho = \frac{(1+x_1^2)^{3/2}}{x_2}, x_2 \neq 0$

where $x_1 = \frac{dx}{dy}$, $x_2 = \frac{d^2x}{dy^2}$ at any point P

- The radius of curvature for the polar curve can be evaluated as,

$\rho = r \frac{dr}{dp}$, where $f(p, r, c)$ is the pedal equation.

V

- Find the radius of curvature of the curve $y = a \log(\sec(x/a))$ at any point

Given,

$$y = a \log(\sec(x/a))$$

differentiate ① w.r.t. x

$$\text{①} \Rightarrow \frac{dy}{dx} = a \cdot \frac{1}{\sec(x/a)} \sec(x/a) \tan(x/a) \cdot 1/a$$

$$\Rightarrow \frac{dy}{dx} = \tan(x/a) = y_1 \rightarrow ②$$

differentiate ② w.r.t. x

$$② \Rightarrow \frac{d^2y}{dx^2} = \sec^2(x/a) \cdot 1/a$$

$$\Rightarrow y_2 = \frac{d^2y}{dx^2} = \frac{1}{a} \sec^2(x/a)$$

∴ W.K.T

$$\rho = \frac{(1 + y_1^2)^{3/2}}{y^2}$$

$$\Rightarrow \rho = \frac{(1 + \tan^2(x/a))^{3/2}}{1/a \sec^2(x/a)}$$

$$\Rightarrow \rho = \frac{a (\sec^2(x/a))^{3/2}}{\sec^2(x/a)}$$

$$\Rightarrow \rho = \frac{a \sec^3(x/a)}{\sec^2(x/a)}$$

$$\Rightarrow \boxed{\rho = a \sec(x/a)}$$

2. Find the radius of the curvature of the curve $x^3 + y^3 = 3axy$ at the point $P(3a/2, 3a/2)$.

Given,

$$x^3 + y^3 = 3axy \rightarrow ①$$

differentiate w.r.t. x

$$① \Rightarrow 3x^2 + 3y^2 \frac{dy}{dx} = 3a [1 \cdot y + x \frac{dy}{dx}]$$

$$\Rightarrow x^2 + y^2 y_1 = a[y + xy_1]$$

$$\Rightarrow x^2 + y^2 y_1 - ay - axy_1 = 0$$

$$\Rightarrow (y^2 - ax)y_1 + (x^2 - ay) = 0$$

$$\Rightarrow (y^2 - ax)y_1 + (x^2 - ay) = 0$$

$$\Rightarrow (y^2 - ax)y_1 = ay - x^2$$

$$\Rightarrow y_1 = \frac{ay - x^2}{y^2 - ax} \longrightarrow ②$$

$$\therefore (y_1)_P = \frac{a\left(\frac{3a}{2}\right) - \left(\frac{3a}{2}\right)^2}{\left(\frac{3a}{2}\right)^2 - a\left(\frac{3a}{2}\right)} = -\frac{\left(\frac{3a}{2}\right)^2 - a\left(\frac{3a}{2}\right)}{\left(\frac{3a}{2}\right)^2 - a\left(\frac{3a}{2}\right)} = -1$$

differentiate ② w.r.t x

$$② \Rightarrow y_2 = \frac{d^2 y}{dx^2} = \frac{(y^2 - ax)(ay_1 - 2x) - (ay - x^2)(2yy_1 - a)}{(y^2 - ax)^2}$$

$$(y_2)_P = \frac{\left[\left(\frac{3a}{2}\right)^2 - 4\left(\frac{3a}{2}\right)\right]a(-1) - 2\left(\frac{3a}{2}\right) - 4\left(\frac{3a}{2}\right) - \left(\frac{3a}{2}\right)^2}{\left(\frac{3a}{2}\right)^2 - a\left(\frac{3a}{2}\right)^2}$$

$$\Rightarrow y_2 = \frac{\left(\frac{3a^2}{4}\right)(-4a) - \left(-\frac{3a^2}{4}\right)(-4a)}{\left(\frac{3a^2}{4}\right)^2}$$

$$\Rightarrow y_2 = \frac{-3a^3 - 3a^3}{\frac{9a^4}{16}}$$

$$\Rightarrow y_2 = -\frac{6a^3 \times 16}{9a^4} = -\frac{32}{3a}$$

$$\therefore \rho = \left| \frac{(1+y_1^2)^{3/2}}{y_2} \right| = \left| \frac{(1+(-1)^2)^{3/2}}{-32/3a} \right|$$

$$\Rightarrow \rho = \frac{2^{3/2}}{32/3a}$$

$$\Rightarrow \rho = \frac{2 \cdot 2^{1/2} \times 3a}{32}$$

$$\Rightarrow \rho = \frac{3\sqrt{2}a}{16}$$

$$\Rightarrow \rho = \frac{3\sqrt{2} \cdot a}{8 \times \sqrt{2} \times \sqrt{2}}$$

$$\boxed{\Rightarrow \rho = \frac{3a}{8\sqrt{2}}}$$

- ③ Find the radius of curvature of the curve $a^2y = x^2 - a^2$ at the point where the curve meets at x -axis.

Given curve $a^2y = x^2 - a^2$ and the curve meet the x axis at the point 'P' and its y coordinate becomes zero.

\therefore the required point $P(a, 0)$

differentiate equation w.r.t x

$$a^2y = x^2 - a^2 \longrightarrow ①$$

$$\textcircled{1} \Rightarrow a^2 y_1 = 2x - 0$$

$$\Rightarrow y_1 = \frac{2x}{a^2} \rightarrow \textcircled{2}$$

$$\therefore (y_1)_p = \frac{2(a)}{a^2} = \frac{2}{a} \rightarrow \textcircled{3}$$

Differentiate $\textcircled{3}$ w.r.t x

$$\textcircled{3} \Rightarrow y_2 = \frac{d^2y}{dx^2} = \frac{2}{a^2}$$

$$(y_2)_p = \frac{2}{a^2} \neq 0$$

$$\rho = \left| \frac{(1-y_1^2)^{3/2}}{y_2} \right|$$

$$\rho = \frac{(1+(2/a)^2)^{3/2}}{2/a^2}$$

$$\Rightarrow \rho = \frac{(1+4/a^2)^{3/2}}{2/a^3}$$

$$\Rightarrow \rho = \frac{a^2 \left(\frac{a^2+4}{a^2} \right)^{3/2}}{2}$$

$$\Rightarrow \rho = \frac{a^2 (a^2 + 4)^{3/2}}{a^3}$$

$$\Rightarrow \boxed{\rho = \frac{(a^2 + 4)^{3/2}}{2a}}$$

④ Find the radius of the curvature of the curve $a^2y = x^3 - a^3$ and meets at x axis.

Given,

$$a^2y = x^3 - a^3 \rightarrow ①$$

differentiate w.r.t. x

$$① \Rightarrow a^2y_1 = 3x^2 \rightarrow ②$$

or

$$y_1 = \frac{3x^2}{a^2} \rightarrow ②$$

$$\therefore (y_1)_p = \frac{3(a^2)}{a^2} = 3$$

differentiate ② w.r.t x

$$② \Rightarrow y_2 = \frac{d^2y}{dx^2} = \frac{6x}{a^3}$$

$$(y_2)_p = \frac{6(a^2)}{a^3} = \frac{6}{a}$$

$$\Rightarrow \rho = \left| \frac{(1+y_1^2)^{3/2}}{y_2} \right|$$

$$\Rightarrow \rho = \frac{(1+3^2)^{3/2}}{6/a}$$

$$\Rightarrow \rho = \frac{(1+9)^{3/2}}{6/a}$$

$$\Rightarrow \rho = \frac{a \cdot 10^{3/2}}{6}$$

$$\Rightarrow \rho = \frac{a \cdot 10 \cdot 10^{1/2}}{6}$$

$$\Rightarrow \boxed{\rho = \frac{5\sqrt{10} \cdot a}{3}}$$

⑤ Find the radius of the curvature of the curve $x^2y = a(x^2 + y^2)$ at Point P(-2a, 2a)

Given,

$$x^2y = a(x^2 + y^2) \quad \dots \textcircled{1}$$

differentiate ① w.r.t y

$$\textcircled{1} \Rightarrow x^2(1) + y \frac{dx}{dy} = a \left(2x \frac{dx}{dy} + 2y \right)$$

$$\Rightarrow x^2 + 2xy \cdot x_1 = 2ax_1 + 2ay$$

$$\Rightarrow 2xy \cdot x_1 - 2ax \cdot x_1 = 2ay - x^2$$

$$\Rightarrow 2x x_1 (y-a) = 2ay - x^2$$

$$\Rightarrow x_1 = \frac{2ay - x^2}{2x(y-a)} = \frac{2ay - x^2}{2xy - 2ax}$$

$$\therefore (x_1)_P = \frac{2a(2a) - (-2a)^2}{2(-2a)(2a-a)}$$

$$\Rightarrow (x_1)_P = 0$$

differentiate ② w.r.t y

$$\textcircled{2} \Rightarrow x_2 = \frac{d^2x}{dy^2} = \frac{(2xy - 2ax)(2a - 2x \cdot x_1) - (2a - 2x \cdot x_1)}{- (2ay - x^2) [2x + 2x_1 y - 2ax_1]} \cdot \frac{1}{[2x(y-a)]^2}$$

$$\therefore (x_2)_p = \frac{[-8a^2 + 4a^2][2a - 0] - 0}{[-4a(a)]^2}$$

$$\Rightarrow x_2 = -\frac{8a^3}{16a^4}$$

$$\Rightarrow x_2 = -\frac{1}{2a} \neq 0$$

$$\therefore p = \left| \frac{(1+x_1^2)^{3/2}}{x_2} \right| = \left| \frac{(1+0)^{3/2}}{-1/2a} \right| = 2a //$$

⑥ Find the radius of curvature of the polar curve $r = a(1 + \cos \theta)$

Given,

$$r = a(1 + \cos \theta) \rightarrow ①$$

differentiate ① w.r.t θ

$$① \Rightarrow \frac{dr}{d\theta} = a(0 - \sin \theta) = -a \sin \theta$$

$$\Rightarrow \frac{1}{r} \frac{dr}{d\theta} = \frac{-a \sin \theta}{a(1 + \cos \theta)}$$

$$\Rightarrow \cot \phi = \frac{-2 \sin \theta / 2 \cos \theta / 2}{2 \cos^2 \theta / 2}$$

$$\Rightarrow \cot \phi = -\tan(\theta/2)$$

w.k.t the pedal equation

$$\frac{1}{P^2} = \frac{1}{r^2} (1 + \cot^2 \phi)$$

$$\Rightarrow \frac{1}{P^2} = \frac{1}{r^2} (1 + \tan^2 \theta/2)$$

$$\Rightarrow \frac{1}{P^2} = \frac{1}{r^2} \sec^2 \theta/2$$

$$\Rightarrow \frac{1}{P^2} = \frac{1}{r^2 \cos^2 \theta/2}$$

$$\Rightarrow P^2 = r^2 \cos^2 \theta/2$$

$$\Rightarrow P^2 = r^2 \left(\frac{1 + \cos \theta}{2} \right)$$

$$\Rightarrow P^2 = \frac{r^2}{2} \left(\frac{\gamma}{\alpha} \right)$$

$$\Rightarrow 2\alpha P^2 = \gamma^3 \rightarrow ②$$

Differentiate ② w.r.t. P

$$② \Rightarrow 2\alpha \cdot 2P = 3\gamma^2 \frac{dP}{dP}$$

$$\Rightarrow 4\alpha P = 3\gamma^2 \frac{dr}{dp}$$

$$\Rightarrow \frac{dr}{dp} = \frac{4\alpha P}{3\gamma^2}$$

$$\Rightarrow \gamma \frac{dr}{dp} = \frac{4\alpha P}{3\gamma^2}$$

$$\Rightarrow \gamma \frac{dr}{dp} = \frac{4\alpha P}{3\gamma}$$

$$\Rightarrow \gamma = \frac{4\alpha P}{3\gamma}$$

$$\Rightarrow \rho^2 = \frac{16a^2 p^2}{q\tau^2}$$

$$\Rightarrow \rho^2 = \frac{16a^2}{q} \cdot \frac{1}{r^2} \left(\frac{\tau^3}{qa} \right)$$

$$\Rightarrow \rho^2 = \left(\frac{8a}{q} \right) r$$

$$\Rightarrow \rho^2 \propto r$$

- ⑦ Show that for the curve $\tau(1 - \cos\theta) = 2a$
or $\frac{2a}{r} = (1 - \cos\theta)$, and ρ^2 varies as τ^3
[$\rho^2 \propto \tau^3$]

Given,

$$\tau(1 - \cos\theta) = 2a \rightarrow ②$$

Differentiate ① w.r.t. θ

$$① \Rightarrow (1 - \cos\theta) \frac{d\tau}{d\theta} + r(\theta + \sin\theta) = 0$$

$$\Rightarrow (1 - \cos\theta) \frac{dr}{d\theta} = -\tau \sin\theta$$

$$\Rightarrow \frac{1}{r} \frac{dr}{d\theta} = \frac{-\sin\theta}{1 - \cos\theta}$$

$$\Rightarrow \cot\phi = \frac{-2\sin\theta/2 \cos\theta/2}{2\sin^2\theta/2}$$

$$\Rightarrow \cot\phi = -\cot(\theta/2)$$

∴ W.K.T the pedal eqn.

$$\frac{1}{P^2} = \frac{1}{r^2} (1 + \omega t^2 \phi)$$

$$\Rightarrow \frac{1}{P^2} = \frac{1}{r^2} (1 + \omega t^2 \Theta/2)$$

$$\Rightarrow \frac{1}{P^2} = \frac{1}{r^2} (\cosec^2 \Theta/2)$$

$$\Rightarrow \frac{1}{P^2} = \frac{1}{r^2 \sin^2 \Theta/2}$$

$$\Rightarrow P^2 = r^2 \sin^2 \Theta/2$$

$$\Rightarrow P^2 = r^2 \left(1 - \frac{\cos \theta}{2} \right)$$

$$\Rightarrow P^2 = \frac{r^2}{2} \left(\frac{2\alpha}{\gamma} \right)$$

$$\Rightarrow P^2 = ar \longrightarrow ②$$

Differentiate ② w.r.t. P

$$② \Rightarrow \frac{\partial P}{\partial r} = a \frac{dr}{dp}$$

$$\Rightarrow \frac{dr}{dp} = \frac{\partial P}{a}$$

$$\Rightarrow r \frac{dr}{dp} = \frac{\partial r p}{a}$$

$$\Rightarrow \rho = \frac{\partial r p}{a}$$

$$\Rightarrow \rho^2 = \frac{4r^2 p^2}{a^2}$$

$$\Rightarrow \rho^2 = \frac{4r^2}{a^2} (ar)$$

$$\Rightarrow \rho^2 = \left(\frac{4}{a}\right)r^3$$

$$\Rightarrow \rho^2 \propto r^3$$

- ⑧ For the Cardioid, $r = a(1 - \cos\theta)$, show that $\frac{\rho^2}{r}$ is constant

Given,

$$r = a(1 - \cos\theta) \rightarrow ①$$

Differentiate ① w.r.t θ

$$\frac{dr}{d\theta} = \sin\theta \cdot a$$

$$\frac{1}{r} \frac{d\gamma}{d\theta} = \frac{a \cdot \frac{1}{2} \sin\theta/2 \cos\theta/2}{a \cdot \frac{1}{2} \sin^2\theta/2}$$

$$\frac{1}{r} \frac{dr}{d\theta} = \cot\theta/2$$

$$\cot\phi = \cot\theta/2$$

$$\phi = \theta/2$$

∴ W.R.T the pedal equation

$$P = r \sin(\theta/2)$$

$$P^2 = r^2 \sin^2(\theta/2)$$

$$P^2 = r^2 \left(\frac{1 - \cos\theta}{2} \right)$$

$$\Rightarrow p^2 = \frac{3}{2} \left(\frac{x}{a} \right)$$

$$\Rightarrow p^2 = \frac{x^3}{2a}$$

then differentiate w.r.t p

$$2p = \frac{3x^2}{2a} \frac{dx}{dp}$$

$$r \frac{dr}{dp} = \frac{2p}{\frac{3x}{2a}}$$

$$\Rightarrow r = \frac{2p \times 2a}{3x}$$

$$\Rightarrow r^2 = \frac{16a}{9x^2} \left(\frac{x^3}{2a} \right)$$

$$\Rightarrow \boxed{\frac{r^2}{r} = \frac{8a}{9}} //$$

⑨ Find the radius of curvature for the curve

$$\theta = \frac{\sqrt{r^2 - a^2}}{a} - \cos^{-1}\left(\frac{a}{r}\right) \text{ at any point on it}$$

Given,

$$\theta = \frac{\sqrt{r^2 - a^2}}{a} - \cos^{-1}\left(\frac{a}{r}\right) \rightarrow ①$$

Differentiate ① w.r.t. r

$$① \Rightarrow \frac{d\theta}{dr} = \frac{1}{a} \frac{1}{2\sqrt{r^2 - a^2}} (2r) - \left(\frac{-1}{\sqrt{1 - (a/r)^2}} \right) \frac{d}{dr} \left(\frac{a}{r} \right)$$

$$\Rightarrow \frac{d\theta}{dr} = \frac{r}{a\sqrt{r^2 - a^2}} + \frac{1}{\sqrt{1 - a^2/r^2}} a \left(-\frac{1}{r^2} \right)$$

$$= \frac{r}{a\sqrt{r^2 - a^2}} - \frac{ar}{\sqrt{r^2 - a^2}} \left(\frac{1}{r^2} \right)$$

$$= \frac{r}{a\sqrt{r^2 - a^2}} - \frac{a}{\sqrt{r^2 - a^2}}$$

$$\Rightarrow \frac{d\theta}{dr} = \frac{1}{\sqrt{r^2 - a^2}} \left[\frac{r}{a} - \frac{a}{r} \right]$$

$$= \frac{1}{\sqrt{r^2 - a^2}} \left[\frac{r^2 - a^2}{ar} \right]$$

$$= \frac{1}{\sqrt{r^2 - a^2}} \frac{\sqrt{r^2 - a^2}}{ar} \frac{\sqrt{r^2 - a^2}}{ar}$$

$$\frac{d\theta}{dr} = \frac{\sqrt{r^2 - a^2}}{ar}$$

$$\Rightarrow \frac{dr}{d\theta} = \frac{ar}{\sqrt{r^2 - a^2}}$$

$$\Rightarrow \frac{1}{\gamma} \frac{d\gamma}{d\theta} = \frac{a}{\sqrt{\gamma^2 - a^2}}$$

$$\Rightarrow \cot \phi = \frac{a}{\sqrt{\gamma^2 - a^2}}$$

$$\Rightarrow \cot \phi = \frac{a^2}{\gamma^2 - a^2}$$

$$\Rightarrow 1 + \cot^2 \phi = \frac{1 + a^2}{\gamma^2 - a^2} = \frac{\gamma^2 - a^2 + a^2}{\gamma^2 - a^2} = \frac{\gamma^2}{\gamma^2 - a^2}$$

∴ the pedal equation

$$\frac{1}{P^2} = \frac{1}{\gamma^2} (1 + \cot^2 \phi) = \frac{1}{\gamma^2} \frac{\gamma^2}{(\gamma^2 - a^2)}$$

$$\Rightarrow \frac{1}{P^2} = \frac{1}{\gamma^2 - a^2}$$

$$\Rightarrow P^2 = \gamma^2 - a^2 \rightarrow ②$$

Differentiate ② w.r.t. P

$$② \Rightarrow 2P = 2\gamma \frac{d\gamma}{dP}$$

$$\Rightarrow \gamma \frac{d\gamma}{dP} = P$$

$$\boxed{\Rightarrow P = \sqrt{\gamma^2 - a^2}}$$

⑤ Find the radius of curvature $\gamma^n = a^n (\sin n\theta)$

Given,

$$\gamma^n = a^n (\sin n\theta) \rightarrow ①$$

Differentiate w.r.t. γ

$$n r^{n-1} \frac{dr}{d\theta} = a^n (\omega \sin \theta)$$

$$\frac{r^n}{r} \frac{dr}{d\theta} = \frac{a^n \cos n\theta}{r}$$

$$\frac{1}{r} \frac{dr}{d\theta} = \frac{a^n \cos n\theta}{a^n \sin n\theta}$$

$$\cot \phi = \cot n\theta$$

∴ The pedal equation

$$\frac{1}{p^2} = \frac{1}{r^2} (1 + \cot^2 \phi)$$

$$\frac{1}{p^2} = \frac{\operatorname{cosec}^2 n\theta}{r^2}$$

$$\Rightarrow \frac{1}{p^2} = \frac{1}{r^2 \sin^2 n\theta}$$

$$\Rightarrow p^2 = r^2 \sin^2 n\theta$$

$$\Rightarrow p^2 = r \sin n\theta$$

$$\Rightarrow p^2 = r \frac{r^n}{a^n} = \frac{r^{n+1}}{a^n}$$

$$\Rightarrow a^n p = r^{n+1} \longrightarrow ②$$

Differentiate ② w.r.t p

$$② \Rightarrow a^n \cdot 1 = (n+1) r^n \frac{dr}{dp}$$

$$\Rightarrow r^n \frac{dr}{dp} = \frac{a^n}{n+1}$$

$$\Rightarrow r \cdot r^{n-1} \frac{dr}{dp} = \frac{a^n}{n+1}$$

$$\Rightarrow r \frac{dr}{d\theta} = \left(\frac{a^n}{n+1} \right) \cdot \left(\frac{1}{r^{n-1}} \right)$$

$$\Rightarrow r = \left(\frac{a^n}{n+1} \right) \left(\frac{1}{r^{n-1}} \right)$$

$$\Rightarrow r \propto \frac{1}{r^{n-1}}$$

ii) Find the radius of curvature $r^n = a^n (\cos n\theta)$
 Given,

$$r^n = a^n (\cos n\theta) \rightarrow ①$$

Differentiate ① w.r.t r

$$① \Rightarrow n r^{n-1} \frac{dr}{d\theta} = -a^n \cdot n \sin^n \theta$$

$$\frac{1}{r} \frac{\frac{dr}{d\theta}}{r} = -\frac{a^n \sin^n \theta}{a^n \cos^n \theta}$$

$$\frac{1}{r} \frac{dr}{d\theta} = -\frac{a^n \sin^n \theta}{a^n \cos^n \theta}$$

$$\cot \phi = -\tan n\theta$$

$$\cot \phi = \cot(\pi/2 + n\theta)$$

$$\phi = \frac{\pi}{2} + n\theta$$

\therefore WKT the pedal equation

$$\Rightarrow p = r \sin \phi$$

$$\Rightarrow p = r \sin (\pi/2 + n\theta)$$

$$\Rightarrow p = r \cos n\theta$$

$$\Rightarrow P = r \left(\frac{r^n}{a^n} \right)$$

$$\Rightarrow p = \frac{r^{n+1}}{a^n}$$

$$\Rightarrow a^n p = r^{n+1} \rightarrow \textcircled{2}$$

Then differentiate w.r.t p

$$a^n = (n+1) r^n \frac{dr}{dp}$$

$$\frac{a^n}{(n+1)} \left(\frac{1}{r^{n-1}} \right) = p$$

$$\Rightarrow p \propto \frac{1}{r^{n-1}}$$

//

The Centre of Curvature , Invaluates and Evaluates

Let C_1 and C_2 be the two smooth curves passing through the same point $P(x, y)$. The centre of the curve C_1 is $C(\alpha, \beta)$ called as the centre of curvature. And it's radius ρ is called circle curvature. The curve C_2 is called as the evaluate of the invaluate C_1 .

The evaluate of an invaluate can be derived by the locus of the centre of the curvature $C(\alpha, \beta)$.


The centre of curvature $C(\alpha, \beta)$ can be evaluated by using the following formulas

$$\alpha = x - \rho \sin \psi$$

$$\beta = y + \rho \cos \psi$$

$$\Rightarrow \alpha = \frac{x - y_1(1 + y_1^2)}{y_2}, \quad \beta = \frac{y + (1 + y_1^2)}{y_2}$$

$$\text{where, } y_1 = \frac{dy}{dx}, \quad y_2 = \frac{dy^2}{dx^2}$$



VIP Show that the evaluate of parabola,

$$y^2 = 4ax \text{ is } \frac{\partial}{\partial x} y^2 = 4(x - 2a)^3$$

Given.

$$y^2 = 4ax \rightarrow ①$$

$$\Rightarrow y = 2a^{1/2} x^{1/2}$$

differentiate ① w.r.t. x

$$① \Rightarrow \frac{\partial y}{\partial x} \frac{dy}{dx} = 4a$$

$$\Rightarrow y \frac{dy}{dx} = 2a$$

$$\Rightarrow \frac{dy}{dx} = y_1 = \frac{2a}{y} \rightarrow ②$$

differentiate ② w.r.t. x

$$② \Rightarrow \frac{d^2y}{dx^2} = y_2 = \frac{y(1) \frac{\partial}{\partial x} y_1}{y^2}$$

$$\Rightarrow y_2 = -\frac{\frac{\partial}{\partial x} y_1}{y^2}$$

$$\Rightarrow y_2 = -\frac{\frac{\partial}{\partial x} (2a/y)}{y^2}$$

$$\Rightarrow y_2 = -\frac{4a^2}{y^3}$$

$$\Rightarrow y_2 = -\frac{4a^2}{(2a^{1/2} \cdot x^{1/2})^3}$$

$$\Rightarrow y_2 = \frac{-4a^3}{8a^{3/2} x^{3/2}}$$

$$\Rightarrow y_2 = \frac{-a^{1/2}}{2x^{3/2}}$$

$$\therefore y_1 = \frac{2a}{2a^{1/3} \chi^{1/2}}$$

$$\Rightarrow y_1 = \frac{a^{1/2}}{\chi^{1/2}}$$

$\therefore w. K. T$

$$\alpha = \frac{x - y_1 (1 + y_1^2)}{y_2}$$

$$\Rightarrow \alpha = x - \frac{\left(\frac{a^{1/2}}{\chi^{1/2}} \right) \left(1 + a/\chi \right)}{-a^{1/2}} \\ \frac{-a^{1/2}}{2\chi^{3/2}}$$

$$\Rightarrow \alpha = \frac{x - 2a^{1/2} \chi^{3/2} \left(\frac{x+a}{\chi} \right)}{-a^{1/2} \chi^{1/2}}$$

$$\Rightarrow \alpha = \frac{x + 2x \cdot \chi^{1/2} \left(\frac{x+a}{\chi} \right)}{\chi^{1/2}}$$

$$\Rightarrow \alpha = x + 2(x+a)$$

$$\Rightarrow \alpha = 3x + 2a$$

$$\Rightarrow 3x = \alpha - 2a$$

$$\Rightarrow x = \frac{(\alpha - 2a)}{3}$$

$$\Rightarrow x^3 = \frac{(\alpha - 2a)^3}{27} \longrightarrow ③$$

$$\Rightarrow \beta = \frac{y + (1 + y_1)^2}{y_2} = \frac{2a^{1/2} \chi^{1/2} + \left(1 + \frac{a}{\chi} \right)}{-\frac{a^{1/2}}{2\chi^{3/2}}}$$

$$\Rightarrow \beta = \frac{2a^{1/2} \chi^{1/2} - 2\chi^{3/2} \left(\chi + \frac{a}{\chi} \right)}{a^{1/2}}$$

$$\Rightarrow \beta = \frac{2a^{1/2} \chi^{1/2} - 2\chi \chi^{1/2} \left(\frac{\chi + a}{\chi} \right)}{a^{1/2}}$$

$$\Rightarrow \beta = \frac{2a^{1/2} \chi^{1/2} - 2\chi^{1/2} (\chi + a)}{a^{1/2}}$$

$$\Rightarrow \beta = \frac{1}{a^{1/2}} \left| 2ax^{1/2} - 2x^{3/2} - 2ax^{1/2} \right|$$

$$\Rightarrow \beta = \frac{-2x^{3/2}}{a^{1/2}}$$

$$\Rightarrow -2x^{3/2} = \beta a^{1/2}$$

$$\Rightarrow x^{3/2} = \frac{-\beta a^{1/2}}{2}$$

$$\Rightarrow (x^{3/2})^2 = \left[\frac{-\beta a^{1/2}}{2} \right]^2$$

$$\Rightarrow x^3 = \frac{\beta^2 a}{4} \rightarrow ④$$

From ③ and ④

$$\frac{(\alpha - 2a)^3}{27} = \frac{\beta^2 a}{4}$$

$$\Rightarrow 4(\alpha - 2a)^3 = 27a\beta^2 \rightarrow ⑤$$

∴ the locus of equation ⑤ is

$$4(x - 2a)^3 = 27ay^2$$

② Show that evaluate of the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$
is $(ax)^{2/3} + (by)^{2/3} = (a^2 - b^2)^{2/3}$

Given.

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1 \rightarrow ①$$

$$\text{Let } x = a\cos\theta, y = b\sin\theta$$

$$\therefore \frac{dx}{d\theta} = -a\sin\theta, \frac{dy}{d\theta} = b\cos\theta$$

$$y_1 = \frac{dy}{dx} = \frac{dy/d\theta}{dx/d\theta} = \frac{b\cos\theta}{-a\sin\theta} = -\frac{b}{a} \cot\theta \rightarrow ②$$

differentiate ② w.r.t x

$$\Rightarrow y_2 = \frac{d^2y}{dx^2} = \frac{d}{dx} \left(\frac{dy}{dx} \right)$$

$$\Rightarrow y_2 = \frac{d}{d\theta} \left(\frac{dy}{dx} \right) \frac{d\theta}{dx}$$

$$\Rightarrow y_2 = \frac{d}{d\theta} \left(-\frac{b}{a} \cot\theta \right) \cdot \frac{1}{dx/d\theta}$$

$$\Rightarrow y_2 = -\frac{b}{a} (-\operatorname{cosec}^2\theta) \cdot \frac{1}{-\sin\theta}$$

$$\Rightarrow y_2 = -\frac{b}{a^2} \frac{1}{\sin^2\theta} \cdot \frac{1}{\sin\theta}$$

$$\Rightarrow y_2 = \frac{-b}{a^2 \sin^3\theta}$$

$\therefore \omega \cdot K.T$

$$\alpha = \frac{x - y_1(1 + y_1^2)}{y_2}$$

$$\alpha = \frac{a \cos \theta - (-b/a \omega t \theta) \left(1 + \frac{b^2}{a^2} \omega t^2 \theta \right)}{-b/a^2 \sin^3 \theta}$$

$$\Rightarrow \alpha = \frac{a \cos \theta - \frac{b}{a} \frac{\cos \theta}{\sin \theta} \left[1 + \frac{b^2}{a^2} \frac{\cos^2 \theta}{\sin^2 \theta} \right]}{\frac{b}{a^2 \sin^3 \theta}}$$

$$\Rightarrow \alpha = a \cos \theta - (a \sin^3 \theta) \left(\frac{\cos \theta}{\sin \theta} \right) \left(\frac{a^2 \sin^2 \theta + b^2 \cos^2 \theta}{a^2 \sin^2 \theta} \right)$$

$$\Rightarrow \alpha = a \cos \theta - \frac{\cos \theta}{a} \left[a^2 (1 - \cos^2 \theta) + b^2 \cos^2 \theta \right]$$

$$\Rightarrow \alpha = a \cos \theta - \frac{\cos \theta}{a} \left[a^2 - a^2 \cos^2 \theta + b^2 \cos^2 \theta \right]$$

$$\Rightarrow \alpha = a \cos \theta - a \cos \theta + \left[a \cos^3 \theta - \frac{b^2}{a} \cos^3 \theta \right]$$

$$\Rightarrow \alpha = \left(a - \frac{b^2}{a} \right) \cos^3 \theta$$

$$\Rightarrow \alpha = \left(\frac{a^2 - b^2}{a} \right) \cos^3 \theta$$

$$\Rightarrow \cos^3 \theta = \frac{a \alpha}{a^2 - b^2}$$

$$\Rightarrow (\cos^3 \theta)^{2/3} = (\alpha \alpha)^{2/3} / (a^2 - b^2)^{2/3}$$

$$\Rightarrow \cos^2 \theta = \frac{(\alpha \alpha)^{2/3}}{(\alpha^2 - b^2)^{2/3}} \longrightarrow ③$$

$$\beta = \frac{y + (1 + y_1^2)}{y_2}$$

$$\Rightarrow \beta = \frac{b \sin \theta + \left(1 + \frac{b^2}{\alpha^2} \omega t^2 \theta \right)}{-b/\alpha^2 \sin^3 \theta}$$

$$\Rightarrow \beta = b \sin \theta - \frac{\alpha^2 \sin^3 \theta}{b} - b \sin \theta + b \sin^3 \theta$$

$$\Rightarrow \beta = \left(b - \frac{\alpha^2}{b} \right) \sin^3 \theta$$

$$\Rightarrow \beta = \left(\frac{b^2 - \alpha^2}{b} \right) \sin^3 \theta$$

$$\Rightarrow \sin^3 \theta = \frac{b \beta}{b^2 - \alpha^2}$$

$$\Rightarrow \sin^3 \theta = \frac{b \beta}{-(\alpha^2 - b^2)}$$

$$\Rightarrow \sin^6 \theta = \frac{(b \beta)^2}{(\alpha^2 - b^2)^{2/3}}$$

$$\Rightarrow \sin^2 \theta = \frac{(b \beta)^{2/3}}{(\alpha^2 - b^2)^{2/3}} \longrightarrow ④$$

$$③ + ④ = \cos^2 \theta + \sin^2 \theta = \frac{(\alpha \alpha)^{2/3}}{(\alpha^2 - b^2)^{2/3}} + \frac{(b \beta)^{2/3}}{(\alpha^2 - b^2)^{2/3}}$$

$$\Rightarrow (ax)^{2/3} + (by)^{2/3} = (a^2 - b^2)^{2/3}$$

\therefore the locus of (α, β)

$$(ax)^{2/3} + (by)^{2/3} = (a^2 - b^2)^{2/3} //$$

③ Show that the evaluate of the hyperbola

$$\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1, \quad (ax)^{2/3} - (by)^{2/3} = (a^2 + b^2)^{2/3}$$

Given,

$$x = a \sec \theta, \quad y = b \tan \theta$$

$$\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1 \rightarrow ①$$

$$\frac{dx}{d\theta} = a \sec \theta \tan \theta, \quad \frac{dy}{d\theta} = b \sec^2 \theta$$

$$\therefore y_1 = \frac{dy}{dx} = \frac{\frac{dy}{d\theta}}{\frac{dx}{d\theta}} = \frac{b \sec^2 \theta}{a \sec \theta \tan \theta} = \frac{b \sec \theta}{a \tan \theta}$$

$$y_1 = \frac{b}{a} \cosec \theta$$

$$y_2 = \frac{d}{dx} \left(\frac{dy}{dx} \right) = \frac{d}{d\theta} \left(\frac{dy}{dx} \right) \frac{d\theta}{dx}$$

$$= \frac{d}{d\theta} \left(\frac{b}{a} \cosec \theta \right) \cdot \frac{1}{\frac{dx}{d\theta}}$$

$$= \frac{b}{a} (-\cosec \theta \cdot \cot \theta) \cdot \frac{1}{a \sec \theta \tan \theta}$$

$$= -\frac{b}{a^2} \frac{1}{\sin \theta} \frac{\cos \theta}{\sin \theta} \cdot \cos \theta \cdot \frac{\cos \theta}{\sin \theta}$$

$$y_2 = -\frac{b}{a^2} \frac{\cos^3 \theta}{\sin^3 \theta}$$

$$\therefore d = \frac{x - y_1(1 + y_1^2)}{y_2}$$

$$d = \frac{a \sec \theta - \left(\frac{b}{a} \csc \theta \right) \left[1 + \frac{b^2}{a^2} \csc^2 \theta \right]}{-\frac{b}{a^2} \frac{\cos^3 \theta}{\sin^3 \theta}}$$

$$d = a \sec \theta + \frac{b}{a} \frac{1}{\sin \theta} \frac{a^2 \tan^3 \theta}{b \cos^3 \theta} \left[1 + \frac{b^2}{a^2} \cdot \frac{1}{\sin^2 \theta} \right]$$

$$d = a \sec \theta + a \frac{\sin^2 \theta}{\cos^3 \theta} \left[\frac{a^2 \sin^2 \theta + b^2}{a^2 \sin^2 \theta} \right]$$

$$d = a \sec \theta + \frac{1}{a \cos^3 \theta} [a^2 \sin^2 \theta + b^2]$$

$$d = a \sec \theta + \frac{a \sin^2 \theta}{\cos^3 \theta} + \frac{b^2}{a \cos^3 \theta}$$

$$d = a \sec \theta + a \tan^2 \theta \sec \theta + \frac{b^2}{a} \sec^3 \theta$$

$$d = a \sec \theta + a \sec \theta (\sec^2 \theta - 1) + \frac{b^2}{a} \sec^3 \theta$$

$$\alpha = a \sec \theta + a \sec^3 \theta - a \sec \theta + \frac{b^2}{a} \sec^3 \theta$$

$$\alpha = \left(a + \frac{b^2}{a} \right) \sec^3 \theta$$

$$\alpha = \left(\frac{a^2 + b^2}{a} \right) \sec^3 \theta$$

$$\sec^3 \theta = \frac{\alpha \ell}{a^2 + b^2}$$

$$\sec^2 \theta = \frac{(a\ell)^{2/3}}{(a^2 + b^2)^{2/3}} \rightarrow ④$$

Similarly $\beta = y + \frac{(1+y_1^2)}{y_2}$

$$\tan^2 \theta = \frac{(b\beta)^{3/2}}{(a^2 + b^2)^{2/3}} \rightarrow ⑤$$

$$④ + ⑤ = 1 = \frac{(a\ell)^{2/3} - (b\beta)^{2/3}}{(a^2 + b^2)^{2/3}}$$

$$(a^2 + b^2)^{2/3} = (a\ell)^{2/3} - (by)^{2/3} //$$

A. Find the pedal equation.

$$\textcircled{1} \quad r^n = a^n \sin n\theta$$

Given,

$$r^n = a^n \sin n\theta \quad \rightarrow \textcircled{1}$$

\textcircled{1} equation differentiate w.r.t. \theta

$$\textcircled{1} \Rightarrow n r^{n-1} \frac{dr}{d\theta} = a^n \cos n\theta \cdot n$$

$$\Rightarrow \frac{r^n}{r} \frac{dr}{d\theta} = a^n \cos n\theta$$

$$\Rightarrow \frac{1}{r} \frac{dr}{d\theta} = \frac{a^n \cos n\theta}{r^n}$$

$$\Rightarrow \omega t \phi = \frac{a^n \cos n\theta}{a^n \sin n\theta}$$

$$\Rightarrow \omega t \phi = \omega t n\theta$$

$$\Rightarrow \phi = n\theta$$

We know that, the pedal equation

$$\frac{1}{P^2} = \frac{1}{r^2} (1 + \cot^2 \phi)$$

$$\Rightarrow \frac{1}{P^2} = \frac{1}{r^2} (1 + \cot^2 n\theta)$$

$$\Rightarrow \frac{1}{P^2} = \frac{1}{r^2} \operatorname{cosec}^2 n\theta$$

$$\Rightarrow P^2 = r^2 \sin^2 n\theta$$

$$\Rightarrow P = r \sin n\theta$$

$$\Rightarrow P = r \left(\frac{r^n}{a^n} \right)$$

$$\boxed{\Rightarrow P = \frac{r^{n+1}}{a^n}}$$

B. Find the pedal equation for the following polar curves.

$$\textcircled{1} \cdot r^m = a^m \cos m\theta + b^m \sin m\theta$$

Given.

$$r^m = a^m \cos m\theta + b^m \sin m\theta \rightarrow \textcircled{1}$$

Differentiate \textcircled{1} w.r.t. \theta

$$m r^{m-1} \frac{dr}{d\theta} = -\sin m\theta \cdot m a^m + \cos m\theta \cdot m b^m$$

$$\frac{r^m}{r} \frac{dr}{d\theta} = b^m \cos m\theta - a^m \sin m\theta$$

$$\frac{1}{r} \frac{dr}{d\theta} = \frac{b^m \cos m\theta - a^m \sin m\theta}{a^m \cos m\theta + b^m \sin m\theta}$$

$$\cot \phi = \frac{b^m \cos m\theta - a^m \sin m\theta}{a^m \cos m\theta + b^m \sin m\theta}$$

$$1 + \cot^2 \phi = 1 + \frac{(b^m \cos m\theta - a^m \sin m\theta)^2}{(a^m \cos m\theta + b^m \sin m\theta)^2}$$

$$= \frac{(a^m \cos m\theta + b^m \sin m\theta)^2 + (b^m \cos m\theta - a^m \sin m\theta)^2}{(a^m \cos m\theta + b^m \sin m\theta)^2}$$

$$1 + \cot^2 \phi = \frac{2(a^2m + b^2m)}{r^{2m}}$$

\therefore W.K.T the pedal equation

$$\frac{1}{P^2} = \frac{1}{r^2} (1 + \cot^2 \phi)$$

$$\frac{1}{P^2} = \frac{1}{r^2} \left(\frac{a^2m + b^2m}{r^{2m}} \right)$$

$$\Rightarrow \frac{1}{P^2} = \frac{a^{2m} + b^{2m}}{\gamma^2 + 2m}$$

$$\Rightarrow \gamma^{2m+2} = P^2(a^{2m} + b^{2m})$$

PRP@SJCIT