

Families of Distributions

Discrete Distributions

- The following are commonly used Discrete Distributions
- They can be defined by means of their Probability Mass Functions $P(x)$
- They are
 - Bernoulli Distribution
 - Binomial Distribution
 - Geometric Distribution
 - Negative Binomial Distribution
 - Poisson Distribution

Bernoulli distribution

- Used to model experiments with a binary outcome (yes/no, pass/fail, true/false)
 - Called Bernoulli Trials
- Random variable can take values 0 (fail) or 1 (pass)
- p = probability of success

$$P(x) = \begin{cases} 1 - p & \text{if } x = 0 \\ p & \text{if } x = 1 \end{cases}$$

$$E(X) = p$$

$$Var(X) = p(1 - p)$$

Binomial Distribution

- It is used to model number of success in a sequence of **independent** Bernoulli trials
- Models the probability of x successes in n trials
- p = probability of success; n = number of trials

$$P(x) = \binom{n}{x} p^x (1 - p)^{n-x}$$

$$E(X) = np$$

$$Var(X) = np(1 - p)$$

Geometric Distribution

- It is used to model number trials needed to achieve the first success in a sequence of **independent** Bernoulli trials
- Models the probability of x^{th} successive trial resulting in a success
- p = probability of success

$$P(x) = (1 - p)^{x-1}p$$

$$E(X) = \frac{1}{p}$$

$$Var(X) = \frac{1 - p}{p^2}$$

Negative Binomial Distribution

- It is used to model number of trials needed to obtain k success in a sequence of **independent** Bernoulli trials
- Models the probability of x^{th} successive trial resulting in the k^{th} success
- p = probability of success; k = number of success

$$P(x) = \binom{x-1}{k-1} (1-p)^{x-k} p^k$$

$$E(X) = \frac{k}{p}$$

$$Var(X) = \frac{k(1-p)}{p^2}$$

Poisson Distribution

- It is used to model number rare events occurring within a fixed period of time
- Models the probability of x rare events occurring in a fixed period of time if we know the frequency at which the events occur on average
- λ = frequency (average number of events in a fixed time period)

$$P(x) = e^{-\lambda} \frac{\lambda^x}{x!}$$

$$E(X) = \lambda$$

$$Var(X) = \lambda$$

Side Note: Poisson Approx. of Binomial

- If the number of trials is large and the probability of success is low, then we can use Poisson Distribution to approximate the Binomial Distribution
 - Also works if probability of failure is very low
- $$\lim_{\substack{n \rightarrow \infty \\ p \rightarrow 0 \\ np \rightarrow \lambda}} \binom{n}{x} p^x (1-p)^{n-x} = e^{-\lambda} \frac{\lambda^x}{x!}$$
- Can use this approximation if $n \geq 30$ and $p \leq 0.05$

Continuous Distributions

- The following are commonly used Continuous Distributions
- They can be defined by means of their Probability Density Functions $f(x)$
- They are
 - Uniform
 - Exponential
 - Gamma
 - Normal

Uniform Distribution

- Used to model scenarios where outcome lies within a given interval (a,b) and all outcomes are equally likely.
- If interval is (0,1) it is called standard uniform distribution

$$f(x) = \frac{1}{b-a}, a < x < b$$

$$E(X) = \frac{a+b}{2}$$

$$Var(X) = \frac{(b-a)^2}{12}$$

Exponential Distribution

- Used to model time (or separation) between events occurring at frequency λ (rate at which the events occur)

$$f(x) = \lambda e^{-\lambda x}, x > 0$$

$$E(X) = \frac{1}{\lambda}$$

$$Var(X) = \frac{1}{\lambda^2}$$

Gamma Distribution

- Used to model total time of multistage processes with α steps (shape parameter) where time of each step can be modeled as a Exponential distribution with frequency λ .

$$f(x) = \frac{\lambda^\alpha}{\Gamma(\alpha)} x^{\alpha-1} e^{-\lambda x} \quad \text{if } \alpha > 0 \ x > 0$$

$$\Gamma(\alpha) = \int_0^\infty x^{\alpha-1} e^{-x} dx \quad \text{if } \alpha > 0$$

$$\text{also } \Gamma(\alpha) = (n-1)! \quad \text{if } \alpha \text{ is a positive integer}$$

$$E(X) = \frac{\alpha}{\lambda}$$

$$Var(X) = \frac{\alpha}{\lambda^2}$$

- Please note that $\text{Gamma}(1, \lambda) = \text{Exponential}(\lambda)$

Side-Note: Gamma-Poisson Formula

- Can be used to simplify calculation of probabilities of RV T with Gamma Distribution.

$$\{T > t\} = \{X < \alpha\}$$

- Where T has Gamma distribution with parameters α (number of events) and λ (frequency of each event).
- X models the number of events that occurs before time t . It has Poisson distribution with parameter λt .

So,

$$P\{T > t\} = P\{X < \alpha\}$$

$$P\{T \leq t\} = P\{X \geq \alpha\}$$

Where T has $\text{Gamma}(\alpha, \lambda)$ distribution and X has $\text{Poisson}(\lambda t)$ distribution

Normal Distribution

- Used to model a large number of scenarios
 - Sums, averages or errors: Mainly due to CLT
 - Naturally occurring phenomena
- Allows you to model a scenario on the basis of expectation μ (location parameter) and standard deviation σ (scale parameter)

$$f(x) = \frac{1}{\sigma\sqrt{2\pi}} e^{\left(\frac{-(x-\mu)^2}{2\sigma^2}\right)} \quad -\infty < x < \infty$$

$$E(X) = \mu$$

$$Var(X) = \sigma^2$$

Central Limit Theorem

- If you a Random variable that is expressed as a sum of large number of independent random variables (usually ≥ 30), then you can use this theorem to model them as a Normal Distribution.

Theorem 1 (CENTRAL LIMIT THEOREM) *Let X_1, X_2, \dots be independent random variables with the same expectation $\mu = \mathbf{E}(X_i)$ and the same standard deviation $\sigma = \text{Std}(X_i)$, and let*

$$S_n = \sum_{i=1}^n X_i = X_1 + \dots + X_n.$$

As $n \rightarrow \infty$, the standardized sum

$$Z_n = \frac{S_n - \mathbf{E}(S_n)}{\text{Std}(S_n)} = \frac{S_n - n\mu}{\sigma\sqrt{n}}$$

converges in distribution to a Standard Normal random variable, that is,

$$F_{Z_n}(z) = \mathbf{P} \left\{ \frac{S_n - n\mu}{\sigma\sqrt{n}} \leq z \right\} \rightarrow \Phi(z)$$

for all z .

Side-Note: Using Normal to Approx. Binomial

- A binomial distribution is a sum of n Bernoulli trials. So if n is large (≥ 30) but p is not small enough (or large enough) to use Poisson approximation ($0.05 \leq p \leq 0.95$) then we can model the binomial as a sum of Bernoulli distributions with mean p and variance $p(1 - p)$.
- So by Central Limit Theorem,

$$\text{Binomial}(n, p) \approx \text{Normal} \left(\mu = np, \sigma = \sqrt{np(1 - p)} \right)$$

- This normal distribution can be calculated by converting it to a standard normal distribution