

Gov 50: 24. More Inference with Mathematical Models

Matthew Blackwell

Harvard University

Roadmap

1. Confidence intervals for experiments
2. Hypothesis testing with the CLT
3. Two-sample tests

1/ Confidence intervals for experiments

Comparison between groups

- More interesting to compare across groups.

Comparison between groups

- More interesting to compare across groups.
 - Differences in public opinion across groups

Comparison between groups

- More interesting to compare across groups.
 - Differences in public opinion across groups
 - Difference between treatment and control groups.

Comparison between groups

- More interesting to compare across groups.
 - Differences in public opinion across groups
 - Difference between treatment and control groups.
- Bedrock of causal inference!

Social pressure experiment

- Back to the Social Pressure Mailer GOTV example.

Social pressure experiment

- Back to the Social Pressure Mailer GOTV example.
 - Primary election in MI 2006

Social pressure experiment

- Back to the Social Pressure Mailer GOTV example.
 - Primary election in MI 2006
- Treatment group: postcards showing their own and their neighbors' voting records.

Social pressure experiment

- Back to the Social Pressure Mailer GOTV example.
 - Primary election in MI 2006
- Treatment group: postcards showing their own and their neighbors' voting records.
 - Sample size of treated group, $n_T = 360$ (artificially reducing sample size to highlight the math)

Social pressure experiment

- Back to the Social Pressure Mailer GOTV example.
 - Primary election in MI 2006
- Treatment group: postcards showing their own and their neighbors' voting records.
 - Sample size of treated group, $n_T = 360$ (artificially reducing sample size to highlight the math)
- Control group: received nothing.

Social pressure experiment

- Back to the Social Pressure Mailer GOTV example.
 - Primary election in MI 2006
- Treatment group: postcards showing their own and their neighbors' voting records.
 - Sample size of treated group, $n_T = 360$ (artificially reducing sample size to highlight the math)
- Control group: received nothing.
 - Sample size of the control group, $n_C = 1890$

Outcomes

- Outcome: $Y_i = 1$ if i voted, 0 otherwise.

Outcomes

- Outcome: $Y_i = 1$ if i voted, 0 otherwise.
- Turnout rate (sample mean) in treated group, $\bar{Y}_T = 0.37$

Outcomes

- Outcome: $Y_i = 1$ if i voted, 0 otherwise.
- Turnout rate (sample mean) in treated group, $\bar{Y}_T = 0.37$
- Turnout rate (sample mean) in control group, $\bar{Y}_C = 0.30$

Outcomes

- Outcome: $Y_i = 1$ if i voted, 0 otherwise.
- Turnout rate (sample mean) in treated group, $\overline{Y}_T = 0.37$
- Turnout rate (sample mean) in control group, $\overline{Y}_C = 0.30$
- Estimated **average treatment effect**

$$\widehat{ATE} = \overline{Y}_T - \overline{Y}_C = 0.07$$

Inference for the difference

- Parameter: **population ATE** $\mu_T - \mu_C$

Inference for the difference

- Parameter: **population ATE** $\mu_T - \mu_C$
 - μ_T : Turnout rate in the population if everyone received treatment.

Inference for the difference

- Parameter: **population ATE** $\mu_T - \mu_C$
 - μ_T : Turnout rate in the population if everyone received treatment.
 - μ_C : Turnout rate in the population if everyone received control.

Inference for the difference

- Parameter: **population ATE** $\mu_T - \mu_C$
 - μ_T : Turnout rate in the population if everyone received treatment.
 - μ_C : Turnout rate in the population if everyone received control.
- Estimator: $\widehat{ATE} = \bar{Y}_T - \bar{Y}_C$

Inference for the difference

- Parameter: **population ATE** $\mu_T - \mu_C$
 - μ_T : Turnout rate in the population if everyone received treatment.
 - μ_C : Turnout rate in the population if everyone received control.
- Estimator: $\widehat{ATE} = \bar{Y}_T - \bar{Y}_C$

Inference for the difference

- Parameter: **population ATE** $\mu_T - \mu_C$
 - μ_T : Turnout rate in the population if everyone received treatment.
 - μ_C : Turnout rate in the population if everyone received control.
- Estimator: $\widehat{ATE} = \bar{Y}_T - \bar{Y}_C$

By the CLT in large samples, we know that:

- $\bar{Y}_T \approx N\left(\mu_T, \frac{\mu_T(1-\mu_T)}{n_C}\right)$

Inference for the difference

- Parameter: **population ATE** $\mu_T - \mu_C$
 - μ_T : Turnout rate in the population if everyone received treatment.
 - μ_C : Turnout rate in the population if everyone received control.
- Estimator: $\widehat{ATE} = \bar{Y}_T - \bar{Y}_C$

By the CLT in large samples, we know that:

- $\bar{Y}_T \approx N\left(\mu_T, \frac{\mu_T(1-\mu_T)}{n_C}\right)$
- $\bar{Y}_C \approx N\left(\mu_C, \frac{\mu_C(1-\mu_C)}{n_C}\right)$

Inference for the difference

- Parameter: **population ATE** $\mu_T - \mu_C$
 - μ_T : Turnout rate in the population if everyone received treatment.
 - μ_C : Turnout rate in the population if everyone received control.
- Estimator: $\widehat{ATE} = \bar{Y}_T - \bar{Y}_C$

By the CLT in large samples, we know that:

- $\bar{Y}_T \approx N\left(\mu_T, \frac{\mu_T(1-\mu_T)}{n_C}\right)$
- $\bar{Y}_C \approx N\left(\mu_C, \frac{\mu_C(1-\mu_C)}{n_C}\right)$
- $\rightsquigarrow \bar{Y}_T - \bar{Y}_C \approx N(\mu_T - \mu_C, SE_{\text{diff}}^2)$

Inference for the difference

- Parameter: **population ATE** $\mu_T - \mu_C$
 - μ_T : Turnout rate in the population if everyone received treatment.
 - μ_C : Turnout rate in the population if everyone received control.
- Estimator: $\widehat{ATE} = \bar{Y}_T - \bar{Y}_C$

By the CLT in large samples, we know that:

- $\bar{Y}_T \approx N\left(\mu_T, \frac{\mu_T(1-\mu_T)}{n_C}\right)$
- $\bar{Y}_C \approx N\left(\mu_C, \frac{\mu_C(1-\mu_C)}{n_C}\right)$
- $\rightsquigarrow \bar{Y}_T - \bar{Y}_C \approx N(\mu_T - \mu_C, SE_{\text{diff}}^2)$

Inference for the difference

- Parameter: **population ATE** $\mu_T - \mu_C$
 - μ_T : Turnout rate in the population if everyone received treatment.
 - μ_C : Turnout rate in the population if everyone received control.
- Estimator: $\widehat{ATE} = \bar{Y}_T - \bar{Y}_C$


By the CLT in large samples, we know that:

- $\bar{Y}_T \approx N\left(\mu_T, \frac{\mu_T(1-\mu_T)}{n_C}\right)$
- $\bar{Y}_C \approx N\left(\mu_C, \frac{\mu_C(1-\mu_C)}{n_C}\right)$
- $\rightsquigarrow \bar{Y}_T - \bar{Y}_C \approx N(\mu_T - \mu_C, SE_{\text{diff}}^2)$

But what is the SE_{diff} in this case?

Spread of a difference in normals

If we take the difference between two independent normal r.v.s, what happens to the spread?



The spread of the difference is **larger** than the spread of the two variables being differenced!

Standard error for the estimated ATE

- SE of a difference in means **adds** the SEs for each group

$$SE_{\text{diff}} = \sqrt{SE_T^2 + SE_C^2}$$

Standard error for the estimated ATE

- SE of a difference in means **adds** the SEs for each group

$$SE_{\text{diff}} = \sqrt{SE_T^2 + SE_C^2}$$

- Using what we know about SEs with binary outcomes:

$$SE_{\text{diff}} = \sqrt{\frac{\mu_T(1 - \mu_T)}{n_t} + \frac{\mu_C(1 - \mu_C)}{n_C}}$$

Standard error for the estimated ATE

- SE of a difference in means **adds** the SEs for each group

$$SE_{\text{diff}} = \sqrt{SE_T^2 + SE_C^2}$$

- Using what we know about SEs with binary outcomes:

$$SE_{\text{diff}} = \sqrt{\frac{\mu_T(1 - \mu_T)}{n_t} + \frac{\mu_C(1 - \mu_C)}{n_C}}$$

- Chance errors $\bar{Y}_T - \bar{Y}_C - (\mu_T - \mu_C) \approx N(0, SE_{\text{diff}})$

Standard error for the estimated ATE

- SE of a difference in means **adds** the SEs for each group

$$SE_{\text{diff}} = \sqrt{SE_T^2 + SE_C^2}$$

- Using what we know about SEs with binary outcomes:

$$SE_{\text{diff}} = \sqrt{\frac{\mu_T(1 - \mu_T)}{n_t} + \frac{\mu_C(1 - \mu_C)}{n_c}}$$

- Chance errors $\bar{Y}_T - \bar{Y}_C - (\mu_T - \mu_C) \approx N(0, SE_{\text{diff}})$
 - We can construct a 95% CI with $\widehat{ATE} \pm 1.96 \times SE_{\text{diff}}$

Confidence intervals

But we don't know μ_T or μ_C ! Plug in our sample proportions to estimate the SE:

$$\begin{aligned}\widehat{SE}_{\text{diff}} &= \sqrt{\frac{\overline{Y}_T(1 - \overline{Y}_T)}{n_t} + \frac{\overline{Y}_C(1 - \overline{Y}_C)}{n_c}} \\ &= \sqrt{\frac{0.37 \times 0.63}{360} + \frac{0.3 \times 0.7}{1890}} = 0.028\end{aligned}$$

Confidence intervals

But we don't know μ_T or μ_C ! Plug in our sample proportions to estimate the SE:

$$\begin{aligned}\widehat{SE}_{\text{diff}} &= \sqrt{\frac{\overline{Y}_T(1 - \overline{Y}_T)}{n_t} + \frac{\overline{Y}_C(1 - \overline{Y}_C)}{n_c}} \\ &= \sqrt{\frac{0.37 \times 0.63}{360} + \frac{0.3 \times 0.7}{1890}} = 0.028\end{aligned}$$

Now we can construct confidence intervals based on the CLT like last time:

$$CI_{95} = \widehat{ATE} \pm 1.96 \times \widehat{SE}_{\text{diff}}$$

Confidence intervals

But we don't know μ_T or μ_C ! Plug in our sample proportions to estimate the SE:

$$\begin{aligned}\widehat{SE}_{\text{diff}} &= \sqrt{\frac{\overline{Y}_T(1 - \overline{Y}_T)}{n_t} + \frac{\overline{Y}_C(1 - \overline{Y}_C)}{n_c}} \\ &= \sqrt{\frac{0.37 \times 0.63}{360} + \frac{0.3 \times 0.7}{1890}} = 0.028\end{aligned}$$

Now we can construct confidence intervals based on the CLT like last time:

$$\begin{aligned}CI_{95} &= \widehat{ATE} \pm 1.96 \times \widehat{SE}_{\text{diff}} \\ &= 0.07 \pm 1.96 \times 0.028\end{aligned}$$

Confidence intervals

But we don't know μ_T or μ_C ! Plug in our sample proportions to estimate the SE:

$$\begin{aligned}\widehat{SE}_{\text{diff}} &= \sqrt{\frac{\overline{Y}_T(1 - \overline{Y}_T)}{n_t} + \frac{\overline{Y}_C(1 - \overline{Y}_C)}{n_c}} \\ &= \sqrt{\frac{0.37 \times 0.63}{360} + \frac{0.3 \times 0.7}{1890}} = 0.028\end{aligned}$$

Now we can construct confidence intervals based on the CLT like last time:

$$\begin{aligned}CI_{95} &= \widehat{ATE} \pm 1.96 \times \widehat{SE}_{\text{diff}} \\ &= 0.07 \pm 1.96 \times 0.028 \\ &= 0.07 \pm 0.054\end{aligned}$$

Confidence intervals

But we don't know μ_T or μ_C ! Plug in our sample proportions to estimate the SE:

$$\begin{aligned}\widehat{SE}_{\text{diff}} &= \sqrt{\frac{\overline{Y}_T(1 - \overline{Y}_T)}{n_t} + \frac{\overline{Y}_C(1 - \overline{Y}_C)}{n_c}} \\ &= \sqrt{\frac{0.37 \times 0.63}{360} + \frac{0.3 \times 0.7}{1890}} = 0.028\end{aligned}$$

Now we can construct confidence intervals based on the CLT like last time:

$$\begin{aligned}CI_{95} &= \widehat{ATE} \pm 1.96 \times \widehat{SE}_{\text{diff}} \\ &= 0.07 \pm 1.96 \times 0.028 \\ &= 0.07 \pm 0.054 \\ &= [0.016, 0.124]\end{aligned}$$

Confidence intervals

But we don't know μ_T or μ_C ! Plug in our sample proportions to estimate the SE:

$$\begin{aligned}\widehat{SE}_{\text{diff}} &= \sqrt{\frac{\overline{Y}_T(1 - \overline{Y}_T)}{n_t} + \frac{\overline{Y}_C(1 - \overline{Y}_C)}{n_c}} \\ &= \sqrt{\frac{0.37 \times 0.63}{360} + \frac{0.3 \times 0.7}{1890}} = 0.028\end{aligned}$$

Now we can construct confidence intervals based on the CLT like last time:

$$\begin{aligned}CI_{95} &= \widehat{ATE} \pm 1.96 \times \widehat{SE}_{\text{diff}} \\ &= 0.07 \pm 1.96 \times 0.028 \\ &= 0.07 \pm 0.054 \\ &= [0.016, 0.124]\end{aligned}$$

Range of possibilities taking into account plausible chance errors.

2/ Hypothesis testing with the CLT

Statistical hypothesis testing

- Statistical hypothesis testing is a **thought experiment**.

Statistical hypothesis testing

- Statistical hypothesis testing is a **thought experiment**.
- What would the world look like **if we knew the truth**?

Statistical hypothesis testing

- Statistical hypothesis testing is a **thought experiment**.
- What would the world look like **if we knew the truth**?
- Conducted with several steps:

Statistical hypothesis testing

- Statistical hypothesis testing is a **thought experiment**.
- What would the world look like **if we knew the truth**?
- Conducted with several steps:
 1. Specify your **null** and **alternative hypotheses**

Statistical hypothesis testing

- Statistical hypothesis testing is a **thought experiment**.
- What would the world look like **if we knew the truth**?
- Conducted with several steps:
 1. Specify your **null** and **alternative hypotheses**
 2. Choose an appropriate **test statistic** and level of test α

Statistical hypothesis testing

- Statistical hypothesis testing is a **thought experiment**.
- What would the world look like **if we knew the truth**?
- Conducted with several steps:
 1. Specify your **null** and **alternative hypotheses**
 2. Choose an appropriate **test statistic** and level of test α
 3. Derive the **reference distribution** of the test statistic under the null.

Statistical hypothesis testing

- Statistical hypothesis testing is a **thought experiment**.
- What would the world look like **if we knew the truth**?
- Conducted with several steps:
 1. Specify your **null** and **alternative hypotheses**
 2. Choose an appropriate **test statistic** and level of test α
 3. Derive the **reference distribution** of the test statistic under the null.
 4. Use this distribution to calculate the **p-value**.

Statistical hypothesis testing

- Statistical hypothesis testing is a **thought experiment**.
- What would the world look like **if we knew the truth**?
- Conducted with several steps:
 1. Specify your **null** and **alternative hypotheses**
 2. Choose an appropriate **test statistic** and level of test α
 3. Derive the **reference distribution** of the test statistic under the null.
 4. Use this distribution to calculate the **p-value**.
 5. Use p-value to decide whether to reject the null hypothesis or not

How popular is Joe Biden?



- What proportion of the public approves of Biden's job as president?

How popular is Joe Biden?



- What proportion of the public approves of Biden's job as president?
- Latest Gallup poll: $\bar{Y} = 0.42$ with $n = 812$

How popular is Joe Biden?



- What proportion of the public approves of Biden's job as president?
- Latest Gallup poll: $\bar{Y} = 0.42$ with $n = 812$
- Could we reject the null that Biden's national support is 50%?

How popular is Joe Biden?



- What proportion of the public approves of Biden's job as president?
- Latest Gallup poll: $\bar{Y} = 0.42$ with $n = 812$
- Could we reject the null that Biden's national support is 50%?
 - Null: $H_0 : p = 0.5$

How popular is Joe Biden?



- What proportion of the public approves of Biden's job as president?
- Latest Gallup poll: $\bar{Y} = 0.42$ with $n = 812$
- Could we reject the null that Biden's national support is 50%?
 - Null: $H_0 : p = 0.5$
 - Alternative: $H_1 : p \neq 0.5$

Sampling distribution, in pictures




population
distribution


The diagram consists of two light gray squares with blue borders, positioned side-by-side on a light gray background. The left square contains the text 'population distribution' and the right square contains the text 'estimator'. There are no arrows or other graphical elements connecting the two boxes.

estimator


Sampling distribution, in pictures




Sampling distribution, in pictures




Sampling distribution, in pictures



Sampling distribution, in pictures



Sampling distribution, in pictures



CLT for hypothesis testing

Under the null, we know the distribution of \bar{Y} :

$$\bar{Y} \approx N\left(p, \frac{p(1-p)}{n}\right) = N\left(0.5, \frac{0.5 \times 0.5}{812}\right)$$

CLT for hypothesis testing

Under the null, we know the distribution of \bar{Y} :

$$\bar{Y} \approx N\left(p, \frac{p(1-p)}{n}\right) = N\left(0.5, \frac{0.5 \times 0.5}{812}\right)$$

Using the rules of normal transformations if $X \sim N(\mu, \sigma^2)$:

$$\frac{X - \mu}{\sigma} \sim N(0, 1)$$

CLT for hypothesis testing

Under the null, we know the distribution of \bar{Y} :

$$\bar{Y} \approx N\left(p, \frac{p(1-p)}{n}\right) = N\left(0.5, \frac{0.5 \times 0.5}{812}\right)$$

Using the rules of normal transformations if $X \sim N(\mu, \sigma^2)$:

$$\frac{X - \mu}{\sigma} \sim N(0, 1)$$

Then under the null, know the distribution of the following test statistic:

$$Z = \frac{\bar{Y} - 0.5}{0.5/\sqrt{812}} \approx N(0, 1)$$

What we observe:

$$\begin{aligned} Z_{\text{obs}} &= \frac{\bar{Y} - 0.5}{0.5/\sqrt{812}} = \frac{0.42 - 0.5}{0.5/\sqrt{812}} \\ &= -\frac{0.08}{0.018} = -4.44 \end{aligned}$$


What we observe:

$$\begin{aligned} Z_{\text{obs}} &= \frac{\bar{Y} - 0.5}{0.5/\sqrt{812}} = \frac{0.42 - 0.5}{0.5/\sqrt{812}} \\ &= -\frac{0.08}{0.018} = -4.44 \end{aligned}$$

Our observed sample proportion is 4.44 SEs away from 0.5 under the null.
What's the probability of being that far away? (**p-value**)

```
pnorm(-4.44, mean = 0, sd = 1) + ## prob being below -4.44  
(1 - pnorm(4.44, mean = 0, sd = 1)) ## prob being above 4.44
```

```
## [1] 0.000009
```



Generalizing hypothesis tests

- Hypothesis testing using the CLT pretty much takes this general form no matter what the estimator of interest is.

Generalizing hypothesis tests

- Hypothesis testing using the CLT pretty much takes this general form no matter what the estimator of interest is.
- Hypotheses: $H_0 : \mu = \mu_0$ (null guess), $H_1 : \mu \neq \mu_0$

Generalizing hypothesis tests

- Hypothesis testing using the CLT pretty much takes this general form no matter what the estimator of interest is.
- Hypotheses: $H_0 : \mu = \mu_0$ (null guess), $H_1 : \mu \neq \mu_0$
- Test statistic:

$$Z = \frac{\text{observed value} - \text{null guess}}{\widehat{SE}} = \frac{\bar{Y} - \mu_0}{\widehat{SE}}$$

Generalizing hypothesis tests

- Hypothesis testing using the CLT pretty much takes this general form no matter what the estimator of interest is.
- Hypotheses: $H_0 : \mu = \mu_0$ (null guess), $H_1 : \mu \neq \mu_0$
- Test statistic:

$$Z = \frac{\text{observed value} - \text{null guess}}{\widehat{SE}} = \frac{\bar{Y} - \mu_0}{\widehat{SE}}$$

- The exact estimator for the standard error \widehat{SE} will depend on the estimator of interest.

Generalizing hypothesis tests

- Hypothesis testing using the CLT pretty much takes this general form no matter what the estimator of interest is.
- Hypotheses: $H_0 : \mu = \mu_0$ (null guess), $H_1 : \mu \neq \mu_0$
- Test statistic:

$$Z = \frac{\text{observed value} - \text{null guess}}{\widehat{SE}} = \frac{\bar{Y} - \mu_0}{\widehat{SE}}$$

- The exact estimator for the standard error \widehat{SE} will depend on the estimator of interest.
- Null distribution: $Z \approx N(0, 1)$ by the CLT


Generalizing hypothesis tests

- Hypothesis testing using the CLT pretty much takes this general form no matter what the estimator of interest is.
- Hypotheses: $H_0 : \mu = \mu_0$ (null guess), $H_1 : \mu \neq \mu_0$
- Test statistic:

$$Z = \frac{\text{observed value} - \text{null guess}}{\widehat{SE}} = \frac{\bar{Y} - \mu_0}{\widehat{SE}}$$


- The exact estimator for the standard error \widehat{SE} will depend on the estimator of interest.
- Null distribution: $Z \approx N(0, 1)$ by the CLT
- p-value: probability of a standard normal being bigger than $|Z_{\text{obs}}|$

Rejecting regions




- Reject if p-value is below α (usually 0.05).

Rejecting regions




- Reject if p-value is below α (usually 0.05).
 - We know 5% of the time Z will be bigger than 1.96.

Rejecting regions



- Reject if p-value is below α (usually 0.05).
 - We know 5% of the time Z will be bigger than 1.96.
 - If $Z_{\text{obs}} > 1.96$ or $Z_{\text{obs}} < -1.96$, then the p-value must be below 0.05

Rejecting regions



- Reject if p-value is below α (usually 0.05).
 - We know 5% of the time Z will be bigger than 1.96.
 - If $Z_{\text{obs}} > 1.96$ or $Z_{\text{obs}} < -1.96$, then the p-value must be below 0.05
 - We can reject if $|Z_{\text{obs}}| > 1.96$

3/ Two-sample tests

Two-sample hypotheses

- Parameter: **population ATE** $\mu_T - \mu_C$

Two-sample hypotheses

- Parameter: **population ATE** $\mu_T - \mu_C$
- Goal: learn about the population difference in means

Two-sample hypotheses

- Parameter: **population ATE** $\mu_T - \mu_C$
- Goal: learn about the population difference in means
- Usual null hypothesis: no difference in population means (ATE = 0)

Two-sample hypotheses

- Parameter: **population ATE** $\mu_T - \mu_C$
- Goal: learn about the population difference in means
- Usual null hypothesis: no difference in population means (ATE = 0)
 - Null: $H_0 : \mu_T - \mu_C = 0$

Two-sample hypotheses

- Parameter: **population ATE** $\mu_T - \mu_C$
- Goal: learn about the population difference in means
- Usual null hypothesis: no difference in population means (ATE = 0)
 - Null: $H_0 : \mu_T - \mu_C = 0$
 - Two-sided alternative: $H_1 : \mu_T - \mu_C \neq 0$

Two-sample hypotheses

- Parameter: **population ATE** $\mu_T - \mu_C$
- Goal: learn about the population difference in means
- Usual null hypothesis: no difference in population means (ATE = 0)
 - Null: $H_0 : \mu_T - \mu_C = 0$
 - Two-sided alternative: $H_1 : \mu_T - \mu_C \neq 0$
- In words: are the differences in sample means just due to chance?

Difference-in-means review

- Sample turnout rates: $\overline{Y}_T = 0.37$, $\overline{Y}_C = 0.30$

Difference-in-means review

- Sample turnout rates: $\bar{Y}_T = 0.37$, $\bar{Y}_C = 0.30$
- Sample sizes: $n_T = 360$, $n_C = 1890$

Difference-in-means review

- Sample turnout rates: $\overline{Y}_T = 0.37$, $\overline{Y}_C = 0.30$
- Sample sizes: $n_T = 360$, $n_C = 1890$
- Estimator is the **sample difference-in-means**:

$$\widehat{ATE} = \overline{Y}_T - \overline{Y}_C = 0.07$$

Difference-in-means review

- Sample turnout rates: $\bar{Y}_T = 0.37$, $\bar{Y}_C = 0.30$
- Sample sizes: $n_T = 360$, $n_C = 1890$
- Estimator is the **sample difference-in-means**:

$$\widehat{ATE} = \bar{Y}_T - \bar{Y}_C = 0.07$$

- Estimated SE for the difference in means:

$$\widehat{SE}_{\text{diff}} = \sqrt{\frac{\bar{Y}_T(1 - \bar{Y}_T)}{n_T} + \frac{\bar{Y}_C(1 - \bar{Y}_C)}{n_C}} = 0.028$$

CLT again and again

Earlier we saw that by the CLT we have:

$$\overline{Y}_T - \overline{Y}_C \approx N(\mu_T - \mu_C, SE_{\text{diff}}^2)$$

CLT again and again

Earlier we saw that by the CLT we have:

$$\overline{Y}_T - \overline{Y}_C \approx N(\mu_T - \mu_C, SE_{\text{diff}}^2)$$

We can use Z-scores to get a test statistic:

$$Z = \frac{(\overline{Y}_T - \overline{Y}_C) - (\mu_T - \mu_C)}{SE_{\text{diff}}} \sim N(0, 1)$$

CLT again and again

Earlier we saw that by the CLT we have:

$$\bar{Y}_T - \bar{Y}_C \approx N(\mu_T - \mu_C, SE_{\text{diff}}^2)$$

We can use Z-scores to get a test statistic:

$$Z = \frac{(\bar{Y}_T - \bar{Y}_C) - (\mu_T - \mu_C)}{SE_{\text{diff}}} \sim N(0, 1)$$

Same general form of the test statistic as with one sample mean/proportion:

$$\frac{\text{observed} - \text{null guess}}{SE}$$

The usual null of no difference

- Null hypothesis: $H_0 : \mu_T - \mu_C = 0$

The usual null of no difference

- Null hypothesis: $H_0 : \mu_T - \mu_C = 0$
- Test statistic:

$$Z = \frac{(\bar{Y}_T - \bar{Y}_C) - (\mu_T - \mu_C)}{SE_{\text{diff}}}$$

The usual null of no difference

- Null hypothesis: $H_0 : \mu_T - \mu_C = 0$
- Test statistic:

$$Z = \frac{(\bar{Y}_T - \bar{Y}_C) - (\mu_T - \mu_C)}{SE_{\text{diff}}} = \frac{(\bar{Y}_T - \bar{Y}_C) - 0}{SE_{\text{diff}}}$$

The usual null of no difference

- Null hypothesis: $H_0 : \mu_T - \mu_C = 0$
- Test statistic:

$$Z = \frac{(\bar{Y}_T - \bar{Y}_C) - (\mu_T - \mu_C)}{SE_{\text{diff}}} = \frac{(\bar{Y}_T - \bar{Y}_C) - 0}{SE_{\text{diff}}}$$

- In large samples, we can replace true SE with an estimate:

$$\widehat{SE}_{\text{diff}} = \sqrt{\widehat{SE}_T^2 + \widehat{SE}_C^2}$$

Calculating p-values

- Finally! Our test statistic in this sample:

$$Z = \frac{\overline{Y}_T - \overline{Y}_C}{\widehat{SE}_{\text{diff}}} = \frac{0.07}{0.028} = 2.5$$

Calculating p-values

- Finally! Our test statistic in this sample:

$$Z = \frac{\overline{Y}_T - \overline{Y}_C}{\widehat{SE}_{\text{diff}}} = \frac{0.07}{0.028} = 2.5$$


- p-value based on a two-sided test: probability of getting a difference in means this big (or bigger) if the null hypothesis were true

Calculating p-values

- Finally! Our test statistic in this sample:

$$Z = \frac{\overline{Y}_T - \overline{Y}_C}{\widehat{SE}_{\text{diff}}} = \frac{0.07}{0.028} = 2.5$$

- p-value based on a two-sided test: probability of getting a difference in means this big (or bigger) if the null hypothesis were true
 - Lower p-values \rightsquigarrow stronger evidence against the null.



```
2 * pnorm(2.5, lower.tail = FALSE)
```

```
## [1] 0.0124
```