Gov 50: 8. Observational Studies

Matthew Blackwell

Harvard University

Roadmap



· Can newspaper endorsements change voters' minds?



- · Can newspaper endorsements change voters' minds?
- Why not compare vote choice of readers of different papers?



- · Can newspaper endorsements change voters' minds?
- · Why not compare vote choice of readers of different papers?
 - Problem: readers choose papers based on their previous beliefs.



- · Can newspaper endorsements change voters' minds?
- · Why not compare vote choice of readers of different papers?
 - Problem: readers choose papers based on their previous beliefs.
 - Liberals \leadsto New York Times, conservatives \leadsto Wall Street Journal.



- · Can newspaper endorsements change voters' minds?
- · Why not compare vote choice of readers of different papers?
 - Problem: readers choose papers based on their previous beliefs.
 - Liberals \leadsto New York Times, conservatives \leadsto Wall Street Journal.
- Study for today: British newspapers switching their endorsements.



- · Can newspaper endorsements change voters' minds?
- · Why not compare vote choice of readers of different papers?
 - Problem: readers choose papers based on their previous beliefs.
 - Liberals \leadsto New York Times, conservatives \leadsto Wall Street Journal.
- Study for today: British newspapers switching their endorsements.
 - Some newspapers endorsing Tories in 1992 switched to Labour in 1997.



- · Can newspaper endorsements change voters' minds?
- · Why not compare vote choice of readers of different papers?
 - · Problem: readers choose papers based on their previous beliefs.
 - Liberals \leadsto New York Times, conservatives \leadsto Wall Street Journal.
- Study for today: British newspapers switching their endorsements.
 - Some newspapers endorsing Tories in 1992 switched to Labour in 1997.
 - Treated group: readers of Tory \rightarrow Labour papers.



- · Can newspaper endorsements change voters' minds?
- · Why not compare vote choice of readers of different papers?
 - Problem: readers choose papers based on their previous beliefs.
 - Liberals \leadsto New York Times, conservatives \leadsto Wall Street Journal.
- Study for today: British newspapers switching their endorsements.
 - Some newspapers endorsing Tories in 1992 switched to Labour in 1997.
 - Treated group: readers of Tory \rightarrow Labour papers.
 - Control group: readers of papers who didn't switch.

Data

Name	Description
to_labour	Read a newspaper that switched endorsement to Labour between 1992 and 1997 (1=Yes, 0=No)?
vote_lab_92	Did respondent vote for Labour in 1992 election (1=Yes, 0=No)?
vote_lab_97	Did respondent vote for Labour in 1997 election (1=Yes, 0=No)?
age	Age of respondent
male	Does the respondent identify as Male (1=Yes, 0=No)?
parent_labour	Did the respondent's parents vote for Labour (1=Yes, 0=No)?
work_class	Does the respondent identify as working class (1=Yes, 0=No)?

library(tidyverse) library(gov50data) newspapers

```
## # A tibble: 1,593 x 7
##
     to labour vote lab 92 vote lab 97 age male
         <fdh>>
                    <dbl>
                               <dbl> <hvn lbll> <dbl>
##
## 1
                                            33
                                                   0
## 2
                                             51
##
   3
                                            46
##
                                            45
                                            29
##
   5
##
   6
                                            47
##
                                            34
##
                                            31
##
                                            24
## 10
                                             48
                                                   0
## # i 1,583 more rows
## # i 2 more variables: parent_labour <dbl>, work_class <dbl>
```

• Example of an observational study:

- Example of an **observational study**:
 - We as researchers observe a naturally assigned treatment

- Example of an observational study:
 - We as researchers observe a naturally assigned treatment
 - Very common: often can't randomize for ethical/logistical reasons.

- Example of an observational study:
 - We as researchers observe a naturally assigned treatment
 - Very common: often can't randomize for ethical/logistical reasons.
- Internal validity: are the causal assumption satisfied? Can we interpret this as a causal effect?


- · Example of an observational study:
 - · We as researchers observe a naturally assigned treatment
 - Very common: often can't randomize for ethical/logistical reasons.
- Internal validity: are the causal assumption satisfied? Can we interpret this as a causal effect?
 - RCTs usually have higher internal validity.

- Example of an observational study:
 - We as researchers observe a naturally assigned treatment
 - Very common: often can't randomize for ethical/logistical reasons.
- Internal validity: are the causal assumption satisfied? Can we interpret
 this as a causal effect?
 - RCTs usually have higher internal validity.
 - Observational studies less so because treatment and control groups may differ in ways that are hard to measure


- Example of an observational study:
 - We as researchers observe a naturally assigned treatment
 - Very common: often can't randomize for ethical/logistical reasons.
- Internal validity: are the causal assumption satisfied? Can we interpret
 this as a causal effect?
 - RCTs usually have higher internal validity.
 - Observational studies less so because treatment and control groups may differ in ways that are hard to measure
- **External validity**: can the conclusions/estimated effects be generalized beyond this study?

- · Example of an observational study:
 - · We as researchers observe a naturally assigned treatment
 - Very common: often can't randomize for ethical/logistical reasons.
- Internal validity: are the causal assumption satisfied? Can we interpret
 this as a causal effect?
 - · RCTs usually have higher internal validity.
 - Observational studies less so because treatment and control groups may differ in ways that are hard to measure
- **External validity**: can the conclusions/estimated effects be generalized beyond this study?
 - RCTs weaker here because often very expensive to conduct on representative samples.


- Example of an observational study:
 - We as researchers observe a naturally assigned treatment
 - Very common: often can't randomize for ethical/logistical reasons.
- Internal validity: are the causal assumption satisfied? Can we interpret
 this as a causal effect?
 - · RCTs usually have higher internal validity.
 - Observational studies less so because treatment and control groups may differ in ways that are hard to measure
- External validity: can the conclusions/estimated effects be generalized beyond this study?
 - RCTs weaker here because often very expensive to conduct on representative samples.
 - Observational studies often have larger/more representative samples that improve external validity.




• **Confounder**: pre-treatment variable affecting treatment & the outcome.




- Confounder: pre-treatment variable affecting treatment & the outcome.
 - Leftists (X) more likely to read newspapers switching to Labour (T).




- Confounder: pre-treatment variable affecting treatment & the outcome.
 - Leftists (X) more likely to read newspapers switching to Labour (T).
 - Leftists (X) also more likely to vote for Labour (Y).



- **Confounder**: pre-treatment variable affecting treatment & the outcome.
 - Leftists (X) more likely to read newspapers switching to Labour (T).
 - Leftists (X) also more likely to vote for Labour (Y).
- Confounding bias in the estimated SATE due to these differences



- **Confounder**: pre-treatment variable affecting treatment & the outcome.
 - Leftists (X) more likely to read newspapers switching to Labour (T).
 - Leftists (X) also more likely to vote for Labour (Y).
- Confounding bias in the estimated SATE due to these differences
 - $\overline{Y}_{control}$ not a good proxy for $\frac{1}{n}\sum_{i=1}^{n}Y_{i}(0)$ in treated group.



- **Confounder**: pre-treatment variable affecting treatment & the outcome.
 - Leftists (X) more likely to read newspapers switching to Labour (T).
 - Leftists (X) also more likely to vote for Labour (Y).
- Confounding bias in the estimated SATE due to these differences
 - $\overline{Y}_{control}$ not a good proxy for $\frac{1}{n} \sum_{i=1}^{n} Y_i(0)$ in treated group.
 - one type: selection bias from self-selection into treatment

· How can we find a good comparison group?


- · How can we find a good comparison group?
- Depends on the data we have available.

- · How can we find a good comparison group?
- Depends on the data we have available.
- Three general types of observational study **reseach designs**:

- · How can we find a good comparison group?
- Depends on the data we have available.
- Three general types of observational study **reseach designs**:
 - 1. **Cross-sectional design**: compare outcomes treated and control units at one point in time.


- · How can we find a good comparison group?
- Depends on the data we have available.
- Three general types of observational study reseach designs:
 - Cross-sectional design: compare outcomes treated and control units at one point in time.
 - 2. **Before-and-after design**: compare outcomes before and after a unit has been treated, but need over-time data on treated group.


- · How can we find a good comparison group?
- · Depends on the data we have available.
- Three general types of observational study reseach designs:
 - Cross-sectional design: compare outcomes treated and control units at one point in time.
 - 2. **Before-and-after design**: compare outcomes before and after a unit has been treated, but need over-time data on treated group.
 - 3. **Difference-in-differences design**: use before/after information for the treated and control group; need over-time on treated & control group.




Cross-sectional design

 Compare treated/control groups after treatment happens.







- Compare treated/control groups after treatment happens.
 - Switching readers vs non-switching readers in 1997.



- Compare treated/control groups after treatment happens.
 - Switching readers vs non-switching readers in 1997.
- Assumption: groups identical on average (like RCTs)




- Compare treated/control groups after treatment happens.
 - Switching readers vs non-switching readers in 1997.
- Assumption: groups identical on average (like RCTs)
 - Sometimes called unconfoundedness or as-if randomized.



- Compare treated/control groups after treatment happens.
 - Switching readers vs non-switching readers in 1997.
- Assumption: groups identical on average (like RCTs)
 - Sometimes called unconfoundedness or as-if randomized.
- · Cross-section estimate:

$$\overline{Y}_{\text{treated}}^{\text{after}} - \overline{Y}_{\text{contro}}^{\text{after}}$$



- Compare treated/control groups after treatment happens.
 - Switching readers vs non-switching readers in 1997.
- Assumption: groups identical on average (like RCTs)
 - Sometimes called unconfoundedness or as-if randomized.
- · Cross-section estimate:

$$\overline{Y}_{\text{treated}}^{\text{after}} - \overline{Y}_{\text{control}}^{\text{after}}$$

· Could there be confounders?

Cross-sectional design in R

```
switched <- newspapers |>
  filter(to_labour == 1) |>
  summarize(mean(vote_lab_97))

no_change <- newspapers |>
  filter(to_labour == 0) |>
  summarize(mean(vote_lab_97))

switched - no_change
```

```
## mean(vote_lab_97)
## 1 0.14
```

• Statistical control: adjust for confounders using statistical procedures.

- Statistical control: adjust for confounders using statistical procedures.
 - Can help to reduce confounding bias.

- Statistical control: adjust for confounders using statistical procedures.
 - · Can help to reduce confounding bias.
- · One type of statistical control: subclassification

- Statistical control: adjust for confounders using statistical procedures.
 - · Can help to reduce confounding bias.
- · One type of statistical control: subclassification
 - · Compare treated and control groups within levels of a confounder.

- Statistical control: adjust for confounders using statistical procedures.
 - · Can help to reduce confounding bias.
- · One type of statistical control: subclassification
 - · Compare treated and control groups within levels of a confounder.
 - Remaining effect can't be due to the confounder.


- Statistical control: adjust for confounders using statistical procedures.
 - · Can help to reduce confounding bias.
- · One type of statistical control: subclassification
 - · Compare treated and control groups within levels of a confounder.
 - Remaining effect can't be due to the confounder.

Statistical control in R


```
newspapers |>
  group_by(parent_labour, to_labour) |>
  summarize(avg_vote = mean(vote_lab_97)) |>
  pivot_wider(
   names_from = to_labour,
   values_from = avg_vote
) |>
  mutate(diff_by_parent = `1` - `0`)
```


```
## # A tibble: 2 x 4
## # Groups: parent_labour [2]
## parent_labour `0` `1` diff_by_parent
## <dbl> <dbl> <dbl> <dbl> <dbl> ## 1
            0 0.279 0.434
            0.155
## 2
            1 0.597 0.698
            0.101
```

 Compare readers of party-switching newspapers before & after switch.




- Compare readers of party-switching newspapers before & after switch.
- Advantage: all person-specific features held fixed






- Compare readers of party-switching newspapers before & after switch.
- Advantage: all person-specific features held fixed
 - · comparing within a person over time.



- Compare readers of party-switching newspapers before & after switch.
- Advantage: all person-specific features held fixed
 - · comparing within a person over time.
- · Before-and-after estimate:


$$\overline{Y}_{\text{treated}}^{\text{after}} - \overline{Y}_{\text{treated}}^{\text{before}}$$



- Compare readers of party-switching newspapers before & after switch.
- Advantage: all person-specific features held fixed
 - comparing within a person over time.
- · Before-and-after estimate:

$$\overline{Y}_{\text{treated}}^{\text{after}} - \overline{Y}_{\text{treated}}^{\text{before}}$$

Assumption: no time-varying confounders



- Compare readers of party-switching newspapers before & after switch.
- Advantage: all person-specific features held fixed
 - · comparing within a person over time.
- · Before-and-after estimate:

$$\overline{Y}_{\text{treated}}^{\text{after}} - \overline{Y}_{\text{treated}}^{\text{before}}$$


- Assumption: no time-varying confounders
 - Time trend: Labour just did better overall in 1997 compared to 1992.


Before and after in R

```
newspapers |>
  mutate(
    vote_change = vote_lab_97 - vote_lab_92
) |>
  summarize(avg_change = mean(vote_change))
```

```
## # A tibble: 1 x 1
## avg_change
## <dbl>
## 1 0.119
```


 Use the before/after difference of control group to infer what would have happened to treatment group without treatment.





- Use the before/after difference of control group to infer what would have happened to treatment group without treatment.
- · DiD estimate:

$$\underbrace{\left(\overline{Y}_{\text{treated}}^{\text{after}} - \overline{Y}_{\text{treated}}^{\text{before}}\right)}_{\text{trend in treated group}} - \underbrace{\left(\overline{Y}_{\text{control}}^{\text{after}} - \overline{Y}_{\text{control}}^{\text{before}}\right)}_{\text{trend in control group}}$$



- Use the before/after difference of control group to infer what would have happened to treatment group without treatment.
- · DiD estimate:

$$\underbrace{\left(\overline{Y}_{\text{treated}}^{\text{after}} - \overline{Y}_{\text{treated}}^{\text{before}}\right)}_{\text{trend in treated group}} - \underbrace{\left(\overline{Y}_{\text{control}}^{\text{after}} - \overline{Y}_{\text{control}}^{\text{before}}\right)}_{\text{trend in control group}}$$


 Change in treated group above and beyond the change in control group.



- Use the before/after difference of control group to infer what would have happened to treatment group without treatment.
- · DiD estimate:

$$\underbrace{\left(\overline{Y}_{\text{treated}}^{\text{after}} - \overline{Y}_{\text{treated}}^{\text{before}}\right)}_{\text{trend in treated group}} - \underbrace{\left(\overline{Y}_{\text{control}}^{\text{after}} - \overline{Y}_{\text{control}}^{\text{before}}\right)}_{\text{trend in control group}}$$

- Change in treated group above and beyond the change in control group.
- · Assumption: parallel trends




 Use the before/after difference of control group to infer what would have happened to treatment group without treatment.

· DiD estimate:

$$\underbrace{\left(\overline{Y}_{\text{treated}}^{\text{after}} - \overline{Y}_{\text{treated}}^{\text{before}}\right)}_{\text{trend in treated group}} - \underbrace{\left(\overline{Y}_{\text{control}}^{\text{after}} - \overline{Y}_{\text{control}}^{\text{before}}\right)}_{\text{trend in control group}}$$

- Change in treated group above and beyond the change in control group.
- · Assumption: parallel trends
 - Changes in vote of readers of non-switching papers roughly the same as changes that readers of switching papers would have been if they read non-switching papers.



 Use the before/after difference of control group to infer what would have happened to treatment group without treatment.

· DiD estimate:

$$\underbrace{\left(\overline{Y}_{\text{treated}}^{\text{after}} - \overline{Y}_{\text{treated}}^{\text{before}}\right)}_{\text{trend in treated group}} - \underbrace{\left(\overline{Y}_{\text{control}}^{\text{after}} - \overline{Y}_{\text{control}}^{\text{before}}\right)}_{\text{trend in control group}}$$

- Change in treated group above and beyond the change in control group.
- Assumption: parallel trends
 - Changes in vote of readers of non-switching papers roughly the same as changes that readers of switching papers would have been if they read non-switching papers.
 - Threat to inference: non-parallel trends.

Difference-in-differences in R

```
newspapers |>
 mutate(
    vote change = vote lab 97 - vote lab 92,
    to_labour = if_else(to_labour == 1, "switched", "unswitched")
 group by(to labour) |>
  summarize(avg change = mean(vote change)) |>
 pivot wider(
   names from = to labour,
    values_from = avg_change
 mutate(DID = switched - unswitched)
```

```
## # A tibble: 1 x 3
## switched unswitched DID
## <dbl> <dbl> <dbl> <dbl>
## 1 0.190 0.110 0.0796
```

1. Cross-sectional comparison

1. Cross-sectional comparison

· Compare treated units with control units after treatment

1. Cross-sectional comparison

- · Compare treated units with control units after treatment
- Assumption: treated and controls units are comparable

1. Cross-sectional comparison

- · Compare treated units with control units after treatment
- · Assumption: treated and controls units are comparable
- Possible confounding

1. Cross-sectional comparison

- · Compare treated units with control units after treatment
- Assumption: treated and controls units are comparable
- · Possible confounding

1. Cross-sectional comparison

- · Compare treated units with control units after treatment
- · Assumption: treated and controls units are comparable
- Possible confounding

2. Before-and-after comparison

Compare the same units before and after treatment

1. Cross-sectional comparison

- · Compare treated units with control units after treatment
- · Assumption: treated and controls units are comparable
- · Possible confounding

- Compare the same units before and after treatment
- · Assumption: no time-varying confounding

1. Cross-sectional comparison

- · Compare treated units with control units after treatment
- · Assumption: treated and controls units are comparable
- Possible confounding

2. Before-and-after comparison

- Compare the same units before and after treatment
- Assumption: no time-varying confounding

1. Cross-sectional comparison

- · Compare treated units with control units after treatment
- · Assumption: treated and controls units are comparable
- · Possible confounding

2. Before-and-after comparison

- · Compare the same units before and after treatment
- Assumption: no time-varying confounding

3. Differences-in-differences

· Assumption: parallel trends assumptions

1. Cross-sectional comparison

- · Compare treated units with control units after treatment
- · Assumption: treated and controls units are comparable
- Possible confounding

2. **Before-and-after comparison**

- Compare the same units before and after treatment
- · Assumption: no time-varying confounding

- · Assumption: parallel trends assumptions
- Under this assumption, it accounts for unit-specific and time-varying confounding.

1. Cross-sectional comparison

- · Compare treated units with control units after treatment
- · Assumption: treated and controls units are comparable
- · Possible confounding

2. Before-and-after comparison

- Compare the same units before and after treatment
- Assumption: no time-varying confounding

- · Assumption: parallel trends assumptions
- Under this assumption, it accounts for unit-specific and time-varying confounding.
- All rely on assumptions that can't be verified to handle confounding.

1. Cross-sectional comparison

- · Compare treated units with control units after treatment
- · Assumption: treated and controls units are comparable
- · Possible confounding

2. Before-and-after comparison

- · Compare the same units before and after treatment
- · Assumption: no time-varying confounding

- · Assumption: parallel trends assumptions
- Under this assumption, it accounts for unit-specific and time-varying confounding.
- All rely on assumptions that can't be verified to handle confounding.
- RCTs handle confounding by design.

Causality understanding check

