

matreex: Simulating European forest dynamics with IPM.

Maxime Jaunatre ^{1¶}, Julien Barrere ^{1,2}, Björn Reineking ¹, Arnaud Guyennon ³, Thomas Cordonnier ⁴, and Georges Kunstler ^{1¶}

¹ Univ. Grenoble Alpes, INRAE, LESSEM, St-Martin-d'Hères, France  ² INRAE, Aix Marseille Université, UMR RECOVER, Aix-en-Provence, France ³ Independent Researcher, France ⁴ ONF, Département Recherche Développement et Innovation, 21 rue du Muguet, 39100 Dole. ¶ Corresponding author

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

Software

- [Review](#) 
- [Repository](#) 
- [Archive](#) 

Editor: [Open Journals](#) 

Reviewers:

- [@openjournals](#)

Submitted: 01 January 1970

Published: unpublished

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#))

Summary

Integral projection models (IPMs) are powerful tools for studying the temporal dynamics of populations structured by continuous traits, allowing for predictions of changes in trait distributions over time ([Ellner et al., 2016](#)). Unlike individual-based or cohort-based models, which represent populations as discrete populations, IPMs describe populations as continuous populations, integrating over the uncertainty of demographic processes. This removes demographic stochasticity and results in fully deterministic simulations, which are complementary to individual-based models (IBMs).

Here, we introduce matreex, an R package specifically designed to build IPMs for European forest tree species. Our package includes pre-fitted species-specific growth, survival and recruitment functions that account for the effect of climate and competition, and functions to efficiently integrate IPMs and run temporal simulations of single-species or multispecies forest communities until equilibrium. This package complements existing R packages for IPMs, such as *ipmr* ([Metcalfe et al., 2013](#)) and *IPMpack* ([Levin et al., 2021](#)), which are not specifically designed for forest ecosystems.

Statement of need

IPMs are rarely applied to forest ecosystems due to the complexity of tree growth kernels, which are challenging to integrate, making the construction of forest IPMs particularly difficult. matreex is an R package specifically designed to build IPMs and run simulations for European forests. These IPMs can be used by the forest ecology research community, to study population dynamic and forest communities trajectories. In matreex IPM simulations, it is also possible to include temporally variable climatic conditions, natural disturbances, harvesting scenarios and regional dispersal affecting population dynaminc depending on tree species sensitivity and stand structure.

State of the field

Other, more generalist, packages exist to build IPMs for various types of organisms (*IPMpack* ([Metcalfe et al., 2013](#)), *ipmr* ([Levin et al., 2021](#))). However they focus on analysis on the matrix itself whereas matreex use the integrated matrices to simulate population dynamics. In addition matreex integrates numerous functions to run simulations for multispecies communities under various harvesting and disturbance scenarios.

Software design

A key feature of the `matreex` package is the development of IPM integration functions designed for complex forest tree growth kernels. As trees have a small annual growth rate, most of the integration effort is concentrated near the diagonal of the matrix. We achieve this by combining different integration methods (Gauss-Legendre and Mid-bin) at different distances of the diagonal, which helps speed up the integration. [Figure 1](#) illustrates how these different methods of integrating the matrix are used. This speed is crucial as we integrate a matrix per competition level (based on species basal area) to account for density dependence. More details about the integration are provided in the [Building IPM vignette](#).




Figure 1: Combination of different integration methods. Dashed line is the identity where $z_t = z_t + 1$ and dark blue distribution is an expected distribution of the growth kernel.

A crucial development objective was to simplify the workflow for ecological researchers, who may work on multispecies models with climate variation, disturbances, and harvesting. To facilitate this, `matreex` provides fitted vital models for European tree species directly in the package ([Barrere et al., 2024](#); [Guyennon et al., 2023](#); [Kunstler et al., 2021](#)), although other new vital models can be used. The object-oriented architecture limits code complexity, allowing users to focus on designing large simulation experiments to tackle their ecological questions. The code below demonstrates a quick way to use the package, resulting in [Figure 2](#), which shows the dynamics of four species in the same environment.

```
library(matreex)
library(dplyr)
library(ggplot2)

# select a climate to run in
data("climate_species")
climate <- subset(
  climate_species,
  climate_label == "optimum" & sp == "Fagus_sylvatica",
  select = -c(sp, climate_label)
)

# integrate ipms and store them in species object
Picea <- species(IPM = make_IPM(
  species = "Picea_abies", fit = fit_Picea_abies,
  climate = climate, clim_lab = "optimum clim",
  mesh = c(m = 700, L = 90, U = get_maxdbh(fit_Picea_abies) * 1.1),
  BA = 0:60
), init_pop = def_initBA(1))
Betula <- species(IPM = make_IPM(
  species = "Betula", fit = fit_Betula,
  climate = climate, clim_lab = "optimum clim",
  mesh = c(m = 700, L = 90, U = get_maxdbh(fit_Betula) * 1.1),
  BA = 0:60
```

```

80 ), init_pop = def_initBA(1))
81 Fagus <- species(IPM = make_IPM(
82   species = "Fagus_sylvatica", fit = fit_Fagus_sylvatica,
83   climate = climate, clim_lab = "optimum clim",
84   mesh = c(m = 700, L = 90, U = get_maxdbh(fit_Fagus_sylvatica) * 1.1),
85   BA = 0:60
86 ), init_pop = def_initBA(1))
87 Abies <- species(IPM = make_IPM(
88   species = "Abies_alba", fit = fit_Abies_alba,
89   climate = climate, clim_lab = "optimum clim",
90   mesh = c(m = 700, L = 90, U = get_maxdbh(fit_Abies_alba) * 1.1),
91   BA = 0:60
92 ), init_pop = def_initBA(1))
93
94 # assemble species in a forest object
95 Forest <- forest(species = list(Picea = Picea, Abies = Abies,
96                               Fagus = Fagus, Betula = Betula))
97 set.seed(20260825) # The seed is here for initial population random functions.
98
99 # Run simulation and plot it
100 Sim <- sim_deter_forest(
101   Forest,
102   tlim = 1000,
103   equil_time = 1000, equil_dist = 50, equil_diff = 1,
104   SurfEch = 0.03,
105   verbose = TRUE
106 )
107 Sim %>%
108   dplyr::filter(var == "BAsp", ! equil) |>
109   ggplot(aes(x = time, y = value, color = species)) +
110   geom_line(linewidth = .4) +
111   ylab("Basal Area (m2)") + xlab("Simulation time (years)")

```




Figure 2: Simulation output for 4 species.

112 Climatic temporal variability

113 Modelling forest dynamics under fluctuating climatic conditions can be computationally
 114 expensive because the IPMs growth kernel must be integrated for every climatic condition. To

avoid this high computation cost, we implement a new method involving the pre-integration of IPM growth matrix blocks for different mean growth rates. The IPM for each climatic condition is then reassembled from these mean growth rate IPM matrix blocks (mu matrix). This allows to speed-up the simulations. This is described in the [matreex climate variation vignette](#).

Disturbance

One key originality of matreex is the possibility to apply storm, fire, biotic or snow disturbances of varying intensity (ranging from 0 to 1) at any time of the IPM simulations. When a disturbance strikes the stand a given year of the simulation, the survival function is replaced by the species-specific equations from Barrere et al. (2024). These equations quantify the annual mortality probability of a tree in a disturbed stand as a function of its species, diameter at breast height, stand structure, nature and intensity of the disturbance. A disturbance striking two different stands with the same intensity will thus result in different mortality rates, depending notably on the sensitivity of the tree species present in the stand. Disturbances in matreex are described in details in Barrere et al. (2024) and in Barrere et al. (in prep). Figure 3 shows an example of multispecies simulations with storm disturbance.




Figure 3: Simulation output for 3 species with a disturbance at $time = 2600$.

Regional dispersal

Most stand-scale forest dynamics models simulate closed systems, where only the tree species already present in the stand contribute to the recruitment of new trees. This limitation preclude the simulation of immigration from external species, which is a key process of forest dynamics, particularly under climate change. To overcome this limitation, we included the possibility to split the recruitment function in two component : (i) within-plot dispersal that depends on the sum of basal area of fecund tree species in the plot, and (ii) external dispersal that depends on the regional pool. This regional pool approach is extensively presented in Barrere et al. (in prep).

Harvesting

Since most temperate forests are managed, it is crucial to incorporate silvicultural interventions into the simulations. We implemented three management strategies:

- First, we implemented a simple constant annual harvesting rates accounting for the effect of the harvesting rates observed in NFI data used for the model calibration (Kunstler et al., 2021).

- Second, we implement an even-aged management. The objective is to apply typical even-aged harvesting, based on a single cohort. Trees are harvested with successive thinning during stand development until the final harvest. Thinning harvest are based on the distance to a self-thinning boundary, based on Aussenac et al. (2021). This is easily connected with management guidelines.
- Third, we implement an uneven-aged harvesting. The uneven-aged harvest scenario consists in selective harvesting across all size classes with the objective to reach a stable size structure with continuous replacement of large mature trees. This scenario depends on the basal area of the stand and the size distribution of the trees (building on Lafond et al. (2014)). These three managements are described in the [matreex harvesting vignette](#).

156 Research impact statement

matreex was designed to be easily adapted to various research ideas and is in continuous development. It has already been used in several scientific publications tackling diverse scientific questions (Baranger et al., in prep; Barrere et al., 2024, in prep; Guyennon et al., 2023; Kunstler et al., 2021). The ability to simulate forests easily with a dedicated R package will help ecologists to analyse the effect of climate change, shifting disturbance regimes, and their interplay with forest management across European forests.

```
163 matreex is an open-source package made available under the MIT license. Installation and
164 usage instructions can be found at the website
```

165 AI usage disclosure

166 No generative AI tools were used in the development of this software, the writing of this
167 manuscript, or the preparation of supporting materials.

168 **Acknowledgements**

169 JB, MJ, BR and GK are funded through the BiodivClim ERA-Net Cofund (joint BiodivERsA
170 Call on “Biodiversity and Climate Change”, 2019-2020) with national co-funding through ANR
171 (France, project ANR-20-EBI5-0005-03). MJ and GK were funded by the ANR DECLIC (grant
172 ANR-1520-CE32-0005-01) and REGE-ADAPT PEPR FORESTTT France 2030 (ANR-24-PEFO-
173 0006). JB, MJ, BR and GK are funded by the RESONATE H2020 project (grant 101000574).
174 GK, BR and AG received support from the REFORCE – EU FP7ERA-NET Sumforest 2016
175 through the call ‘Sustainable forests for the society of the future’, with the ANR as national
176 funding agency (grant ANR-16-SUMF-0002).

177 References

- Aussenac, R., Pérot, T., Fortin, M., Coligny, F. de, Monnet, J.-M., & Vallet, P. (2021). The
salem simulator version 2.0: A tool for predicting the productivity of pure and mixed
forest stands and simulating management operations. *Open Research Europe*, 1, 61.
<https://doi.org/10.12688/openreseurope.13671.2>
- Baranger, A., Cordonnier, T., Barrere, J., & Kunstler, G. (in prep). *Direct and indirect effects
of climate on competition drive species population performance across their climatic ranges.*
- Barrere, J., Grünig, M., Jabot, F., Jaunatre, M., Rammer, W., Reineking, B., Seidl, R., Senf,
C., & Kunstler, G. (in prep). *Competition, dispersal and disturbances drive the transient
response of forest composition to climate change across europe.*

- 187 Barrere, J., Reineking, B., Jaunatre, M., & Kunstler, G. (2024). Forest storm resilience depends
188 on the interplay between functional composition and climate—insights from european-
189 scale simulations. *Functional Ecology*, 38(3), 500–516. <https://doi.org/10.1111/1365-2435.14489>
190
- 191 Ellner, S. P., Childs, D. Z., & Rees, M. (2016). *Data-driven Modelling of Structured Populations: A Practical Guide to the Integral Projection Model*. Springer International Publishing.
192 <https://doi.org/10.1007/978-3-319-28893-2>
193
- 194 Guyennon, A., Reineking, B., Salguero-Gomez, R., Dahlgren, J., Lehtonen, A., Ratcliffe, S.,
195 Ruiz-Benito, P., Zavala, M. A., & Kunstler, G. (2023). Beyond mean fitness: Demographic
196 stochasticity and resilience matter at tree species climatic edges. *Global Ecology and Biogeography*, n/a(n/a). <https://doi.org/10.1111/geb.13640>
197
- 198 Kunstler, G., Guyennon, A., Ratcliffe, S., Rüger, N., Ruiz-Benito, P., Childs, D. Z., Dahlgren,
199 J., Lehtonen, A., Thuiller, W., Wirth, C., Zavala, M. A., & Salguero-Gomez, R. (2021).
200 Demographic performance of european tree species at their hot and cold climatic edges.
201 *Journal of Ecology*, 109(2), 1041–1054. <https://doi.org/10.1111/1365-2745.13533>
- 202 Lafond, V., Lagarrigues, G., Cordonnier, T., & Courbaud, B. (2014). Uneven-aged management
203 options to promote forest resilience for climate change adaptation: Effects of group
204 selection and harvesting intensity. *Annals of Forest Science*, 71(2), 173–186. <https://doi.org/10.1007/s13595-013-0291-y>
205
- 206 Levin, S. C., Childs, D. Z., Compagnoni, A., Evers, S., Knight, T. M., & Salguero-Gómez, R. (2021). Ipmr: Flexible implementation of integral projection models in r. *Methods in Ecology and Evolution*, 12(10), 1826–1834. <https://doi.org/10.1111/2041-210X.13683>
207
- 208 Metcalf, C. J. E., McMahon, S. M., Salguero-Gómez, R., & Jongejans, E. (2013). Pack: An r
209 package for integral projection models. *Methods in Ecology and Evolution*, 4(2), 195–200.
210 <https://doi.org/10.1111/2041-210x.12001>
211