

# Action constrained quasi-Newton methods

Robert Gower



**EUROPT 2014 Workshop on Advances in  
Continuous Optimization**

July 10, 2014

---

What and why quasi-Newton?

Classic quasi-Newton

New Action Constrained quasi-Newton

Implementing a Preconditioned Newton-CG with new metric


## What's new

- ▶ A few Hessian-vector products are cheap  $\Rightarrow$  use a handful to build Hessian approximation <sup>1</sup>
- ▶ Framework for “tracking” inverses of matrix fields
- ▶ General purpose Newton-CG preconditioners
- ▶ Good results on regularized logistic regression (compared to BFGS or Newton-CG)

---

<sup>1</sup>Walther, A. (2008). Computing sparse Hessians with automatic differentiation. ACM Trans. Math. Software, 34(1), Art. 3, 15.

## Solving sequences of linear systems




$x_0$

min  $f(x)$  ( $C^2$  – diffeomorphism)

$$f(x_k + d) \approx f_k + \nabla f_k^T d + \frac{1}{2} d^T \nabla^2 f_k d$$

## Solving sequences of linear systems




$$\min \quad f(x)$$

$$f(x_k + d) \approx f_k + \nabla f_k^T d + \frac{1}{2} d^T \nabla^2 f_k d$$


$x_0$

## Solving sequences of linear systems



$$\min \quad f(x)$$

$$f(x_k + d) \approx f_k + \nabla f_k^T d + \frac{1}{2} d^T \nabla^2 f_k d$$




$x_0$

$$d_0 = -\nabla^2 f_0^{-1} \nabla f_0$$


$x_1$

## Solving sequences of linear systems



The quadratic model “slowly” changes; how to take advantage?

## Solving sequences of linear systems



The quadratic model “slowly” changes; how to take advantage?


Maintain an approximation of the inverse:  $H_{k+1} \approx \nabla^2 f_{k+1}^{-1}$ .

“Slowly” update with cheap low rank matrices:  $H_{k+1} = H_k + E_k$ .

$H_k$  is a *metric* matrix.



## Solving sequences of linear systems



The quadratic model “slowly” changes; how to take advantage?

Maintain an approximation of the inverse:  $H_{k+1} \approx \nabla^2 f_{k+1}^{-1}$ .

“Slowly” update with cheap low rank matrices:  $H_{k+1} = H_k + E_k$ .

$H_k$  is a *metric* matrix.

## Classic approach: imitating action of Hessian with the secant equation

$$\mathcal{A}\nabla^2 f_{k+1} = \int_0^1 \nabla^2 f(x_k + td_k) dt, \quad \gamma_k = (\nabla f_{k+1} - \nabla f_k).$$

Fundamental Theorem of Calculus says

$$\begin{aligned}\mathcal{A}\nabla^2 f_{k+1} \cdot d_k &= \gamma_k \Rightarrow \\ d_k &= (\mathcal{A}\nabla^2 f_{k+1})^{-1} \gamma_k \approx (\nabla^2 f_{k+1})^{-1} \gamma_k\end{aligned}$$

---

<sup>2</sup>Fletcher, B. R., & Powell, M. J. D. (1960). A rapidly convergent descent method for minimization, (1).

## Classic approach: imitating action of Hessian with the secant equation

$$\mathcal{A}\nabla^2 f_{k+1} = \int_0^1 \nabla^2 f(x_k + td_k) dt, \quad \gamma_k = (\nabla f_{k+1} - \nabla f_k).$$

Fundamental Theorem of Calculus says

$$\begin{aligned} \mathcal{A}\nabla^2 f_{k+1} \cdot d_k &= \gamma_k \Rightarrow \\ d_k &= (\mathcal{A}\nabla^2 f_{k+1})^{-1} \gamma_k \approx (\nabla^2 f_{k+1})^{-1} \gamma_k \end{aligned}$$

Choose a metric that satisfies

### The Secant Equation

$$d_k = H_{k+1} \gamma_k = H_{k+1} \mathcal{A}\nabla^2 f_{k+1} d_k$$

Still under-determined. Least squares idea + slowly changing <sup>2</sup>

## Classic approach: imitating action of Hessian with the secant equation

$$\mathcal{A}\nabla^2 f_{k+1} = \int_0^1 \nabla^2 f(x_k + td_k) dt, \quad \gamma_k = (\nabla f_{k+1} - \nabla f_k).$$

Fundamental Theorem of Calculus says

$$\begin{aligned} \mathcal{A}\nabla^2 f_{k+1} \cdot d_k &= \gamma_k \Rightarrow \\ d_k &= (\mathcal{A}\nabla^2 f_{k+1})^{-1} \gamma_k \approx (\nabla^2 f_{k+1})^{-1} \gamma_k \end{aligned}$$

Choose a metric that satisfies

### The Secant Equation

$$d_k = H_{k+1} \gamma_k = H_{k+1} \mathcal{A}\nabla^2 f_{k+1} d_k$$

Still under-determined. Least squares idea + slowly changing <sup>2</sup>

---

<sup>2</sup>Fletcher, B. R., & Powell, M. J. D. (1960). A rapidly convergent descent method for minimization, (1).

## Classic approach: imitating action of Hessian with the secant equation

$$\mathcal{A}\nabla^2 f_{k+1} = \int_0^1 \nabla^2 f(x_k + td_k) dt, \quad \gamma_k = (\nabla f_{k+1} - \nabla f_k).$$

Fundamental Theorem of Calculus says

$$\begin{aligned} \mathcal{A}\nabla^2 f_{k+1} \cdot d_k &= \gamma_k \Rightarrow \\ d_k &= (\mathcal{A}\nabla^2 f_{k+1})^{-1} \gamma_k \approx (\nabla^2 f_{k+1})^{-1} \gamma_k \end{aligned}$$

Choose a metric that satisfies

### The Secant Equation

$$d_k = H_{k+1} \gamma_k = H_{k+1} \mathcal{A}\nabla^2 f_{k+1} d_k$$

Still under-determined. Least squares idea + slowly changing <sup>2</sup>

---

<sup>2</sup>Fletcher, B. R., & Powell, M. J. D. (1960). A rapidly convergent descent method for minimization, (1).

## Designing classic quasi-Newton

$$\begin{aligned} \min_{H_{k+1}} \quad & \|H_{k+1} - H_k\|_{Frobenius(W)}^2 = \|E_k\|_{Frobenius(W)}^2 = \sum_{i,j} E_{i,j}^2 W_{i,j}^2 \\ \text{s.t.} \quad & H_{k+1} \mathcal{A} \nabla^2 f_{k+1} d_k = d_k, \quad H_{k+1} = H_{k+1}^T. \end{aligned}$$

- Iteratively updating metric; changes “slowly”<sup>3</sup>

---

<sup>3</sup>Goldfarb, D. (1970). A Family of Variable-Metric Methods Derived by Variational Means. *Mathematics of Computation*, 24(109), 23.

## Designing classic quasi-Newton

$$\begin{aligned} \min_{H_{k+1}} \quad & \|H_{k+1} - H_k\|_{Frobenius(W)}^2 = \|E_k\|_{Frobenius(W)}^2 = \sum_{i,j} E_{i,j}^2 W_{i,j}^2 \\ \text{s.t.} \quad & H_{k+1} \mathcal{A} \nabla^2 f_{k+1} d_k = d_k, \quad H_{k+1} = H_{k+1}^T. \end{aligned}$$

- ▶ Iteratively updating metric; changes “slowly”<sup>3</sup>
- ▶ Secant equation Must be symmetric

<sup>3</sup>Goldfarb, D. (1970). A Family of Variable-Metric Methods Derived by Variational Means. Mathematics of Computation, 24(109), 23.

## Designing classic quasi-Newton

$$\begin{aligned} \min_{H_{k+1}} \quad & \|H_{k+1} - H_k\|_{Frobenius(W)}^2 = \|E_k\|_{Frobenius(W)}^2 = \sum_{i,j} E_{i,j}^2 W_{i,j}^2 \\ \text{s.t.} \quad & H_{k+1} \mathcal{A} \nabla^2 f_{k+1} d_k = d_k, \quad H_{k+1} = H_{k+1}^T. \end{aligned}$$

- ▶ Iteratively updating metric; changes “slowly”<sup>3</sup>
- ▶ Secant equation Must be symmetric

BFGS by choosing  $W = \mathcal{A} \nabla^2 f_{k+1}$ ,

$$H_{k+1} = \frac{d_k d_k^T}{d_k^T \gamma_k} + \left( I - \frac{d_k \gamma_k^T}{d_k^T \gamma_k} \right) H_k \left( I - \frac{\gamma_k d_k^T}{d_k^T \gamma_k} \right).$$

<sup>3</sup>Goldfarb, D. (1970). A Family of Variable-Metric Methods Derived by Variational Means. Mathematics of Computation, 24(109), 23.



## Designing classic quasi-Newton

$$\begin{aligned} \min_{H_{k+1}} \quad & \|H_{k+1} - H_k\|_{Frobenius(W)}^2 = \|E_k\|_{Frobenius(W)}^2 = \sum_{i,j} E_{i,j}^2 W_{i,j}^2 \\ \text{s.t.} \quad & H_{k+1} \mathcal{A} \nabla^2 f_{k+1} d_k = d_k, \quad H_{k+1} = H_{k+1}^T. \end{aligned}$$

- ▶ Iteratively updating metric; changes “slowly”<sup>3</sup>
- ▶ Secant equation Must be symmetric

BFGS by choosing  $W = \mathcal{A} \nabla^2 f_{k+1}$ ,

$$H_{k+1} = \frac{d_k d_k^T}{d_k^T \gamma_k} + \left( I - \frac{d_k \gamma_k^T}{d_k^T \gamma_k} \right) H_k \left( I - \frac{\gamma_k d_k^T}{d_k^T \gamma_k} \right).$$

<sup>3</sup>Goldfarb, D. (1970). A Family of Variable-Metric Methods Derived by Variational Means. Mathematics of Computation, 24(109), 23.

## Designing classic quasi-Newton

$$\begin{aligned} \min_{H_{k+1}} \quad & \|H_{k+1} - H_k\|_{Frobenius(W)}^2 = \|E_k\|_{Frobenius(W)}^2 = \sum_{i,j} E_{i,j}^2 W_{i,j}^2 \\ \text{s.t.} \quad & H_{k+1} \mathcal{A} \nabla^2 f_{k+1} d_k = d_k, \quad H_{k+1} = H_{k+1}^T. \end{aligned}$$

- Iteratively updating metric; changes “slowly”<sup>3</sup>
- Secant equation Must be symmetric

BFGS by choosing  $W = \mathcal{A} \nabla^2 f_{k+1}$ ,

$$H_{k+1} = \text{proj}_{d_k}^{\mathcal{A} \nabla^2 f_k} + \left( I - \text{proj}_{d_k}^{\mathcal{A} \nabla^2 f_k} \mathcal{A} \nabla^2 f_k \right) H_k \left( I - \mathcal{A} \nabla^2 f_k \text{proj}_{d_k}^{\mathcal{A} \nabla^2 f_k} \right).$$

$\text{proj}_d^A := d(d^T A d)^{-1} d^T A$  = oblique  $A$ -projection onto  $\text{span}(d)$ .

<sup>3</sup>Goldfarb, D. (1970). A Family of Variable-Metric Methods Derived by Variational Means. Mathematics of Computation, 24(109), 23.

## Designing classic quasi-Newton

$$\begin{aligned} \min_{H_{k+1}} \quad & \|H_{k+1} - H_k\|_{Frobenius(W)}^2 = \|E_k\|_{Frobenius(W)}^2 = \sum_{i,j} E_{i,j}^2 W_{i,j}^2 \\ \text{s.t.} \quad & H_{k+1} \mathcal{A} \nabla^2 f_{k+1} d_k = d_k, \quad H_{k+1} = H_{k+1}^T. \end{aligned}$$

- Iteratively updating metric; changes “slowly”<sup>3</sup>
- Secant equation Must be symmetric

BFGS by choosing  $W = \mathcal{A} \nabla^2 f_{k+1}$ ,

$$H_{k+1} = \text{proj}_{d_k}^{\mathcal{A} \nabla^2 f_k} + \left( I - \text{proj}_{d_k}^{\mathcal{A} \nabla^2 f_k} \mathcal{A} \nabla^2 f_k \right) H_k \left( I - \mathcal{A} \nabla^2 f_k \text{proj}_{d_k}^{\mathcal{A} \nabla^2 f_k} \right).$$

$\text{proj}_d^A := d(d^T A d)^{-1} d^T A$  = oblique  $A$ -projection onto  $\text{span}(d)$ .

<sup>3</sup>Goldfarb, D. (1970). A Family of Variable-Metric Methods Derived by Variational Means. Mathematics of Computation, 24(109), 23.

## Designing classic quasi-Newton

$$\begin{aligned} \min_{H_{k+1}} \quad & \|H_{k+1} - H_k\|_{Frobenius(W)}^2 = \|E_k\|_{Frobenius(W)}^2 = \sum_{i,j} E_{i,j}^2 W_{i,j}^2 \\ \text{s.t.} \quad & H_{k+1} \mathcal{A} \nabla^2 f_{k+1} d_k = d_k, \quad H_{k+1} = H_{k+1}^T. \end{aligned}$$

- Iteratively updating metric; changes “slowly”<sup>3</sup>
- Secant equation Must be symmetric

BFGS by choosing  $W = \mathcal{A} \nabla^2 f_{k+1}$ ,

$$H_{k+1} = \text{proj}_{d_k}^{\mathcal{A} \nabla^2 f_k} + \left( I - \text{proj}_{d_k}^{\mathcal{A} \nabla^2 f_k} \mathcal{A} \nabla^2 f_k \right) H_k \left( I - \mathcal{A} \nabla^2 f_k \text{proj}_{d_k}^{\mathcal{A} \nabla^2 f_k} \right).$$

$\text{proj}_d^A := d(d^T A d)^{-1} d^T A$  = oblique  $A$ -projection onto  $\text{span}(d)$ .

<sup>3</sup>Goldfarb, D. (1970). A Family of Variable-Metric Methods Derived by Variational Means. Mathematics of Computation, 24(109), 23.

## Designing the action constrained metrics

$$\min H_{k+1}$$

$$\|H_{k+1} - H_k\|_{Frobenius(W)}^2$$

s.t.

$$H_{k+1} \nabla^2 f_{k+1} D_k = D_k,$$

$$H_{k+1} = H_{k+1}^T.$$

- ▶ Iteratively updating metric; changes “slowly”
- ▶ Same action of  $\nabla^2 f_{k+1}^{-1}$  and  $H_{k+1}$  over  $\nabla^2 f_{k+1} D_k$  where  $D_k \in \mathbb{R}^{n \times p}$ ,  $p \ll n$  is a tall thin matrix. Must be symmetric

$$H_{k+1} = H_k + W^{-1} \text{proj}_{\nabla^2 f_{k+1} D_k}^{W^{-1}} (I - H_k \nabla^2 f_{k+1}) (I - \text{proj}_{\nabla^2 f_{k+1} D_k}^{W^{-1}} W^{-1}) \\ + (I - H_k \nabla^2 f_{k+1}) \text{proj}_{\nabla^2 f_{k+1} D_k}^{W^{-1}} W^{-1}$$

## Designing the action constrained metrics

$$\min H_{k+1}$$

$$\|H_{k+1} - H_k\|_{Frobenius(W)}^2$$

s.t.

$$H_{k+1} \nabla^2 f_{k+1} D_k = D_k,$$

$$H_{k+1} = H_{k+1}^T.$$

- ▶ Iteratively updating metric; changes “slowly”
- ▶ Same action of  $\nabla^2 f_{k+1}^{-1}$  and  $H_{k+1}$  over  $\nabla^2 f_{k+1} D_k$  where  $D_k \in \mathbb{R}^{n \times p}$ ,  $p \ll n$  is a tall thin matrix. Must be symmetric

$$H_{k+1} = H_k + W^{-1} \text{proj}_{\nabla^2 f_{k+1} D_k}^{W^{-1}} (I - H_k \nabla^2 f_{k+1}) \left( I - \text{proj}_{\nabla^2 f_{k+1} D_k}^{W^{-1}} W^{-1} \right) \\ + (I - H_k \nabla^2 f_{k+1}) \text{proj}_{\nabla^2 f_{k+1} D_k}^{W^{-1}} W^{-1}$$

$$\text{proj}_D^A A := D(D^T A D)^{-1} D^T A.$$

## Designing the action constrained metrics

$$\min H_{k+1}$$

$$\|H_{k+1} - H_k\|_{Frobenius(W)}^2$$

s.t.

$$H_{k+1} \nabla^2 f_{k+1} D_k = D_k,$$

$$H_{k+1} = H_{k+1}^T.$$

- ▶ Iteratively updating metric; changes “slowly”
- ▶ Same action of  $\nabla^2 f_{k+1}^{-1}$  and  $H_{k+1}$  over  $\nabla^2 f_{k+1} D_k$  where  $D_k \in \mathbb{R}^{n \times p}$ ,  $p \ll n$  is a tall thin matrix. Must be symmetric

$$H_{k+1} = H_k + W^{-1} \text{proj}_{\nabla^2 f_{k+1} D_k}^{W^{-1}} (I - H_k \nabla^2 f_{k+1}) \left( I - \text{proj}_{\nabla^2 f_{k+1} D_k}^{W^{-1}} W^{-1} \right) \\ + (I - H_k \nabla^2 f_{k+1}) \text{proj}_{\nabla^2 f_{k+1} D_k}^{W^{-1}} W^{-1}$$

$$\text{proj}_D^A A := D(D^T A D)^{-1} D^T A.$$

Reverse Automatic Differentiation  $\nabla^2 f_{k+1}(D_k)$  costs  $O(p)$ !

## Designing the action constrained metrics

$$\min H_{k+1}$$

$$\|H_{k+1} - H_k\|_{\text{Frobenius}(W)}^2$$

s.t.

$$H_{k+1} \nabla^2 f_{k+1} D_k = D_k,$$

$$H_{k+1} = H_{k+1}^T.$$

- ▶ Iteratively updating metric; changes “slowly”
- ▶ Same action of  $\nabla^2 f_{k+1}^{-1}$  and  $H_{k+1}$  over  $\nabla^2 f_{k+1} D_k$  where  $D_k \in \mathbb{R}^{n \times p}$ ,  $p \ll n$  is a tall thin matrix. Must be symmetric

$$H_{k+1} = H_k + W^{-1} \text{proj}_{\nabla^2 f_{k+1} D_k}^{W^{-1}} (I - H_k \nabla^2 f_{k+1}) \left( I - \text{proj}_{\nabla^2 f_{k+1} D_k}^{W^{-1}} W^{-1} \right) \\ + (I - H_k \nabla^2 f_{k+1}) \text{proj}_{\nabla^2 f_{k+1} D_k}^{W^{-1}} W^{-1}$$

$$\text{proj}_D^A A := D(D^T A D)^{-1} D^T A.$$

Reverse Automatic Differentiation  $\nabla^2 f_{k+1}(D_k)$  costs  $O(p)$ !

A small rank- $3p$  update.



## Designing the action constrained metrics

$$\min H_{k+1}$$

$$\|H_{k+1} - H_k\|_{Frobenius(W)}^2$$

s.t.

$$H_{k+1} \nabla^2 f_{k+1} D_k = D_k,$$

$$H_{k+1} = H_{k+1}^T.$$

- ▶ Iteratively updating metric; changes “slowly”
- ▶ Same action of  $\nabla^2 f_{k+1}^{-1}$  and  $H_{k+1}$  over  $\nabla^2 f_{k+1} D_k$  where  $D_k \in \mathbb{R}^{n \times p}$ ,  $p \ll n$  is a tall thin matrix. Must be symmetric

$$H_{k+1} = H_k + W^{-1} \text{proj}_{\nabla^2 f_{k+1} D_k}^{W^{-1}} (I - H_k \nabla^2 f_{k+1}) \left( I - \text{proj}_{\nabla^2 f_{k+1} D_k}^{W^{-1}} W^{-1} \right) \\ + (I - H_k \nabla^2 f_{k+1}) \text{proj}_{\nabla^2 f_{k+1} D_k}^{W^{-1}} W^{-1}$$

$$\text{proj}_D^A A := D(D^T A D)^{-1} D^T A.$$

Reverse Automatic Differentiation  $\nabla^2 f_{k+1}(D_k)$  costs  $O(p)$ !

A small rank- $3p$  update.  $W = ?$  and  $D_k = ?$

## Designing the action constrained metrics

$$\min H_{k+1}$$

$$\|H_{k+1} - H_k\|_{\text{Frobenius}(W)}^2$$

s.t.

$$H_{k+1} \nabla^2 f_{k+1} D_k = D_k,$$

$$H_{k+1} = H_{k+1}^T.$$

- ▶ Iteratively updating metric; changes “slowly”
- ▶ Same action of  $\nabla^2 f_{k+1}^{-1}$  and  $H_{k+1}$  over  $\nabla^2 f_{k+1} D_k$  where  $D_k \in \mathbb{R}^{n \times p}$ ,  $p \ll n$  is a tall thin matrix. Must be symmetric

$$H_{k+1} = H_k + W^{-1} \text{proj}_{\nabla^2 f_{k+1} D_k}^{W^{-1}} (I - H_k \nabla^2 f_{k+1}) \left( I - \text{proj}_{\nabla^2 f_{k+1} D_k}^{W^{-1}} W^{-1} \right) \\ + (I - H_k \nabla^2 f_{k+1}) \text{proj}_{\nabla^2 f_{k+1} D_k}^{W^{-1}} W^{-1}$$


$$\text{proj}_D^A A := D(D^T A D)^{-1} D^T A.$$

Reverse Automatic Differentiation  $\nabla^2 f_{k+1}(D_k)$  costs  $O(p)$ !

A small rank- $3p$  update.  $W = ?$  and  $D_k = ?$


## Overall idea of $D_k$

Build metric  $H$  that captures  $D = [d_1, \dots, d_p]$  directions of positive curvature





## Overall idea of $D_k$

Build metric  $H$  that captures  $D = [d_1, \dots, d_p]$  directions of positive curvature



## Overall idea of $D_k$

Build metric  $H$  that captures  $D = [d_1, \dots, d_p]$  directions of positive curvature





$$\nabla^2 f(x) D = H^{-1} D$$

$$H \nabla^2 f(x) D = D$$

Newton-CG produces  
directions of positive  
curvature!

## Overall idea of $D_k$

Build metric  $H$  that captures  $D = [d_1, \dots, d_p]$  directions of positive curvature



Newton-CG produces  
directions of positive  
curvature!

## Choosing $D_k$ and $W$

Analogous to BFGS by choosing  $W = \nabla^2 f_k$

quNac: quasi-Newton action constrained

$$H_{k+1} = \text{proj}_{D_k}^{\nabla^2 f_{k+1}} + (I - \text{proj}_{D_k}^{\nabla^2 f_{k+1}} \nabla^2 f_{k+1}) H_k (I - \nabla^2 f_{k+1} \text{proj}_{D_k}^{\nabla^2 f_{k+1}}).$$

- ▶ A rank- $2p$  update
- ▶ Only need to calculate the  $p$  columns  $\nabla^2 f_{k+1}(D_k)$ .
- ▶  $\text{proj}_{D_k}^{\nabla^2 f_{k+1}} = D_k (D_k^T \nabla^2 f_{k+1} D_k)^{-1} D_k^T$  is expensive?

Choose  $D_k$  according to

- ▶ **Hereditary** property  $\Rightarrow$  for local convergence
- ▶ **Descent property**  $\Rightarrow$  for Global stability

### Quadratic Hereditary $\nabla^2 f(x) \equiv Q$ property

If  $[D_1, \dots, D_k] \in \mathbb{R}^{n \times n}$  and  $D_i^T Q D_j = 0$  for  $1 \leq j < i \leq k$  then

$$H_{k+1} Q D_i = D_i, \quad \text{for } i = 1, \dots, k.$$

Lemma:  $H_{k+1} Q = I \Rightarrow H_{k+1} = Q^{-1}$ .



## Quadratic Hereditary $\nabla^2 f(x) \equiv Q$ property

If  $[D_1, \dots, D_k] \in \mathbb{R}^{n \times n}$  and  $D_i^T Q D_j = 0$  for  $1 \leq j < i \leq k$  then

$$H_{k+1} Q D_i = D_i, \quad \text{for } i = 1, \dots, k.$$

**Lemma:**  $H_{k+1} Q = I \Rightarrow H_{k+1} = Q^{-1}$ .

**proof:**

$$\begin{aligned} H_{k+1} &= H_k + E_k \\ &= H_i + E_i + \dots + E_k. \end{aligned}$$

If  $D_i^T Q D_j = 0$  for  $j < i$  then

$$\begin{aligned} \left( \text{proj}_{D_j}^Q Q D_i = 0 \right) &\Rightarrow (E_j Q D_i = 0). \\ &\Rightarrow H_{k+1} Q D_i = H_{i+1} Q D_i = D_i. \quad \square \end{aligned}$$

Choose  $D_k$  conjugate to  $\nabla^2 f_{k+1}$  as an approximation!

## Quadratic Hereditary $\nabla^2 f(x) \equiv Q$ property

If  $[D_1, \dots, D_k] \in \mathbb{R}^{n \times n}$  and  $D_i^T Q D_j = 0$  for  $1 \leq j < i \leq k$  then

$$H_{k+1} Q D_i = D_i, \quad \text{for } i = 1, \dots, k.$$

**Lemma:**  $H_{k+1} Q = I \Rightarrow H_{k+1} = Q^{-1}$ .

**proof:**

$$\begin{aligned} H_{k+1} &= H_k + E_k \\ &= H_i + E_i + \dots + E_k. \end{aligned}$$

## Quadratic Hereditary $\nabla^2 f(x) \equiv Q$ property

If  $[D_1, \dots, D_k] \in \mathbb{R}^{n \times n}$  and  $D_i^T Q D_j = 0$  for  $1 \leq j < i \leq k$  then

$$H_{k+1} Q D_i = D_i, \quad \text{for } i = 1, \dots, k.$$

**Lemma:**  $H_{k+1} Q = I \Rightarrow H_{k+1} = Q^{-1}$ .

**proof:**

$$\begin{aligned} H_{k+1} &= H_k + E_k \\ &= H_i + E_i + \dots + E_k. \end{aligned}$$

If  $D_i^T Q D_j = 0$  for  $j < i$  then

## Quadratic Hereditary $\nabla^2 f(x) \equiv Q$ property

If  $[D_1, \dots, D_k] \in \mathbb{R}^{n \times n}$  and  $D_i^T Q D_j = 0$  for  $1 \leq j < i \leq k$  then

$$H_{k+1} Q D_i = D_i, \quad \text{for } i = 1, \dots, k.$$

**Lemma:**  $H_{k+1} Q = I \Rightarrow H_{k+1} = Q^{-1}$ .

**proof:**

$$\begin{aligned} H_{k+1} &= H_k + E_k \\ &= H_i + E_i + \dots + E_k. \end{aligned}$$

If  $D_i^T Q D_j = 0$  for  $j < i$  then

$$\begin{aligned} \left( \text{proj}_{D_j}^Q Q D_i = 0 \right) &\Rightarrow (E_j Q D_i = 0). \\ &\Rightarrow H_{k+1} Q D_i = H_{i+1} Q D_i = D_i. \quad \square \end{aligned}$$

## Quadratic Hereditary $\nabla^2 f(x) \equiv Q$ property

If  $[D_1, \dots, D_k] \in \mathbb{R}^{n \times n}$  and  $D_i^T Q D_j = 0$  for  $1 \leq j < i \leq k$  then

$$H_{k+1} Q D_i = D_i, \quad \text{for } i = 1, \dots, k.$$

**Lemma:**  $H_{k+1} Q = I \Rightarrow H_{k+1} = Q^{-1}$ .

**proof:**

$$\begin{aligned} H_{k+1} &= H_k + E_k \\ &= H_i + E_i + \dots + E_k. \end{aligned}$$

If  $D_i^T Q D_j = 0$  for  $j < i$  then

$$\begin{aligned} \left( \text{proj}_{D_j}^Q Q D_i = 0 \right) &\Rightarrow (E_j Q D_i = 0). \\ &\Rightarrow H_{k+1} Q D_i = H_{i+1} Q D_i = D_i. \quad \square \end{aligned}$$

Choose  $D_k$  conjugate to  $\nabla^2 f_{k+1}$  as an approximation!

## Quadratic Hereditary $\nabla^2 f(x) \equiv Q$ property

If  $[D_1, \dots, D_k] \in \mathbb{R}^{n \times n}$  and  $D_i^T Q D_j = 0$  for  $1 \leq j < i \leq k$  then

$$H_{k+1} Q D_i = D_i, \quad \text{for } i = 1, \dots, k.$$

**Lemma:**  $H_{k+1} Q = I \Rightarrow H_{k+1} = Q^{-1}$ .

**proof:**

$$\begin{aligned} H_{k+1} &= H_k + E_k \\ &= H_i + E_i + \dots + E_k. \end{aligned}$$

If  $D_i^T Q D_j = 0$  for  $j < i$  then

$$\begin{aligned} \left( \text{proj}_{D_j}^Q Q D_i = 0 \right) &\Rightarrow (E_j Q D_i = 0). \\ &\Rightarrow H_{k+1} Q D_i = H_{i+1} Q D_i = D_i. \quad \square \end{aligned}$$

Choose  $D_k$  conjugate to  $\nabla^2 f_{k+1}$  as an approximation!

## Quadratic Hereditary $\nabla^2 f(x) \equiv Q$ property

If  $[D_1, \dots, D_k] \in \mathbb{R}^{n \times n}$  and  $D_i^T QW^{-1}QD_j = 0$  for  $1 \leq j < i \leq k$  then

$$H_{k+1}QD_i = D_i, \quad \text{for } i = 1, \dots, k.$$

**Lemma:**  $H_{k+1}Q = I \Rightarrow H_{k+1} = Q^{-1}$ .

**proof:**

$$\begin{aligned} H_{k+1} &= H_k + E_k \\ &= H_i + E_i + \dots + E_k. \end{aligned}$$

If  $D_i^T QW^{-1}QD_j = 0$  for  $j < i$  then

$$\begin{aligned} \left( \text{proj}_{QD_j}^W W^{-1}QD_i = 0 \right) &\Rightarrow (E_jQD_i = 0) \\ &\Rightarrow H_{k+1}QD_i = H_{i+1}QD_i = D_i. \quad \square \end{aligned}$$

Choose  $D_k$  conjugate to  $\nabla^2 f_{k+1}$  as an approximation!

## Descent and Positive Definiteness

**Descent:**  $d_k = -H_k \nabla f_k$  is less than  $90^\circ$  with  $-\nabla f_k$ .

### Sufficient Descent condition

If  $H_k \succ 0$  then  $-d_k^T \nabla f_k = \nabla f_k^T H_k \nabla f_k > 0$ .

### Classic quasi-Newton

If  $H_k \succ 0$  and  $\gamma_k^T d_k = d_k^T \mathcal{A} \nabla^2 f_{k+1} d_k > 0$  then  $H_{k+1} \succ 0$ .



## Descent and Positive Definiteness

**Descent:**  $d_k = -H_k \nabla f_k$  is less than  $90^\circ$  with  $-\nabla f_k$ .

### Sufficient Descent condition

If  $H_k \succ 0$  then  $-d_k^T \nabla f_k = \nabla f_k^T H_k \nabla f_k > 0$ .

### Classic quasi-Newton

If  $H_k \succ 0$  and  $\gamma_k^T d_k = d_k^T \mathcal{A} \nabla^2 f_{k+1} d_k > 0$  then  $H_{k+1} \succ 0$ .

### quNac Descent condition

If  $H_k \succ 0$  and  $D_k$  columns are directions of positive curvature  
 $D_k^T \nabla^2 f_{k+1} D_k \succ 0$  then  $H_{k+1} \succ 0$ .

## Descent and Positive Definiteness

**Descent:**  $d_k = -H_k \nabla f_k$  is less than  $90^\circ$  with  $-\nabla f_k$ .

### Sufficient Descent condition

If  $H_k \succ 0$  then  $-d_k^T \nabla f_k = \nabla f_k^T H_k \nabla f_k > 0$ .

### Classic quasi-Newton

If  $H_k \succ 0$  and  $\gamma_k^T d_k = d_k^T \mathcal{A} \nabla^2 f_{k+1} d_k > 0$  then  $H_{k+1} \succ 0$ .

### quNac Descent condition

If  $H_k \succ 0$  and  $D_k$  columns are directions of positive curvature  
 $D_k^T \nabla^2 f_{k+1} D_k \succ 0$  then  $H_{k+1} \succ 0$ .

## Implementing a Preconditioned Newton-CG with quNac

- ▶ Call Conjugate gradients on  $\nabla^2 f_{k+1} d = -\nabla f_{k+1}$  to get  $D_k$  directions of positive curvature
- ▶  $\text{proj}_{D_k}^{\nabla^2 f_{k+1}} = D_k (D_k^T \nabla^2 f_{k+1} D_k)^{-1} D_k^T$  is  $O(p^2)$  because  $D_k^T \nabla^2 f_{k+1} D_k = \text{diagonal}!$

## Implementing a Preconditioned Newton-CG with quNac

- ▶ Call Conjugate gradients on  $\nabla^2 f_{k+1} d = -\nabla f_{k+1}$  to get  $D_k$  directions of positive curvature
- ▶  $\text{proj}_{D_k}^{\nabla^2 f_{k+1}} = D_k (D_k^T \nabla^2 f_{k+1} D_k)^{-1} D_k^T$  is  $O(p^2)$  because  $D_k^T \nabla^2 f_{k+1} D_k = \text{diagonal!}$
- ▶  $H_{k+1}$  is positive definite and  $H_{k+1} \nabla^2 f_{k+1} D_k = D_k$ , thus  $H_{k+1} \nabla^2 f_{k+1}$  has concentrated Eigen-values  $\Rightarrow$  Good preconditioner CG!

## Implementing a Preconditioned Newton-CG with quNac

- ▶ Call Conjugate gradients on  $\nabla^2 f_{k+1} d = -\nabla f_{k+1}$  to get  $D_k$  directions of positive curvature
- ▶  $\text{proj}_{D_k}^{\nabla^2 f_{k+1}} = D_k (D_k^T \nabla^2 f_{k+1} D_k)^{-1} D_k^T$  is  $O(p^2)$  because  $D_k^T \nabla^2 f_{k+1} D_k = \text{diagonal!}$
- ▶  $H_{k+1}$  is positive definite and  $H_{k+1} \nabla^2 f_{k+1} D_k = D_k$ , thus  $H_{k+1} \nabla^2 f_{k+1}$  has concentrated Eigen-values  $\Rightarrow$  Good preconditioner CG!
- ▶ Use  $H_k \approx \nabla^2 f_{k+1}$  as a preconditioner in a Newton-CG framework!

## Implementing a Preconditioned Newton-CG with quNac

- ▶ Call Conjugate gradients on  $\nabla^2 f_{k+1} d = -\nabla f_{k+1}$  to get  $D_k$  directions of positive curvature
- ▶  $\text{proj}_{D_k}^{\nabla^2 f_{k+1}} = D_k (D_k^T \nabla^2 f_{k+1} D_k)^{-1} D_k^T$  is  $O(p^2)$  because  $D_k^T \nabla^2 f_{k+1} D_k = \text{diagonal!}$
- ▶  $H_{k+1}$  is positive definite and  $H_{k+1} \nabla^2 f_{k+1} D_k = D_k$ , thus  $H_{k+1} \nabla^2 f_{k+1}$  has concentrated Eigen-values  $\Rightarrow$  Good preconditioner CG!
- ▶ Use  $H_k \approx \nabla^2 f_{k+1}$  as a preconditioner in a Newton-CG framework!

## quNac quasi-Newton action Constrained

►  $H_0, x_0 \in \mathbb{R}, k = 0$

## quNac quasi-Newton action Constrained

- ▶  $H_0, x_0 \in \mathbb{R}, k = 0$
- ▶ While  $|\nabla f_k|/|\nabla f_0| > \epsilon$



## quNac quasi-Newton action Constrained

- ▶  $H_0, x_0 \in \mathbb{R}, k = 0$
- ▶ While  $|\nabla f_k|/|\nabla f_0| > \epsilon$ 
  - ▶ If  $k = 0$

## quNac quasi-Newton action Constrained

- ▶  $H_0, x_0 \in \mathbb{R}, k = 0$
- ▶ While  $|\nabla f_k|/|\nabla f_0| > \epsilon$ 
  - ▶ **If  $k = 0$** 
    - ▶  $s_0 = -H_0 \nabla f_0$

## quNac quasi-Newton action Constrained

- ▶  $H_0, x_0 \in \mathbb{R}, k = 0$
- ▶ While  $|\nabla f_k|/|\nabla f_0| > \epsilon$ 
  - ▶ **If**  $k = 0$ 
    - ▶  $s_0 = -H_0 \nabla f_0$
  - ▶ **Else**

## quNac quasi-Newton action Constrained

- ▶  $H_0, x_0 \in \mathbb{R}, k = 0$
- ▶ While  $|\nabla f_k|/|\nabla f_0| > \epsilon$ 
  - ▶ **If**  $k = 0$ 
    - ▶  $s_0 = -H_0 \nabla f_0$
  - ▶ **Else**
    - ▶  $s_0 = d_{CG}$

## quNac quasi-Newton action Constrained

- ▶  $H_0, x_0 \in \mathbb{R}, k = 0$
- ▶ While  $|\nabla f_k|/|\nabla f_0| > \epsilon$ 
  - ▶ **If**  $k = 0$ 
    - ▶  $s_0 = -H_0 \nabla f_0$
  - ▶ **Else**
    - ▶  $s_0 = d_{CG}$
- ▶ Select step-size  $a_k$

## quNac quasi-Newton action Constrained

- ▶  $H_0, x_0 \in \mathbb{R}, k = 0$
- ▶ While  $|\nabla f_k|/|\nabla f_0| > \epsilon$ 
  - ▶ **If**  $k = 0$ 
    - ▶  $s_0 = -H_0 \nabla f_0$
  - ▶ **Else**
    - ▶  $s_0 = d_{CG}$
  - ▶ Select step-size  $a_k$
  - ▶  $x_{k+1} = x_k + a_k s_k$

## quNac quasi-Newton action Constrained

- ▶  $H_0, x_0 \in \mathbb{R}, k = 0$
- ▶ While  $|\nabla f_k|/|\nabla f_0| > \epsilon$ 
  - ▶ **If**  $k = 0$ 
    - ▶  $s_0 = -H_0 \nabla f_0$
  - ▶ **Else**
    - ▶  $s_0 = d_{CG}$
  - ▶ Select step-size  $a_k$
  - ▶  $x_{k+1} = x_k + a_k s_k$
  - ▶ Calculate  $d_{CG}$  by applying CG to preconditioned system

## quNac quasi-Newton action Constrained

- ▶  $H_0, x_0 \in \mathbb{R}, k = 0$
- ▶ While  $|\nabla f_k|/|\nabla f_0| > \epsilon$ 
  - ▶ **If**  $k = 0$ 
    - ▶  $s_0 = -H_0 \nabla f_0$
  - ▶ **Else**
    - ▶  $s_0 = d_{CG}$
  - ▶ Select step-size  $a_k$
  - ▶  $x_{k+1} = x_k + a_k s_k$
  - ▶ Calculate  $d_{CG}$  by applying CG to preconditioned system
    - ▶  $H_k \nabla^2 f_{k+1} d = -H_k \nabla f_{k+1}.$



## quNac quasi-Newton action Constrained

- ▶  $H_0, x_0 \in \mathbb{R}, k = 0$
- ▶ While  $|\nabla f_k|/|\nabla f_0| > \epsilon$ 
  - ▶ **If**  $k = 0$ 
    - ▶  $s_0 = -H_0 \nabla f_0$
  - ▶ **Else**
    - ▶  $s_0 = d_{CG}$
  - ▶ Select step-size  $a_k$
  - ▶  $x_{k+1} = x_k + a_k s_k$
  - ▶ Calculate  $d_{CG}$  by applying CG to preconditioned system
    - ▶  $H_k \nabla^2 f_{k+1} d = -H_k \nabla f_{k+1}$ .
    - ▶ Store conjugate directions in  $D$  and store  $\nabla^2 f_{k+1} D$ .

## quNac quasi-Newton action Constrained

- ▶  $H_0, x_0 \in \mathbb{R}, k = 0$
- ▶ While  $|\nabla f_k|/|\nabla f_0| > \epsilon$ 
  - ▶ **If**  $k = 0$ 
    - ▶  $s_0 = -H_0 \nabla f_0$
  - ▶ **Else**
    - ▶  $s_0 = d_{CG}$
  - ▶ Select step-size  $a_k$
  - ▶  $x_{k+1} = x_k + a_k s_k$
  - ▶ Calculate  $d_{CG}$  by applying CG to preconditioned system
    - ▶  $H_k \nabla^2 f_{k+1} d = -H_k \nabla f_{k+1}$ .
    - ▶ Store conjugate directions in  $D$  and store  $\nabla^2 f_{k+1} D$ .
  - ▶  $H_{k+1} = H_k + E_k (D, \nabla^2 f_{k+1}(D))$

## LquNac quasi-Newton action Constrained

- ▶  $H_0, x_0 \in \mathbb{R}, k = 0$
- ▶ While  $|\nabla f_k|/|\nabla f_0| > \epsilon$ 
  - ▶ **If**  $k = 0$ 
    - ▶  $s_0 = -H_0 \nabla f_0$
  - ▶ **Else**
    - ▶  $s_0 = d_{CG}$
  - ▶ Select step-size  $a_k$
  - ▶  $x_{k+1} = x_k + a_k s_k$
  - ▶ Calculate  $d_{CG}$  by applying CG to preconditioned system
    - ▶  $H_k \nabla^2 f_{k+1} d = -H_k \nabla f_{k+1}$ .
    - ▶ Store conjugate directions in  $D$  and store  $\nabla^2 f_{k+1} D$ .
  - ▶  $H_{k+1} = H_k + E_k (D, \nabla^2 f_{k+1}(D))$

## LquNac Limited quasi-Newton action Constrained

- ▶  $H_0, x_0 \in \mathbb{R}, k = 0$
- ▶ While  $|\nabla f_k|/|\nabla f_0| > \epsilon$ 
  - ▶ **If**  $k = 0$ 
    - ▶  $s_0 = -H_0 \nabla f_0$
  - ▶ **Else**
    - ▶  $s_0 = d_{CG}$
  - ▶ Select step-size  $a_k$
  - ▶  $x_{k+1} = x_k + a_k s_k$
  - ▶ Calculate  $d_{CG}$  by applying CG to preconditioned system
    - ▶  $OpH_k \nabla^2 f_{k+1} d = -OpH_k \nabla f_{k+1}$ .
    - ▶ Store conjugate directions in  $D$  and store  $\nabla^2 f_{k+1} D$ .
  - ▶  $OpH_{k+1} = H_0^k + E_k(D, \nabla^2 f_{k+1}(D))$

## Logistic L2 Regression tests:

$$\min_w L_w(y, X) + \|w\|_2^2$$

$$L_w(y, X) = \sum_{i=1}^m \ln(1 + \exp(-y_i \langle x^i, w \rangle)) .$$

quNac	vs	BFGS
41		3
quNac	vs	Newton_CG
31		12
LquNac	vs	LBFGS
27		17
LquNac	vs	Newton_CG
14		29

Table: # fastest runs on 44 binary classifications problems from LibSVM

## Logistic pseudo-Huber Regression tests:

$$\min_w L_w(y, X) + R_\mu(w) := \mu \sum_{i=1}^n \left( \sqrt{1 + x_i^2 / \mu^2} - 1 \right).^4$$

quNac vs	BFGS
32	10
quNac vs	Newton.CG
37	4
LquNac vs	LBFGS
18	25
LquNac vs	Newton.CG
24	16

**Table:** # fastest runs on 44 binary classifications problems from LibSVM

<sup>4</sup>Fountoulakis, K., & Gondzio, J. (2013). A Second-Order Method for Strongly Convex l1-regularization Problems.

## Conclusion

- ▶ Framework for approximating a changing inverse Hessian.
- ▶ Has good properties: **Hereditary** and **Descent**.
- ▶ Variable amount of curvature information at each iteration (depends on CG error).
- ▶ Developed Newton-CG Preconditioner for smooth functions.

### In the works:

- ▶ Full and limited memory implementations for non-convex unconstrained.
- ▶ How does extra flexibility help mesh into globalization strategies: Interior point, Trust region, Sequential Quadratic?
- ▶ Connections to other problems: Matrix completion?

## Conclusion

- ▶ Framework for approximating a changing inverse Hessian.
- ▶ Has good properties: **Hereditary** and **Descent**.
- ▶ Variable amount of curvature information at each iteration (depends on CG error).
- ▶ Developed Newton-CG Preconditioner for smooth functions.

### In the works:

- ▶ Full and limited memory implementations for non-convex unconstrained.
- ▶ How does extra flexibility help mesh into globalization strategies: Interior point, Trust region, Sequential Quadratic?
- ▶ Connections to other problems: Matrix completion?



## References



Gower, R. M., (2014).

New quasi-Newton family through action constraints (*in preparation*)



Fletcher, B. R., Powell, M. J. D. (1960).

A rapidly convergent descent method for minimization.



Davidon, W. C. (1959). Variable metric method for minimization.



Goldfarb, D. (1970).

A Family of Variable-Metric Methods Derived by Variational Means.

Mathematics of Computation, 24(109), 23.



Shanno, D. F. (1971).

Conditioning of Quasi-Newton Methods for Function Minimization.

Mathematics of Computation, 24(111), 647656.

## Underlying Matrix Optimization Problem

$$\begin{aligned} \min_E & \|E\|_{Frobenius(W)}^2 \\ \text{s.t.} \quad & ED = RD \\ & E = E^T \end{aligned}$$

Which has a low rank- $3p$  solution.

$$E = W^{-1} \text{proj}_D^{W^{-1}} R \left( I - \text{proj}_D^{W^{-1}} W^{-1} \right) + R \text{proj}_D^{W^{-1}} W^{-1}.$$

This is a matrix completion problem where one knows the desired matrix is symmetric and can only observe its action on a small subspace.