

# Randomized iterative methods for linear systems and inverting matrices

Robert Gower  
joint work with Peter Richtárik



Optimization and Big Data 2015, 7th of May, Edinburgh.



Gower, Robert M., Richtárik, Peter, April 2015.

Randomized Iterative Methods for Linear Systems (in progress)



Gower, Robert M., Richtárik, Peter, April 2015.

Randomized Iterative Methods for Inverting Matrices (in progress)

# The Problem

## The Problem

Solve a consistent linear system  $Ax_* = b$ , where  $A \in \mathbb{R}^{m \times n}$ ,  $m \geq n$ .

$$\begin{bmatrix} \text{---} A_1: \text{---} \\ \text{---} A_2: \text{---} \\ \vdots \\ \vdots \\ \text{---} A_{m-1}: \text{---} \\ \text{---} A_m: \text{---} \end{bmatrix} \begin{bmatrix} x_*^1 \\ \vdots \\ x_*^n \end{bmatrix} = \begin{bmatrix} b^1 \\ b^2 \\ \vdots \\ \vdots \\ b^{m-1} \\ b^m \end{bmatrix}$$

Solve with an iterative method

$$x_{k+1} = \text{update\_formula}(A, x_k)$$

such that  $x_k \rightarrow x_*$ .

# The Problem

## The Problem

Solve a consistent linear system  $Ax_* = b$ , where  $A \in \mathbb{R}^{m \times n}$ ,  $m \geq n$ .

$$\begin{bmatrix} \text{---} A_{1:} \text{---} \\ \text{---} A_{2:} \text{---} \\ \vdots \\ \vdots \\ \text{---} A_{m-1:} \text{---} \\ \text{---} A_{m:} \text{---} \end{bmatrix} \begin{bmatrix} x_*^1 \\ \vdots \\ x_*^n \end{bmatrix} = \begin{bmatrix} b^1 \\ b^2 \\ \vdots \\ \vdots \\ b^{m-1} \\ b^m \end{bmatrix}$$

Solve with an iterative method

$$x_{k+1} = \text{update\_formula}(A, x_k)$$

such that  $x_k \rightarrow x_*$ .

**Notation:** Let  $\|x\|_B^2 := x^T B x$  for  $B \succ 0$ .

# The Problem

## The Problem

Solve a consistent linear system  $Ax_* = b$ , where  $A \in \mathbb{R}^{m \times n}$ ,  $m \geq n$ .

$$\begin{bmatrix} \text{---} A_{1:} \text{---} \\ \text{---} A_{2:} \text{---} \\ \vdots \\ \vdots \\ \text{---} A_{m-1:} \text{---} \\ \text{---} A_{m:} \text{---} \end{bmatrix} \begin{bmatrix} x_*^1 \\ \vdots \\ x_*^n \end{bmatrix} = \begin{bmatrix} b^1 \\ b^2 \\ \vdots \\ \vdots \\ b^{m-1} \\ b^m \end{bmatrix}$$

Solve with an iterative method

$$x_{k+1} = \text{update\_formula}(A, x_k)$$

such that  $x_k \rightarrow x_*$ .

**Notation:** Let  $\|x\|_B^2 := x^T B x$  for  $B \succ 0$ .

# Table of Contents

## Old methods return randomized

The Kaczmarz method (Stochastic gradient)

The Coordinate Descent method (Gauss Seidel)

## Framework for randomized methods

Geometry and Duality

Convergence analysis

New Gaussian methods

## Numerical tests

## Iteratively Inverting matrices

Future work: Randomized Preconditioning?

## The Return of old methods

- ▶ Old methods (Kaczmarz 1937, Gauss-Seidel 1823) make a randomized return, why?
- ▶ Often suitable for Big Data problems (short recurrence, low memory,...etc)
- ▶ Easy to implement
- ▶ Easy to analyse, good complexity
- ▶ Often fits in parallel architecture

- └ Old methods return randomized
- └ The Kaczmarz method (Stochastic gradient)

## Kaczmarz method



Choose the  $i$ th row then iterate

$$x_{k+1} = \arg \min_{x \in \mathbb{R}^n} \|x - x_k\|_2^2 \quad \text{subject to} \quad A_{i:}x = b^i.$$

$$x_{k+1} = x_k - \frac{A_{i:}x_k - b^i}{\|A_{i:}\|_2^2} A_{i:}^T$$

- ▶ Developed in 1937 Kaczmarz
- ▶ Implemented in the first CT scanner 1972

1




<sup>1</sup>G.N. Hounsfield. Computerized transverse axial scanning (tomography): Part I. description of the system. British Journal Radiology. 1973



- └ Old methods return randomized
- └ The Kaczmarz method (Stochastic gradient)

## Kaczmarz Interpretation

$$x_{k+1} = \arg \min_{x \in \mathbb{R}^n} \|x - x_k\|_2^2 \quad \text{subject to} \quad A_i: x = b^i.$$




$$\begin{aligned} \{x \mid A_i: x = b^i\} &= \{x \mid A_i: (x - x_*) = 0\} \\ &= x_* + \{x \mid A_i: x = 0\} \\ &= x_* + \mathbf{Null}(A_i) \end{aligned}$$

- └ Old methods return randomized
- └ The Kaczmarz method (Stochastic gradient)

## Kaczmarz Interpretation

$$x_{k+1} = \arg \min_{x \in \mathbb{R}^n} \|x - x_k\|_2^2 \quad \text{subject to} \quad A_i: x = b^i.$$



$$\begin{aligned} \{x \mid A_i: x = b^i\} &= \{x \mid A_i: (x - x_*) = 0\} \\ &= x_* + \{x \mid A_i: x = 0\} \\ &= x_* + \mathbf{Null}(A_i) \end{aligned}$$

- └ Old methods return randomized
- └ The Kaczmarz method (Stochastic gradient)

## How to choose $i$

$$x_{k+1} = \arg \min_{x \in \mathbb{R}^n} \|x - x_k\|_2^2 \quad \text{subject to} \quad A_i x = b^i.$$

- ▶ Traditional Kaczmarz: Cycle  $i = 1, 2, \dots, m$ . Slow in practice + difficult to interpret complexity
- ▶ Pick  $i$  with probability  $p_i = 1/m$ . Better in practice + difficult to interpret complexity
- ▶ **Break-Through (Strohmer & Vershynin, 2009)**: pick  $i$  with probability  $p_i = \|A_{i:}\|_2^2 / \|A\|_F^2$ .

$$\mathbf{E} [\|x_k - x_*\|_2^2] \leq \left(1 - \frac{\lambda_{\min}(A^T A)}{\|A\|_F^2}\right)^k \|x_0 - x_*\|_2^2.$$

$$\lambda_{\min}(A^T A) / \|A\|_F^2 = 1 / \|A\|_F^2 \|A^\dagger\|_2^2$$

- └ Old methods return randomized
- └ The Kaczmarz method (Stochastic gradient)

## How to choose $i$

$$x_{k+1} = \arg \min_{x \in \mathbb{R}^n} \|x - x_k\|_2^2 \quad \text{subject to} \quad A_i x = b^i.$$

- ▶ Traditional Kaczmarz: Cycle  $i = 1, 2, \dots, m$ . Slow in practice + difficult to interpret complexity
- ▶ Pick  $i$  with probability  $p_i = 1/m$ . Better in practice + difficult to interpret complexity
- ▶ **Break-Through (Strohmer & Vershynin, 2009)**: pick  $i$  with probability  $p_i = \|A_i\|_2^2 / \|A\|_F^2$ .

$$\mathbf{E} [\|x_k - x_*\|_2^2] \leq \left(1 - \frac{\lambda_{\min}(A^T A)}{\|A\|_F^2}\right)^k \|x_0 - x_*\|_2^2.$$

$$\lambda_{\min}(A^T A) / \|A\|_F^2 = 1 / \|A\|_F^2 \|A^\dagger\|_2^2$$

└ Old methods return randomized

└ The Coordinate Descent method (Gauss Seidel)

## Coordinate Descent (Gauss-Seidel)

Choose the  $i$ th coordinate then

$$x_{k+1} = \arg \min_{x \in \mathbb{R}^n} \|Ax - b\|_2^2 \quad \text{subject to} \quad x = x_k + te_i, \quad t \in \mathbb{R}.$$

$$x_{k+1} = x_k - \frac{(A_{:i})^T (Ax_k - b)}{\|A_{:i}\|_2^2} e_i$$

Note that  $\|Ax - b\|_2^2 = \|A(x - x_*)\|_2^2 = \|x - x_*\|_{A^T A}^2$

## Convergence (Leventhal & Lewis, 2010)


Pick  $i$  with probability  $p_i = \|A_{:i}\|_2^2 / \|A\|_F^2$ .

$$\mathbf{E} [\|x_k - x_*\|_{A^T A}^2] \leq \left(1 - \frac{\lambda_{\min}(A^T A)}{\|A\|_F^2}\right)^k \|x_0 - x_*\|_{A^T A}^2.$$


- └ Old methods return randomized
- └ The Coordinate Descent method (Gauss Seidel)

## Coordinate Descent Interpretation

$$x_{k+1} = \arg \min_{x \in \mathbb{R}^n} \|x - x_*\|_{A^T A}^2 \quad \text{subject to} \quad x = x_k + t e_i, \quad t \in \mathbb{R}.$$




Kaczmarz Interpretation




- └ Old methods return randomized
- └ The Coordinate Descent method (Gauss Seidel)

## Coordinate Descent Interpretation

$$x_{k+1} = \arg \min_{x \in \mathbb{R}^n} \|x - x_*\|_{A^T A}^2 \quad \text{subject to} \quad x = x_k + t e_i, \quad t \in \mathbb{R}.$$



Kaczmarz Interpretation



# Table of Contents

## Old methods return randomized

The Kaczmarz method (Stochastic gradient)

The Coordinate Descent method (Gauss Seidel)

## Framework for randomized methods

Geometry and Duality

Convergence analysis

New Gaussian methods

## Numerical tests

## Iteratively Inverting matrices

Future work: Randomized Preconditioning?



## Framework for designing randomized methods

Choose  $B \succ 0 \in \mathbb{R}^{n \times n}$  and a **random** matrix  $S$  independently drawn at each iteration  $k$ . **Two** viewpoints of the **same** method.

$$(I) \quad x_{k+1} = \arg \min_{x \in \mathbb{R}^n} \|x - x_k\|_B^2 \quad \text{s. t.} \quad S^T A x = S^T b,$$

$$(II) \quad x_{k+1} = \arg \min_{x \in \mathbb{R}^n} \|x - x_k\|_B^2 \quad \text{s. t.} \quad x \in x_k + B^{-1} \text{Range} \left( A^T S \right)$$

(I) : Project  $x_k$  onto a **randomly compacted** system.

$$\boxed{S^T} \begin{bmatrix} A \\ x \end{bmatrix} = \boxed{S^T A} \begin{bmatrix} x \end{bmatrix}$$

**Kaczmarz** fits nicely with  $B = I$  and  $S = e_i$ .

**Block Kaczmarz** choose  $B = I$  and  $S = I_{:,C}$  a subset of columns of identity.

## Framework for designing randomized methods

Choose  $B \succ 0 \in \mathbb{R}^{n \times n}$  and a **random** matrix  $S$  independently drawn at each iteration  $k$ . **Two** viewpoints of the **same** method.

$$(I) \quad x_{k+1} = \arg \min_{x \in \mathbb{R}^n} \|x - x_k\|_B^2 \quad \text{s. t.} \quad S^T A x = S^T b,$$

$$(II) \quad x_{k+1} = \arg \min_{x \in \mathbb{R}^n} \|x - x_k\|_B^2 \quad \text{s. t.} \quad x \in x_k + B^{-1} \text{Range} \left( A^T S \right)$$

(I) : Project  $x_k$  onto a **randomly compacted** system.

$$\boxed{S^T} \quad \boxed{A} \quad \boxed{x} = \boxed{S^T A} \quad \boxed{x}$$

**Kaczmarz** fits nicely with  $B = I$  and  $S = e_i$ .

**Block Kaczmarz** choose  $B = I$  and  $S = I_{:,C}$  a subset of columns of identity.

## Coordinate descent methods fit (II)

$$(II) \quad x_{k+1} = \arg \min_{x \in \mathbb{R}^n} \|x - x_*\|_B^2 \quad \text{subject to} \quad x \in x_k + \text{Range} \left( B^{-1} A^T S \right)$$

- **Least-Squares Coord. Desc:** With  $B = A^T A$  and  $S = A e_i = A_{:,i}$

$$x_{k+1} = \arg \min_{x \in \mathbb{R}^n} \|Ax - b\|_2^2 \quad \text{subject to} \quad x = x_k + t e_i, \quad t \in \mathbb{R}.$$

**Stochastic Newton (SDNA<sup>2</sup> Method 1)** Let  $S = A_{:,C} = A_{:,C}$  subset of columns of  $A$

$$x_{k+1} = \arg \min_{x \in \mathbb{R}^n} \|Ax - b\|_2^2 \quad \text{subject to} \quad x = x_k + t l_{:,C}, \quad t \in \mathbb{R}^{|C|}.$$

- **Positive Definite Coord. Desc:** When  $A \succ 0$ ,  $B = A$  and  $S = l_{:,C}$  then

$$x_{k+1} = \arg \min_{x \in \mathbb{R}^n} \underbrace{\frac{1}{2} x^T A x - x^T b}_{= \|x - x_*\|_A^2} \quad \text{subject to} \quad x = x_k + t l_{:,C}, \quad t \in \mathbb{R}^{|C|},$$


---

<sup>2</sup>Qu, Z., Richtárik, P., Takáč, M., & Fercoq, O. (2015). SDNA: Stochastic Dual Newton Ascent for Empirical Risk Minimization.

Let  $\langle x, y \rangle = x^T B y$  in  $\mathbb{R}^n$ .  $\{x \mid S^T A x = S^T b\} = x_* + \{x \mid S^T A x = 0\}$  and



Let  $\langle x, y \rangle = x^T B y$  in  $\mathbb{R}^n$ .  $\{x \mid S^T A x = S^T b\} = x_* + \{x \mid S^T A x = 0\}$  and

$$\text{Null}(S^T A) \oplus B^{-1} \text{Range}(A^T S) = \mathbb{R}^n.$$




Let  $\langle x, y \rangle = x^T B y$  in  $\mathbb{R}^n$ .  $\{x \mid S^T A x = S^T b\} = x_* + \{x \mid S^T A x = 0\}$  and

$$\text{Null}(S^T A) \oplus B^{-1} \text{Range}(A^T S) = \mathbb{R}^n.$$



Let  $\langle x, y \rangle = x^T B y$  in  $\mathbb{R}^n$ .  $\{x \mid S^T A x = S^T b\} = x_* + \{x \mid S^T A x = 0\}$  and

$$\text{Null}(S^T A) \oplus B^{-1} \text{Range}(A^T S) = \mathbb{R}^n.$$




| Project  $x_k$  onto  $x_* + \text{Null}(S^T A)$

$$x_{k+1} = \arg \min_{x \in \mathbb{R}^n} \|x - x_k\|_B^2 \quad \text{subject to} \quad S^T A x = S^T b$$

Let  $\langle x, y \rangle = x^T B y$  in  $\mathbb{R}^n$ .  $\{x \mid S^T A x = S^T b\} = x_* + \{x \mid S^T A x = 0\}$  and

$$\text{Null}(S^T A) \oplus B^{-1} \text{Range}(A^T S) = \mathbb{R}^n.$$



I Project  $x_k$  onto  $x_* + \text{Null}(S^T A)$

$$x_{k+1} = \arg \min_{x \in \mathbb{R}^n} \|x - x_k\|_B^2 \quad \text{subject to} \quad S^T A x = S^T b$$


II Project  $x_*$  onto  $x_k + B^{-1} \text{Range}(A^T S)$

$$x_{k+1} = \arg \min_{x \in \mathbb{R}^n} \|x - x_*\|_B^2 \quad \text{subject to} \quad x \in x_k + B^{-1} \text{Range}(A^T S)$$



Let  $\langle x, y \rangle = x^T B y$  in  $\mathbb{R}^n$ .  $\{x \mid S^T A x = S^T b\} = x_* + \{x \mid S^T A x = 0\}$  and

$$\text{Null}(S^T A) \oplus B^{-1} \text{Range}(A^T S) = \mathbb{R}^n.$$



I Project  $x_k$  onto  $x_* + \text{Null}(S^T A)$

$$x_{k+1} = \arg \min_{x \in \mathbb{R}^n} \|x - x_k\|_B^2 \quad \text{subject to} \quad S^T A x = S^T b$$

II Project  $x_*$  onto  $x_k + B^{-1} \text{Range}(A^T S)$

$$x_{k+1} = \arg \min_{x \in \mathbb{R}^n} \|x - x_*\|_B^2 \quad \text{subject to} \quad x \in x_k + B^{-1} \text{Range}(A^T S)$$

# The Solution

Assuming  $A^T S$  has full column rank  $\Rightarrow$  closed form solution

## II Project $x_*$ onto $x_k + B^{-1}\text{Range}(A^T S)$

$$\begin{aligned} x_{k+1} &= x_k + \text{proj}_{B^{-1}\text{Range}(A^T S)}(x_* - x_k) \\ &= x_k + B^{-1}A^T S(S^T A B^{-1}A^T S)^{-1}S^T A(x_k - x_*) \\ &= x_k + B^{-1}A^T S \underbrace{(S^T A B^{-1}A^T S)^{-1}S^T A}_{\text{Solve small system.}}(x_k - x_*) \end{aligned}$$

## I Project $x_k$ onto $x_* + \text{Null}(S^T A)$

$$\begin{aligned} x_{k+1} &= x_* + \text{proj}_{\text{Null}(A^T S)}(x_k - x_*) \\ &= x_* + \underbrace{(I - B^{-1}A^T S(S^T A B^{-1}A^T S)^{-1}S^T A)}_Z(x_k - x_*) \\ &= x_* + (I - B^{-1}Z)(x_k - x_*). \end{aligned}$$

# The Solution

Assuming  $A^T S$  has full column rank  $\Rightarrow$  closed form solution

II Project  $x_*$  onto  $x_k + B^{-1}\text{Range}(A^T S)$

$$\begin{aligned} x_{k+1} &= x_k + \text{proj}_{B^{-1}\text{Range}(A^T S)}(x_* - x_k) \\ &= x_k + B^{-1}A^T S(S^T A B^{-1}A^T S)^{-1}S^T A(x_k - x_*) \\ &= x_k + B^{-1}A^T S \underbrace{(S^T A B^{-1}A^T S)^{-1}S^T A}_{\text{Solve small system.}}(x_k - x_*) \end{aligned}$$

I Project  $x_k$  onto  $x_* + \text{Null}(S^T A)$

$$\begin{aligned} x_{k+1} &= x_* + \text{proj}_{\text{Null}(A^T S)}(x_k - x_*) \\ &= x_* + (I - B^{-1}A^T S \underbrace{(S^T A B^{-1}A^T S)^{-1}S^T A}_Z)(x_k - x_*) \\ &= x_* + (I - B^{-1}Z)(x_k - x_*). \end{aligned}$$

## The Fixed point form

All randomness is in the range space projection  $B^{-1}Z$

$$Z \stackrel{\text{def}}{=} A^T S (S^T A B^{-1} A^T S)^{-1} S^T A.$$

For analysis, fixed point form

$$x_{k+1} - x_* = (I - B^{-1}Z)(x_k - x_*).$$

# The Fixed point form

All randomness is in the range space projection  $B^{-1}Z$

$$Z \stackrel{\text{def}}{=} A^T S (S^T A B^{-1} A^T S)^{-1} S^T A.$$

For analysis, fixed point form

$$\begin{aligned} \mathbf{E}[x_{k+1} - x_* \mid x_k] &= (I - B^{-1} \mathbf{E}[Z])(x_k - x_*). \\ \mathbf{E}[\mathbf{E}[x_{k+1} - x_* \mid x_k]] &= \mathbf{E}[x_{k+1} - x_*] \\ &= \mathbf{E}[(I - B^{-1} \mathbf{E}[Z])(x_k - x_*)] \\ &= (I - B^{-1} \mathbf{E}[Z]) \mathbf{E}[x_k - x_*]. \end{aligned}$$

## The Fixed point form

All randomness is in the range space projection  $B^{-1}Z$

$$Z \stackrel{\text{def}}{=} A^T S (S^T A B^{-1} A^T S)^{-1} S^T A.$$

For analysis, fixed point form

$$\begin{aligned} \mathbf{E}[x_{k+1} - x_* | x_k] &= (I - B^{-1} \mathbf{E}[Z])(x_k - x_*). \\ \mathbf{E}[\mathbf{E}[x_{k+1} - x_* | x_k]] &= \mathbf{E}[x_{k+1} - x_*] \\ &= \mathbf{E}[(I - B^{-1} \mathbf{E}[Z])(x_k - x_*)] \\ &= (I - B^{-1} \mathbf{E}[Z]) \mathbf{E}[x_k - x_*]. \end{aligned}$$

### Convergence Theorems

$$\|\mathbf{E}[x_k] - x_*\| \leq \left(1 - \lambda_{\min}(B^{-1/2} \mathbf{E}[Z] B^{-1/2})\right)^k \|x_0 - x_*\|$$

and when  $\mathbf{E}[Z]$  nonsingular

$$\mathbf{E}[\|x_k - x_*\|] \leq \left(1 - \lambda_{\min}(B^{-1/2} \mathbf{E}[Z] B^{-1/2})\right)^k \|x_0 - x_*\|$$

## The Fixed point form

All randomness is in the range space projection  $B^{-1}Z$

$$Z \stackrel{\text{def}}{=} A^T S (S^T A B^{-1} A^T S)^{-1} S^T A.$$

For analysis, fixed point form

$$\begin{aligned} \mathbf{E}[x_{k+1} - x_* | x_k] &= (I - B^{-1} \mathbf{E}[Z])(x_k - x_*). \\ \mathbf{E}[\mathbf{E}[x_{k+1} - x_* | x_k]] &= \mathbf{E}[x_{k+1} - x_*] \\ &= \mathbf{E}[(I - B^{-1} \mathbf{E}[Z])(x_k - x_*)] \\ &= (I - B^{-1} \mathbf{E}[Z]) \mathbf{E}[x_k - x_*]. \end{aligned}$$

### Convergence Theorems

$$\|\mathbf{E}[x_k] - x_*\| \leq \left(1 - \lambda_{\min}(B^{-1/2} \mathbf{E}[Z] B^{-1/2})\right)^k \|x_0 - x_*\|$$

and when  $\mathbf{E}[Z]$  nonsingular

$$\mathbf{E}[\|x_k - x_*\|] \leq \left(1 - \lambda_{\min}(B^{-1/2} \mathbf{E}[Z] B^{-1/2})\right)^k \|x_0 - x_*\|$$

## Theorem (General S)

Let  $S$  be a random matrix such that  $A^T S$  full column rank. Then for all  $k \geq 0$ ,

$$\|\mathbf{E}[x_k] - x_*\| \leq \rho^k \|x_0 - x_*\|,$$

where

$$\rho = 1 - \lambda_{\min}(B^{-1/2} \mathbf{E}[Z] B^{-1/2}) \quad \text{and} \quad 0 \leq \rho \leq 1.$$

## Proof.

Taking conditional expectation with respect to  $x_k$ , we get

$$\mathbf{E}[x_{k+1} - x_* \mid x_k] = (I - B^{-1} \mathbf{E}[Z])(x_k - x_*). \quad (1)$$

Taking full expectation, we get

$$\begin{aligned} \mathbf{E}[x_{k+1} - x_*] &= \mathbf{E}[\mathbf{E}[x_{k+1} - x_* \mid x_k]] \\ &\stackrel{(1)}{=} \mathbf{E}[(I - B^{-1} \mathbf{E}[Z])(x_k - x_*)] \\ &= (I - B^{-1} \mathbf{E}[Z]) \mathbf{E}[x_k - x_*]. \end{aligned}$$

Now unroll the recurrence and apply the operator norm. As  $B^{-1}Z$  is a projection, by Jensen's inequality with the convex functions  $\lambda_{\max}$  and  $-\lambda_{\min}$ , we have

$$0 \leq \lambda_{\max}(B^{-1} \mathbf{E}[Z]) \leq \lambda_{\max}(B^{-1} Z) \leq 1. \quad \square$$



## Unifying previous methods & analysis

### Theorem (Discrete random vector)

Let  $S$  be discrete r.v. such  $S = s_i \in \mathbb{R}^n$  (for concreteness, think of  $s_i = e_i$ ) with probability  $p_i > 0$ , for  $i = 1, \dots, m$ , and let

$$\mathbf{S} = [s_1, \dots, s_m].$$

Then

$$x_{k+1} = x_k + \frac{s_i^T (Ax_k - b)}{s_i^T AB^{-1}A^T s_i} B^{-1}A^T s_i, \quad \text{with prob } p_i.$$

If we choose

$$p_i = \frac{s_i^T AB^{-1}A^T s_i}{\|B^{-1/2}A^T \mathbf{S}\|_F^2}, \quad \text{for } i = 1, \dots, m,$$

then

$$\mathbf{E}[Z] = \frac{A^T \mathbf{S} \mathbf{S}^T A}{\|B^{-1/2}A^T \mathbf{S}\|_F^2} \quad \text{and} \quad \rho = 1 - \frac{\lambda_{\min}(B^{-1/2}A^T \mathbf{S} \mathbf{S}^T AB^{-1/2})}{\|B^{-1/2}A^T \mathbf{S}\|_F^2}.$$

Furthermore, if  $\mathbf{S}^T A$  has full column rank then  $\rho < 1$ .

## Proof.

$$\begin{aligned}\mathbf{E}[Z] &= \sum_{i=1}^m A^T s_i (s_i^T A B^{-1} A^T s_i)^{-1} s_i^T A p_i \\ &= \frac{1}{\|B^{-1/2} A^T \mathbf{S}\|_F^2} \sum_{i=1}^m A^T s_i s_i^T A \\ &= \frac{1}{\|B^{-1/2} A^T \mathbf{S}\|_F^2} A^T \mathbf{S} \mathbf{S}^T A.\end{aligned}$$

Thus the  $\rho$  is given by

$$\rho = 1 - \lambda_{\min} \left( B^{-1/2} \mathbf{E}[Z] B^{-1/2} \right) = 1 - \frac{\lambda_{\min} \left( B^{-1/2} A^T \mathbf{S} \mathbf{S}^T A B^{-1/2} \right)}{\|B^{-1/2} A^T \mathbf{S}\|_F^2}.$$

As  $\mathbf{S}^T A$  has full column rank,  $\mathbf{E}[Z]$  is positive definite and  $\rho < 1$ . □

## Unifying previous methods & analysis

$$p_i = \frac{s_i^T A B^{-1} A^T s_i}{\|B^{-1/2} A^T \mathbf{S}\|_F^2} \quad \text{with} \quad \rho = 1 - \frac{\lambda_{\min}(B^{-1/2} A^T \mathbf{S} \mathbf{S}^T A B^{-1/2})}{\|B^{-1/2} A^T \mathbf{S}\|_F^2}.$$

Name	$B$	$S$	$\mathbf{S}$	$p_i$	$1 - \rho$
Kaczmarz	$I$	$e_i$	$I$	$\ A_{i:}\ _2^2 / \ A\ _F^2$	$\lambda_{\min}(A^T A) / \ A\ _F^2$
CD $\ Ax - b\ _2^2$	$A^T A$	$A_{:i}$	$A$	$\ A_{:i}\ _2^2 / \ A\ _F^2$	$\lambda_{\min}(A^T A) / \ A\ _F^2$
CD $x^T A x / 2 - x^T b$	$A$	$e_i$	$I$	$A_{ii} / \text{Tr}(A)$	$\lambda_{\min}(A) / \text{Tr}(A)$

New possibilities suggested:

- Covers new cases, e.g.,  $S = \alpha_i e_i + \alpha_j e_j$

## Unifying previous methods & analysis

$$p_i = \frac{s_i^T A B^{-1} A^T s_i}{\|B^{-1/2} A^T \mathbf{S}\|_F^2} \quad \text{with} \quad \rho = 1 - \frac{\lambda_{\min}(B^{-1/2} A^T \mathbf{S} \mathbf{S}^T A B^{-1/2})}{\|B^{-1/2} A^T \mathbf{S}\|_F^2}.$$

Name	$B$	$S$	$\mathbf{S}$	$p_i$	$1 - \rho$
Kaczmarz	$I$	$e_i$	$I$	$\ A_{i:}\ _2^2 / \ A\ _F^2$	$\lambda_{\min}(A^T A) / \ A\ _F^2$
CD $\ Ax - b\ _2^2$	$A^T A$	$A_{:,i}$	$A$	$\ A_{:,i}\ _2^2 / \ A\ _F^2$	$\lambda_{\min}(A^T A) / \ A\ _F^2$
CD $x^T A x / 2 - x^T b$	$A$	$e_i$	$I$	$A_{ii} / \text{Tr}(A)$	$\lambda_{\min}(A) / \text{Tr}(A)$

New possibilities suggested:

- Covers new cases, e.g.,  $S = \alpha_i e_i + \alpha_j e_j$
- For  $B = I$ , then ideally  $\mathbf{S}^T \approx A^\dagger$  then  $\rho \approx 1 - 1/n$ . If we have a preconditioner  $P \approx A^\dagger$  then  $S = \text{sample rows of } P$ .

## Unifying previous methods & analysis

$$p_i = \frac{s_i^T A B^{-1} A^T s_i}{\|B^{-1/2} A^T \mathbf{S}\|_F^2} \quad \text{with} \quad \rho = 1 - \frac{\lambda_{\min}(B^{-1/2} A^T \mathbf{S} \mathbf{S}^T A B^{-1/2})}{\|B^{-1/2} A^T \mathbf{S}\|_F^2}.$$

Name	$B$	$S$	$\mathbf{S}$	$p_i$	$1 - \rho$
Kaczmarz	$I$	$e_i$	$I$	$\ A_{i:}\ _2^2 / \ A\ _F^2$	$\lambda_{\min}(A^T A) / \ A\ _F^2$
CD $\ Ax - b\ _2^2$	$A^T A$	$A_{:,i}$	$A$	$\ A_{:,i}\ _2^2 / \ A\ _F^2$	$\lambda_{\min}(A^T A) / \ A\ _F^2$
CD $x^T A x / 2 - x^T b$	$A$	$e_i$	$I$	$A_{ii} / \text{Tr}(A)$	$\lambda_{\min}(A) / \text{Tr}(A)$

New possibilities suggested:

- Covers new cases, e.g.,  $S = \alpha_i e_i + \alpha_j e_j$
- For  $B = I$ , then ideally  $\mathbf{S}^T \approx A^\dagger$  then  $\rho \approx 1 - 1/n$ . If we have a preconditioner  $P \approx A^\dagger$  then  $S = \text{sample rows of } P$ .

## Gaussian based sampling

**Why not make  $S$  a continuous random matrix?**

Sample  $S = \xi \sim N(0, \Sigma)$  a normal random variable then

$$Z = A^T S (S^T A B^{-1} A^T S)^{-1} S^T A = \frac{A^T \xi \xi^T A}{\xi^T A B^{-1} A^T \xi}.$$

$$x_{k+1} = x_k - \frac{\xi^T (A x_k - b)}{\xi^T A B^{-1} A^T \xi} B^{-1} A^T \xi.$$

Iteration cost  $O(\text{product } A^T \cdot \xi)$ .

The convergence rate determined by

$$\rho = 1 - \lambda_{\min}(B^{-1/2} \mathbf{E}[Z] B^{-1/2}) = 1 - \lambda_{\min}\left(\mathbf{E}\left[\frac{\bar{\xi} \bar{\xi}^T}{\bar{\xi}^T \bar{\xi}}\right]\right),$$

where  $\bar{\xi} = B^{-1/2} A^T \xi \sim N(0, \Omega)$ , and  $\Omega = B^{-1/2} A \Sigma A^T B^{-1/2}$ .

## Gaussian based sampling

**Why not make  $S$  a continuous random matrix?**

Sample  $S = \xi \sim N(0, \Sigma)$  a normal random variable then

$$Z = A^T S (S^T A B^{-1} A^T S)^{-1} S^T A = \frac{A^T \xi \xi^T A}{\xi^T A B^{-1} A^T \xi}.$$

$$x_{k+1} = x_k - \frac{\xi^T (A x_k - b)}{\xi^T A B^{-1} A^T \xi} B^{-1} A^T \xi.$$

Iteration cost  $O(\text{product } A^T \cdot \xi)$ .

The convergence **rate** determined by

$$\rho = 1 - \lambda_{\min}(B^{-1/2} \mathbf{E}[Z] B^{-1/2}) = 1 - \lambda_{\min} \left( \mathbf{E} \left[ \frac{\bar{\xi} \bar{\xi}^T}{\bar{\xi}^T \bar{\xi}} \right] \right),$$

where  $\bar{\xi} = B^{-1/2} A^T \xi \sim N(0, \Omega)$ , and  $\Omega = B^{-1/2} A \Sigma A^T B^{-1/2}$ .

## New Gaussian Methods

Sample  $S = \xi \sim N(0, \Sigma)$ . Let  $\eta \sim N(0, I)$ .

**Gauss. Kaczmarz**  $B = I$  and  $\Sigma = I$

$$x_{k+1} = \arg \min_{x \in \mathbb{R}^n} \|x - x_k\|_2^2 \quad \text{subject to} \quad \eta^T (Ax - b) = 0.$$

**Gauss Least-squares**  $B = A^T A$  and  $\Sigma = AA^T$

$$x_{k+1} = \arg \min_{x \in \mathbb{R}^n} \|Ax - b\|_2^2 \quad \text{subject to} \quad x = x_k + t\eta, \quad t \in \mathbb{R}.$$

**Gauss. Pos. Def.**  $B = A$  and  $\Sigma = I$

$$x_{k+1} = \arg \min_{x \in \mathbb{R}^n} 1/2 x^T A x - x^T b \quad \text{subject to} \quad x = x_k + t\eta, \quad t \in \mathbb{R}.$$



# Table of Contents

## Old methods return randomized

The Kaczmarz method (Stochastic gradient)

The Coordinate Descent method (Gauss Seidel)

## Framework for randomized methods

Geometry and Duality

Convergence analysis

New Gaussian methods

## Numerical tests

## Iteratively Inverting matrices

Future work: Randomized Preconditioning?

# Dense Overdetermined Gaussian Matrix




Figure :  $m \times n = 500 \times 1000$ ,  $A = \text{randn}(m, n)$

Dense matrix  $\Rightarrow$  High iteration cost of Gaussian methods  $O(A \cdot \eta)$ .

# Sparse Square Gaussian Matrix





Figure :  $m \times n = 10000 \times 1000$ , density =  $1/\sqrt{m} = 1\%$ ;  $\kappa = \sqrt{n}$ ;  $A = \text{sprandsym}(n, \text{density}, \text{rc})$

Sparse matrices  $\Rightarrow$  Gauss methods become competitive.

# Regression SUSY

## The SUSY<sup>3</sup> Classification problem



Solving least-squares regression  $\min \|Ax - b\|_2^2$  with  $m = 5 \cdot 10^6$  and  $n = 18$

<sup>3</sup>Baldi, P., P. Sadowski, and D. Whiteson. Searching for Exotic Particles in High-energy Physics with Deep Learning. Nature Communications 5 (July 2, 2014)

# Table of Contents

## Old methods return randomized

The Kaczmarz method (Stochastic gradient)

The Coordinate Descent method (Gauss Seidel)

## Framework for randomized methods

Geometry and Duality

Convergence analysis

New Gaussian methods

## Numerical tests

## Iteratively Inverting matrices

Future work: Randomized Preconditioning?

## Why iteratively invert a matrix $A \in \mathbb{R}^{n \times n}$ ?

- ▶ Needed to calculate Schur complements, a projection operator...etc
- ▶ Iterative is good when we can tolerate an error
- ▶ Iterative is good when we have an initial guess  $X_0 \approx A^{-1}$ .
- ▶ Staging for randomized variable metric methods and randomized Preconditioning.

New context:  $A \in \mathbb{R}^{n \times n}$  non-singular.

# Framework

- ▶ Assume we observe  $S^T A$  where  $S$  is random.
- ▶ Given  $X_k \approx A \in \mathbb{R}^{n \times n}$ , we want to iteratively calculate

$$X_{k+1} = \text{update\_formula}(S^T A, X_k)$$

such that  $X_{k+1} \rightarrow A^{-1}$ .

$$X_{k+1} = \arg \min_{X \in \mathbb{R}^{n \times n}} \|X - X_k\|_{\text{Frobenius}(B)}^2 \quad \text{s.t.} \quad S^T A X = X,$$

The solution

$$\begin{aligned} X_{k+1} &= X_k + \text{proj}_{B^{-1}\text{Range}(A^T S)}(A^{-1} - X_k) \\ &= X_k + B^{-1} A^T S (S^T A B^{-1} A^T S)^{-1} S^T A (A^{-1} - X_k) \\ &= X_k + B^{-1} A^T S \underbrace{(S^T A B^{-1} A^T S)^{-1}}_{\text{Invert small matrix}} S^T (I - A X_k). \end{aligned}$$

What about the symmetric case  $A^T = A$ ?

# Framework

- ▶ Assume we observe  $S^T A$  where  $S$  is random.
- ▶ Given  $X_k \approx A \in \mathbb{R}^{n \times n}$ , we want to iteratively calculate

$$X_{k+1} = \text{update\_formula}(S^T A, X_k)$$

such that  $X_{k+1} \rightarrow A^{-1}$ .

$$X_{k+1} = \arg \min_{X \in \mathbb{R}^{n \times n}} \|X - X_k\|_{\text{Frobenius}(B)}^2 \quad \text{s.t.} \quad S^T A X = X,$$

The solution

$$\begin{aligned} X_{k+1} &= X_k + \text{proj}_{B^{-1} \text{Range}(A^T S)}(A^{-1} - X_k) \\ &= X_k + B^{-1} A^T S (S^T A B^{-1} A^T S)^{-1} S^T A (A^{-1} - X_k) \\ &= X_k + B^{-1} A^T S \underbrace{(S^T A B^{-1} A^T S)^{-1}}_{\text{Invert small matrix}} S^T (I - A X_k). \end{aligned}$$

What about the symmetric case  $A^T = A$ ?




# Symmetric matrices

$$X_{k+1} = \arg \min_{X \in \mathbb{R}^{n \times n}} \|X - X_k\|_{\text{Frobenius}(B)}^2$$

s.t.  $S^T A X = X$

$X = X^T$




# Symmetric matrices

$$X_{k+1} = \arg \min_{X \in \mathbb{R}^{n \times n}} \|X - X_k\|_{\text{Frobenius}(B)}^2$$

$$\text{s.t. } S^T A X = X$$

$$X = X^T$$



Solution:<sup>4</sup>

$$X_{k+1} = X_k + \text{proj}_{B^{-1}\text{Range}(AS)}(X_k - A^{-1})\text{proj}_{B^{-1}\text{Range}(AS)} \\ - (X_k - A^{-1})\text{proj}_{B^{-1}\text{Range}(AS)} - \text{proj}_{B^{-1}\text{Range}(AS)}(X_k - A^{-1})$$


<sup>4</sup>Gower and Gondzio 2014

# Symmetric matrices

$$X_{k+1} = \arg \min_{X \in \mathbb{R}^{n \times n}} \|X - X_k\|_{\text{Frobenius}(B)}^2$$

$$\text{s.t. } S^T A X = X$$

$$X = X^T$$



**Solution:**<sup>4</sup>

$$X_{k+1} = X_k + \text{proj}_{B^{-1}\text{Range}(AS)}(X_k - A^{-1})\text{proj}_{B^{-1}\text{Range}(AS)}$$

$$- (X_k - A^{-1})\text{proj}_{B^{-1}\text{Range}(AS)} - \text{proj}_{B^{-1}\text{Range}(AS)}(X_k - A^{-1})$$

<sup>4</sup>Gower and Gondzio 2014

## Theorem (Convergence)

Let  $S$  be equal to a column of a full rank matrix  $\mathbf{S} := [s_1, \dots, s_n]$  with probability  $\|B^{-1/2}As_i\|^2 / \|B^{-1/2}A\mathbf{S}\|_F^2$ . Then from a given  $X_0 \in \mathbb{R}^{n \times n}$ , the iteration

$$X_{k+1} = X_k + \text{proj}_{B^{-1}\text{Range}(AS)}(X_k - A^{-1})\text{proj}_{B^{-1}\text{Range}(AS)} \\ - (X_k - A^{-1})\text{proj}_{B^{-1}\text{Range}(AS)} - \text{proj}_{B^{-1}\text{Range}(AS)}(X_k - A^{-1})$$

converges with

$$\mathbf{E} \left[ \|X_k - A^{-1}\|_{\text{Frob}(B)}^2 \right] = \left( I - \frac{1}{\kappa_F^2(B^{-1/2}A\mathbf{S})} \right)^k \|X_0 - A^{-1}\|_{\text{Frob}(B)}^2,$$

where  $\kappa_F(B^{-1/2}A\mathbf{S}) = \|B^{-1/2}A\mathbf{S}\|_F \|\mathbf{S}^{-1}A^{-1}B^{1/2}\|_F$ .

**Self-preconditioning Method:** This suggests that  $\mathbf{S} \approx A^{-1}$ .  
But  $X_k \approx A^{-1}$  so try  $S = \text{sample columns of } X_k$ .

## Theorem (Convergence)

Let  $S$  be equal to a column of a full rank matrix  $\mathbf{S} := [s_1, \dots, s_n]$  with probability  $\|B^{-1/2}As_i\|^2 / \|B^{-1/2}A\mathbf{S}\|_F^2$ . Then from a given  $X_0 \in \mathbb{R}^{n \times n}$ , the iteration

$$X_{k+1} = X_k + \text{proj}_{B^{-1}\text{Range}(AS)}(X_k - A^{-1})\text{proj}_{B^{-1}\text{Range}(AS)} \\ - (X_k - A^{-1})\text{proj}_{B^{-1}\text{Range}(AS)} - \text{proj}_{B^{-1}\text{Range}(AS)}(X_k - A^{-1})$$

converges with

$$\mathbf{E} \left[ \|X_k - A^{-1}\|_{\text{Frob}(B)}^2 \right] = \left( I - \frac{1}{\kappa_F^2(B^{-1/2}A\mathbf{S})} \right)^k \|X_0 - A^{-1}\|_{\text{Frob}(B)}^2,$$

where  $\kappa_F(B^{-1/2}A\mathbf{S}) = \|B^{-1/2}A\mathbf{S}\|_F \|\mathbf{S}^{-1}A^{-1}B^{1/2}\|_F$ .

**Self-preconditioning Method:** This suggests that  $\mathbf{S} \approx A^{-1}$ . But  $X_k \approx A^{-1}$  so try  $S = \text{sample columns of } X_k$ .

## Initial experiments $A$ positive definite

**Newton-Schulz:**  $X_0 = A^T / (0.99 \|A^T A\|_2)$ ,  $X_{k+1} = 2X_k - X_k A X_k$ .

**Self-preconditioning Method:**  $B = A$ ,  $X_0 = I$ ,  $X_{k+1} = \text{proj}_S + (I - \text{proj}_S A) X_k (I - A \text{proj}_S)$ ,  
where  $S$  = sample columns of  $X_k$ .




Figure:  $n = 10'000$ , with  $nnz = 10^8$ ,  $A = \text{randn}(n, n)$ ,  $A = (A') * A$ ;

## Towards Randomized Preconditioning

**Initialize**  $X_0 \in \mathbb{R}^{n \times n}$  and  $x_0 \in \mathbb{R}^n$ .

**While** (stopping\_criteria)

$S_k = \text{sample\_function}(A, X_k)$

$x_{k+1} = \arg \min_{x \in \mathbb{R}^n} \|x - x_k\|_B^2 \text{ s.t. } S_k^T A x = S_k^T b$

$X_{k+1} = \arg \min_{X \in \mathbb{R}^{n \times n}} \|X - X_k\|_{\text{Frobenius}(B)} \text{ s.t. } S_k^T A X = S_k^T, X = X^T.$

$k = k + 1$

**end**

What if  $(A_k)_k$  and  $(b_k)$  are slowly changing (like a Hessian matrix)?

# Conclusion

- ▶ A natural framework for designing and analysing randomized iterative methods
- ▶ Analyse previous methods through one Theorem.
- ▶ New Gaussian methods, with potential on sparse problems
- ▶ New randomized matrix inversion methods.
- ▶ Paving a path towards randomized preconditioning.

Thank you for your attention!







# Conclusion

- ▶ A natural framework for designing and analysing randomized iterative methods
- ▶ Analyse previous methods through one Theorem.
- ▶ New Gaussian methods, with potential on sparse problems
- ▶ New randomized matrix inversion methods.
- ▶ Paving a path towards randomized preconditioning.

Thank you for your attention!

- └ Iteratively Inverting matrices
- └ Future work: Randomized Preconditioning?

# References

-  Gower, Robert M., Richtárik, Peter, April 2015.  
Randomized Iterative Methods for Linear Systems (in progress)
-  Gower, Robert M., Richtárik, Peter, April 2015.  
Randomized Iterative Methods for Inverting Matrices (in progress)
-  Qu, Z., Richtárik, P., Takáč, M., & Fercoq, O. (2015).  
SDNA: Stochastic Dual Newton Ascent for Empirical Risk Minimization.
-  Gower, Robert M., Gondzio, Jacek (2014).  
Action constrained quasi-Newton methods (in progress)