

Optimization and Numerical Analysis: Nonlinear programming without constraints

Robert Gower



October 9, 2018

Table of Contents

Unconstrained Nonlinear Programming

Multivariate Calculus

Local Descent Lemma and Local Optimality

Sufficient and Necessary conditions for local optimality

Convex functions

The Problem: Nonlinear programming

Minimize a nonlinear differentiable function $f : x \in \mathbb{R}^n \mapsto f(x) \in \mathbb{R}$

$$x^* = \arg \min_{x \in \mathbb{R}^n} f(x). \quad (1)$$

Warning: This problem is often impossible. First check there **exists** a minimum. Even linear programming does not always have a maximum! Develop iterative methods x^1, \dots, x^k, \dots , such that

$$\lim_{k \rightarrow \infty} x^k = x^*.$$

Template method

$$x^{k+1} = x^k + s_k d^k,$$

where $s_k > 0$ is a *step size* and $d^k \in \mathbb{R}^n$ is *search direction*. Satisfy the *descent condition*

$$f(x^{k+1}) < f(x^k).$$

Local and Global Minima

Definition of Local Minima

The point $x^* \in \mathbb{R}^n$ is a *local minima* of $f(x)$ if there exists $r > 0$ such that

$$f(x^*) \leq f(x), \quad \forall \|x - x^*\|_2 < r. \quad (2)$$

Definition of Global Minima

The point $x^* \in \mathbb{R}^n$ is a *global minima* of $f(x)$ if

$$f(x^*) \leq f(x), \quad \forall x. \quad (3)$$

In general finding global minima impossible. Finding local is good enough.

Multivariate Calculus

For a differentiable function $f : x \in \mathbb{R}^n \mapsto f(x) \in \mathbb{R}$, we refer to $\nabla f(x)$ as the gradient evaluated at x defined by

$$\nabla f(x) = \left[\frac{\partial f(x)}{\partial x_1}, \dots, \frac{\partial f(x)}{\partial x_n} \right]^\top.$$

Note that $\nabla f(x)$ is a column-vector. For any $F : x \in \mathbb{R}^n \rightarrow F(x) = [f_1(x), \dots, f_n(x)]^\top \in \mathbb{R}^n$ define the *Jacobian matrix* by

$$\begin{aligned} \nabla F(x) &\stackrel{\text{def}}{=} \begin{bmatrix} \frac{\partial f_1(z)}{\partial x_1} & \frac{\partial f_2(z)}{\partial x_1} & \frac{\partial f_3(z)}{\partial x_1} & \cdots & \frac{\partial f_n(z)}{\partial x_1} \\ \frac{\partial f_1(z)}{\partial x_2} & \frac{\partial f_2(z)}{\partial x_2} & \frac{\partial f_3(z)}{\partial x_2} & \cdots & \frac{\partial f_n(z)}{\partial x_2} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \frac{\partial f_1(z)}{\partial x_n} & \frac{\partial f_2(z)}{\partial x_n} & \frac{\partial f_3(z)}{\partial x_n} & \cdots & \frac{\partial f_n(z)}{\partial x_n} \end{bmatrix} \\ &= [\nabla f_1(x), \nabla f_2(x), \nabla f_3(x), \dots, \nabla f_n(x)] \end{aligned}$$

Multivariate Calculus

The gradient is useful because of 1st order Taylor expansion

$$f(x^0 + d) = f(x^0) + \nabla f(x^0)^\top d + \epsilon(d)\|d\|_2, \quad (4)$$

where $\epsilon(d)$ is a real valued such that

$$\lim_{d \rightarrow 0} \epsilon(d) = 0. \quad (5)$$

Example (The $\epsilon(d)$ function)

If $f(x) = \|x\|_2^2$ or $f(x) = x^\top Ax$ what is $\epsilon(d)$? Name three functions that satisfy (5).

The Hessian Matrix

If $f \in C^2$, we refer to $\nabla^2 f(x)$ as the Hessian matrix:

$$\nabla^2 f(x) \stackrel{\text{def}}{=} \begin{bmatrix} \frac{\partial^2 f(z)}{\partial x_1 \partial x_1} & \frac{\partial^2 f(z)}{\partial x_1 \partial x_2} & \frac{\partial^2 f(z)}{\partial x_1 \partial x_3} & \cdots & \frac{\partial^2 f(z)}{\partial x_1 \partial x_n} \\ \frac{\partial^2 f(z)}{\partial x_1 \partial x_2} & \frac{\partial^2 f(z)}{\partial x_2 \partial x_2} & \frac{\partial^2 f(z)}{\partial x_2 \partial x_3} & \cdots & \frac{\partial^2 f(z)}{\partial x_2 \partial x_n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \frac{\partial^2 f(z)}{\partial x_n \partial x_1} & \frac{\partial^2 f(z)}{\partial x_n \partial x_2} & \frac{\partial^2 f(z)}{\partial x_n \partial x_3} & \cdots & \frac{\partial^2 f(z)}{\partial x_n \partial x_n} \end{bmatrix}$$

If $f \in C^2$ then

$$\frac{\partial^2 f(z)}{\partial x_i \partial x_j} = \frac{\partial^2 f(z)}{\partial x_j \partial x_i}, \quad \forall i, j \in \{1, \dots, n\}, \quad \Leftrightarrow \quad \nabla^2 f(x) = \nabla^2 f(x)^\top.$$

Hessian matrix useful for 2nd order Taylor expansion.

$$f(x^0 + d) = f(x^0) + \nabla f(x^0)^\top d + \frac{1}{2} d^\top \nabla^2 f(x^0) d + \epsilon(d) \|d\|_2^2. \quad (6)$$

Example

If $f(x) = x^3$ or $f(x) = x^\top A x$ what is $\epsilon(d)$?

The Product-rule

The vector valued version of the product rule

- ▶ For any function $F(x) : \mathbb{R}^n \rightarrow \mathbb{R}^n$ and matrix $A \in \mathbb{R}^{n \times n}$ we have

$$\nabla(F(x)^\top A) = \nabla F(x)^\top A. \quad (7)$$

- ▶ For any two vector valued functions F_1 and F_2 we have that

$$\nabla(F_1(x)^\top F_2(x)) = \nabla F_1(x) F_2(x) + \nabla F_2(x) F_1(x). \quad (8)$$

Example

Let $f(x) = \frac{1}{2}x^\top Ax$, where $A \in \mathbb{R}^{n \times n}$ is a symmetric positive definite matrix. Calculate the gradient and the Hessian of $f(x)$.

The Product-rule

The vector valued version of the product rule

- ▶ For any function $F(x) : \mathbb{R}^n \rightarrow \mathbb{R}^n$ and matrix $A \in \mathbb{R}^{n \times n}$ we have

$$\nabla(F(x)^\top A) = \nabla F(x)^\top A. \quad (7)$$

- ▶ For any two vector valued functions F_1 and F_2 we have that

$$\nabla(F_1(x)^\top F_2(x)) = \nabla F_1(x) F_2(x) + \nabla F_2(x) F_1(x). \quad (8)$$

Example

Let $f(x) = \frac{1}{2}x^\top Ax$, where $A \in \mathbb{R}^{n \times n}$ is a symmetric positive definite matrix. Calculate the gradient and the Hessian of $f(x)$.

Let $F_1(x) = A^\top x$ and $F_2(x) = x$ then

$\nabla f(x) = \frac{1}{2}\nabla(A^\top x)x + \frac{1}{2}\nabla(x)A^\top x = \frac{1}{2}(A + A^\top)x$ since

$\nabla(A^\top x) = A\nabla(x) = A$. Differentiating again

$\nabla(\nabla f(x)) = \nabla(Ax) = \nabla(A)x + \nabla(x)A = A$.