Satellite imagery for rural infrastructure

"Satellite imagery for rural infrastructure mapping" *Generated by LeapSpace*

Satellite Imagery for Rural Infrastructure Mapping at a Glance

Key Findings Table

Key finding	Infrastructure / setting	Practical implication	Citations
High-resolution optical imagery is the strongest-supported data source for mapping rural roads, buildings, and settlements, especially when paired with deep learning.	Roads, rooftops, dispersed settlements	Best suited for fine delineation where sub-field features matter.	[1] [2] [3]
Data fusion consistently improves separability of rural built features from cropland, bare soil, and other confusing backgrounds.	Built-up areas, roads, mixed rural landscapes	Combine optical, SAR, texture, indices, and/or multi-season inputs where possible.	[4] [5] [6] [7] [8] [9]
Deep learning generally outperforms classical segmentation and OBIA, but gains depend on context and are weaker for hard rural targets such as unpaved tracks.	Road and building extraction	Modern segmentation/detection is preferable, but rural-specific tuning remains necessary.	[1] [10] [11] [12] [13]
SAR is essential in cloudy and inaccessible rural regions, but usually works best as a complement to optical imagery rather than a full replacement for detailed infrastructure delineation.	Cloud-prone tropics, flood-prone corridors, inaccessible regions	Use SAR for continuity, flooding, and temporal coverage; fuse with optical for finer feature mapping.	[14] [15] [16] [17] [18] [19] [20]
Validation is a major bottleneck because many studies rely on manual labels, benchmark datasets, map products, or sampled ROIs rather than field GPS.	Remote rural settings	Reported accuracies should be interpreted with caution for operational deployment.	[15] [16] [21] [22] [23] [24] [25] [26] [27]

Key finding	Infrastructure / setting	Practical implication	Citations
The literature is shifting from feature extraction to decision support, especially electrification, accessibility, cadastral assessment, and rural development monitoring.	Planning and policy use cases	Highest-value outputs may be accessibility gaps, service coverage, or electrification indicators rather than feature masks alone.	[28] [29] [30] [31] [32] [33]

Direct Answer

Satellite imagery is already a practical tool for rural infrastructure mapping, with the strongest evidence favoring **high-resolution optical imagery plus deep-learning segmentation/detection** for roads, buildings, and settlements, and **SAR** as a critical all-weather complement in cloud-prone or hard-to-access regions [1] [2] [3] [14] [15] [16]. Across studies, performance improves when workflows use **fusion**—for example optical plus SAR, multispectral plus texture, or multi-season stacks—because rural built features are often visually similar to bare soil, cropland, or flood-affected surfaces [4] [5] [6] [7] [8].

For practice, the evidence supports a **tiered workflow**: use broad-coverage Sentinel-1/2-style time series for screening and persistence under clouds, then deploy high-resolution optical deep-learning models over candidate settlements, roads, or service corridors [1] [14] [16] [20]. The main constraint is not only model accuracy but **validation quality**: many reported results depend on manual annotations or secondary reference products rather than field GPS, which limits certainty about transferability across regions and seasons [21] [22] [25] [27].

Confidence level: High - consistent cross-source support exists for optical dominance in fine extraction, the value of fusion, the importance of SAR under clouds, and validation as a bottleneck [1] [2] [4] [14] [15] [26].

Study Scope

This report covers evidence summarized from studies spanning roughly **2015-2025**, with emphasis on **2018-2025** applications and benchmarks relevant to rural and remote settings [2] [33] [34]. It includes work from **remote sensing, computer vision, land-cover mapping, flood/wetland monitoring, rural development, electrification planning, and geospatial accessibility analysis** [31] [35] [36]. Methods represented include **classical classification, object-based image analysis, image segmentation, deep-learning detection/segmentation, SAR time-series analysis, and planning-oriented geospatial integration** [8] [31] [37] [38] [39].

Assumptions & Limitations

- The evidence base is much stronger for **optical/deep-learning workflows** than for **hyperspectral** rural infrastructure mapping or exact rural-specific spatial-resolution thresholds [34] [40] [41].

- Reported accuracies are often derived from **manual annotations, benchmark labels, external map products, or sampled validation points**, not systematic field GPS for roads and buildings in remote rural areas [21] [25] [26].
- Benchmark success does not automatically guarantee **cross-region or cross-season transferability** in low-density rural landscapes [27] [42] [43] [44].

Suggested Further Research

- Build **rural-specific benchmark datasets** for unpaved roads, dispersed housing, utilities, and small service facilities [1] [36].
- Test **minimum useful spatial resolution** for each infrastructure class using consistent rural validation protocols [2] [34].
- Expand studies using **field GPS or cadastral-quality rural reference data** at operational scale [26] [32].
- Evaluate **hyperspectral and spectral-unmixing** methods more directly for rural built-feature mapping [40] [41].
- Measure how extraction errors propagate into **accessibility, electrification, and service-planning decisions** [29] [31] [33].

Full Report

1. Introduction and Scope

Rural infrastructure mapping from satellite imagery focuses on features that are spatially sparse, heterogeneous, and often difficult to validate: minor roads and tracks, small buildings, dispersed settlements, flood-prone access corridors, and service-related built features [1] [31]. The practical motivation is strong. Satellite-derived infrastructure products are increasingly used for **electrification planning**, **monitoring settlement change**, **accessibility analysis**, and **land-governance support**, particularly where conventional mapping is expensive or incomplete [28] [29] [30] [32] [33].

Taken together, these findings suggest that rural infrastructure mapping is no longer just a technical image-analysis problem; it is becoming a decision-support pipeline in which extraction quality matters because it feeds planning and policy outputs [31] [33] [36].

Confidence level: High - the shift toward planning applications appears across multiple independent studies and the meta-synthesis [28] [29] [31].

2. Foundational Context

2.1 What makes rural infrastructure hard to map?

Rural targets are often **small, elongated, low-contrast, and spectrally confused** with natural or agricultural surfaces [1] [4]. Unpaved roads can resemble bare soil or harvested fields, while scattered rooftops may be difficult to distinguish from bright ground materials without sufficient spatial detail or contextual features [1] [34]. This is why simple single-date classification often struggles in rural environments [17] [37].

2.2 Why sensor choice matters

Three broad data families dominate the evidence:

- **High-resolution optical imagery**: best for fine boundary delineation of roads, buildings, and settlements because it preserves visible shape and edge information [1] [2] [3].
- **Multispectral/hyperspectral imagery**: useful when spectral richness helps separate built-up areas from cropland, bare soil, or other land covers, especially when enhanced by fusion or pan-sharpening [4] [5] [7] [45] [46].
- **SAR/radar imagery**: valuable in cloudy regions because it provides all-weather imaging and multi-temporal monitoring, though not always the best standalone source for fine infrastructure edges [14] [15] [16] [17] [18].

2.3 Why workflows increasingly use fusion

Fusion is repeatedly reported as useful because rural infrastructure classes often require **both structural and contextual cues**: geometry from high-resolution imagery, spectral separation from multispectral bands, and weather-robust temporal consistency from SAR [4] [5] [6] [8]. This is one of the clearest patterns in the evidence base [7] [9].

Confidence level: High - multiple sources converge on the same mechanism: rural separability improves when complementary signals are combined [4] [5] [17] [18].

3. Satellite Data Types and Spatial Requirements

3.1 High-resolution optical imagery for roads, buildings, and settlements

The strongest direct evidence for rural infrastructure extraction comes from **high-resolution optical imagery** [1] [2] [3]. In building extraction, one benchmark reported **DeepLabV3+** with Accuracy **96.77%**, IoU **89.85%**, and F1 **94.53%**, outperforming several related models on high-resolution imagery [2]. A separate building study using **Mask R-CNN** on Pleiades imagery reported **F1 = 0.85** [3].

For rural roads, optical orthoimagery remains difficult because unpaved tracks blend into agricultural background, but recent tiled-inference pipelines using **YOLOv11 plus SAHI** were reported to detect rural roads and related infrastructure with high mean average precision [1]. Even so, the evidence also indicates that “high resolution” alone does not remove rural ambiguity; model design and context remain important [1] [22].

A notable gap is that the provided studies do **not** establish exact rural class-specific resolution thresholds for roads versus rooftops versus hamlets [34] [40]. What the evidence does support is the general proposition that finer optical detail improves delineation of continuous roads and building boundaries compared with coarser imagery [34].

3.2 Multispectral and hyperspectral imagery

Multispectral imagery helps identify built-up features when combined with **spectral indices, texture, and seasonal context** [4]. Sentinel-2A impervious-surface mapping at **10 m** reportedly showed clearer continuous roads and building boundaries than a **30 m** Landsat 8 product, although the excerpt did not provide a separability metric for those classes [34].

Fusion methods recur throughout this literature. **Pan-sharpening** and **hyperspectral-multispectral fusion** improve spatial and spectral detail and have been reported to boost road extraction, land-use segmentation, and urban material mapping [5] [6] [7] [45] [46]. For hyperspectral work, the evidence is thinner and more urban-material-oriented than rural-infrastructure-specific; for example, one AVIRIS-NG study argued that **MESMA** was better suited than linear unmixing for heterogeneous built-up scenes, but did not provide rural infrastructure separability metrics [41].

Taken together, these findings suggest multispectral methods are already practical for rural screening and broad built-up mapping, while hyperspectral methods remain under-evaluated for operational rural infrastructure tasks [4] [41].

3.3 SAR and radar for cloud-prone rural regions

SAR is consistently valuable in **cloud-prone tropical and remote settings**, supporting land-cover and monitoring tasks where optical imagery is interrupted by weather [14] [15] [16]. A dry/wet-season Sentinel-1 classification reported **94.81% overall accuracy** and **Kappa 0.92** using multi-temporal SAR and texture features, illustrating the benefit of temporal context [14]. Other studies show that full-year or multi-scene SAR stacks outperform single-date imagery, such as forest/non-forest accuracy improving from **77% to 85%**, with annual mean and standard deviation features raising mean accuracy to **87%** [8].

For flood-affected access corridors and built-up flooding, SAR can detect open water and inundated built areas, but performance may be modest: PALSAR-2 amplitude plus coherence achieved **Kappa 0.583** for open water and **0.429** for built-up flooding [19]. This supports a recurring conclusion: SAR is indispensable for robustness and repeat coverage, but SAR-only delineation is often weaker than fused optical+SAR approaches for fine infrastructure products [17] [18] [47].

Confidence level: High - the all-weather role of SAR and the benefit of temporal stacking are strongly supported, though fine-feature infrastructure extraction is less consistently strong with SAR alone [8] [14] [15].

4. Mapping Methods and Extraction Workflows

4.1 Preprocessing and feature engineering

Common preprocessing steps include **pan-sharpening**, **multisensor fusion**, **texture extraction**, **edge-based features**, and **multi-season compositing** [5] [6] [14] [45]. These steps are especially important in rural landscapes because pixel-level spectral signals alone may not distinguish roads from adjacent soil or built-up patches from dry fields [4] [6]. Multi-temporal or multi-season inputs repeatedly improve class separation in both optical and SAR workflows [4] [8] [9].

4.2 Classical classification and object-based image analysis

Earlier and still-relevant workflows include **maximum-likelihood classification**, **pixel-based methods**, **object-based image analysis (OBIA)**, and classical segmentation [6] [37] [38]. In one road-network study, **spectral difference segmentation** outperformed **multi-resolution segmentation** and classical maximum-likelihood classification for road detection [6] [37].

OBIA and segmentation-plus-classical-ML remain useful where object shape and contextual grouping matter, but comparative evidence generally places them below the best deep segmenters for current infrastructure tasks [38]. In one comparative framework, U-Net achieved the best IoU (**0.763**), ahead of DeepLabV3 and Mask R-CNN in that experiment [38].

4.3 Deep learning for infrastructure delineation

Deep learning now dominates road and building extraction workflows [10] [11] [12] [13]. Foundational architectures such as **U-Net**, **DeepLab**, **SegNet**, and **LinkNet** established the encoder-decoder and multi-scale segmentation patterns widely reused for satellite imagery [42] [43] [48] [49]. Residual learning also influenced later road-extraction models, including **ResU-Net** variants [50] [51].

Comparative findings show meaningful architecture effects. **ResU-Net** outperformed Reconstruction Bias U-Net, and **U-Net++** was reported as the strongest road extractor in one comparative study of U-Net variants [39]. In building extraction, **DeepLabV3+** led one benchmark, ahead of Attention U-Net, U-Net, UNet++, and U2Net [2]. For large-area road extraction, one **UU-Net** study outperformed single-source optical/SAR baselines whose average F1 and IoU were **0.5502** and **0.4021** respectively [22].

Benchmark normalization through datasets and platforms such as **DeepGlobe** and **Google Earth Engine** has helped standardize development and scaling of these methods [52] [53]. However, benchmark performance does not guarantee success in sparse rural environments [36] [44].

Confidence level: High - deep learning's dominance is well supported, but rural gains are explicitly context-dependent [1] [10] [44].

5. Accuracy, Validation, and Transferability

5.1 Reported accuracy patterns across methods

The broad pattern is clear: **deep-learning methods usually outperform classical methods**, but accuracy varies widely with target type, sensor, and validation design [10] [11] [38]. Strong building-extraction results were reported for DeepLabV3+ and Mask R-CNN [2] [3]. For SAR-based thematic mapping, multi-temporal workflows often improve OA and Kappa relative to single-date inputs [8] [14]. In crop-related SAR comparisons, TerraSAR-X outperformed Sentinel-1 with **86% OA, Kappa 0.71** versus **80-82% OA, Kappa 0.60** for Sentinel-1 VV and lower for VH [54].

This pattern implies that “best method” depends on whether the goal is **fine feature delineation**, **weather-robust monitoring**, or **broad thematic screening** [2] [14] [20].

5.2 Reference data quality and validation design

Validation quality is one of the most important limitations in the evidence base [21] [25]. Many optical extraction studies relied on **manual annotations, benchmark labels, or map products such as OSM**, rather than field GPS [21] [22]. In rural built-up mapping, one study used labor-intensive manual reference grids plus crowdsourcing and visual interpretation, while still depending on WSF-2015 as a key reference product [25].

More rigorous designs do appear in some studies. A Mekong Delta land-cover study used **270 independent, spatially distributed validation points**, stratified by land cover and supported by official maps and high-resolution imagery, with separate training and validation data and explicit error matrices [26]. Another framework without direct field surveys combined internal cross-validation, external product comparison, statistical checks, and regional generalization tests before transfer to Romania and the USA [27].

The practical consequence is that reported performance often measures agreement with a **reference representation**, not necessarily with on-the-ground infrastructure reality [21] [27].

5.3 Transferability across regions and seasons

Transferability is a frontier issue. Some newer models report stronger cross-dataset generalization, including a **SAM2-based road model** performing better than baselines across DeepGlobe, SpaceNet, and Massachusetts [44]. Similarly, a non-infrastructure agricultural framework explicitly tested regional transfer and maintained high overall accuracy after transfer [27].

But the evidence also warns against overgeneralizing benchmark success to rural settings with different road materials, settlement morphologies, and annotation scarcity [36] [42] [43]. This tension between benchmark performance and real-world rural deployment is one of the clearest cross-source themes [27] [44].

Confidence level: Medium-High - the validation bottleneck is strongly supported, while transferability evidence is growing but still limited [25] [27] [44].

6. Applications for Planning and Development Monitoring

6.1 Energy access and electrification planning

Planning applications are among the strongest rural use cases. A satellite-imagery dataset for Uganda, Kenya, and Sierra Leone mapped **578,010 geolocated homes** for electrical infrastructure planning, showing home-level extraction at policy-relevant scale [28]. In rural Uganda, satellite imagery plus machine learning was used to estimate local livelihood impacts of **grid expansion**, linking electrification to village-level asset wealth for low-cost policy evaluation [29]. In Kenya, multi-temporal high-resolution imagery across **1,000 communities** was used to detect settlement growth around electrification, finding that communities receiving electricity access grew **15% more slowly** than those without access [30].

6.2 Connectivity, accessibility, and service coverage

Mapped infrastructure is increasingly combined with other geospatial layers to create **accessibility descriptors**, **service coverage maps**, and **connectivity graphs** [31]. In rural Ukraine, a geospatial framework combined OSM and humanitarian datasets to map access to healthcare, education, transit, and other services, identifying underserved areas for targeted revitalization [31]. This is an important evolution from feature extraction toward infrastructure analytics [31] [33].

6.3 Cadastral and development monitoring uses

Satellite imagery has also supported **cadastral fitness assessment** in Nigerian rural, peri-urban, and urban settlements, with evidence that imagery resolution matters more in informal rural settlements for land-governance planning [32]. More broadly, imagery-based monitoring of land-use change from **2000-2020** has been combined with population data to identify socioeconomic change and support local planning and sustainable development [33].

A caution remains: planning-oriented settlement products can overlook vulnerable populations if ground truth is weak or biased [55].

Confidence level: High - planning and monitoring applications are directly evidenced across multiple countries and use cases [28] [29] [31] [33].

7. Practical Synthesis: What Works Best in Rural Mapping?

Based on the evidence, the most defensible operational strategy is:

1. **Use SAR or multispectral time series for broad-area screening and persistence under cloud cover** [14] [20].

2. **Use high-resolution optical imagery for detailed delineation of roads, buildings, and settlement edges** [1] [2].
3. **Prefer deep-learning segmentation/detection over classical pipelines**, while tuning specifically for sparse, low-contrast rural features [1] [10].
4. **Fuse modalities and temporal views whenever feasible** [4] [5] [8].
5. **Invest in validation design**, especially targeted field checks or high-quality reference construction in uncertain zones [25] [26].

Inferred: a cost-effective rural workflow may be a **tiered system** in which Sentinel-1/2-like data identify candidate settlements and corridors, followed by targeted high-resolution optical inference only where finer mapping is needed. This inference is consistent with the documented strengths of broad-coverage all-weather screening plus high-resolution deep learning, but it is not directly evaluated as a unified pipeline in the supplied evidence [1] [14] [20].

Confidence level: Medium - each component is directly supported, but the integrated tiered workflow is a synthesis across studies rather than a single validated design [1] [14].

8. Conclusions

Satellite imagery is a mature and increasingly policy-relevant tool for rural infrastructure mapping [28] [31]. The clearest evidence favors **high-resolution optical imagery plus deep learning** for detailed extraction, **SAR** for all-weather continuity, and **data fusion** for difficult rural separability problems [1] [2] [4] [14]. The major unresolved issue is not whether rural infrastructure can be mapped, but how reliably results transfer across regions and whether reference data are strong enough for operational decisions [25] [27] [44].

The field is also changing in purpose: instead of stopping at roads or buildings as map layers, recent studies increasingly convert mapped infrastructure into **electrification, accessibility, cadastral, and development-monitoring products** [29] [32] [33]. That shift is likely to define the next phase of rural satellite infrastructure mapping, provided validation practices improve [26] [36].

Overall confidence level: High - core conclusions are supported by repeated patterns across the provided evidence, though exact rural spatial-resolution thresholds and field-validated transferability remain insufficiently resolved [1] [27] [34].

References

1. Sanita, M., Nepi, L., Malinverni, E.S., (...), Balawejder, M. (2026) Automatic Extraction and Multi-Class Instance Segmentation of Rural Road Networks from Orthoimagery using YOLOv11 and SAHI Sliced Inference for Cadastral Update. *International Archives of the Photogrammetry, Remote Sensing and Spatial Information Sciences - ISPRS Archives*. https://doi.org/10.5194/isprs-archives-XLIX-B2-2026-1397-2026

2. Ekiz, S., Acar, U. (2025) Improving building extraction from high-resolution aerial images: Error correction and performance enhancement using deep learning on the Inria dataset. *Science Progress*. https://doi.org/10.1177/00368504251318202
3. Lin, Y.H., Tsai, F. (2025) Precise Building Boundary Extraction Using Deep Learning Method. *46th Asian Conference on Remote Sensing, ACRS 2025 - Harnessing Remote Sensing for Global Sustainability and Innovation*. https://www.scopus.com/pages/publications/105031785767
4. Kamusoko, C. (2022) Improving Urban Land Cover Mapping. *Springer Geography*. https://doi.org/10.1007/978-981-16-5149-6_5
5. Irfan, A. (2024) Fusion of Hyperspectral and Multispectral Images for Land Use Segmentation. *Lecture Notes in Networks and Systems*. https://doi.org/10.1007/978-981-97-3556-3_13
6. Wang, R., Hu, Y., Zhang, X. (2005) Extraction of road networks using pan-sharpened multispectral and panchromatic quickbird images. *Geomatica*. https://www.scopus.com/pages/publications/28444460786
7. Salvini, R., Garzelli, A., Rindinella, A., (...), Candela, L. (2025) Joint processing of Sentinel-2 and PRISMA data for monitoring urban areas (TUS:CAN Project). *Proceedings of SPIE - The International Society for Optical Engineering*. https://doi.org/10.1117/12.3069875
8. Hansen, J.N., Mitchard, E.T.A., King, S. (2020) Assessing forest/non-forest separability using sentinel-1 C-band synthetic aperture radar. *Remote Sensing*. https://doi.org/10.3390/rs12111899
9. San Martín, L., Morandeira, N.S., Grimson, R., (...), Kandus, P. (2020) The contribution of ALOS/PALSAR-1 multi-temporal data to map permanently and temporarily flooded coastal wetlands. *International Journal of Remote Sensing*. https://doi.org/10.1080/01431161.2019.1673915
10. Na, H.S., Chen, C.-B. (2022) Deep Learning Methods for Road Extraction from Remote Sensing Imagery. *IISE Annual Conference and Expo 2022*. https://www.scopus.com/pages/publications/85137180674
11. Neupane, B., Aryal, J., Rajabifard, A. (2024) CNNs for remote extraction of urban features: A survey-driven benchmarking. *Expert Systems with Applications*. https://doi.org/10.1016/j.eswa.2024.124751
12. Soni, T.K., Pujari, P. (2025) Emerging deep learning approaches for urban satellite image analysis: a survey on classification, segmentation, and change detection. *Evolutionary Intelligence*. https://doi.org/10.1007/s12065-025-01090-2
13. Deressu, T.F., Kumsa, A.K., Debelee, T.G., (...), Gebissa, K.W. (2025) Enhancing land use and land cover classification with deep learning-based satellite imagery segmentation. *International Journal of Applied Earth Observation and Geoinformation*. https://doi.org/10.1016/j.jag.2025.104839
14. Ngo, K.D., Lechner, A.M., Vu, T.T. (2020) Land cover mapping of the Mekong Delta to support natural resource management with multi-temporal Sentinel-1A synthetic aperture radar imagery. *Remote Sensing Applications: Society and Environment*. https://doi.org/10.1016/j.rsase.2019.100272

15. Schultz, F.M., Lodwick, G.D. (1989) Information extraction from SAR imagery for tropical land cover evaluation. *Digest - International Geoscience and Remote Sensing Symposium (IGARSS)*. https://www.scopus.com/pages/publications/0024895519

16. Hütt, C., Waldhoff, G., Bareth, G. (2020) Fusion of sentinel-1 with official topographic and cadastral geodata for crop-type enriched LULC mapping using FOSS and open data. *ISPRS International Journal of Geo-Information*. https://doi.org/10.3390/ijgi9020120

17. Xu, F., Shi, Y., Yang, W., (...), Zhu, X.X. (2024) CloudSeg: A multi-modal learning framework for robust land cover mapping under cloudy conditions. *ISPRS Journal of Photogrammetry and Remote Sensing*. https://doi.org/10.1016/j.isprsjprs.2024.06.001

18. Ling, J., Zhang, H., Lin, Y. (2021) Improving urban land cover classification in cloud-prone areas with polarimetric sar images. *Remote Sensing*. https://doi.org/10.3390/rs13224708

19. Ohki, M., Tadono, T., Itoh, T., (...), Shimada, M. (2019) Flood area detection using PALSAR-2 amplitude and coherence data: The case of the 2015 heavy rainfall in Japan. *IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing*. https://doi.org/10.1109/JSTARS.2019.2911596

20. Caballero, G., Pezzola, A., Winschel, C., (...), Delegido, J. (2022) Quantifying Irrigated Winter Wheat LAI in Argentina Using Multiple Sentinel-1 Incidence Angles. *Remote Sensing*. https://doi.org/10.3390/rs14225867

21. Lee, H., Kim, K., Lee, K. (2024) Application of Segment Anything Model (SAM) for Optical-SAR Fusion Images to Surface Water Body Object Extraction: A Case Study using KOMPSAT-3/3A and KOMPSAT-5 Images. *Proceedings of SPIE - The International Society for Optical Engineering*. https://doi.org/10.1117/12.3045140

22. Lin, Y., Wan, L., Zhang, H., (...), Zhao, Z. (2021) Leveraging optical and SAR data with a UU-Net for large-scale road extraction. *International Journal of Applied Earth Observation and Geoinformation*. https://doi.org/10.1016/j.jag.2021.102498

23. Gstaiger, V., Huth, J., Gebhardt, S., (...), Kuenzer, C. (2012) Multi-sensoral and automated derivation of inundated areas using TerraSAR-X and ENVISAT ASAR data. *International Journal of Remote Sensing*. https://doi.org/10.1080/01431161.2012.700421

24. Iyyappan, I., Shanmugam, R.S., Srinivasa Raju, S.R. (2017) SAR polarimetric decomposition with ALOS PALSAR-1 for agricultural land and other land use/cover classification: case study in Rajasthan, India. *Environmental Earth Sciences*. https://doi.org/10.1007/s12665-017-6783-6

25. Li, M., Wang, Y., Rosier, J.F., (...), Van Vliet, J. (2022) Global maps of 3D built-up patterns for urban morphological analysis. *International Journal of Applied Earth Observation and Geoinformation*. https://doi.org/10.1016/j.jag.2022.103048

26. Ngo, K.D., Lechner, A.M., Vu, T.T. (2020) Land cover mapping of the Mekong Delta to support natural resource management with multi-temporal Sentinel-1A synthetic aperture radar imagery. *Remote Sensing Applications: Society and Environment*. https://doi.org/10.1016/j.rsase.2019.100272

27. Qiu, J., Hu, Y., Peng, D., (...), Xu, J. (2026) Analyzing the disruption of agricultural systems by conflict: A case study of sunflower production in eastern Ukraine. *Agricultural Systems*. https://doi.org/10.1016/j.agsy.2026.104636

28. Leonard, A., Wheeler, S., McCulloch, M. (2022) Rural Home Annotation Dataset Mapped by Citizen Scientists in Satellite Imagery. *Data in Brief*. https://doi.org/10.1016/j.dib.2022.108262

29. Ratledge, N., Cadamuro, G., de la Cuesta, B., (...), Burke, M. (2022) Using machine learning to assess the livelihood impact of electricity access. *Nature*. https://doi.org/10.1038/s41586-022-05322-8

30. Correa, S., Pérez, G., Jaramillo, P., Taneja, J. (2023) Taking the Long View: Enhancing Learning On Multi-Temporal, High-Resolution, and Disparate Remote Sensing Data. *BuildSys 2023 - Proceedings of the10th ACM International Conference on Systems for Energy-Efficient Buildings, Cities, and Transportation*. https://doi.org/10.1145/3600100.3623722

31. Kussul, N., Svirsh, V., Potuzhnyi, B. (2024) Integrated Geospatial Analysis for Rural Development Metrics. *CEUR Workshop Proceedings*. https://www.scopus.com/pages/publications/85191465860

32. Taiwo, I.O., Ibitoye, M.O., Oladejo, S.O., Koeva, M. (2024) Fitness of Multi-Resolution Remotely Sensed Data for Cadastral Mapping in Ekiti State, Nigeria. *Remote Sensing*. https://doi.org/10.3390/rs16193670

33. Grabska, E., Hashemvand Khiabani, P., Pesoa-Marcilla, M., (...), de Vries, W.T. (2024) Exploring land use dynamics in rural areas. An analysis of eight cases in the Global North. *Land Use Policy*. https://doi.org/10.1016/j.landusepol.2024.107246

34. Xu, Rudong, Liu, Jin, Xu, Jianhui (2018) Extraction of High-Precision Urban Impervious Surfaces from Sentinel-2 Multispectral Imagery via Modified Linear Spectral Mixture Analysis. *Sensors (Basel, Switzerland)*. https://doi.org/10.3390/s18092873

35. Wang, W., Yang, N., Zhang, Y., (...), Eklund, P. (2016) A review of road extraction from remote sensing images. *Journal of Traffic and Transportation Engineering (English Edition)*. https://doi.org/10.1016/j.jtte.2016.05.005

36. Hargreaves, P.K., Watmough, G.R. (2021) Satellite Earth observation to support sustainable rural development. *International Journal of Applied Earth Observation and Geoinformation*. https://doi.org/10.1016/j.jag.2021.102466

37. Leena, C., Saxena, M.R., Dwivedi, R.S. (2021) Road extraction from cartosat-2F multispectral data with object-oriented analysis. *Photogrammetric Engineering and Remote Sensing*. https://doi.org/10.14358/PERS.87.3.171

38. Toma, A., Şandric, I., Mihai, B.-A. (2024) Flooded area detection and mapping from Sentinel-1 imagery. Complementary approaches and comparative performance evaluation. *European Journal of Remote Sensing*. https://doi.org/10.1080/22797254.2024.2414004

39. Nebili, B., Louali, R., Boukerche, H., (...), Boulahmar, M. (2025) AI-Driven Road Extraction from Satellite Imagery: Evaluating the Impact of U-Net Symmetry on Segmentation Accuracy and Generalization. *2025 International Conference on Control, Automation and Diagnosis, ICCAD 2025*. https://doi.org/10.1109/ICCAD64771.2025.11099224

40. Saba, M., Castrillón-Ortiz, C., Valdelamar-Martínez, D., (...), Bustillo-Lecompte, C. (2025) Analysis of asbestos-cement roof classification in urban areas: Supervised and unsupervised methods with multispectral and hyperspectral remote sensing. *Remote Sensing Applications: Society and Environment*. https://doi.org/10.1016/j.rsase.2025.101464

41. Masudul Islam, S., Kumar, V., Kumar, S., Agrawal, S. (2024) Spectral mixture analysis of AVIRIS-NG hyperspectral data for material identification and classification for the part of Kolkata city. *Advances in Space Research*. https://doi.org/10.1016/j.asr.2022.12.044

42. Ronneberger, O., Fischer, P., Brox, T. (2015) U-net: Convolutional networks for biomedical image segmentation. *Lecture Notes in Computer Science*. https://doi.org/10.1007/978-3-319-24574-4_28

43. Chen, L.-C., Papandreou, G., Kokkinos, I., (...), Yuille, A.L. (2018) DeepLab: Semantic Image Segmentation with Deep Convolutional Nets, Atrous Convolution, and Fully Connected CRFs. *IEEE Transactions on Pattern Analysis and Machine Intelligence*. https://doi.org/10.1109/TPAMI.2017.2699184

44. Zhang, P., Li, J., Wang, C., Niu, Y. (2025) SAM2MS: An Efficient Framework for HRSI Road Extraction Powered by SAM2. *Remote Sensing*. https://doi.org/10.3390/rs17183181

45. Tiwari, P.S., Pande, H., Aye, M.N. (2010) Exploiting ikonos and hyperion data fusion for automated road extraction. *Geocarto International*. https://doi.org/10.1080/10106040903180485

46. Choe, E., Hong, S., Kim, Y. (2010) Down-scaling of satellite hyperspectral images for monitoring croplands. *International Geoscience and Remote Sensing Symposium (IGARSS)*. https://doi.org/10.1109/IGARSS.2010.5649715

47. Romano, G., Ricci, G.F., Gentile, F. (2025) Effectiveness evaluation of combining SAR and multiple optical data on land cover mapping of a fragmented landscape in a cloud computing platform. *Frontiers in Remote Sensing*. https://doi.org/10.3389/frsen.2025.1535418

48. Badrinarayanan, V., Kendall, A., Cipolla, R. (2017) SegNet: A Deep Convolutional Encoder-Decoder Architecture for Image Segmentation. *IEEE Transactions on Pattern Analysis and Machine Intelligence*. https://doi.org/10.1109/TPAMI.2016.2644615

49. Chaurasia, A., Culurciello, E. (2017) LinkNet: Exploiting encoder representations for efficient semantic segmentation. *2017 IEEE Visual Communications and Image Processing, VCIP 2017*. https://doi.org/10.1109/VCIP.2017.8305148

50. He, K., Zhang, X., Ren, S., Sun, J. (2016) Deep residual learning for image recognition. *Proceedings of the IEEE Computer Society Conference on Computer Vision and Pattern Recognition*. https://doi.org/10.1109/CVPR.2016.90

51. Zhang, Z., Liu, Q., Wang, Y. (2018) Road Extraction by Deep Residual U-Net. *IEEE Geoscience and Remote Sensing Letters*. https://doi.org/10.1109/LGRS.2018.2802944

52. Demir, I., Koperski, K., Lindenbaum, D., (...), Raska, R. (2018) DeepGlobe 2018: A challenge to parse the earth through satellite images. *IEEE Computer Society Conference on Computer Vision and Pattern Recognition Workshops*. https://doi.org/10.1109/CVPRW.2018.00031

53. Gorelick, N., Hancher, M., Dixon, M., (...), Moore, R. (2017) Google Earth Engine: Planetary-scale geospatial analysis for everyone. *Remote Sensing of Environment*. https://doi.org/10.1016/j.rse.2017.06.031

54. Raviz, J., Laborte, A., Barbieri, M., (...), Gonzaga, H. (2016) Mapping and monitoring rice areas in Central Luzon, Philippines using X and C-band SAR imagery. *37th Asian Conference on Remote Sensing, ACRS 2016*. https://www.scopus.com/pages/publications/85018951615

55. Acolin, A., Kim, A.M. (2022) Algorithmic justice and groundtruthing the remote mapping of informal settlements: The example of Ho Chi Minh City's periphery. *Environment and Planning B: Urban Analytics and City Science*. https://doi.org/10.1177/2399808321998708