



Rural Healthcare in India: Research for a Real Telemedicine Product

India's rural healthcare problem is not simply "there are fewer hospitals." It is a **last-mile access problem** involving distance, absent specialists, medicine stock-outs, weak diagnostics, poor referral coordination, emergency transport, affordability, and unreliable connectivity. Your product should therefore not be just a doctor video-call app; it should function as a **low-bandwidth care-coordination system connecting patients, frontline workers, PHCs, specialists, pharmacies, laboratories, ambulances, and referral hospitals.**

The strongest hackathon argument is:

"Telemedicine should not replace rural healthcare infrastructure. It should make the existing infrastructure more reachable, better coordinated, and more accountable."

1. Current healthcare availability

India's rural public-health system is designed as a three-level structure:

Facility	Intended role	Official population norm	Rural facilities functioning, 2021–22	Average rural population covered
Sub-Centre / Health and Wellness Centre	First contact, prevention, maternal-child care, screening, basic treatment	5,000 in plains; 3,000 in hilly/tribal areas	157,935	5,691
Primary Health Centre	Medical officer, outpatient care, basic diagnostics, referral	30,000 in plains; 20,000 in hilly/tribal areas	24,935	36,049
Community Health Centre	Referral hospital and specialist care	120,000 in plains; 80,000 in hilly/tribal areas	5,480	164,027

The official norms are therefore being exceeded at the national average: one rural PHC covers approximately 36,049 people against a plains norm of 30,000, while one CHC covers approximately 164,027 people against a plains norm of 120,000. The gap becomes more serious in difficult terrain, where distance and travel time matter more than the simple number of facilities. ^[1] ^[2]

The rural system also has:

- 1,275 functioning sub-divisional or sub-district hospitals.
- 767 district hospitals.

- Approximately 134,224 beds at PHCs, 184,175 at CHCs, 109,937 at sub-divisional hospitals, and 162,749 at district hospitals.
- An average radial distance of 6.28 km to a PHC and 13.39 km to a CHC.
- A PHC serving an average of 27 villages and a CHC serving approximately 121 villages. ^[1]

These are **straight-line averages**, not actual travel distances. Roads, rivers, hills, monsoon flooding, lack of public transport, and ambulance delays can make the effective journey much longer.

The 24×7 problem

Of the 24,935 rural PHCs functioning in 2021–22, only 11,250 were operating on a 24×7 basis—approximately 45.1%. This means that more than half of rural PHCs were not recorded as 24-hour facilities, which is especially important for childbirth, accidents, fever emergencies, snakebites, and acute respiratory conditions. ^[3]

A centre may exist on paper but still be functionally unavailable at night because of:

- No doctor after working hours.
- No nurse or pharmacist.
- No electricity backup.
- No laboratory technician.
- No ambulance.
- No essential medicines.
- No capacity to admit or refer a patient safely.

This distinction is important for your product: your app should show “**service availability now**”, not merely the location of the nearest hospital.

2. Availability of doctors and specialists

General doctors

The number of allopathic doctors at rural PHCs increased from 20,308 in 2005 to 30,640 in 2022, a rise of approximately 50.9%. At the national level, the official report recorded only a 3.1% shortfall against the overall requirement for PHC doctors. ^[1]

However, this number can hide serious state-level and local-level variation. A doctor may also be:

- Posted at the PHC but absent on a particular day.
- Responsible for several facilities.
- Available only during limited outpatient hours.
- Unable to provide services because laboratory staff or medicines are missing.
- Not trained or equipped for complicated cases.

Specialist doctors

The specialist shortage is much more severe at CHCs. A CHC is expected to provide four core specialists:

- Surgeon.
- Physician.
- Obstetrician and gynaecologist.
- Paediatrician.

In 2021–22, rural CHCs required approximately 21,920 specialist doctors but had only 4,485 available. This represents a shortfall of approximately 17,435 specialists. ^[4] ^[5]

Specialist	Shortfall at rural CHCs
Surgeons	83.2%
Obstetricians and gynaecologists	74.2%
Physicians	79.1%
Paediatricians	81.6%
Overall specialists	79.5%

Only a small fraction of CHCs had the complete four-specialist team. One report based on the same Rural Health Statistics data noted that only 541 of 5,480 functioning CHCs had all four required specialists. ^[6]

This creates several consequences:

- Pregnant women may need to travel to district hospitals for complications.
- Children with serious infections may not receive paediatric review locally.
- Surgical cases are referred late or directly to distant private hospitals.
- General doctors handle cases beyond their training.
- Patients may repeatedly visit a PHC without receiving a definitive diagnosis.
- Specialist consultation becomes expensive because patients turn to private providers.

What telemedicine can realistically solve

Telemedicine can reduce the specialist gap for cases that do not require immediate physical intervention. It can help with:

- Dermatology and visible skin conditions.
- Hypertension and diabetes follow-up.
- Medication review.
- Maternal-health risk screening.
- Paediatric triage.

- Mental-health counselling.
- ECG, oxygen, blood-pressure, blood-sugar, and temperature interpretation.
- Second opinions.
- Referral decisions.

It cannot replace:

- Surgery.
- Emergency childbirth procedures.
- Blood transfusion.
- Imaging that is unavailable locally.
- Physical examination when essential.
- Critical care.
- Ambulance transport.

Therefore, the app must include a **triage and referral layer**, not only a consultation screen.

3. Medicine and medical-stock availability

Medicine availability is one of the most important gaps because a consultation is useless if the patient cannot obtain the prescribed treatment.

A recent ICMR–WHO-supported assessment examined 415 facilities across 19 districts in seven Indian states. The study found that overall readiness for diabetes and hypertension care was lowest at:

- Sub-Centres: 61%.
- Community Health Centres: 59%.
- PHCs: 73%.
- Government tertiary facilities: generally above 70%.

The researchers considered a readiness score of at least 70% as the threshold for being prepared to provide these services.^[7]

Reported medicine stock-outs

Among 105 assessed sub-centres:

Medicine	Facilities reporting stock-out	Median stock-out duration
Glimepiride	50.5%	3 months
Metformin	35.2%	7 months
Atenolol	68.5%	4 months
Amlodipine	44.8%	7 months
Enalapril	66.7%	5 months

Medicine	Facilities reporting stock-out	Median stock-out duration
Frusemide	31.4%	1 month
Chlorothiazide	41.9%	Not reported

These were stock-outs reported during the previous year, not necessarily stock-outs on the survey day. The duration data still shows the scale of the problem: medicine gaps can last for months rather than a few days.^[7]

The same study reported that medicines were generally more available at government medical colleges than at peripheral public facilities. The medicine domain was the weakest part of facility readiness, even when equipment availability was better.^[7]

Why medicine stock-outs harm patients

Stock-outs cause patients to:

- Buy medicines from private pharmacies.
- Stop treatment until money becomes available.
- Split tablets or reduce doses.
- Travel repeatedly to another facility.
- Miss follow-up visits.
- Depend on informal providers.
- Experience preventable complications.

For chronic diseases, the damage accumulates slowly. A hypertensive patient who misses medication may not feel symptoms immediately but remains at risk of stroke, heart failure, kidney disease, and other complications.

A good product must therefore answer:

“Can this facility provide the medicine today?”

rather than merely:

“Was this medicine theoretically prescribed?”

4. Diagnostics and service readiness

Rural healthcare is also constrained by weak diagnostic capacity. A PHC may have a doctor but lack:

- Blood tests.
- Urine tests.
- Pregnancy tests.
- Blood-sugar testing.
- Haemoglobin testing.

- ECG.
- X-ray.
- Ultrasound.
- Trained laboratory staff.
- Reliable sample transportation.

The ICMR-WHO assessment found that public facilities were not equally prepared across all service domains. PHCs performed better than sub-centres and CHCs for diabetes and hypertension care, but staff, medicines, guidelines, and diagnostics still required improvement. ^[7]

This means a telemedicine application should not force doctors to diagnose from symptoms alone. It should support:

- Structured symptom capture.
- Vitals entry.
- Photo or document upload when possible.
- Point-of-care test entry.
- “Test required before consultation” workflows.
- Laboratory referral.
- Repeat-test reminders.
- Doctor visibility into previous readings.

For example, a hypertension consultation should ideally include:

- Blood pressure readings from multiple days.
- Pulse rate.
- Current medicines.
- Symptoms such as chest pain, weakness, breathlessness, or severe headache.
- Diabetes status.
- Kidney-related history if available.
- Whether the prescribed medicines are currently in stock.

5. How patients benefit or suffer

Benefits when care works

Rural patients benefit when services are available close to home:

- Lower travel expenses.
- Less wage loss for daily-wage workers.
- Earlier diagnosis.

- Faster specialist advice.
- Better continuity for chronic diseases.
- Reduced crowding at district hospitals.
- Lower dependence on private hospitals.
- Improved maternal and child-health follow-up.
- Better monitoring of elderly and disabled patients.

Digital connectivity creates a genuine opportunity. As of March 2024, India had approximately 398.35 million rural internet subscribers. As of April 2024, 612,952 of 644,131 villages—approximately 95.15%—had 3G or 4G mobile connectivity according to government data.^[8]

But village-level mobile coverage does not guarantee:

- Strong signal inside homes.
- Affordable data.
- Smartphone ownership.
- Digital literacy.
- Private space for consultation.
- Reliable electricity.
- Ability to upload video.
- Local-language support.

This is why your product should be designed for **low bandwidth and assisted use**, not for urban smartphone users alone.

How patients suffer

The major patient burdens are:

- Long travel to PHCs, CHCs, district hospitals, or private hospitals.
- Transport delays during emergencies.
- Repeated visits because the doctor, specialist, medicine, or test is unavailable.
- Out-of-pocket expenditure on medicines and private consultations.
- Loss of wages and school attendance.
- Delayed referral.
- Poor follow-up after the initial consultation.
- Language and literacy barriers.
- Fear of travelling alone, especially for women, elderly people, and children.

One geographic-accessibility study in a rural and remote Indian region found that delivery and inpatient care could require approximately three hours of walking time in some scenarios. It also found villages without ambulatory, delivery, or inpatient care within practical distances.

This is not a national average, but it illustrates why physical distance can become a clinical risk. ^[9]

For emergency care, the benchmark is not simply whether an ambulance exists. The National Health Mission has used a quality benchmark of ambulance arrival within 30 minutes in rural areas and arrival at a health facility within the next 30 minutes. ^[10]

6. What your product should actually solve

Product concept

Build an **offline-first rural care coordination platform** with three interfaces:

1. Patient interface

- Works through Android app, WhatsApp-style interface, IVR, or assisted kiosk.
- Supports local languages.
- Uses text and audio before video.
- Stores patient information offline.
- Shows consultation status and next steps.

2. Frontline-worker interface

- Used by ASHA workers, ANMs, village health workers, pharmacists, and rural volunteers.
- Captures vitals, symptoms, medicines, photos, and location.
- Creates referrals.
- Tracks whether the patient reached the facility.
- Records medicine and diagnostic availability.

3. Doctor and facility interface

- Allows remote doctors to review structured case summaries.
- Shows facility readiness and current stock.
- Enables scheduled audio consultations.
- Supports referral acceptance or rejection.
- Creates follow-up tasks.

The core workflow

A realistic workflow could be:

1. A patient visits a village health worker.
2. The worker records symptoms, vitals, age, pregnancy status, and existing conditions.
3. The app works offline and stores the case locally.

4. When any network becomes available, it synchronises a small text-based clinical summary.
5. A remote doctor reviews the case asynchronously or joins by audio.
6. The system checks whether the required medicine and test are available nearby.
7. If available, the patient receives a prescription and follow-up date.
8. If unavailable, the app suggests a verified alternative facility or pharmacy.
9. If dangerous symptoms are detected, the system escalates to a referral hospital and ambulance workflow.
10. The frontline worker receives a reminder to confirm whether the patient completed the referral.

This is much more valuable than a video call because it closes the gaps between **consultation, treatment, medicine availability, referral, and follow-up**.

Features that create real-world value

Problem	Product feature	Measurable outcome
Specialist shortage	Asynchronous doctor review and scheduled audio consultation	Time to specialist advice
Poor connectivity	Offline-first sync, compressed text, audio fallback	Consultation completion rate
Medicine stock-outs	Facility-level stock dashboard and substitution approval workflow	Prescription fulfilment rate
Unclear referral	Referral ticket with receiving-facility confirmation	Referral completion rate
Emergency delays	Danger-sign triage and ambulance escalation	Time from red flag to transport
Language barriers	Tamil, Hindi, Telugu, Bengali, and local-language audio prompts	Patient comprehension
Poor follow-up	SMS, IVR, and health-worker reminders	Follow-up completion rate
Fragmented records	Longitudinal patient record with consent	Repeat-visit continuity
Low digital literacy	Assisted mode for ASHA/ANM workers	Successful assisted consultations
Weak accountability	Facility availability and stock timestamp	Data freshness and reliability

7. Low-bandwidth design

Do not make video the default. Use a communication hierarchy:

1. Offline structured form.
2. Store-and-forward text summary.
3. Compressed image.
4. Voice note or audio call.
5. Video only when the network supports it.

A rural consultation record should be small enough to synchronise over an unstable mobile connection. The app can send:

- Patient age and sex.
- Village and facility.
- Main symptoms.
- Duration.
- Vital signs.
- Red-flag responses.
- Existing diseases.
- Current medicines.
- One or two compressed images where clinically useful.
- Requested specialist.
- Required urgency.

The doctor does not always need a live video call. In many cases, a structured case packet reviewed within a defined time is more reliable.

Suggested network states

- **Offline:** collect data, provide first-aid and triage guidance, queue consultation.
- **Weak network:** synchronise text, vitals, and compressed images.
- **Moderate network:** audio consultation.
- **Strong network:** video consultation and document exchange.

This design also protects patients in areas where the 95% village connectivity figure does not represent continuous usable internet.

8. Safety and medical governance

This is a healthcare product, so clinical safety should be central to your pitch.

The app should not

- Diagnose emergencies solely through an AI chatbot.
- Give prescription suggestions without doctor supervision.
- Promise that a nearby facility has stock unless the information has a timestamp.
- Replace in-person examination where it is medically necessary.
- Hide uncertainty from the patient.
- Allow unverified providers to provide clinical advice.
- Treat a video consultation as proof that treatment is safe.

The app should include

- Red-flag screening.
- Doctor verification.
- Consent before sharing records.
- Role-based access.
- Audit logs.
- Encrypted data transfer and storage.
- Clear escalation instructions.
- Timestamped medicine and facility information.
- Patient confirmation in the local language.
- Human review for high-risk cases.

Examples of red flags requiring urgent in-person escalation include:

- Severe breathing difficulty.
- Chest pain.
- Unconsciousness or confusion.
- Stroke-like weakness or speech difficulty.
- Severe bleeding.
- Pregnancy with heavy bleeding, seizures, or prolonged labour.
- Severe dehydration in a child.
- Snakebite.
- Serious injury.
- Very high fever with altered mental status.

The exact clinical rules should be validated with licensed doctors and public-health experts before deployment.

9. How to validate it beyond a hackathon

Do not claim that the prototype has improved health outcomes. Instead, propose a pilot with measurable indicators.

Pilot design

Choose:

- One district.
- Three to five PHCs.
- Ten to twenty sub-centres or village workers.
- One referral hospital.

- One laboratory or sample-collection partner.
- One ambulance or emergency coordination partner.

Start with a limited disease area, such as:

- Hypertension and diabetes.
- Maternal risk screening.
- Paediatric fever and respiratory symptoms.
- Dermatology.
- Elderly-care follow-up.

Do not begin with every disease. A narrow, measurable use case is more credible.

Metrics

Measure before and after deployment:

- Average travel trips avoided.
- Time from patient registration to doctor advice.
- Percentage of consultations completed despite weak connectivity.
- Percentage of prescriptions fulfilled.
- Medicine stock accuracy.
- Referral acceptance rate.
- Referral completion rate.
- Follow-up completion rate.
- Percentage of patients needing escalation.
- Patient-reported cost saved.
- Patient and health-worker satisfaction.
- Number of cases incorrectly escalated or missed.

A strong pilot target could be:

“Within three months, demonstrate that at least 80% of scheduled remote consultations are completed, at least 90% of stock records are updated within seven days, and at least 70% of high-priority referrals receive confirmation from the receiving facility.”

These are proposed product targets, not existing national statistics. Label them clearly as pilot goals.

10. Your hackathon pitch

Use this simple structure:

Problem

“Rural India has a large network of public facilities, but availability is uneven. A rural PHC serves about 36,049 people on average, a CHC about 164,027, and nearly 80% of required specialists at CHCs are missing. Even where doctors are available, medicines and diagnostics may not be.”^[1]

Insight

“The problem is not only the absence of doctors. It is the failure to connect the patient to the right person, facility, medicine, test, transport, and follow-up at the right time.”

Solution

“Our platform is an offline-first, low-bandwidth care-coordination system. It enables frontline workers to capture cases, remote doctors to review them, facilities to report real-time readiness, patients to receive local-language instructions, and referral hospitals to confirm whether patients actually arrived.”

Differentiator

“Most telemedicine apps assume that the patient has a smartphone, good internet, a private room, and a nearby pharmacy. Our product assumes the opposite: shared phones, weak networks, limited literacy, missing medicines, and the need for an intermediary health worker.”

Impact

“We measure care completion, not app downloads: consultation completion, prescription fulfilment, referral completion, follow-up adherence, travel avoided, and cost saved.”

Evidence limitations

Use the data carefully in your presentation:

- The latest fully accessible nationwide Rural Health Statistics source used here is for 2021–22, with data up to 31 March 2022.^[1]
- The medicine-stock study covered 415 facilities in 19 districts across seven states, so its stock-out percentages should not be presented as the exact national rate.^[7]
- Internet connectivity figures refer mainly to village-level 3G/4G coverage or subscriber counts; they do not guarantee reliable indoor connectivity or smartphone access for every household.^[8]
- Facility “availability” does not necessarily mean that the facility is open, staffed, stocked, and clinically capable at the moment the patient arrives.

Your strongest product position is therefore not “we will digitally cure rural healthcare.” It is:

“We make rural healthcare actionable: find a functioning provider, transmit a case on weak networks, verify medicines and tests, coordinate referrals, and ensure follow-up.”

That is a real-world problem, a credible technology solution, and a much stronger hackathon story than a generic teleconsultation app.

✱✱

1. <https://ruralhospitalnetwork.org/wp-content/uploads/2023/01/RHS-2021-22.pdf>
2. <https://www.pib.gov.in/PressReleaseIframePage.aspx?PRID=1896950>
3. <https://sansad.in/getFile/loksabhaquestions/annex/1711/AU1553.pdf>
4. <https://www.thehindu.com/data/data-shortfall-of-surgeons-gynaecologists-and-paediatricians-in-rural-india-was-80-in-2022/article66452879.ece>
5. <https://www.etvbharat.com/english/sukhibhava/sukhibhava-news/indias-rural-health-sector-facing-shortage-of-specialists-union-health-ministry-report/na20230113104327913913549>
6. <https://www.downtoearth.org.in/health/poor-infrastructure-staff-crunch-continue-to-plague-healthcare-in-rural-india-centre-87250>
7. <https://ijmr.org.in/preparedness-of-public-private-health-facilities-for-management-of-diabetes-hypertension-in-19-districts-in-india/>
8. <https://www.pib.gov.in/PressReleaseIframePage.aspx?PRID=2040566>
9. <https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0239326>
10. <https://pmc.ncbi.nlm.nih.gov/articles/PMC10692338/>
11. <https://sansad.in/getFile/loksabhaquestions/annex/1715/AU226.pdf?source=pqals>
12. <https://medicaldialogues.in/news/health/80-shortfall-of-specialist-doctors-in-chcs-health-ministry-report-105783>
13. <https://ruralindiaonline.org/en/library/resource/rural-health-statistics-2021-22/>
14. <https://m.thewire.in/article/health/eight-charts-to-explain-the-state-of-rural-healthcare-in-india>
15. <https://www.tribuneindia.com/news/editorials/ailing-rural-healthcare-470130/>
16. <https://theprint.in/health/primary-health-centres-see-over-50-increase-in-doctors-but-shortfall-of-80-specialists-at-chcs/1310905/>
17. <https://pdfs.semanticscholar.org/4486/842098378fcb4b346fc583b4cd7106257b7.pdf>
18. <https://www.tribuneindia.com/news/nation/80-shortfall-of-specialists-in-rural-india-469835/>
19. <https://www.ris.org.in/sites/default/files/Publication/Policy-Brief-No-129-Monika-Kochar.pdf>
20. <https://vajiramandravi.com/current-affairs/rural-health-statistics-report-2021-22/>
21. <https://www.indiaherald.com/VidhyaKiVaidhyam/Read/994895407/India-Doctor-Shortage-MBBS-Seats-vs-Rural-Health-care-Gap>
22. <https://www.mohfw.gov.in/documents/staistics>
23. <https://www.mohfw.gov.in/documents/publication/archives>
24. <https://ijmr.org.in/?view-pdf=1&embedded=true&article=99953280a738a111f2389dd3aa90b5b1TxZeVfXxK2TeUA==>

25. <https://ijmr.org.in/?view-pdf=1&embedded=true&article=03ec7df6976e31f827ee3c0e113b1734rgsGPLxBrppwUg==>
26. <https://mohfw.gov.in/documents/publication/archives>
27. [https://www.mohfw.gov.in/sites/default/files/induction material.pdf](https://www.mohfw.gov.in/sites/default/files/induction%20material.pdf)
28. <https://www.mohfw.gov.in/documents/statistics/archives>
29. <https://journals.sagepub.com/doi/pdf/10.1177/09720634221078064>
30. <https://pmc.ncbi.nlm.nih.gov/articles/PMC9107149/>
31. https://www.nmji.in/content/141/2021/34/1/Images/NatlMedJIndia_2021_34_1_29_323661.pdf
32. <https://nmji.in/?view-pdf=1&embedded=true&article=e0d7668b87ded599d6b239755e10797849HzDDCGoGA=>
33. <https://pdfs.semanticscholar.org/1dcf/15d009841714cbea3c1cb91fc7cf1ba9a594.pdf>
34. https://cbhidghs.mohfw.gov.in/WriteReadData/l892s/Final_Central Bureau of Health Intelligence July 2024.pdf
35. <https://health.economictimes.indiatimes.com/news/industry/critical-medicine-shortages-uncovered-for-diabetes-and-hypertension-in-rural-india/121589284>
36. <https://www.pib.gov.in/PressReleasePage.aspx?PRID=1947691>
37. https://cag.gov.in/uploads/download_audit_report/2024/08.-Chapter-V-0678f47ad07bc24.87026435.pdf
38. https://cag.gov.in/uploads/download_audit_report/2024/I-CHAPTER-V--Healthcare-Infrastructure-067640dd62a5453.38034004.pdf
39. <https://www.pib.gov.in/PressReleasePage.aspx?PRID=1896948>
40. <https://www.pib.gov.in/PressReleasePage.aspx?PRID=1943657>
41. <https://hal.science/hal-04384064/document>
42. <https://so03.tci-thaijo.org/index.php/jpss/article/download/274918/185291/1184544>
43. <https://www.frontiersin.org/journals/public-health/articles/10.3389/fpubh.2023.1160088/full>
44. <https://archive.org/details/PARI.national-family-health-survey-nfhs-5-2019-20-fact-sheets-key-indicators--->
45. <https://ruralindiaonline.org/en/library/resource/national-family-health-survey-nfhs-5-2019-21-compendium-of-fact-sheets-key-indicators---india-and-14-statesuts-phase-ii/>
46. https://www.medicalbuyer.co.in/wp-content/uploads/2023/07/RHS-2021-22_compressed-1.pdf
47. <https://static.pib.gov.in/WriteReadData/specificdocs/documents/2025/jun/doc2025630578601.pdf>
48. <https://static.pib.gov.in/WriteReadData/specificdocs/documents/2025/jun/doc2025612568801.pdf>
49. https://www.trai.gov.in/sites/default/files/2025-04/QPIR-Dec-24_24_04.pdf
50. https://www.trai.gov.in/sites/default/files/2024-09/Report_14082024.pdf
51. <https://static.pib.gov.in/WriteReadData/specificdocs/documents/2024/dec/doc20241226477201.pdf>
52. <https://static.pib.gov.in/WriteReadData/specificdocs/documents/2024/oct/doc20241021421101.pdf>
53. <https://static.pib.gov.in/WriteReadData/specificdocs/documents/2024/aug/doc202486366801.pdf>
54. <https://www.lightreading.com/digital-divide/indian-telcos-eye-rural-expansion-as-data-consumption-and-digital-adoption-grow>
55. <https://telecomtalk.info/india-achieves-95percent-rural-internet-connectivity-3g4g/979718/>
56. <https://telecom.economictimes.indiatimes.com/news/policy/95-15-of-villages-in-india-have-internet-access-today-telcom-ministry/112215646>

57. <https://economictimes.indiatimes.com/tech/technology/4g-network-penetrates-95-of-the-population-internet-connections-jump-to-over-97-crore-in-june-2024/articleshow/121860412.cms>
58. <https://ddnews.gov.in/en/centre-boosts-internet-connectivity-across-india-under-digital-india-initiative/>
59. <https://www.indiatvnews.com/technology/news/internet-access-in-india-over-95-percent-villages-covered-under-3g-4g-connectivity-2024-08-02-944893>
60. <https://myjiostatic.cdn.jio.com/cms/assets/iv3/Analysys Mason - Industry Report for Jio 2026-06-17.pdf>
61. https://journals.lww.com/ijcm/fulltext/2024/49020/can_the_emergency_medical_service_ems_system.32.aspx
62. [https://www.unescap.org/sites/default/files/Bulletin 84 Article5.pdf](https://www.unescap.org/sites/default/files/Bulletin%2084%20Article5.pdf)
63. [https://journals.lww.com/jfmpc/fulltext/2021/10050/insights from national survey on household.20.aspx](https://journals.lww.com/jfmpc/fulltext/2021/10050/insights_from_national_survey_on_household.20.aspx)
64. <https://d-nb.info/124132235X/34>
65. <https://pmc.ncbi.nlm.nih.gov/articles/PMC12645763/>
66. <https://pmc.ncbi.nlm.nih.gov/articles/PMC12230409/>
67. <https://www.nirdprojms.in/index.php/jrd/article/download/172462/116669/415596>
68. <https://www.tandfonline.com/doi/full/10.1080/23288604.2024.2380251>
69. <https://pure.jgu.edu.in/id/eprint/1672/1/Mukhopadhyay2021.pdf>
70. <https://www.iimb.ac.in/sites/default/files/inline-files/Emergency Healthcare v3 May 2013.pdf>
71. <https://www.downtoearth.org.in/environment/ambulance-emergency-40605>
72. <https://hss.iitm.ac.in/ctap/wp-content/uploads/2023/01/108-Final-Report.pdf>
73. <https://hsrii.org/wp-content/uploads/2016/06/PLOS-One-2014-ERS-Punjab.pdf>