

ON THE LARGE-SCALE SUBLINEAR GEOMETRY OF HEINTZE SPACES

Gabriel Pallier - Paris-Sud University, Orsay

March 19th, 2019

Analysis and Geometry Seminar, Bristol

Outline

Heintze spaces

Large-scale geometry and the boundary

Invariants

HEINTZE SPACES

Heintze space: Definition

Definition

Let Y be a connected Riemannian manifold of **negative sectional curvature**. If the isometry group $\text{Isom}(Y)$ acts transitively on Y , then Y is called a Heintze space.


Heintze space: Definition

Definition

Let Y be a connected Riemannian manifold of **negative sectional curvature**. If the isometry group $\text{Isom}(Y)$ acts transitively on Y , then Y is called a Heintze space.

Example: Y is $\mathbf{R} \times \mathbf{R}^2$ with coordinates (t, x, y) and the Riemannian metric $ds^2 = dt^2 + e^{-2t}(dx^2 + dy^2)$.

Y is the hyperbolic 3-space (constant curvature -1) and (x, y, t) are **horospherical coordinates**.




Isometries of \mathbb{H}^3 seen as a Heintze space

$$ds^2 = dt^2 + e^{-t}(dx^2 + dy^2).$$

Let us check that $\text{Isom}(Y)$ is transitive.

1. For all $(u, v) \in \mathbb{R}^2$,
 $(t, x, y) \mapsto (t, x + u, y + v)$ is an isometry.




Isometries of \mathbb{H}^3 seen as a Heintze space

$$ds^2 = dt^2 + e^{-t}(dx^2 + dy^2).$$

Let us check that $\text{Isom}(Y)$ is transitive.

1. For all $(u, v) \in \mathbb{R}^2$,
 $(t, x, y) \mapsto (t, x + u, y + v)$ is an isometry.




Isometries of \mathbb{H}^3 seen as a Heintze space

$$ds^2 = dt^2 + e^{-t}(dx^2 + dy^2).$$

Let us check that $\text{Isom}(Y)$ is transitive.

1. For all $(u, v) \in \mathbb{R}^2$,
 $(t, x, y) \mapsto (t, x + u, y + v)$ is an isometry.




Isometries of \mathbb{H}^3 seen as a Heintze space

$$ds^2 = dt^2 + e^{-t}(dx^2 + dy^2).$$

Let us check that $\text{Isom}(Y)$ is transitive.

1. For all $(u, v) \in \mathbb{R}^2$,
 $(t, x, y) \mapsto (t, x + u, y + v)$ is an isometry.




Isometries of \mathbb{H}^3 seen as a Heintze space

$$ds^2 = dt^2 + e^{-t}(dx^2 + dy^2).$$

Let us check that $\text{Isom}(Y)$ is transitive.

1. For all $(u, v) \in \mathbb{R}^2$,
 $(t, x, y) \mapsto (t, x + u, y + v)$ is an isometry.




Isometries of \mathbb{H}^3 seen as a Heintze space

$$ds^2 = dt^2 + e^{-t}(dx^2 + dy^2).$$

Let us check that $\text{Isom}(Y)$ is transitive.

1. For all $(u, v) \in \mathbb{R}^2$,
 $(t, x, y) \mapsto (t, x + u, y + v)$ is an isometry.




Isometries of \mathbb{H}^3 seen as a Heintze space

$$ds^2 = dt^2 + e^{-t}(dx^2 + dy^2).$$

Let us check that $\text{Isom}(Y)$ is transitive.

1. For all $(u, v) \in \mathbb{R}^2$,
 $(t, x, y) \mapsto (t, x + u, y + v)$ is an isometry.
2. For all $s \in \mathbb{R}$,
 $(t, x, y) \mapsto (t + s, e^s x, e^s y)$ is an isometry.




Isometries of \mathbb{H}^3 seen as a Heintze space

$$ds^2 = dt^2 + e^{-t}(dx^2 + dy^2).$$

Let us check that $\text{Isom}(Y)$ is transitive.

1. For all $(u, v) \in \mathbb{R}^2$,
 $(t, x, y) \mapsto (t, x + u, y + v)$ is an isometry.
2. For all $s \in \mathbb{R}$,
 $(t, x, y) \mapsto (t + s, e^s x, e^s y)$ is an isometry.




Isometries of \mathbb{H}^3 seen as a Heintze space

$$ds^2 = dt^2 + e^{-t}(dx^2 + dy^2).$$

Let us check that $\text{Isom}(Y)$ is transitive.

1. For all $(u, v) \in \mathbb{R}^2$,
 $(t, x, y) \mapsto (t, x + u, y + v)$ is an isometry.
2. For all $s \in \mathbb{R}$,
 $(t, x, y) \mapsto (t + s, e^s x, e^s y)$ is an isometry.



$G = \mathbb{R} \ltimes \mathbb{R}^2$ acts **simply transitively** by isometries on Y .

Other Heintze spaces

To form G , we used $\mathbf{R} \curvearrowright \mathbf{R}^2$ by $s.(x,y) = \delta^s(x,y) = (e^s x, e^s y)$. More general:

Other Heintze spaces

To form G , we used $\mathbf{R} \curvearrowright \mathbf{R}^2$ by $s.(x,y) = \delta^s(x,y) = (e^s x, e^s y)$. More general:

1. **Diagonal** action: Let $\mu \geq 1$ be a parameter

$$\delta^s(x,y) = (e^s x, e^{\mu s} y) = \exp \left[s \begin{pmatrix} 1 & 0 \\ 0 & \mu \end{pmatrix} \right] \begin{pmatrix} x \\ y \end{pmatrix}$$

a left inv. metric is $ds^2 = dt^2 + e^{-2t}(dx^2 + e^{2(1-\mu)t}dy^2)$.

2. **Unipotent** action:

$$\delta^s(x,y) = (e^s x + s e^s y, e^s y) = \exp \left[s \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \right] \begin{pmatrix} x \\ y \end{pmatrix}$$


a left inv. metric is $ds^2 = dt^2 + e^{-2t}(dx^2 + (1 + t^2)dy^2 - 2tdxdy)$.

Fact (Consequence of Heintze's theorem)

The corresponding groups G , G_μ , G' with all their left invariant metrics are **all the 3-dim Heintze spaces**.


The boundary of a 3-dimensional Heintze space

All the isometries described fix a **special boundary point** of Y that we denote ω . Horospheres centered at ω are **Euclidean planes** (if $\dim Y = 3$).




The boundary of a 3-dimensional Heintze space

All the isometries described fix a **special boundary point** of Y that we denote ω . Horospheres centered at ω are **Euclidean planes** (if $\dim Y = 3$). The boundary minus ω is denoted $\partial_\infty^* Y$. Geodesic lines ending at ω have form $\{(t, x, y)\}_{t \in \mathbb{R}}$.



The boundary of a 3-dimensional Heintze space

All the isometries described fix a **special boundary point** of Y that we denote ω . Horospheres centered at ω are **Euclidean planes** (if $\dim Y = 3$). The boundary minus ω is denoted $\partial_\infty^* Y$. Geodesic lines ending at ω have form $\{(t, x, y)\}_{t \in \mathbb{R}}$. One can identify $\partial_\infty^* Y$ and \mathbb{R}^2 .




A quasidistance on the boundary minus ω

$$\rho((-\infty, x, y), (-\infty, x', y')) := \exp \left(-\frac{1}{2} \lim_{t \rightarrow -\infty} d_Y((-t, x_0, y_0), (-t, (x_1, y_1)) + 2t) \right)$$

The boundary of a 3-dimensional Heintze space

All the isometries described fix a **special boundary point** of Y that we denote ω . Horospheres centered at ω are **Euclidean planes** (if $\dim Y = 3$). The boundary minus ω is denoted $\partial_\infty^* Y$. Geodesic lines ending at ω have form $\{(t, x, y)\}_{t \in \mathbb{R}}$. One can identify $\partial_\infty^* Y$ and \mathbb{R}^2 .




A quasidistance on the boundary minus ω

$$\rho((-\infty, x, y), (-\infty, x', y')) := \exp \left(-\frac{1}{2} \lim_{t \rightarrow -\infty} d_Y((-t, x_0, y_0), (-t, (x_1, y_1)) + 2t) \right)$$


Why such a quasidistance on the boundary?

Isometries of Y extend to ∂_∞^* , by homeomorphisms.



Why such a quasidistance on the boundary?

Isometries of Y extend to ∂_∞^* , by homeomorphisms. But there is no invariant distance inducing the topology.




Why such a quasidistance on the boundary?

Isometries of Y extend to ∂_∞^* , by homeomorphisms. But there is no invariant distance inducing the topology. Nevertheless,

$$\forall \xi_0, \xi_1 \in \partial_\infty^* Y, \rho(\delta^s \xi_0, \delta^s \xi_1) = e^s \rho(\xi_0, \xi_1).$$

Equipped with ρ , ∂_∞^* is a self-similar space. Identified with \mathbf{R}^2 , self-similarities are the δ^s .




How do the small balls of ρ look like?

$$\begin{array}{l} \text{Group} \\ \text{self-similarities} \end{array} \left\| \left\{ \begin{array}{c} G \\ \left(\begin{array}{cc} e^s & 0 \\ 0 & e^s \end{array} \right) \end{array} \right\} \middle| \left\{ \begin{array}{c} G' \\ \left(\begin{array}{cc} e^s & se^s \\ 0 & e^s \end{array} \right) \end{array} \right\} \middle| \left\{ \begin{array}{c} G_\mu \ (\mu > 1) \\ \left(\begin{array}{cc} e^s & 0 \\ 0 & e^{\mu s} \end{array} \right) \end{array} \right\} \right.$$

How do the small balls of ρ look like?

Group self-similarities


G	G'	$G_\mu (\mu > 1)$
$\left\{ \begin{pmatrix} e^s & 0 \\ 0 & e^s \end{pmatrix} \right\}$	$\left\{ \begin{pmatrix} e^s & se^s \\ 0 & e^s \end{pmatrix} \right\}$	$\left\{ \begin{pmatrix} e^s & 0 \\ 0 & e^{\mu s} \end{pmatrix} \right\}$



How do the small balls of ρ look like?

Group self-similarities


G	G'	$G_\mu (\mu > 1)$
$\left\{ \begin{pmatrix} e^s & 0 \\ 0 & e^s \end{pmatrix} \right\}$	$\left\{ \begin{pmatrix} e^s & se^s \\ 0 & e^s \end{pmatrix} \right\}$	$\left\{ \begin{pmatrix} e^s & 0 \\ 0 & e^{\mu s} \end{pmatrix} \right\}$



How do the small balls of ρ look like?


Group self-similarities $\left\| \left\{ \begin{pmatrix} e^s & 0 \\ 0 & e^s \end{pmatrix} \right\} \middle| \left\{ \begin{pmatrix} e^s & se^s \\ 0 & e^s \end{pmatrix} \right\} \middle| \left\{ \begin{pmatrix} e^s & 0 \\ 0 & e^{\mu s} \end{pmatrix} \right\} \right\|$

$G \qquad G' \qquad G_\mu (\mu > 1)$



How do the small balls of ρ look like?

Group self-similarities $\left\| \left\{ \begin{pmatrix} e^s & 0 \\ 0 & e^s \end{pmatrix} \right\} \middle| \left\{ \begin{pmatrix} e^s & se^s \\ 0 & e^s \end{pmatrix} \right\} \middle| \left\{ \begin{pmatrix} e^s & 0 \\ 0 & e^{\mu s} \end{pmatrix} \right\} \right\|$



LARGE-SCALE GEOMETRY AND THE BOUNDARY

Quasiisometry and Sublinearly biLipschitz Equivalence

Y, Y' are pointed metric spaces, $\lambda \geq 1$.

$f : Y \rightarrow Y'$ is a **quasiisometry** (QI) if $\exists c \geq 0$ s.t. $\forall y_1, y_2 \in Y, \forall y' \in Y'$,

$$\begin{cases} \lambda^{-1}d(y_1, y_2) - c \leq d(f(y_1), f(y_2)) \leq \lambda d(y_1, y_2) + c \\ d(y', f(Y)) \leq c. \end{cases}$$

Quasiisometry and Sublinearly biLipschitz Equivalence

Y, Y' are pointed metric spaces, $\lambda \geq 1$.

$f : Y \rightarrow Y'$ is a **quasiisometry** (QI) if $\exists c \geq 0$ s.t. $\forall y_1, y_2 \in Y, \forall y' \in Y'$,

$$\begin{cases} \lambda^{-1}d(y_1, y_2) - c \leq d(f(y_1), f(y_2)) \leq \lambda d(y_1, y_2) + c \\ d(y', f(Y)) \leq c. \end{cases}$$

$f : Y \rightarrow Y'$ is a **sublinearly biLipschitz equivalence** (SBE) if there exists a sublinear $\nu : \mathbf{R}_{\geq 0} \rightarrow \mathbf{R}_{\geq 1}$ s.t. $\forall y_1, y_2 \in Y$ and $\forall y' \in Y'$,

$$\begin{cases} \lambda^{-1}d(y_1, y_2) - \nu(|y_1| + |y_2|) \leq d(f(y_1), f(y_2)) \leq \lambda d(y_1, y_2) + \nu(|y_1| + |y_2|) \\ d(y', f(Y)) \leq \nu(|y'|), \end{cases}$$

where $|\cdot|$ denotes the distance to base-point.

Large-scale geometries of Heintze spaces

For a pair of Heintze spaces Y, Y' :

Large-scale geometries of Heintze spaces

For a pair of Heintze spaces Y, Y' :

Y, Y' isometric

$\Downarrow (1)$

$\text{Isom}(Y), \text{Isom}(Y')$ isomorphic (up to taking cocompact subgroups and quotients
by compact subgroups)

$\Downarrow (2)$

Y, Y' quasiisometric

$\Downarrow (3)$

Y, Y' sublinearly biLipschitz equivalent

Large-scale geometries of Heintze spaces

For a pair of Heintze spaces Y, Y' :

Y, Y' isometric

\Downarrow (1)

$\text{Isom}(Y), \text{Isom}(Y')$ isomorphic (up to taking cocompact subgroups and quotients
by compact subgroups)

\Downarrow (2)

Y, Y' quasiisometric

\Downarrow (3)

Y, Y' sublinearly biLipschitz equivalent

Question

Can (2), or even (2) and (3) be reversed? If not, how? Positive partial answers can be obtained through the search of QI or SBE **invariants**.

Quasiisometries and the boundary

Let $t \geq 1$. A pair of subsets (a, a^+) of a quasimetric space is a **t -ring** if there is a ball B such that $B \subseteq a \subseteq a^+ \subseteq tB$. $\text{radius}(B)$ is an **inner radius** and $\tau = \log t$ is called an **asphericity** for (a, a^+) . If $a = a^+$, **round set**.

A homeomorphism f is called quasisymmetric if f and its inverse **preserve bounded asphericity**.

Quasiisometries and the boundary

Let $t \geq 1$. A pair of subsets (a, a^+) of a quasimetric space is a **t -ring** if there is a ball B such that $B \subseteq a \subseteq a^+ \subseteq tB$. $\text{radius}(B)$ is an **inner radius** and $\tau = \log t$ is called an **asphericity** for (a, a^+) . If $a = a^+$, **round set**.

A homeomorphism f is called **quasisymmetric** if f and its inverse **preserve bounded asphericity**.

Theorem (known under some form since the 70s)

Let Y and Y' be Heintze spaces. Assume that there exists a quasiisometry $f : Y \rightarrow Y'$. Then $\partial_\infty f$ extends to a homeomorphism $\partial_\infty f : \partial_\infty Y \rightarrow \partial_\infty Y'$. Further one can assume that f preserves maps the distinguished points one to another, and $\partial_\infty f : \partial_\infty^* Y \rightarrow \partial_\infty^* Y'$ is **quasisymmetric**.

SBEs and the boundary

Let $s_n \rightarrow -\infty$. A family of rings (a_n, a_n^+) on $(\partial_\infty^*, \rho)$ with inner radii e^{s_n} and asphericities τ_n is said to have **sublinear asphericity** if $\tau_n \ll |s_n|$.

SBEs and the boundary

Let $s_n \rightarrow -\infty$. A family of rings (a_n, a_n^+) on $(\partial_\infty^*, \rho)$ with inner radii e^{s_n} and asphericities τ_n is said to have **sublinear asphericity** if $\tau_n \ll |s_n|$.

A homeomorphism is sublinearly quasymmetric if it is **biHölder continuous** and **preserves sublinear asphericity** for families of rings.

SBEs and the boundary

Let $s_n \rightarrow -\infty$. A family of rings (a_n, a_n^+) on $(\partial_\infty^*, \rho)$ with inner radii e^{s_n} and asphericities τ_n is said to have **sublinear asphericity** if $\tau_n \ll |s_n|$.

A homeomorphism is sublinearly quasymmetric if it is **biHölder continuous** and **preserves sublinear asphericity** for families of rings.

Theorem

1. (Cornulier 2016) SBEs induce biHölder homeomorphisms between the boundaries [of Heintze spaces, and more generally Gromov-hyperbolic spaces].

SBEs and the boundary


Let $s_n \rightarrow -\infty$. A family of rings (a_n, a_n^+) on $(\partial_\infty^*, \rho)$ with inner radii e^{s_n} and asphericities τ_n is said to have **sublinear asphericity** if $\tau_n \ll |s_n|$.

A homeomorphism is sublinearly quasymmetric if it is **biHölder continuous** and **preserves sublinear asphericity** for families of rings.


Theorem

1. (Cornulier 2016) SBEs induce biHölder homeomorphisms between the boundaries [of Heintze spaces, and more generally Gromov-hyperbolic spaces].
2. (P. 2018) SBEs induce sublinearly quasymmetric homeomorphisms between the boundaries.


Sublinear asphericity is preserved




Sublinear asphericity is preserved




Sublinear asphericity is preserved




$$\tau_n = O(\sqrt{n})$$



Sublinear asphericity is preserved



$$\tau_n = O(\sqrt{n})$$



$$\tau'_n = O(\sqrt{n})$$

Figure: Round sets and their images in Euclidean \mathbf{R}^2 .

G and G' are SBE

Let Y and Y' be the Heintze spaces associated to G and G'

G and G' are SBE

Let Y and Y' be the Heintze spaces associated to G and G'

Observation (Cornulier 2008, 2011)

Y and Y' are SBE (via the identity map in horospherical coordinates).

G and G' are SBE

Let Y and Y' be the Heintze spaces associated to G and G'

Observation (Cornulier 2008, 2011)

Y and Y' are SBE (via the identity map in horospherical coordinates).

On the boundary: the identity map through identification with \mathbf{R}^2 is a sublinearly quasisymmetric homeomorphism. Precisely $\nu = O(\log)$.

G and G' are SBE

Let Y and Y' be the Heintze spaces associated to G and G'

Observation (Cornulier 2008, 2011)

Y and Y' are SBE (via the identity map in horospherical coordinates).

On the boundary: the identity map through identification with \mathbf{R}^2 is a sublinearly quasisymmetric homeomorphism. Precisely $\nu = O(\log)$.

Negative answer to the second part of the Question (even restricted to 3-dim Heintze groups).

A method to produce sublin-q.s. homeos (I)

Ingredients

The Lebesgue measure λ on $[0, 1]$,

A method to produce sublin-q.s. homeos (I)

Ingredients

The Lebesgue measure λ on $[0, 1]$,

A decreasing $(\epsilon_n)_{n \geq 0}$ in $(0, 1)$ going to 0 **but not in ℓ^1** ,

A method to produce sublin-q.s. homeos (I)



Ingredients

The Lebesgue measure λ on $[0, 1]$,

A decreasing $(\epsilon_n)_{n \geq 0}$ in $(0, 1)$ going to 0 **but not in** ℓ^1 ,

An infinite rooted binary tree,

\aleph_0 independent random variables uniformly distributed in $\{\leftarrow, \rightarrow\}$.



A method to produce sublin-q.s. homeos (I)

Ingredients



The Lebesgue measure λ on $[0, 1]$,

A decreasing $(\epsilon_n)_{n \geq 0}$ in $(0, 1)$ going to 0 **but not in** ℓ^1 ,

An infinite rooted binary tree,

\aleph_0 independent random variables uniformly distributed in $\{\leftarrow, \rightarrow\}$.

1st step: Obtain a random measure M on $[0, 1]$.



A method to produce sublin-q.s. homeos (I)

Ingredients



The Lebesgue measure λ on $[0, 1]$,

A decreasing $(\epsilon_n)_{n \geq 0}$ in $(0, 1)$ going to 0 **but not in** ℓ^1 ,

An infinite rooted binary tree,

\aleph_0 independent random variables uniformly distributed in $\{\leftarrow, \rightarrow\}$.

1st step: Obtain a random measure M on $[0, 1]$.



A method to produce sublin-q.s. homeos (I)

Ingredients



The Lebesgue measure λ on $[0, 1]$,

A decreasing $(\epsilon_n)_{n \geq 0}$ in $(0, 1)$ going to 0 **but not in** ℓ^1 ,

An infinite rooted binary tree,

\aleph_0 independent random variables uniformly distributed in $\{\leftarrow, \rightarrow\}$.

1st step: Obtain a random measure M on $[0, 1]$.



A method to produce sublin-q.s. homeos (I)

Ingredients



The Lebesgue measure λ on $[0, 1]$,

A decreasing $(\epsilon_n)_{n \geq 0}$ in $(0, 1)$ going to 0 **but not in** ℓ^1 ,

An infinite rooted binary tree,

\aleph_0 independent random variables uniformly distributed in $\{\leftarrow, \rightarrow\}$.

1st step: Obtain a random measure M on $[0, 1]$.



A method to produce sublin-q.s. homeos (I)

Ingredients


The Lebesgue measure λ on $[0, 1]$,

A decreasing $(\epsilon_n)_{n \geq 0}$ in $(0, 1)$ going to 0 **but not in** ℓ^1 ,


An infinite rooted binary tree,

\aleph_0 independent random variables uniformly distributed in $\{\leftarrow, \rightarrow\}$.

1st step: Obtain a random measure M on $[0, 1]$.



$$M = \lim_n \lambda_n$$



A method to produce sublin-q.s. homeos (II)

2nd step: Take the primitive $\phi : [0, 1] \rightarrow [0, 1]$ in the distributional sense.

ϕ is not absolutely continuous. The derivative is λ -a.e. 0. The modulus of continuity deviates sublinearly from that of a Lipschitz function:

$\log |\phi(x) - \phi(y)| \leq \log |x - y| + \nu(\log |x - y|)$, sublinear ν .

A method to produce sublin-q.s. homeos (II)

2nd step: Take the primitive $\phi : [0, 1] \rightarrow [0, 1]$ in the distributional sense.

ϕ is not absolutely continuous. The derivative is λ -a.e. 0. The modulus of continuity deviates sublinearly from that of a Lipschitz function:

$$\log |\phi(x) - \phi(y)| \leq \log |x - y| + \nu(\log |x - y|), \text{ sublinear } \nu.$$

3rd step: To get a sublinearly quasisymmetric homeomorphism of the square (torus), consider a product map $\Phi = \phi_1 \times \phi_2$, where ϕ_1 and ϕ_2 are as previously. It is not ACL.

A method to produce sublin-q.s. homeos (II)

2nd step: Take the primitive $\phi : [0, 1] \rightarrow [0, 1]$ in the distributional sense.

ϕ is not absolutely continuous. The derivative is λ -a.e. 0. The modulus of continuity deviates sublinearly from that of a Lipschitz function:

$$\log |\phi(x) - \phi(y)| \leq \log |x - y| + \nu(\log |x - y|), \text{ sublinear } \nu.$$

3rd step: To get a sublinearly quasisymmetric homeomorphism of the square (torus), consider a product map $\Phi = \phi_1 \times \phi_2$, where ϕ_1 and ϕ_2 are as previously. It is not ACL.

Proposition

ϕ and Φ are sublinearly quasisymmetric. The asphericity distortions at scale s for ϕ and Φ are bounded by $(\sum_{n < \log_2 s} \epsilon_n)$ (in fact they are a.e. much lower).

INVARIANTS

Conformal dimensions

The **topological dimension**, the **conformal dimension** are invariant under quasimetric homeomorphisms.

Conformal dimensions

The **topological dimension**, the **conformal dimension** are invariant under quasimetric homeomorphisms.

$$\text{Cdim}(Z) = \inf \left\{ p > 0 : \text{mod}_p^{\{\tau_j\}}(\text{nonconstant curves in } Z) = 0 \right\}.$$

$\{\tau_j\}$ are asphericity parameters for packing measures; $\text{mod}_p^{\{\tau_j\}}$ are the combinatorial moduli. Can be sublinearized and **sometimes computed** using the diffusivity lemma from conf. dim theory.

Conformal dimensions

The **topological dimension**, the **conformal dimension** are invariant under quasisymmetric homeomorphisms.

$$\text{Cdim}(Z) = \inf \left\{ p > 0 : \text{mod}_p^{\{\tau_j\}}(\text{nonconstant curves in } Z) = 0 \right\}.$$

$\{\tau_j\}$ are asphericity parameters for packing measures; $\text{mod}_p^{\{\tau_j\}}$ are the combinatorial moduli. Can be sublinearized and **sometimes computed** using the diffusivity lemma from conf. dim theory.

Self-similar space	Sublinear conformal dimension
\mathbb{R}^2 with scalar or unipotent δ	2
\mathbb{R}^2 with $\delta = \text{diag}(1, \mu)$	$1 + \mu$
General (nilpotent)	trace of the generator of dilations δ

Large-scale classifications of 3-dim Heintze spaces

Csq of Xie 2011 and of Carrasco Piaggio 2014

The 3-dimensionsal Heintze spaces Y and Y' are quasiisometric if and only if $\text{Isom}(Y)$ and $\text{Isom}(Y')$ are isomorphic.

Large-scale classifications of 3-dim Heintze spaces

Csq of Xie 2011 and of Carrasco Piaggio 2014

The 3-dimensional Heintze spaces Y and Y' are quasiisometric if and only if $\text{Isom}(Y)$ and $\text{Isom}(Y')$ are isomorphic.

Conformal dimension does not suffice. Two methods: Q -capacity, or ℓ^ϕ cohomology.

Large-scale classifications of 3-dim Heintze spaces

Csq of Xie 2011 and of Carrasco Piaggio 2014

The 3-dimensional Heintze spaces Y and Y' are quasiisometric if and only if $\text{Isom}(Y)$ and $\text{Isom}(Y')$ are isomorphic.

Conformal dimension does not suffice. Two methods: Q -capacity, or ℓ^ϕ cohomology.

With the sublinear conformal dimension

The 3-dimensional Heintze spaces Y and Y' are SBE if and only if

1. Either $\text{Isom}(Y)$ and $\text{Isom}(Y')$ are isomorphic.
2. Or Y and Y' are isometric to left-invariant Riemannian metrics on G and G' .

Ongoing work on higher-dimensional Heintze groups

Rk1. Conformal changes of metrics
preserve the Dirichlet **energy**
of functions.

Ongoing work on higher-dimensional Heintze groups

Rk1. Conformal changes of metrics
preserve the Dirichlet **energy**
of functions.

Rk2. Lower bound on moduli \leftrightarrow
lower bound on energies.

Ongoing work on higher-dimensional Heintze groups

Rk1. Conformal changes of metrics preserve the Dirichlet **energy** of functions.


Rk2. Lower bound on moduli \leftrightarrow
lower bound on energies.



Ongoing work on higher-dimensional Heintze groups

Rk1. Conformal changes of metrics preserve the Dirichlet **energy** of functions.

Rk2. Lower bound on moduli \leftrightarrow lower bound on energies.



A diagram of a rectangle with horizontal lines, representing a domain. The left vertical boundary is labeled $f \equiv 0$ and the right vertical boundary is labeled $f \equiv 1$. Below the rectangle, the equation $E_p^{\{\tau_j\}}(f) \geq \text{mod}_p^{\{\tau_j\}}(\Gamma)$ is written.


$$E_p^{\{\tau_j\}}(f) \geq \text{mod}_p^{\{\tau_j\}}(\Gamma)$$

One can define **functions of locally bounded p -energy** $\mathcal{W}_{\text{loc}}^p$; if φ is a sublin-q.s. homeo then $\mathcal{W}_{\text{loc}}^p(\Omega) \xrightarrow{\sim} \mathcal{W}_{\text{loc}}^p(\varphi^{-1}\Omega)$ for Ω an open in the target. $\mathcal{W}_{\text{loc}}^p(\Omega)$ is a Fréchet algebra whose **spectrum** is a quotient of Ω , the largest space of leaves that it separates.

Ongoing work on higher-dimensional Heintze groups

Rk1. Conformal changes of metrics preserve the Dirichlet **energy** of functions.

Rk2. Lower bound on moduli \leftrightarrow lower bound on energies.


$$E_p^{\{\tau_j\}}(f) \geq \text{mod}_p^{\{\tau_j\}}(\Gamma)$$

One can define **functions of locally bounded p -energy** $\mathcal{W}_{\text{loc}}^p$; if φ is a sublin-q.s. homeo then $\mathcal{W}_{\text{loc}}^p(\Omega) \xrightarrow{\sim} \mathcal{W}_{\text{loc}}^p(\varphi^{-1}\Omega)$ for Ω an open in the target. $\mathcal{W}_{\text{loc}}^p(\Omega)$ is a Fréchet algebra whose **spectrum** is a quotient of Ω , the largest space of leaves that it separates.

Work in progress

$\mathbf{R} \ltimes_{\delta_1} \mathbf{R}^n$ and $\mathbf{R} \ltimes_{\delta_2} \mathbf{R}^n$ with invariant metrics are SBE if and only if $\chi_{\delta_1} = \chi_{\delta_2}$.

General Heintze spaces

Theorem (Heintze 1974)

Every Heintze space is the left-invariant Riemannian metric of a $\mathbf{R} \ltimes_{\delta} N$ where N is a connected nilpotent Lie group and δ is a derivation of its Lie algebra with positive eigenvalues.

General Heintze spaces

Theorem (Heintze 1974)

Every Heintze space is the left-invariant Riemannian metric of a $\mathbf{R} \ltimes_{\delta} N$ where N is a connected nilpotent Lie group and δ is a derivation of its Lie algebra with positive eigenvalues.

Examples (different from already seen) include:

General Heintze spaces

Theorem (Heintze 1974)

Every Heintze space is the left-invariant Riemannian metric of a $\mathbf{R} \ltimes_{\delta} N$ where N is a connected nilpotent Lie group and δ is a derivation of its Lie algebra with positive eigenvalues.

Examples (different from already seen) include:

1. **Carnot type:** N is a Carnot gradable group and δ is a grading derivation.

General Heintze spaces

Theorem (Heintze 1974)

Every Heintze space is the left-invariant Riemannian metric of a $\mathbf{R} \ltimes_{\delta} N$ where N is a connected nilpotent Lie group and δ is a derivation of its Lie algebra with positive eigenvalues.

Examples (different from already seen) include:

1. **Carnot type:** N is a Carnot gradable group and δ is a grading derivation.
2. **Rank one symmetric space of non-compact type:** N is a (general) Heisenberg group and δ is a grading derivation.

General Heintze spaces

Theorem (Heintze 1974)

Every Heintze space is the left-invariant Riemannian metric of a $\mathbf{R} \ltimes_{\delta} N$ where N is a connected nilpotent Lie group and δ is a derivation of its Lie algebra with positive eigenvalues.

Examples (different from already seen) include:

1. **Carnot type:** N is a Carnot gradable group and δ is a grading derivation.
2. **Rank one symmetric space of non-compact type:** N is a (general) Heisenberg group and δ is a grading derivation.

Conjecture (Hamenstädt 1980s) — known for Carnot type (Pansu 1989)

If two Heintze spaces are quasiisometric then the underlying groups under purely real form are isomorphic.

Theorem (Cornulier 2008)

Let N be a nilpotent Lie group. Let δ_1, δ_2 be Heintze derivations with semisimple parts σ_1, σ_2 . If $G_1 = \mathbf{R} \ltimes_{\sigma_1} N$ and $G_2 = \mathbf{R} \ltimes_{\sigma_2} N$ are isomorphic then $G_1 = \mathbf{R} \ltimes_{\delta_1} N$ and $G_2 = \mathbf{R} \ltimes_{\delta_2} N$ with left-invariant metrics are SBE.

Cornulier reduction

Theorem (Cornulier 2008)

Let N be a nilpotent Lie group. Let δ_1, δ_2 be Heintze derivations with semisimple parts σ_1, σ_2 . If $G_1 = \mathbf{R} \ltimes_{\sigma_1} N$ and $G_2 = \mathbf{R} \ltimes_{\sigma_2} N$ are isomorphic then $G_1 = \mathbf{R} \ltimes_{\delta_1} N$ and $G_2 = \mathbf{R} \ltimes_{\delta_2} N$ with left-invariant metrics are SBE.

Replacement of the former question: do the converse hold?

Cornulier reduction

Theorem (Cornulier 2008)

Let N be a nilpotent Lie group. Let δ_1, δ_2 be Heintze derivations with semisimple parts σ_1, σ_2 . If $G_1 = \mathbf{R} \ltimes_{\sigma_1} N$ and $G_2 = \mathbf{R} \ltimes_{\sigma_2} N$ are isomorphic then $G_1 = \mathbf{R} \ltimes_{\delta_1} N$ and $G_2 = \mathbf{R} \ltimes_{\delta_2} N$ with left-invariant metrics are SBE.

Replacement of the former question: do the converse hold? Very partial answers:

1. Yes in the 3-dimensional case.


Theorem (Cornulier 2008)

Let N be a nilpotent Lie group. Let δ_1, δ_2 be Heintze derivations with semisimple parts σ_1, σ_2 . If $G_1 = \mathbf{R} \ltimes_{\sigma_1} N$ and $G_2 = \mathbf{R} \ltimes_{\sigma_2} N$ are isomorphic then $G_1 = \mathbf{R} \ltimes_{\delta_1} N$ and $G_2 = \mathbf{R} \ltimes_{\delta_2} N$ with left-invariant metrics are SBE.


Replacement of the former question: do the converse hold? Very partial answers:

1. Yes in the 3-dimensional case.
2. Ongoing work to a positive answer for abelian N .

Overview



Overview



Thank you for your attention!