

ECE 68000: MODERN AUTOMATIC CONTROL

Professor Stan Žak

Quadratic Lyapunov's Functions

Quadratic Forms

- $V = \mathbf{x}^T \mathbf{P} \mathbf{x}$, where $\mathbf{P} = \mathbf{P}^T$
- If \mathbf{P} not symmetric, need to symmetrize it
- First observe that because the transposition of a scalar equals itself, we have

$$\left(\mathbf{x}^T \mathbf{P} \mathbf{x} \right)^T = \mathbf{x}^T \mathbf{P}^T \mathbf{x} = \mathbf{x}^T \mathbf{P} \mathbf{x}$$

Symmetrizing Quadratic Form

□ Perform manipulations

$$\begin{aligned}x^T P x &= \frac{1}{2} x^T P x + \frac{1}{2} x^T P x \\&= \frac{1}{2} x^T P x + \frac{1}{2} x^T P^T x \\&= x^T \left(\frac{P + P^T}{2} \right) x\end{aligned}$$

□ Note that

$$\left(\frac{P + P^T}{2} \right)^T = \frac{P + P^T}{2}$$

Tests for Positive and Positive Semi-Definiteness of Quadratic Form

- $V = \mathbf{x}^T \mathbf{P} \mathbf{x}$, where $\mathbf{P} = \mathbf{P}^T$, is positive definite if and only if all eigenvalues of \mathbf{P} are positive
- $V = \mathbf{x}^T \mathbf{P} \mathbf{x}$, where $\mathbf{P} = \mathbf{P}^T$, is positive semi-definite if and only if all eigenvalues of \mathbf{P} are non-negative

Comments on the Eigenvalue Tests

- ❑ These tests are only good for the case when $\mathbf{P} = \mathbf{P}^T$. You must symmetrize \mathbf{P} before applying the above tests
- ❑ Other tests, the Sylvester's criteria, involve checking the signs of principal minors of \mathbf{P}

Negative Definite Quadratic Form

$V = \mathbf{x}^T \mathbf{P} \mathbf{x}$ is negative definite if and only if

$$-\mathbf{x}^T \mathbf{P} \mathbf{x} = \mathbf{x}^T (-\mathbf{P}) \mathbf{x}$$

is positive definite

Negative Semi-Definite Quadratic Form

$V = \mathbf{x}^T \mathbf{P} \mathbf{x}$ is negative semi-definite if and only if

$$-\mathbf{x}^T \mathbf{P} \mathbf{x} = \mathbf{x}^T (-\mathbf{P}) \mathbf{x}$$

is positive semi-definite

Example: Checking the Sign Definiteness of a Quadratic Form

- Is \mathbf{P} , equivalently, is the associated quadratic form, $V = \mathbf{x}^T \mathbf{P} \mathbf{x}$, pd, psd, nd, nsd, or neither?

$$\mathbf{P} = \begin{bmatrix} 2 & -6 \\ 0 & 2 \end{bmatrix}$$

- The associated quadratic form

$$V = \mathbf{x}^T \mathbf{P} \mathbf{x} = 2x_1^2 - 6x_1x_2 + 2x_2^2$$

Example: Symmetrizing the Underlying Matrix of the Quadratic Form

- ❑ Applying the eigenvalue test to the given quadratic form would seem to indicate that the quadratic form is pd, which turns out to be false
- ❑ Need to symmetrize the underlying matrix first and then can apply the eigenvalue test

Example: Symmetrized Matrix

- Symmetrizing manipulations

$$\begin{aligned}\frac{1}{2}(\mathbf{P} + \mathbf{P}^T) &= \frac{1}{2} \left(\begin{bmatrix} 2 & -6 \\ 0 & 2 \end{bmatrix} + \begin{bmatrix} 2 & 0 \\ -6 & 2 \end{bmatrix} \right) \\ &= \begin{bmatrix} 2 & -3 \\ -3 & 2 \end{bmatrix}\end{aligned}$$

- The eigenvalues of the symmetrized matrix are: 5 and -1
- The quadratic form is indefinite!

Example: Further Analysis

□ Direct check that the quadratic form is indefinite

□ Take $\mathbf{x} = [1 \ 0]^T$. Then

$$\mathbf{x}^T \mathbf{P} \mathbf{x} = \begin{bmatrix} 1 & 0 \end{bmatrix} \begin{bmatrix} 2 & -6 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \end{bmatrix} = 2 > 0$$

□ Take $\mathbf{x} = [1 \ 1]^T$. Then

$$\mathbf{x}^T \mathbf{P} \mathbf{x} = \begin{bmatrix} 1 & 1 \end{bmatrix} \begin{bmatrix} 2 & -6 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = -2 < 0$$

Stability Test for $\mathbf{x}_e = \mathbf{0}$ of $d\mathbf{x}/dt = \mathbf{A}\mathbf{x}$

- Let $V = \mathbf{x}^T \mathbf{P} \mathbf{x}$ where $\mathbf{P} = \mathbf{P}^T > 0$
- For V to be a Lyapunov function, that is, for $\mathbf{x}_e = \mathbf{0}$ to be a.s.,

$$\dot{V}(\mathbf{x}(t)) < 0$$

- Evaluate the time derivative of V on the solution of the system $d\mathbf{x}/dt = \mathbf{A}\mathbf{x}$ --- Lyapunov derivative

Lyapunov Derivative for $\dot{\mathbf{x}} = \mathbf{A}\mathbf{x}$

- Note that $V(\mathbf{x}(t)) = \mathbf{x}(t)^T \mathbf{P} \mathbf{x}(t)$
- Use the chain rule

$$\begin{aligned}\dot{V}(\mathbf{x}(t)) &= \dot{\mathbf{x}}^T \mathbf{P} \mathbf{x} + \mathbf{x}^T \mathbf{P} \dot{\mathbf{x}} \\ &= \mathbf{x}^T \mathbf{A}^T \mathbf{P} \mathbf{x} + \mathbf{x}^T \mathbf{P} \mathbf{A} \mathbf{x} \\ &= \mathbf{x}^T (\mathbf{A}^T \mathbf{P} + \mathbf{P} \mathbf{A}) \mathbf{x}\end{aligned}$$

- We used $\dot{\mathbf{x}}^T = \mathbf{x}^T \mathbf{A}^T$

Lyapunov Matrix Equation

□ Denote

$$A^T P + P A = -Q$$

□ Then the Lyapunov derivative can be represented as

$$\dot{V} = \frac{d}{dt}V = -x^T Q x$$

where

$$Q = Q^T > 0$$

Terms to Our Vocabulary

- Theorem---a major result of independent interest
- Lemma---an auxiliary result that is used as a stepping stone toward a theorem
- Corollary---a direct consequence of a theorem, or even a lemma

Lyapunov's Theorem

The real matrix \mathbf{A} is a.s., that is, all eigenvalues of \mathbf{A} have negative real parts if and only if for any $\mathbf{Q} = \mathbf{Q}^T > 0$ the solution \mathbf{P} of the continuous matrix Lyapunov equation

$$\mathbf{A}^T \mathbf{P} + \mathbf{P} \mathbf{A} = -\mathbf{Q}$$

is (symmetric) positive definite

How Do We Use the Lyapunov Theorem?

- ❑ Select an arbitrary symmetric positive definite \mathbf{Q} , for example, an identity matrix, \mathbf{I}_n
- ❑ Solve the Lyapunov equation for $\mathbf{P} = \mathbf{P}^T$
- ❑ If \mathbf{P} is positive definite, the matrix \mathbf{A} is a.s. If \mathbf{P} is not p.d. then \mathbf{A} is not a.s.

How NOT to Use the Lyapunov Theorem

- ❑ It would be no use choosing **P** to be positive definite and then calculating **Q**
- ❑ For unless **Q** turns out to be positive definite, nothing can be said about a.s. of **A** from the Lyapunov equation

Example: How NOT to Use the Lyapunov Theorem

- Consider an a.s. matrix

$$A = \begin{bmatrix} -1 & 3 \\ 0 & -1 \end{bmatrix}$$

- Try

$$P = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

- Compute $Q = - (A^T P + P A)$

Example: Computing Q

$$\begin{aligned} Q &= -\left(A^T P + P A\right) \\ &= -\left(\begin{bmatrix} -1 & 0 \\ 3 & -1 \end{bmatrix} + \begin{bmatrix} -1 & 3 \\ 0 & -1 \end{bmatrix}\right) \\ &= \begin{bmatrix} 2 & -3 \\ -3 & 2 \end{bmatrix} \end{aligned}$$

The matrix Q is indefinite!---
recall the previous example

Solving the Continuous Matrix Lyapunov Equation Using MATLAB

- Use the MATLAB's command `lyap`

- Example:

$$A = \begin{bmatrix} -1 & 3 \\ 0 & -1 \end{bmatrix}$$

- $Q = I_2$

- $P = \text{lyap}(A', Q)$

$$P = \begin{bmatrix} 0.50 & 0.75 \\ 0.75 & 2.75 \end{bmatrix}$$

- Eigenvalues of P are positive: 0.2729 and 2.9771; P is positive definite

Benefits of the Lyapunov Theory

- ❑ Solution to differential equation are not needed to infer about stability properties of equilibrium state of interest



Lyapunov functions are useful in designing robust and adaptive controllers