

Harmony, Stability, and Identity

On the possibility of an intensional account
in proof-theoretic semantics

Alberto Naibo
(joint work with Luca Tranchini)

Université Paris 1 Panthéon-Sorbonne, IHPST (UMR 8590)

October 30 & November 1, 2017
PUC-Rio

Table of contents

Part I. Harmony and Stability

Part II. Stability and Identity

Table of contents

Part I. Harmony and Stability

Part II. Stability and Identity

Truth-theoretic *vs* proof-theoretic semantics

- ▶ Proof-theoretic semantics is usually conceived in opposition to truth-theoretic semantics.
- ▶ In truth-theoretic semantics, truth is considered as a primitive and non-analyzed notion, and meaning is then explained in terms of it.
- ▶ In proof-theoretic semantics, meaning is explained in terms of (our) inferential abilities, and truth is then explained in terms of proofs (where a proof is a concatenation of inferences).

Truth-theoretic *vs* proof-theoretic semantics

- ▶ Proof-theoretic semantics is usually conceived in opposition to truth-theoretic semantics.
- ▶ In truth-theoretic semantics, truth is considered as a primitive and non-analyzed notion, and meaning is then explained in terms of it.
- ▶ In proof-theoretic semantics, meaning is explained in terms of (our) inferential abilities, and truth is then explained in terms of proofs (where a proof is a concatenation of inferences).

Truth-theoretic *vs* proof-theoretic semantics

- ▶ Proof-theoretic semantics is usually conceived in opposition to truth-theoretic semantics.
- ▶ In truth-theoretic semantics, truth is considered as a primitive and non-analyzed notion, and meaning is then explained in terms of it.
- ▶ In proof-theoretic semantics, meaning is explained in terms of (our) inferential abilities, and truth is then explained in terms of proofs (where a proof is a concatenation of inferences).

Truth-theoretic *vs* proof-theoretic semantics

- In truth-theoretic semantics the interpretation of a sentence A corresponds to a truth-value: either the true or the false.

$$\llbracket A \rrbracket = \mathcal{B}, \text{ where } \mathcal{B} \in \{\text{T}, \text{F}\}$$

- In proof-theoretic semantics the interpretation of a sentence A corresponds to a set of proofs sharing a bunch of properties \mathcal{P} saying that these proofs possess some *common inferential structure* (e.g. canonicity or canonisability).

$$\llbracket A \rrbracket = \{\pi \mid \mathcal{P}(\pi)\}$$

- But what happens if the principles used for obtaining these properties \mathcal{P} are somehow mutually incompatible and lead to the identification all proofs of A ?

$$\llbracket A \rrbracket = \mathcal{B}, \text{ where } \mathcal{B} \in \{\mathfrak{P}_A, \emptyset\} \text{ and } \mathfrak{P}_A \text{ is the set of all proofs of } A$$

Truth-theoretic *vs* proof-theoretic semantics

- ▶ In truth-theoretic semantics the interpretation of a sentence A corresponds to a truth-value: either the true or the false.

$$\llbracket A \rrbracket = \mathcal{B}, \text{ where } \mathcal{B} \in \{\mathbf{T}, \mathbf{F}\}$$

- ▶ In proof-theoretic semantics the interpretation of a sentence A corresponds to a set of proofs sharing a bunch of properties \mathcal{P} saying that these proofs possess some *common inferential structure* (e.g. canonicity or canonisability).

$$\llbracket A \rrbracket = \{\pi \mid \mathcal{P}(\pi)\}$$

- ▶ But what happens if the principles used for obtaining these properties \mathcal{P} are somehow mutually incompatible and lead to the identification all proofs of A ?

$$\llbracket A \rrbracket = \mathcal{B}, \text{ where } \mathcal{B} \in \{\mathfrak{P}_A, \emptyset\} \text{ and } \mathfrak{P}_A \text{ is the set of all proofs of } A$$

Truth-theoretic *vs* proof-theoretic semantics

- ▶ In truth-theoretic semantics the interpretation of a sentence A corresponds to a truth-value: either the true or the false.

$$\llbracket A \rrbracket = \mathcal{B}, \text{ where } \mathcal{B} \in \{\mathbf{T}, \mathbf{F}\}$$

- ▶ In proof-theoretic semantics the interpretation of a sentence A corresponds to a set of proofs sharing a bunch of properties \mathcal{P} saying that these proofs possess some *common inferential structure* (e.g. canonicity or canonisability).

$$\llbracket A \rrbracket = \{\pi \mid \mathcal{P}(\pi)\}$$

- ▶ But what happens if the principles used for obtaining these properties \mathcal{P} are somehow mutually incompatible and lead to the identification all proofs of A ?

$$\llbracket A \rrbracket = \mathcal{B}, \text{ where } \mathcal{B} \in \{\mathfrak{P}_A, \emptyset\} \text{ and } \mathfrak{P}_A \text{ is the set of all proofs of } A$$

Truth-theoretic *vs* proof-theoretic semantics

- ▶ In truth-theoretic semantics the interpretation of a sentence A corresponds to a truth-value: either the true or the false.

$$\llbracket A \rrbracket = \mathcal{B}, \text{ where } \mathcal{B} \in \{\mathbf{T}, \mathbf{F}\}$$

- ▶ In proof-theoretic semantics the interpretation of a sentence A corresponds to a set of proofs sharing a bunch of properties \mathcal{P} saying that these proofs possess some *common inferential structure* (e.g. canonicity or canonisability).

$$\llbracket A \rrbracket = \{\pi \mid \mathcal{P}(\pi)\}$$

- ▶ But what happens if the principles used for obtaining these properties \mathcal{P} are somehow mutually incompatible and lead to the identification all proofs of A ?

$$\llbracket A \rrbracket = \mathcal{B}, \text{ where } \mathcal{B} \in \{\mathfrak{P}_A, \emptyset\} \text{ and } \mathfrak{P}_A \text{ is the set of all proofs of } A$$

Truth-theoretic *vs* proof-theoretic semantics

- ▶ In truth-theoretic semantics the interpretation of a sentence A corresponds to a truth-value: either the true or the false.

$$\llbracket A \rrbracket = \mathcal{B}, \text{ where } \mathcal{B} \in \{\mathbf{T}, \mathbf{F}\}$$

- ▶ In proof-theoretic semantics the interpretation of a sentence A corresponds to a set of proofs sharing a bunch of properties \mathcal{P} saying that these proofs possess some *common inferential structure* (e.g. canonicity or canonisability).

$$\llbracket A \rrbracket = \{\pi \mid \mathcal{P}(\pi)\}$$

- ▶ But what happens if the principles used for obtaining these properties \mathcal{P} are somehow mutually incompatible and lead to the identification all proofs of A ?

$$\llbracket A \rrbracket = \mathcal{B}, \text{ where } \mathcal{B} \in \{\mathfrak{P}_A, \emptyset\} \text{ and } \mathfrak{P}_A \text{ is the set of all proofs of } A$$

Truth-theoretic *vs* proof-theoretic semantics

- ▶ In truth-theoretic semantics the interpretation of a sentence A corresponds to a truth-value: either the true or the false.

$$\llbracket A \rrbracket = \mathcal{B}, \text{ where } \mathcal{B} \in \{\mathbf{T}, \mathbf{F}\}$$

- ▶ In proof-theoretic semantics the interpretation of a sentence A corresponds to a set of proofs sharing a bunch of properties \mathcal{P} saying that these proofs possess some *common inferential structure* (e.g. canonicity or canonisability).

$$\llbracket A \rrbracket = \{\pi \mid \mathcal{P}(\pi)\}$$

- ▶ But what happens if the principles used for obtaining these properties \mathcal{P} are somehow mutually incompatible and lead to the identification all proofs of A ?

$$\llbracket A \rrbracket = \mathcal{B}, \text{ where } \mathcal{B} \in \{\mathfrak{P}_A, \emptyset\} \text{ and } \mathfrak{P}_A \text{ is the set of all proofs of } A$$

Proof-theoretic semantics *vs* provability

- ▶ In this case, proof-theoretic semantics does not necessarily boil down to truth-theoretic semantics.
- ▶ However, the meaning of a certain sentence A would be explained without any reference to the inferential structure of its proofs: it is sufficient that *there exists* a proof of A , but it is not necessary to know how this proof is built.
- ▶ In other words, what counts for the meaning of a sentence A is not the way in which it is proved, but just the fact it is *provable*.
- ▶ Provability is not enough for having a genuine proof-theoretic semantics:
 - ▶ a theory based on provability is *extensional*: what counts is just the *relation* between premisses and conclusions;
 - ▶ a theory based on proofs is *intentional*: what counts is the way in which premisses and conclusions are related (in principle, there could be different proofs relating the same premisses and conclusions).

Proof-theoretic semantics *vs* provability

- ▶ In this case, proof-theoretic semantics does not necessarily boil down to truth-theoretic semantics.
- ▶ However, the meaning of a certain sentence A would be explained without any reference to the inferential structure of its proofs: it is sufficient that *there exists* a proof of A , but it is not necessary to know how this proof is built.
- ▶ In other words, what counts for the meaning of a sentence A is not the way in which it is proved, but just the fact it is *provable*.
- ▶ Provability is not enough for having a genuine proof-theoretic semantics:
 - ▶ a theory based on provability is *extensional*: what counts is just the *relation* between premisses and conclusions;
 - ▶ a theory based on proofs is *intentional*: what counts is the way in which premisses and conclusions are related (in principle, there could be different proofs relating the same premisses and conclusions).

Proof-theoretic semantics *vs* provability

- ▶ In this case, proof-theoretic semantics does not necessarily boil down to truth-theoretic semantics.
- ▶ However, the meaning of a certain sentence A would be explained without any reference to the inferential structure of its proofs: it is sufficient that *there exists* a proof of A , but it is not necessary to know how this proof is built.
- ▶ In other words, what counts for the meaning of a sentence A is not the way in which it is proved, but just the fact it is *provable*.
- ▶ Provability is not enough for having a genuine proof-theoretic semantics:
 - ▶ a theory based on provability is *extensional*: what counts is just the *relation* between premisses and conclusions;
 - ▶ a theory based on proofs is *intentional*: what counts is the way in which premisses and conclusions are related (in principle, there could be different proofs relating the same premisses and conclusions).

Proof-theoretic semantics *vs* provability

- ▶ In this case, proof-theoretic semantics does not necessarily boil down to truth-theoretic semantics.
- ▶ However, the meaning of a certain sentence A would be explained without any reference to the inferential structure of its proofs: it is sufficient that *there exists* a proof of A , but it is not necessary to know how this proof is built.
- ▶ In other words, what counts for the meaning of a sentence A is not the way in which it is proved, but just the fact it is *provable*.
- ▶ Provability is not enough for having a genuine proof-theoretic semantics:
 - ▶ a theory based on provability is *extensional*: what counts is just the *relation* between premisses and conclusions;
 - ▶ a theory based on proofs is *intentional*: what counts is the way in which premisses and conclusions are related (in principle, there could be different proofs relating the same premisses and conclusions).

Proof-theoretic semantics *vs* provability

- ▶ In this case, proof-theoretic semantics does not necessarily boil down to truth-theoretic semantics.
- ▶ However, the meaning of a certain sentence A would be explained without any reference to the inferential structure of its proofs: it is sufficient that *there exists* a proof of A , but it is not necessary to know how this proof is built.
- ▶ In other words, what counts for the meaning of a sentence A is not the way in which it is proved, but just the fact it is *provable*.
- ▶ Provability is not enough for having a genuine proof-theoretic semantics:
 - ▶ a theory based on provability is *extensional*: what counts is just the *relation* between premisses and conclusions;
 - ▶ a theory based on proofs is *intentional*: what counts is the way in which premisses and conclusions are related (in principle, there could be different proofs relating the same premisses and conclusions).

Proof-theoretic semantics *vs* provability

- ▶ In this case, proof-theoretic semantics does not necessarily boil down to truth-theoretic semantics.
- ▶ However, the meaning of a certain sentence A would be explained without any reference to the inferential structure of its proofs: it is sufficient that *there exists* a proof of A , but it is not necessary to know how this proof is built.
- ▶ In other words, what counts for the meaning of a sentence A is not the way in which it is proved, but just the fact it is *provable*.
- ▶ Provability is not enough for having a genuine proof-theoretic semantics:
 - ▶ a theory based on provability is *extensional*: what counts is just the *relation* between premisses and conclusions;
 - ▶ a theory based on proofs is *intentional*: what counts is the way in which premisses and conclusions are related (in principle, there could be different proofs relating the same premisses and conclusions).

Proof-theoretic semantics *vs* provability

- ▶ In this case, proof-theoretic semantics does not necessarily boil down to truth-theoretic semantics.
- ▶ However, the meaning of a certain sentence A would be explained without any reference to the inferential structure of its proofs: it is sufficient that *there exists* a proof of A , but it is not necessary to know how this proof is built.
- ▶ In other words, what counts for the meaning of a sentence A is not the way in which it is proved, but just the fact it is *provable*.
- ▶ Provability is not enough for having a genuine proof-theoretic semantics:
 - ▶ a theory based on provability is *extensional*: what counts is just the *relation* between premisses and conclusions;
 - ▶ a theory based on proofs is *intentional*: what counts is the way in which premisses and conclusions are related (in principle, there could be different proofs relating the same premisses and conclusions).

Proof-theoretic semantics *vs* provability

- ▶ In this case, proof-theoretic semantics does not necessarily boil down to truth-theoretic semantics.
- ▶ However, the meaning of a certain sentence A would be explained without any reference to the inferential structure of its proofs: it is sufficient that *there exists* a proof of A , but it is not necessary to know how this proof is built.
- ▶ In other words, what counts for the meaning of a sentence A is not the way in which it is proved, but just the fact it is *provable*.
- ▶ Provability is not enough for having a genuine proof-theoretic semantics:
 - ▶ a theory based on provability is *extensional*: what counts is just the *relation* between premisses and conclusions;
 - ▶ a theory based on proofs is *intentional*: what counts is the way in which premisses and conclusions are related (in principle, there could be different proofs relating the same premisses and conclusions).

Proof-theoretic semantics *vs* provability

- ▶ In this case, proof-theoretic semantics does not necessarily boil down to truth-theoretic semantics.
- ▶ However, the meaning of a certain sentence A would be explained without any reference to the inferential structure of its proofs: it is sufficient that *there exists* a proof of A , but it is not necessary to know how this proof is built.
- ▶ In other words, what counts for the meaning of a sentence A is not the way in which it is proved, but just the fact it is *provable*.
- ▶ Provability is not enough for having a genuine proof-theoretic semantics:
 - ▶ a theory based on provability is *extensional*: what counts is just the *relation* between premisses and conclusions;
 - ▶ a theory based on proofs is *intentional*: what counts is the way in which premisses and conclusions are related (in principle, there could be different proofs relating the same premisses and conclusions).

Two principles under analysis: harmony and stability

- ▶ The proof-theoretic debate is centered around the study of the principles used for obtaining the properties \mathcal{P} (harmony, conservativeness, uniqueness, stability, deducibility of identicals, etc.).
- ▶ However, little is said about whether these principles entail the collapse of certain sets of proofs or not.
- ▶ Our aim is to focus the attention on two of these principles – *harmony* and *stability* – and to show that even if they are very natural to demand, they could lead to the collapse of the set of proofs of certain specific sentences.
- ▶ We will not reduce our analysis to sentences constructed with propositional operators; we will also look at first-order predicates, like the identity predicate.

Two principles under analysis: harmony and stability

- ▶ The proof-theoretic debate is centered around the study of the principles used for obtaining the properties \mathcal{P} (harmony, conservativeness, uniqueness, stability, deducibility of identicals, etc.).
- ▶ However, little is said about whether these principles entail the collapse of certain sets of proofs or not.
- ▶ Our aim is to focus the attention on two of these principles – *harmony* and *stability* – and to show that even if they are very natural to demand, they could lead to the collapse of the set of proofs of certain specific sentences.
- ▶ We will not reduce our analysis to sentences constructed with propositional operators; we will also look at first-order predicates, like the identity predicate.

Two principles under analysis: harmony and stability

- ▶ The proof-theoretic debate is centered around the study of the principles used for obtaining the properties \mathcal{P} (harmony, conservativeness, uniqueness, stability, deducibility of identicals, etc.).
- ▶ However, little is said about whether these principles entail the collapse of certain sets of proofs or not.
- ▶ Our aim is to focus the attention on two of these principles – *harmony* and *stability* – and to show that even if they are very natural to demand, they could lead to the collapse of the set of proofs of certain specific sentences.
- ▶ We will not reduce our analysis to sentences constructed with propositional operators; we will also look at first-order predicates, like the identity predicate.

Two principles under analysis: harmony and stability

- ▶ The proof-theoretic debate is centered around the study of the principles used for obtaining the properties \mathcal{P} (harmony, conservativeness, uniqueness, stability, deducibility of identicals, etc.).
- ▶ However, little is said about whether these principles entail the collapse of certain sets of proofs or not.
- ▶ Our aim is to focus the attention on two of these principles – *harmony* and *stability* – and to show that even if they are very natural to demand, they could lead to the collapse of the set of proofs of certain specific sentences.
- ▶ We will not reduce our analysis to sentences constructed with propositional operators; we will also look at first-order predicates, like the identity predicate.

Harmony and the inversion principle

- ▶ Harmony is usually understood as the idea that there should be a balance between the conditions for the (correct) assertion of a certain (complex) sentence and what can be drawn from this assertion.
- ▶ Usually, this is understood in terms of Prawitz's *inversion principle*, according to which,
[...] by an application of an elimination rule one essentially only restores what had been established if the major premiss of the application was inferred by an application of an introduction rule. (Prawitz 1965: 33)

Harmony and the inversion principle

- ▶ Harmony is usually understood as the idea that there should be a balance between the conditions for the (correct) assertion of a certain (complex) sentence and what can be drawn from this assertion.
- ▶ Usually, this is understood in terms of Prawitz's *inversion principle*, according to which,

[...] by an application of an elimination rule one essentially only restores what had been established if the major premiss of the application was inferred by an application of an introduction rule. (Prawitz 1965: 33)

Harmony and the inversion principle

- ▶ Harmony is usually understood as the idea that there should be a balance between the conditions for the (correct) assertion of a certain (complex) sentence and what can be drawn from this assertion.
- ▶ Usually, this is understood in terms of Prawitz's *inversion principle*, according to which,
[...] by an application of an elimination rule one essentially only restores what had been established if the major premiss of the application was inferred by an application of an introduction rule. (Prawitz 1965: 33)

Harmonious rules

- ▶ The inference rules of standard *intuitionistic* connectives are harmonious.

- ▶ Consider, for example, the rules of conjunction:

$$\frac{A_1 \quad A_2}{A_1 \wedge A_2} \wedge_I \qquad \frac{A_1 \wedge A_2}{A_i} \wedge_{E_i} \ (i \in \{1, 2\})$$

- ▶ These rules satisfy Prawitz's inversion principle, since a *hillock*, or (local) *complexity peak*, of the following kind

$$\frac{\frac{\mathcal{D}_1 \quad \mathcal{D}_2}{A_1 \quad A_2} \wedge_I}{\frac{A_1 \wedge A_2}{A_i} \wedge_{E_i}} \text{ reduces to } \frac{\mathcal{D}_i}{A_i}$$

Harmonious rules

- ▶ The inference rules of standard *intuitionistic* connectives are harmonious.
- ▶ Consider, for example, the rules of conjunction:

$$\frac{A_1 \quad A_2}{A_1 \wedge A_2} \wedge_I \qquad \frac{A_1 \wedge A_2}{A_i} \wedge_{E_i} \ (i \in \{1, 2\})$$

- ▶ These rules satisfy Prawitz's inversion principle, since a *hillock*, or (local) *complexity peak*, of the following kind

$$\frac{\frac{\mathcal{D}_1 \quad \mathcal{D}_2}{A_1 \quad A_2} \wedge_I}{\frac{A_1 \wedge A_2}{A_i} \wedge_{E_i}} \text{ reduces to } \frac{\mathcal{D}_i}{A_i}$$

Harmonious rules

- ▶ The inference rules of standard *intuitionistic* connectives are harmonious.
- ▶ Consider, for example, the rules of conjunction:

$$\frac{A_1 \quad A_2}{A_1 \wedge A_2} \wedge_I \qquad \frac{A_1 \wedge A_2}{A_i} \wedge_{E_i} \ (i \in \{1, 2\})$$

- ▶ These rules satisfy Prawitz's inversion principle, since a *hillock*, or (local) *complexity peak*, of the following kind

$$\frac{\frac{\mathcal{D}_1 \quad \mathcal{D}_2}{A_1 \quad A_2} \wedge_I}{A_i} \wedge_{E_i} \qquad \text{reduces to} \qquad \frac{\mathcal{D}_i}{A_i}$$

Harmonious rules

- ▶ The inference rules of standard *intuitionistic* connectives are harmonious.
- ▶ Consider, for example, the rules of conjunction:

$$\frac{A_1 \quad A_2}{A_1 \wedge A_2} \wedge_I \qquad \frac{A_1 \wedge A_2}{A_i} \wedge_{E_i} \ (i \in \{1, 2\})$$

- ▶ These rules satisfy Prawitz's inversion principle, since a *hillock*, or (local) *complexity peak*, of the following kind

$$\frac{\frac{\mathcal{D}_1 \quad \mathcal{D}_2}{A_1 \quad A_2} \wedge_I}{\frac{A_1 \wedge A_2}{A_i} \wedge_{E_i}} \text{ reduces to } \frac{\mathcal{D}_i}{A_i}$$

Not harmonious rules

- ▶ A typical example of inference rules which fail to satisfy harmony are those of **tonk** (Prior 1960):

$$\frac{A_1}{A_1 \text{ tonk } A_2} \text{tonk}_I \qquad \frac{A_1 \text{ tonk } A_2}{A_2} \text{tonk}_E$$

- ▶ These rules do not satisfy Prawitz's inversion principle since they do not allow one to reduce the following hillock, or (local) complexity peak:

$$\frac{\frac{\mathcal{D}}{A_1} \text{tonk}_I}{A_1 \text{ tonk } A_2} \text{tonk}_E$$

Not harmonious rules

- ▶ A typical example of inference rules which fail to satisfy harmony are those of **tonk** (Prior 1960):

$$\frac{A_1}{A_1 \text{ tonk } A_2} \text{tonk}_I \qquad \frac{A_1 \text{ tonk } A_2}{A_2} \text{tonk}_E$$

- ▶ These rules do not satisfy Prawitz's inversion principle since they do not allow one to reduce the following hillock, or (local) complexity peak:

$$\frac{\mathcal{D} \quad \frac{A_1}{A_1 \text{ tonk } A_2} \text{tonk}_I}{A_2} \text{tonk}_E$$

No more and no less conditions

- ▶ However, as Dummett remarks, « harmony is an excessively modest demand » (1991: 287).

No more and no less conditions

- ▶ However, as Dummett remarks, « harmony is an excessively modest demand » (1991: 287).
- ▶ Harmony guarantees that what can be drawn from a complex sentence A by means of the elimination rules for its main connective \dagger is *no more* than what can be drawn from the premisses of the introduction rules for \dagger (used in order to obtain A).

No more and no less conditions

- ▶ However, as Dummett remarks, « harmony is an excessively modest demand » (1991: 287).
- ▶ Harmony guarantees that what can be drawn from a complex sentence A by means of the elimination rules for its main connective \dagger is *no more* than what has to be established in order to infer A using the introduction rules for \dagger .

No more and no less conditions

- ▶ However, as Dummett remarks, « harmony is an excessively modest demand » (1991: 287).
- ▶ Harmony guarantees that what can be drawn from a complex sentence A by means of the elimination rules for its main connective \dagger is *no more* than what has to be established in order to infer A using the introduction rules for \dagger .
- ▶ In order to fully capture the idea that the rules of inference *define* the meaning of linguistic expressions, it is natural to ask also to fulfill the *no less* condition.

No more and no less conditions

- ▶ However, as Dummett remarks, « harmony is an excessively modest demand » (1991: 287).
- ▶ Harmony guarantees that what can be drawn from a complex sentence A by means of the elimination rules for its main connective \dagger is *no more* than what has to be established in order to infer A using the introduction rules for \dagger .
- ▶ In order to fully capture the idea that the rules of inference *define* the meaning of linguistic expressions, it is natural to ask also to fulfill the *no less* condition.
- ▶ A principle that has been proposed in order to capture this condition is the principle of *deducibility of identicals* (see Hacking 1979, Pfenning & Davies 2001, Francez & Dyckhoff 2012, Naibo & Petrolo 2015).

Deducibility of identicals: An example

- ▶ Deducibility of identicals says that given a complex sentence A , it is possible to derive A from itself using only the inference rules of the principal connective of A .
- ▶ The idea is that what can be drawn from the elimination rules of a certain connective \dagger is *enough* in order to re-establish the conclusion of the introduction rules of \dagger .
- ▶ The inference rules of standard intuitionistic connectives satisfy the deducibility of identicals. Consider, for example, the rules of conjunction. We have that:

$$\begin{array}{ccc} \mathcal{D} & & \mathcal{D} \\ A_1 \wedge A_2 & \text{expands to} & \frac{\frac{A_1 \wedge A_2}{A_1} \wedge_{E_1} \quad \frac{A_1 \wedge A_2}{A_2} \wedge_{E_2}}{A_1 \wedge A_2} \wedge_I \end{array}$$

generating a *valley*, or (local) *complexity riff*.

Deducibility of identicals: An example

- ▶ Deducibility of identicals says that given a complex sentence A , it is possible to derive A from itself using only the inference rules of the principal connective of A .
- ▶ The idea is that what can be drawn from the elimination rules of a certain connective \dagger is *enough* in order to re-establish the conclusion of the introduction rules of \dagger .
- ▶ The inference rules of standard intuitionistic connectives satisfy the deducibility of identicals. Consider, for example, the rules of conjunction. We have that:

$$\begin{array}{c} \mathcal{D} \\ A_1 \wedge A_2 \end{array} \quad \text{expands to} \quad \frac{\frac{\mathcal{D}}{A_1 \wedge A_2} \wedge_{E_1} \quad \frac{\frac{\mathcal{D}}{A_1 \wedge A_2} \wedge_{E_2}}{A_2} \wedge_I}{A_1 \wedge A_2}$$

generating a *valley*, or (local) *complexity riff*.

Deducibility of identicals: An example

- ▶ Deducibility of identicals says that given a complex sentence A , it is possible to derive A from itself using only the inference rules of the principal connective of A .
- ▶ The idea is that what can be drawn from the elimination rules of a certain connective \dagger is *enough* in order to re-establish the conclusion of the introduction rules of \dagger .
- ▶ The inference rules of standard intuitionistic connectives satisfy the deducibility of identicals. Consider, for example, the rules of conjunction. We have that:

$$\begin{array}{c} \mathcal{D} \\ A_1 \wedge A_2 \end{array} \quad \text{expands to} \quad \frac{\frac{\mathcal{D}}{A_1 \wedge A_2} \wedge_{E_1} \quad \frac{\frac{\mathcal{D}}{A_1 \wedge A_2} \wedge_{E_2}}{A_2} \wedge_I}{A_1 \wedge A_2}$$

generating a *valley*, or (local) *complexity riff*.

Deducibility of identicals: An example

- ▶ Deducibility of identicals says that given a complex sentence A , it is possible to derive A from itself using only the inference rules of the principal connective of A .
- ▶ The idea is that what can be drawn from the elimination rules of a certain connective \dagger is *enough* in order to re-establish the conclusion of the introduction rules of \dagger .
- ▶ The inference rules of standard intuitionistic connectives satisfy the deducibility of identicals. Consider, for example, the rules of conjunction. We have that:

$$\begin{array}{c} \mathcal{D} \\ A_1 \wedge A_2 \end{array} \quad \text{expands to} \quad \frac{\frac{\mathcal{D}}{A_1 \wedge A_2} \wedge_{E_1} \quad \frac{\frac{\mathcal{D}}{A_1 \wedge A_2} \wedge_{E_2}}{A_2} \wedge_I}{A_1 \wedge A_2}$$

generating a *valley*, or (local) *complexity riff*.

Deducibility of identicals: An example

- ▶ Deducibility of identicals says that given a complex sentence A , it is possible to derive A from itself using only the inference rules of the principal connective of A .
- ▶ The idea is that what can be drawn from the elimination rules of a certain connective \dagger is *enough* in order to re-establish the conclusion of the introduction rules of \dagger .
- ▶ The inference rules of standard intuitionistic connectives satisfy the deducibility of identicals. Consider, for example, the rules of conjunction. We have that:

$$\begin{array}{c} \mathcal{D} \\ A_1 \wedge A_2 \end{array} \quad \text{expands to} \quad \frac{\frac{\frac{\mathcal{D}}{A_1 \wedge A_2} \wedge_{E_1}}{A_1} \quad \frac{\frac{\frac{\mathcal{D}}{A_1 \wedge A_2} \wedge_{E_2}}{A_2} \wedge_I}{A_1 \wedge A_2}}$$

generating a *valley*, or (local) *complexity riff*.

Deducibility of identicals: An example

- ▶ Deducibility of identicals says that given a complex sentence A , it is possible to derive A from itself using only the inference rules of the principal connective of A .
- ▶ The idea is that what can be drawn from the elimination rules of a certain connective \dagger is *enough* in order to re-establish the conclusion of the introduction rules of \dagger .
- ▶ The inference rules of standard intuitionistic connectives satisfy the deducibility of identicals. Consider, for example, the rules of conjunction. We have that:

$$\begin{array}{ccc} \mathcal{D} & & \mathcal{D} \\ A_1 \wedge A_2 & \text{expands to} & \frac{\frac{A_1 \wedge A_2}{A_1} \wedge_{E_1} \quad \frac{A_1 \wedge A_2}{A_2} \wedge_{E_2}}{A_1 \wedge A_2} \wedge_I \end{array}$$

generating a *valley*, or (local) *complexity riff*.

Deducibility of identicals: An example

- ▶ Deducibility of identicals says that given a complex sentence A , it is possible to derive A from itself using only the inference rules of the principal connective of A .
- ▶ The idea is that what can be drawn from the elimination rules of a certain connective \dagger is *enough* in order to re-establish the conclusion of the introduction rules of \dagger .
- ▶ The inference rules of standard intuitionistic connectives satisfy the deducibility of identicals. Consider, for example, the rules of conjunction. We have that:

$$\begin{array}{ccc} \mathcal{D} & & \mathcal{D} \\ A_1 \wedge A_2 & \text{expands to} & \frac{\frac{A_1 \wedge A_2}{A_1} \wedge_{E_1} \quad \frac{A_1 \wedge A_2}{A_2} \wedge_{E_2}}{A_1 \wedge A_2} \wedge_I \end{array}$$

generating a *valley*, or (local) *complexity riff*.

Deducibility of identicals: A counter-example

- Consider a binary connective \star having the following inference rules:

$$\frac{A \quad B}{A \star B} \star_I \qquad \frac{A \star B \quad A}{B} \star_E$$

- These rules satisfy harmony, since the hillock, or (local) complexity peak,

$$\frac{\frac{\mathcal{D}_1 \quad \mathcal{D}_2}{A \quad B} \star_I \quad \mathcal{D}_3}{B} \star_E \quad \text{reduces to} \quad \frac{\mathcal{D}_2}{B}$$

Deducibility of identicals: A counter-example

- Consider a binary connective \star having the following inference rules:

$$\frac{A \quad B}{A \star B} \star_I \qquad \frac{A \star B \quad A}{B} \star_E$$

- These rules satisfy harmony, since the hillock, or (local) complexity peak,

$$\frac{\frac{\mathcal{D}_1 \quad \mathcal{D}_2}{A \quad B} \star_I \quad \mathcal{D}_3}{B} \star_E \quad \text{reduces to} \quad \mathcal{D}_2$$

Deducibility of identicals: A counter-example

- Consider a binary connective \star having the following inference rules:

$$\frac{A \quad B}{A \star B} \star_I \qquad \frac{A \star B \quad A}{B} \star_E$$

- These rules satisfy harmony, since the hillock, or (local) complexity peak,

$$\frac{\frac{\mathcal{D}_1 \quad A}{A} \quad \frac{\mathcal{D}_2 \quad B}{B} \star_I \quad \frac{\mathcal{D}_3 \quad A}{A} \star_E}{B} \text{ reduces to } \frac{\mathcal{D}_2 \quad B}{B}$$

Deducibility of identicals: A counter-example

- Consider a binary connective \star having the following inference rules:

$$\frac{A \quad B}{A \star B} \star_I \qquad \frac{A \star B \quad A}{B} \star_E$$

- These rules satisfy harmony, since the hillock, or (local) complexity peak,

$$\frac{\frac{\mathcal{D}_1 \quad A}{A} \quad \frac{\mathcal{D}_2 \quad B}{B} \star_I \quad \frac{\mathcal{D}_3 \quad A}{A} \star_E}{B} \text{ reduces to } \frac{\mathcal{D}_2 \quad B}{B}$$

- However, they do not satisfy the deducibility of identicals, since from

$$\frac{\mathcal{D} \quad A \star B}{A \star B} \text{ we get } \frac{A \quad \frac{\frac{\mathcal{D} \quad A \star B}{A \star B} \quad A}{B} \star_E}{A \star B} \star_I$$

Deducibility of identicals: A counter-example

- Consider a binary connective \star having the following inference rules:

$$\frac{A \quad B}{A \star B} \star_I \qquad \frac{A \star B \quad A}{B} \star_E$$

- These rules satisfy harmony, since the hillock, or (local) complexity peak,

$$\frac{\frac{\mathcal{D}_1 \quad \mathcal{D}_2}{A \quad B} \star_I \quad \mathcal{D}_3}{B} \star_E \quad \text{reduces to} \quad \frac{\mathcal{D}_2}{B}$$

- However, they do not satisfy the deducibility of identicals, since from

$$\frac{\mathcal{D}}{A \star B} \quad \text{we get} \quad \frac{\frac{\mathcal{D}}{A \star B} \quad \frac{A}{A}}{A \star B} \star_I$$

(Note: In the original image, the A in the top right of the second fraction and the A in the bottom left of the second fraction are circled in red.)

But an extra-assumption A has been used.

Deducibility of identicals: A counter-example

- Consider a binary connective \star having the following inference rules:

$$\frac{A \quad B}{A \star B} \star_I \qquad \frac{A \star B \quad A}{B} \star_E$$

- These rules satisfy harmony, since the hillock, or (local) complexity peak,

$$\frac{\frac{\mathcal{D}_1 \quad \mathcal{D}_2}{A \quad B} \star_I \quad \mathcal{D}_3}{B} \star_E \quad \text{reduces to} \quad \frac{\mathcal{D}_2}{B}$$

- However, they do not satisfy the deducibility of identicals, since from

$$\frac{\mathcal{D}}{A \star B} \quad \text{we get} \quad \frac{\frac{\mathcal{D}}{A \star B} \quad \frac{\mathcal{D}}{A}}{A \star B} \star_I$$

(Note: In the original image, the A in the top right of the second fraction and the A in the bottom left of the second fraction are circled in red.)

This means that what can be drawn from the elimination rule of \star is less than what has to be established in order to infer $A \star B$ using the introduction rules of \star .

Univocity and stability

- ▶ However, deducibility of identicals is not yet enough in order to *univocally* determine the set of elimination rules associated to the introduction rules of a certain given connective.
- ▶ In order to solve this problem, Dummett asks for an additional principle, that of *stability*.

Univocity and stability

- ▶ However, deducibility of identicals is not yet enough in order to *univocally* determine the set of elimination rules associated to the introduction rules of a certain given connective.
- ▶ In order to solve this problem, Dummett asks for an additional principle, that of *stability*.

Univocity and stability

- ▶ However, deducibility of identicals is not yet enough in order to *univocally* determine the set of elimination rules associated to the introduction rules of a certain given connective.
- ▶ In order to solve this problem, Dummett asks for an additional principle, that of *stability*.
- ▶ Consider the rules for standard disjunction:

$$\frac{A_i}{A_1 \vee A_2} \vee_{I_i} \ (i \in \{1, 2\}) \qquad \frac{A_1 \vee A_2 \quad \begin{array}{cc} [A_1] & [A_2] \\ C & C \end{array}}{C} \vee_E$$

Univocity and stability

- ▶ However, deducibility of identicals is not yet enough in order to *univocally* determine the set of elimination rules associated to the introduction rules of a certain given connective.
- ▶ In order to solve this problem, Dummett asks for an additional principle, that of *stability*.
- ▶ Consider then the rules for *quantum disjunction*:

$$\frac{A_i}{A_1 \bar{\vee} A_2} \vee_{I_i} \ (i \in \{1, 2\}) \qquad \frac{A_1 \bar{\vee} A_2 \quad \frac{[A_1] \quad C}{C} \quad \frac{[A_2] \quad C}{C}}{C} \vee_E$$

Univocity and stability

- ▶ However, deducibility of identicals is not yet enough in order to *univocally* determine the set of elimination rules associated to the introduction rules of a certain given connective.
- ▶ In order to solve this problem, Dummett asks for an additional principle, that of *stability*.
- ▶ Consider then the rules for *quantum disjunction*:

$$\frac{A_i}{A_1 \bar{\vee} A_2} \nabla_{I_i} \ (i \in \{1, 2\}) \qquad \frac{A_1 \bar{\vee} A_2 \quad \frac{[A_1] \quad [A_2]}{C} \quad C}{C} \nabla_E$$

Differently from the standard disjunction, the elimination rule of the quantum disjunction can be applied only in certain conditions, i.e. when there is *no side assumptions* in the sub-derivations of the two minor premisses.

Standard disjunction and quantum disjunction: Analogies

- Standard disjunction satisfies both harmony and deducibility of identicals:

$$\frac{\frac{\mathcal{D}}{A_i} \vee_{I_i} \quad \frac{\frac{[A_1]^m}{\mathcal{D}_1} \quad \frac{[A_2]^n}{\mathcal{D}_2}}{C} \vee_E(m, n)}{C} \quad \text{reduces to} \quad \frac{\mathcal{D}}{[A_i]} \mathcal{D}_i C$$

and

$$\frac{\mathcal{D}}{A_1 \vee A_2} \quad \text{expands to} \quad \frac{\frac{\mathcal{D}}{A_1 \vee A_2} \quad \frac{\frac{A_1^m}{A_1 \vee A_2} \vee_{I_1}}{A_1 \vee A_2} \quad \frac{\frac{A_2^n}{A_1 \vee A_2} \vee_{I_2}}{A_1 \vee A_2} \vee_E(m, n)}{A_1 \vee A_2}$$

Standard disjunction and quantum disjunction: Analogies

- Also quantum disjunction satisfies both harmony and deducibility of identicals:

$$\frac{\frac{\mathcal{D}}{A_i} \quad \bar{\vee}_{I_i} \quad \frac{\frac{^m[A_1]}{\mathcal{D}_1} \quad C}{C} \quad \frac{\frac{^n[A_2]}{\mathcal{D}_2} \quad C}{C}}{C} \bar{\vee}_E(m, n) \quad \text{reduces to} \quad \frac{\mathcal{D}}{[A_i]} \quad \frac{\mathcal{D}_i}{C}$$

and

$$\frac{\mathcal{D}}{A_1 \bar{\vee} A_2} \quad \text{expands to} \quad \frac{\frac{\mathcal{D}}{A_1 \bar{\vee} A_2} \quad \frac{\frac{^m A_1}{A_1 \bar{\vee} A_2} \quad \bar{\vee}_{I_1}}{A_1 \bar{\vee} A_2} \quad \frac{\frac{^n A_2}{A_1 \bar{\vee} A_2} \quad \bar{\vee}_{I_2}}{A_1 \bar{\vee} A_2}}{A_1 \bar{\vee} A_2} \bar{\vee}_E(m, n)$$

Standard disjunction and quantum disjunction: Analogies

- Also quantum disjunction satisfies both harmony and deducibility of identicals:

$$\frac{\frac{\mathcal{D}}{A_i} \quad \bar{\vee}_{I_i} \quad \frac{\frac{^m[A_1]}{\mathcal{D}_1} \quad C}{C} \quad \frac{\frac{^n[A_2]}{\mathcal{D}_2} \quad C}{C} \quad \bar{\vee}_E(m, n)}{C} \quad \text{reduces to} \quad \frac{\mathcal{D}}{[A_i]} \quad \frac{\mathcal{D}_i}{C}$$

and

$$\frac{\mathcal{D}}{A_1 \bar{\vee} A_2} \quad \text{expands to} \quad \frac{\frac{\mathcal{D}}{A_1 \bar{\vee} A_2} \quad \frac{\frac{^m A_1}{A_1 \bar{\vee} A_2} \quad \bar{\vee}_{I_1}}{A_1 \bar{\vee} A_2} \quad \frac{\frac{^n A_2}{A_1 \bar{\vee} A_2} \quad \bar{\vee}_{I_2}}{A_1 \bar{\vee} A_2} \quad \bar{\vee}_E(m, n)}{A_1 \bar{\vee} A_2}$$

- Since standard disjunction and quantum disjunction have the same set of introduction rules, what we showed is that this set is compatible with two different elimination rules.

Standard disjunction and quantum disjunction: Analogies

- Also quantum disjunction satisfies both harmony and deducibility of identicals:

$$\frac{\frac{\mathcal{D}}{A_i} \quad \bar{\vee}_{I_i} \quad \frac{\frac{^m[A_1]}{\mathcal{D}_1} \quad C}{C} \quad \frac{\frac{^n[A_2]}{\mathcal{D}_2} \quad C}{C} \quad \bar{\vee}_E(m, n)}{C} \quad \text{reduces to} \quad \frac{\mathcal{D}}{[A_i]} \quad \bar{\vee}_i \quad C$$

and

$$\frac{\mathcal{D}}{A_1 \bar{\vee} A_2} \quad \text{expands to} \quad \frac{\mathcal{D}}{A_1 \bar{\vee} A_2} \quad \frac{\frac{^m A_1}{A_1 \bar{\vee} A_2} \quad \bar{\vee}_{I_1}}{A_1 \bar{\vee} A_2} \quad \frac{\frac{^n A_2}{A_1 \vee A_2} \quad \bar{\vee}_{I_2}}{\bar{\vee}_E(m, n)} \quad A_1 \bar{\vee} A_2$$

- Since standard disjunction and quantum disjunction have the same set of introduction rules, what we showed is that this set is compatible with two different elimination rules. This could give rise to ambiguities concerning the use of these rules.

Generalized expansion

- However, consider now the case in which the expansion of a disjunctive sentence is operated in the middle of a derivation, and not just at the conclusion of it:

$$\begin{array}{c} \mathcal{D} \\ [A_1 \vee A_2] \\ \mathcal{D}' \\ C \end{array} \quad \text{expands to} \quad \frac{\frac{\mathcal{D}}{A_1 \vee A_2} \quad \frac{\frac{A_1^m}{A_1 \vee A_2} \vee_{I_1} \quad \frac{A_2^n}{A_1 \vee A_2} \vee_{I_2}}{[A_1 \vee A_2]} \vee_E(m, n)}{\mathcal{D}' \quad C}$$

Generalized expansion

- ▶ However, consider now the case in which the expansion of a disjunctive sentence is operated in the middle of a derivation, and not just at the conclusion of it:

$$\begin{array}{c} \mathcal{D} \\ [A_1 \vee A_2] \\ \mathcal{D}' \\ C \end{array} \quad \text{expands to} \quad \frac{\frac{\mathcal{D}}{A_1 \vee A_2} \quad \frac{\frac{A_1^m}{A_1 \vee A_2} \vee_{I_1} \quad \frac{A_2^n}{A_1 \vee A_2} \vee_{I_2}}{[A_1 \vee A_2]} \vee_E(m, n)}{\mathcal{D}' \quad C}$$

the red highlighted part can be *permuted* upward in the sub-derivations of the minor premisses.

Generalized expansion

- ▶ However, consider now the case in which the expansion of a disjunctive sentence is operated in the middle of a derivation, and not just at the conclusion of it:

$$\begin{array}{c}
 \mathcal{D} \\
 [A_1 \vee A_2] \\
 \mathcal{D}' \\
 C
 \end{array}
 \quad \text{expands to} \quad
 \begin{array}{c}
 \mathcal{D} \\
 A_1 \vee A_2 \\
 \hline
 C
 \end{array}
 \quad \vee_E(m, n)$$

$\frac{\frac{m}{A_1} \quad \vee_{I_1} \quad \frac{n}{A_2} \quad \vee_{I_2}}{\frac{[A_1 \vee A_2] \quad \mathcal{D}' \quad C}{[A_1 \vee A_2] \quad \mathcal{D}' \quad C} \vee_{I_1} \vee_{I_2}}$

Generalized expansion

- ▶ However, consider now the case in which the expansion of a disjunctive sentence is operated in the middle of a derivation, and not just at the conclusion of it:

$$\begin{array}{ccc} \begin{array}{c} \mathcal{D} \\ [A_1 \vee A_2] \\ \mathcal{D}' \\ C \end{array} & \text{expands to} & \begin{array}{c} \mathcal{D} \qquad \mathcal{D}' \qquad \mathcal{D}' \\ A_1 \vee A_2 \quad C \quad C \\ \hline C \end{array} \end{array}$$

$\frac{\frac{A_1^m}{[A_1 \vee A_2]} \vee_{I_1} \quad \frac{A_2^n}{[A_1 \vee A_2]} \vee_{I_2}}{C} \vee_E(m, n)$

This can be considered as a form of *generalized expansion*.

A failure of generalized expansion

- This generalized expansion fails in the case of quantum disjunction:

$$\begin{array}{c} \mathcal{D} \\ [A_1 \bar{\vee} A_2] \\ \mathcal{D}' \\ C \end{array} \text{ does not expand to } \frac{\frac{\frac{\mathcal{D}}{A_1 \bar{\vee} A_2}}{\mathcal{D}'} \quad \frac{\frac{\frac{\mathcal{D}'}{C}}{A_1 \bar{\vee} A_2}}{\mathcal{D}'} \quad \frac{\frac{\frac{\mathcal{D}'}{C}}{A_1 \bar{\vee} A_2}}{\mathcal{D}'}}{\frac{\mathcal{D}'}{C}} \bar{\vee}_{I_1} \quad \frac{\frac{\frac{\mathcal{D}'}{C}}{A_1 \bar{\vee} A_2}}{\mathcal{D}'}}{\frac{\mathcal{D}'}{C}} \bar{\vee}_{I_2}}{\frac{\mathcal{D}'}{C}} \bar{\vee}_E(m, n)$$

since the sub-derivation \mathcal{D}' could contain side assumptions, and thus the application of the rule $\bar{\vee}_E$ would not be correct.

- The generalized expansion can be seen as a way of capturing Dummett's stability (Jacinto & Read 2016, Tranchini 2016).
- Moreover, the no less condition can be generalized: what can be drawn from a complex sentence A by means of the elimination rules for its main connective \dagger is no less than what can be drawn from the premisses of the introduction rules for \dagger (used in order to obtain A).

A failure of generalized expansion

- ▶ This generalized expansion fails in the case of quantum disjunction:

$$\begin{array}{c} \mathcal{D} \\ [A_1 \bar{\vee} A_2] \\ \mathcal{D}' \\ C \end{array} \text{ does not expand to } \frac{\frac{\frac{\mathcal{D}}{A_1 \bar{\vee} A_2}}{\frac{\frac{A_1^m}{[A_1 \bar{\vee} A_2]} \bar{\vee}_{I_1}}{\frac{A_2^n}{[A_1 \bar{\vee} A_2]} \bar{\vee}_{I_2}}} \mathcal{D}'}{C} \bar{\vee}_E(m, n)$$

since the sub-derivation \mathcal{D}' could contain side assumptions, and thus the application of the rule $\bar{\vee}_E$ would not be correct.

- ▶ The generalized expansion can be seen as a way of capturing Dummett's stability (Jacinto & Read 2016, Tranchini 2016).
- ▶ Moreover, the no less condition can be generalized:
what can be drawn from a complex sentence A by means of the elimination rules for its main connective \dagger is no less than what can be drawn from the premisses of the introduction rules for \dagger (used in order to obtain A).

A failure of generalized expansion

- ▶ This generalized expansion fails in the case of quantum disjunction:

$$\begin{array}{c} \mathcal{D} \\ [A_1 \bar{\vee} A_2] \\ \mathcal{D}' \\ C \end{array} \text{ does not expand to } \frac{\frac{\frac{\mathcal{D}}{A_1 \bar{\vee} A_2}}{\frac{\frac{\overline{A_1}^m}{[A_1 \bar{\vee} A_2]} \bar{\vee}_{I_1}}{\frac{\mathcal{D}'}{C}}} \quad \frac{\frac{\overline{A_2}^n}{[A_1 \bar{\vee} A_2]} \bar{\vee}_{I_2}}{\frac{\mathcal{D}'}{C}} \bar{\vee}_E(m, n)$$

since the sub-derivation \mathcal{D}' could contain side assumptions, and thus the application of the rule $\bar{\vee}_E$ would not be correct.

- ▶ The generalized expansion can be seen as a way of capturing Dummett's stability (Jacinto & Read 2016, Tranchini 2016).
- ▶ Moreover, the no less condition can be generalized:

what can be drawn from a complex sentence A by means of the elimination rules for its main connective \dagger is no less than what can be drawn from the premisses of the introduction rules for \dagger (used in order to obtain A).

A failure of generalized expansion

- ▶ This generalized expansion fails in the case of quantum disjunction:

$$\begin{array}{c} \mathcal{D} \\ [A_1 \bar{\vee} A_2] \\ \mathcal{D}' \\ C \end{array} \text{ does not expand to } \frac{\frac{\frac{\mathcal{D}}{A_1 \bar{\vee} A_2}}{\frac{\frac{A_1^m}{[A_1 \bar{\vee} A_2]} \bar{\vee}_{I_1}}{\frac{A_2^n}{[A_1 \bar{\vee} A_2]} \bar{\vee}_{I_2}}} \mathcal{D}'}{C} \bar{\vee}_E(m, n)$$

since the sub-derivation \mathcal{D}' could contain side assumptions, and thus the application of the rule $\bar{\vee}_E$ would not be correct.

- ▶ The generalized expansion can be seen as a way of capturing Dummett's stability (Jacinto & Read 2016, Tranchini 2016).
- ▶ Moreover, the no less condition can be generalized:
what can be drawn from a complex sentence A by means of the elimination rules for its main connective \dagger is no less than what can be drawn from the premisses of the introduction rules for \dagger (used in order to obtain A).

Generalized expansion and general elimination rules

- ▶ The generalized expansion works only in the case of general elimination rules, but not when they are *flattened* elimination rules *à la* Negri & von Plato (2001).
- ▶ Consider the flattened rules for implication:

$$\frac{[A] \quad B}{A \rightarrow B} \rightarrow_I \qquad \frac{A \rightarrow B \quad A \quad C}{C} \rightarrow_E$$

Generalized expansion and general elimination rules

- ▶ The generalized expansion works only in the case of general elimination rules, but not when they are *flattened* elimination rules *à la* Negri & von Plato (2001).
- ▶ Consider the flattened rules for implication:

$$\frac{[A] \quad B}{A \rightarrow B} \rightarrow_I \qquad \frac{A \rightarrow B \quad A \quad C}{C} \rightarrow_E$$

Generalized expansion and general elimination rules

- ▶ The generalized expansion works only in the case of general elimination rules, but not when they are *flattened* elimination rules *à la* Negri & von Plato (2001).
- ▶ Consider the flattened rules for implication:

$$\frac{[A] \quad B}{A \rightarrow B} \rightarrow_I \qquad \frac{A \rightarrow B \quad A \quad [B] \quad C}{C} \rightarrow_E$$

Generalized expansion and general elimination rules

- ▶ The generalized expansion works only in the case of general elimination rules, but not when they are *flattened* elimination rules *à la* Negri & von Plato (2001).
- ▶ Consider the flattened rules for implication:

$$\frac{[A] \quad B}{A \rightarrow B} \rightarrow_I \qquad \frac{A \rightarrow B \quad A \quad [B] \quad C}{C} \rightarrow_E$$

- ▶ If we try to expand

$$\begin{array}{c} \mathcal{D} \\ [A \rightarrow B] \\ \mathcal{D}' \\ C \end{array} \quad \text{we get}$$

Generalized expansion and general elimination rules

- ▶ The generalized expansion works only in the case of general elimination rules, but not when they are *flattened* elimination rules *à la* Negri & von Plato (2001).
- ▶ Consider the flattened rules for implication:

$$\frac{[A] \quad B}{A \rightarrow B} \rightarrow_I$$

$$\frac{A \rightarrow B \quad A \quad [B] \quad C}{C} \rightarrow_E$$

- ▶ If we try to expand

$$\begin{array}{c} \mathcal{D} \\ [A \rightarrow B] \\ \mathcal{D}' \\ C \end{array}$$

we get

$$\frac{\mathcal{D} \quad A \rightarrow B \quad A \quad \frac{\mathcal{D}' \quad C}{[A \rightarrow B]} \rightarrow_I}{C} \rightarrow_E (n)$$

Generalized expansion and general elimination rules

- ▶ The generalized expansion works only in the case of general elimination rules, but not when they are *flattened* elimination rules *à la* Negri & von Plato (2001).
- ▶ Consider the flattened rules for implication:

$$\frac{[A] \quad B}{A \rightarrow B} \rightarrow_I \qquad \frac{A \rightarrow B \quad A \quad C}{C} \rightarrow_E$$

- ▶ If we try to expand

$$\begin{array}{ccc} \mathcal{D} & & \frac{\overset{n}{B}}{[A \rightarrow B]} \rightarrow_I \\ [A \rightarrow B] & \text{we get} & \mathcal{D}' \\ \mathcal{D}' & & A \rightarrow B \quad \textcircled{A} \quad C \\ C & & \hline C \end{array} \rightarrow_E (n)$$

But this is not a generalized expansion, since we added an extra-hypothesis A .

Generalized expansion and general elimination rules

- ▶ On the contrary, general expansions can be obtained by working with general elimination rules having the form of *higher level* rules *à la* Schroeder-Heister (1984).
- ▶ Consider the higher level rules for implication:

$$\frac{[A] \quad B}{A \rightarrow B} \rightarrow_I \qquad \frac{A \rightarrow B \quad C}{C} \rightarrow_E$$

- ▶ With these rules we have that:

$$\begin{array}{c} \mathcal{D} \\ [A \rightarrow B] \\ \mathcal{D}' \\ C \end{array} \quad \text{expands to} \quad \frac{\frac{\frac{m}{A}}{B} A \overset{n}{\Rightarrow} B}{[A \rightarrow B]} \rightarrow_I (m) \quad \frac{\mathcal{D} \quad \mathcal{D}' \quad C}{A \rightarrow B \quad C} \rightarrow_E (n)$$

Generalized expansion and general elimination rules

- ▶ On the contrary, general expansions can be obtained by working with general elimination rules having the form of *higher level* rules *à la* Schroeder-Heister (1984).
- ▶ Consider the higher level rules for implication:

$$\frac{[A] \quad B}{A \rightarrow B} \rightarrow_I \qquad \frac{A \rightarrow B \quad C}{C} \rightarrow_E \quad [A \Rightarrow B]$$

- ▶ With these rules we have that:

$$\begin{array}{c} \mathcal{D} \\ [A \rightarrow B] \\ \mathcal{D}' \\ C \end{array} \quad \text{expands to} \quad \frac{\frac{\frac{m}{A}}{B} A \Rightarrow B}{[A \rightarrow B]} \rightarrow_I (m) \quad \frac{\mathcal{D} \quad \mathcal{D}' \quad C}{A \rightarrow B \quad C} \rightarrow_E (n)$$

Generalized expansion and general elimination rules

- ▶ On the contrary, general expansions can be obtained by working with general elimination rules having the form of *higher level* rules *à la* Schroeder-Heister (1984).
- ▶ Consider the higher level rules for implication:

$$\frac{[A] \quad B}{A \rightarrow B} \rightarrow_I \qquad \frac{A \rightarrow B \quad C}{C} \rightarrow_E \quad [A \Rightarrow B]$$

- ▶ With these rules we have that:

$$\begin{array}{c} \mathcal{D} \\ [A \rightarrow B] \\ \mathcal{D}' \\ C \end{array} \quad \text{expands to} \quad \frac{\frac{\frac{m}{A}}{B} A \Rightarrow B}{[A \rightarrow B]} \rightarrow_I (m) \quad \frac{\mathcal{D} \quad \mathcal{D}' \quad C}{A \rightarrow B \quad C} \rightarrow_E (n)$$

Generalized expansion and general elimination rules

- ▶ On the contrary, general expansions can be obtained by working with general elimination rules having the form of *higher level* rules *à la* Schroeder-Heister (1984).
- ▶ Consider the higher level rules for implication:

$$\frac{[A] \quad B}{A \rightarrow B} \rightarrow_I \qquad \frac{A \rightarrow B \quad C}{C} \rightarrow_E \quad [A \Rightarrow B]$$

- ▶ With these rules we have that:

$$\begin{array}{c} \mathcal{D} \\ [A \rightarrow B] \\ \mathcal{D}' \\ C \end{array} \quad \text{expands to} \quad \frac{\frac{\frac{m}{A} \quad A \xRightarrow{n} B}{[A \rightarrow B]} \rightarrow_I (m)}{\frac{\mathcal{D} \quad \mathcal{D}'}{A \rightarrow B \quad C} \rightarrow_E (n)}$$

A limit case

- ▶ A limit case of the satisfaction of harmony and stability is represented by the 0-ary connective \perp :

no introduction rules $\frac{\perp}{C} \perp_E$

- ▶ The elimination rule allows one to obtain:
 - (i) no more than what has to be established in order to infer \perp from its introduction rules, since nothing is enough to infer \perp by an introduction rule.
 - (ii) no less than what has to be established in order to infer any C obtainable from \perp , since

A limit case

- ▶ A limit case of the satisfaction of harmony and stability is represented by the 0-ary connective \perp :

$$\text{no introduction rules} \qquad \frac{\perp}{C} \perp_E$$

- ▶ The elimination rule allows one to obtain:
 - (i) no more than what has to be established in order to infer \perp from its introduction rules, since nothing is enough to infer \perp by an introduction rule.
 - (ii) no less than what has to be established in order to infer any C obtainable from \perp , since

A limit case

- ▶ A limit case of the satisfaction of harmony and stability is represented by the 0-ary connective \perp :

no introduction rules $\frac{\perp}{C} \perp_E$

- ▶ The elimination rule allows one to obtain:
 - (i) no more than what has to be established in order to infer \perp from its introduction rules, since nothing is enough to infer \perp by an introduction rule.
 - (ii) no less than what has to be established in order to infer any C obtainable from \perp , since

A limit case

- ▶ A limit case of the satisfaction of harmony and stability is represented by the 0-ary connective \perp :

no introduction rules $\frac{\perp}{C} \perp_E$

- ▶ The elimination rule allows one to obtain:
 - (i) no more than what has to be established in order to infer \perp from its introduction rules, since nothing is enough to infer \perp by an introduction rule.
 - (ii) no less than what has to be established in order to infer any C obtainable from \perp , since

A limit case

- ▶ A limit case of the satisfaction of harmony and stability is represented by the 0-ary connective \perp :

$$\text{no introduction rules} \qquad \frac{\perp}{C} \perp_E$$

- ▶ The elimination rule allows one to obtain:
 - (i) no more than what has to be established in order to infer \perp from its introduction rules, since nothing is enough to infer \perp by an introduction rule.
 - (ii) no less than what has to be established in order to infer any C obtainable from \perp , since

A limit case

- ▶ A limit case of the satisfaction of harmony and stability is represented by the 0-ary connective \perp :

$$\text{no introduction rules} \qquad \frac{\perp}{C} \perp_E$$

- ▶ The elimination rule allows one to obtain:
 - (i) no more than what has to be established in order to infer \perp from its introduction rules, since nothing is enough to infer \perp by an introduction rule.
 - (ii) no less than what has to be established in order to infer any C obtainable from \perp , since

$$\frac{\mathcal{D}}{[\perp]} \quad \text{“expands” to} \quad \frac{\mathcal{D}}{\frac{[\perp]}{C} \perp_E}$$

A limit case

- ▶ A limit case of the satisfaction of harmony and stability is represented by the 0-ary connective \perp :

$$\text{no introduction rules} \qquad \frac{\perp}{C} \perp_E$$

- ▶ The elimination rule allows one to obtain:
 - (i) no more than what has to be established in order to infer \perp from its introduction rules, since nothing is enough to infer \perp by an introduction rule.
 - (ii) no less than what has to be established in order to infer any C obtainable from \perp , since

$$\begin{array}{c} \mathcal{D} \\ [\perp] \\ \mathcal{D}' \\ C \end{array} \quad \text{“expands” to} \quad \frac{\mathcal{D}}{\frac{[\perp]}{C} \perp_E}$$

This is a *sui generis* expansion, since the derivation on the right corresponds to a sort of simplified version of the derivation on the left.

A limit case

- ▶ A limit case of the satisfaction of harmony and stability is represented by the 0-ary connective \perp :

$$\text{no introduction rules} \qquad \frac{\perp}{C} \perp_E$$

- ▶ The elimination rule allows one to obtain:
 - (i) no more than what has to be established in order to infer \perp from its introduction rules, since nothing is enough to infer \perp by an introduction rule.
 - (ii) no less than what has to be established in order to infer any C obtainable from \perp , since

$$\begin{array}{c} \mathcal{D} \\ [\perp] \\ \mathcal{D}' \\ C \end{array} \quad \text{“expands” to} \quad \frac{\mathcal{D}}{\frac{[\perp]}{C} \perp_E} \perp_E$$

This is due to the fact that \perp_E has no minor premisses where to permute upward $\begin{array}{c} \mathcal{D}' \\ C \end{array}$.

A limit case

- ▶ A limit case of the satisfaction of harmony and stability is represented by the 0-ary connective \perp :

$$\text{no introduction rules} \qquad \frac{\perp}{C} \perp_E$$

- ▶ The elimination rule allows one to obtain:
 - (i) no more than what has to be established in order to infer \perp from its introduction rules, since nothing is enough to infer \perp by an introduction rule.
 - (ii) no less than what has to be established in order to infer any C obtainable from \perp , since

$$\frac{\mathcal{D}}{\perp} \quad \text{expands to} \quad \frac{\perp}{\perp} \perp_E$$

The sense of the word ‘expansion’ becomes clear when we consider the simple expansion (i.e. deducibility of identicals).

A limit case

- ▶ A limit case of the satisfaction of harmony and stability is represented by the 0-ary connective \perp :

$$\text{no introduction rules} \qquad \frac{\perp}{C} \perp_E$$

- ▶ The elimination rule allows one to obtain:
 - (i) no more than what has to be established in order to infer \perp from its introduction rules, since nothing is enough to infer \perp by an introduction rule.
 - (ii) no less than what has to be established in order to infer any C obtainable from \perp , since

$$\begin{array}{c} \mathcal{D} \\ [\perp] \\ \mathcal{D}' \\ \perp \end{array} \quad \text{expands to} \quad \frac{\mathcal{D}}{[\perp]} \perp_E$$

In fact, the same expanded proof is obtained when C is \perp .

Inductive definitions: natural numbers

- ▶ Harmony and stability are satisfied by those expressions defined by (generalized) inductive definitions (see Martin-Löf 1971).
- ▶ Consider the rules for the predicate N for natural numbers (Martin-Löf 1984: 71):

$$\frac{}{N0} N_{I_1} \quad \frac{Nt}{Nt'} N_{I_2} \quad \frac{Nt \quad C(0/y) \quad [Nx/y], [C(x/y)] \quad C(x'/y)}{C(t/y)} N_E$$

where x is an eigenvariable.

Inductive definitions: natural numbers

- ▶ Harmony and stability are satisfied by those expressions defined by (generalized) inductive definitions (see Martin-Löf 1971).
- ▶ Consider the rules for the predicate N for natural numbers (Martin-Löf 1984: 71):

$$\frac{}{N0} N_{I_1} \quad \frac{Nt}{Nt'} N_{I_2} \quad \frac{Nt \quad C(0/y) \quad [Nx/y], [C(x/y)] \quad C(x'/y)}{C(t/y)} N_E$$

where x is an eigenvariable.

Inductive definitions: natural numbers

- ▶ Harmony and stability are satisfied by those expressions defined by (generalized) inductive definitions (see Martin-Löf 1971).
- ▶ Consider the rules for the predicate N for natural numbers (Martin-Löf 1984: 71):

$$\frac{}{N0} N_{I_1} \quad \frac{Nt}{Nt'} N_{I_2} \quad \frac{Nt \quad C(0/y) \quad [Nx/y], [C(x/y)] \quad C(x'/y)}{C(t/y)} N_E$$

where x is an eigenvariable.

Inductive definitions: natural numbers

- Harmony and stability are satisfied by those expressions defined by (generalized) inductive definitions (see Martin-Löf 1971).
- Consider the rules for the predicate N for natural numbers (Martin-Löf 1984: 71):

$$\frac{}{N0} N_{I_1} \quad \frac{Nt}{Nt'} N_{I_2} \quad \frac{Nt \quad C(0/y) \quad \frac{[Nx/y], [C(x/y)]}{C(x'/y)} N_E}{C(t/y)} N_E$$

where x is an eigenvariable.

- These rules satisfy harmony:

$$\frac{\frac{}{N0} N_{I_1} \quad \frac{\mathcal{D}_1(0/y) \quad \mathcal{D}_2(x/y)}{C(0/y)} \quad \frac{[Nx/y]^m, [C(x'/y)]^n}{C(x'/y)} N_E(m,n)}{C(0/y)} \text{ reduces to } \frac{\mathcal{D}_1(0/y)}{C(0/y)}$$

Inductive definitions: natural numbers

- ▶ Harmony and stability are satisfied by those expressions defined by (generalized) inductive definitions (see Martin-Löf 1971).
- ▶ Consider the rules for the predicate N for natural numbers (Martin-Löf 1984: 71):

$$\frac{}{N0} N_{I_1} \quad \frac{Nt}{Nt'} N_{I_2} \quad \frac{Nt \quad C(0/y) \quad \frac{[Nx/y], [C(x/y)]}{C(x'/y)} N_E}{C(t/y)} N_E$$

where x is an eigenvariable.

- ▶ These rules satisfy harmony:

$$\frac{\mathcal{D} \quad \frac{Nt}{Nt'} N_{I_2} \quad \frac{\mathcal{D}_1(0/y) \quad \mathcal{D}_2(x/y)}{C(0/y) \quad C(x'/y)} \quad \frac{[Nx/y], [C(x/y)]}{C(t'/y)} N_E}{C(t'/y)} \text{ reduces to } \mathcal{D} \quad \frac{Nt \quad \mathcal{D}_1(0/y) \quad \mathcal{D}_2(x/y)}{C(0/y) \quad C(x'/y)} N_E \quad \frac{[Nx/y], [C(x/y)]}{C(t'/y)} N_E$$

Inductive definitions: natural numbers

- ▶ Harmony and stability are satisfied by those expressions defined by (generalized) inductive definitions (see Martin-Löf 1971).
- ▶ Consider the rules for the predicate N for natural numbers (Martin-Löf 1984: 71):

$$\frac{}{N0} N_{I_1} \quad \frac{Nt}{Nt'} N_{I_2} \quad \frac{Nt \quad C(0/y) \quad \frac{[Nx/y], [C(x/y)]}{C(x'/y)} N_E}{C(t/y)} N_E$$

where x is an eigenvariable.

- ▶ And they satisfy stability:

$$\begin{array}{l} \mathcal{D} \\ [Nt] \\ \mathcal{D}'(t/y) \\ C(t/y) \end{array} \quad \text{expands to} \quad \frac{\mathcal{D} \quad Nt \quad \frac{\frac{}{N0} N_{I_1} \quad \frac{\frac{^n Nx}{[Nx']} N_{I_2}}{C(x'/y)} N_E(n)}{C(0/y)} N_E(n)}{C(t/y)} N_E(n)$$

Inductive definitions: natural numbers

- ▶ Harmony and stability are satisfied by those expressions defined by (generalized) inductive definitions (see Martin-Löf 1971).
- ▶ Consider the rules for the predicate N for natural numbers (Martin-Löf 1984: 71):

$$\frac{}{N0} N_{I_1} \quad \frac{Nt}{Nt'} N_{I_2} \quad \frac{Nt \quad C(0/y) \quad \frac{[Nx/y], [C(x/y)]}{C(x'/y)} N_E}{C(t/y)} N_E$$

where x is an eigenvariable.

- ▶ And they satisfy stability:

$$\begin{array}{ccc} \mathcal{D} & & \frac{N^nx}{[Nx']} N_{I_2} \\ [Nt] & & \frac{}{N0} N_{I_1} \\ \mathcal{D}'(t/y) & \text{expands to} & \mathcal{D} \quad \mathcal{D}'(0/y) \quad \mathcal{D}'(x'/y) \\ C(t/y) & & \frac{Nt \quad C(0/y) \quad C(x'/y)}{C(t/y)} N_{E(n)} \end{array}$$

Here, it is crucial the fact that we presented a N_E as corresponding to the operation of *recursion*.

Inductive definitions: natural numbers

- ▶ Harmony and stability are satisfied by those expressions defined by (generalized) inductive definitions (see Martin-Löf 1971).
- ▶ Consider the rules for the predicate N for natural numbers (Martin-Löf 1984: 71):

$$\frac{}{N0} N_{I_1} \quad \frac{Nt}{Nt'} N_{I_2} \quad \frac{Nt \quad C(0/y) \quad [Nx/y], [C(x/y)] \quad C(x'/y)}{C(t/y)} N_E$$

where x is an eigenvariable.

- ▶ And they satisfy stability:

$$\frac{\mathcal{D}}{Nt} \quad \text{expands to} \quad \frac{\mathcal{D} \quad \frac{}{N0} N_{I_1} \quad \frac{\frac{Nt}{Nt'} N_{I_2}}{Nt} N_E(n)}{Nt}$$

Inductive definitions: natural numbers

- ▶ Harmony and stability are satisfied by those expressions defined by (generalized) inductive definitions (see Martin-Löf 1971).
- ▶ Consider the rules for the predicate N for natural numbers (Martin-Löf 1984: 71):

$$\frac{}{N0} N_{I_1} \quad \frac{Nt}{Nt'} N_{I_2} \quad \frac{Nt \quad C(0/y) \quad C(x'/y)}{C(t/y)} N_E$$

where x is an eigenvariable.

- ▶ And they satisfy stability:

$$\frac{\mathcal{D}}{Nt} \quad \text{expands to} \quad \frac{\mathcal{D}}{Nt} \quad \frac{\frac{Nt}{N0/y} N_{I_1} \quad \frac{Nt}{Nx'/y} N_{I_2}}{Nt/y} N_{E(n)}$$

Take C as N and \mathcal{D}' as a rule of N introduction.

Inductive definitions: natural numbers

- ▶ Harmony and stability are satisfied by those expressions defined by (generalized) inductive definitions (see Martin-Löf 1971).
- ▶ Consider the rules for the predicate N for natural numbers (Martin-Löf 1984: 71):

$$\frac{}{N0} N_{I_1} \quad \frac{Nt}{Nt'} N_{I_2} \quad \frac{Nt \quad C(0/y) \quad C(x'/y)}{C(t/y)} N_E$$

$[Nx/y], [C(x/y)]$

where x is an eigenvariable.

- ▶ And they satisfy stability:

$$\frac{\mathcal{D}}{Nt} \quad \text{expands to} \quad \frac{\mathcal{D}}{Nt} \quad \frac{\frac{}{N0/y} N_{I_1} \quad \frac{Nx/y}{Nx'/y} N_{I_2}}{Nt/y} N_{E(n)}$$

Take C as N and \mathcal{D}' as a rule of N introduction.

For satisfying only deducibility of identical it is sufficient to consider a N_E which corresponds to the operation of *iteration*.

Inductive definitions: identity

- Consider the following rules of identity (Martin-Löf 1971):

$$\frac{}{t = t} =_I \qquad \frac{t = s \quad C(x/y, x/z)}{C(t/y, s/z)} =_E$$

- The usual rule of indiscernibility of identicals

$$\frac{t = s \quad A(t/z)}{A(s/z)}$$

can be obtained by taking $C(x/y, x/z)$ as $A(x/y) \rightarrow A(x/z)$ and considering then the following derivation:

$$\frac{t = s \quad \frac{\frac{A(x/z)}{A(x/y) \rightarrow A(x/z)} \rightarrow_I (1)}{A(t/y) \rightarrow A(s/z)} =_E \quad A(t/z)}{A(s/z)} \rightarrow_E$$

Inductive definitions: identity

- Consider the following rules of identity (Martin-Löf 1971):

$$\frac{}{t = t} =_I \qquad \frac{t = s \quad C(x/y, x/z)}{C(t/y, s/z)} =_E$$

- The usual rule of indiscernibility of identicals

$$\frac{t = s \quad A(t/z)}{A(s/z)}$$

can be obtained by taking $C(x/y, x/z)$ as $A(x/y) \rightarrow A(x/z)$ and considering then the following derivation:

$$\frac{\frac{t = s \quad \frac{\overset{1}{A(x/z)}}{A(x/y) \rightarrow A(x/z)} \rightarrow_I (1)}{A(t/y) \rightarrow A(s/z)} =_E \quad A(t/z)}{A(s/z)} \rightarrow_E$$

Inductive definitions: identity

- Consider the following rules of identity (Martin-Löf 1971):

$$\frac{}{t = t} =_I \qquad \frac{t = s \quad C(x/y, x/z)}{C(t/y, s/z)} =_E$$

- The usual rule of indiscernibility of identicals

$$\frac{t = s \quad A(t/z)}{A(s/z)}$$

can be obtained by taking $C(x/y, x/z)$ as $A(x/y) \rightarrow A(x/z)$ and considering then the following derivation:

$$\frac{\frac{t = s \quad \frac{\frac{A(x/z)}{A(x/y) \rightarrow A(x/z)} \rightarrow_I (1)}{A(t/y) \rightarrow A(s/z)} =_E \quad A(t/z)}{A(s/z)} \rightarrow_E$$

Inductive definitions: identity

- Consider the following rules of identity (Martin-Löf 1971):

$$\frac{}{t = t} =_I \qquad \frac{t = s \quad C(x/y, x/z)}{C(t/y, s/z)} =_E$$

- The usual rule of indiscernibility of identicals

$$\frac{t = s \quad A(t/z)}{A(s/z)}$$

can be obtained by taking $C(x/y, x/z)$ as $A(x/y) \rightarrow A(x/z)$ and considering then the following derivation:

$$\frac{\frac{t = s \quad \frac{\frac{A(x/z)}{A(x/y) \rightarrow A(x/z)} \rightarrow_I (1)}{A(t/y) \rightarrow A(s/z)} =_E \quad A(t/z)}{A(s/z)} \rightarrow_E$$

Inductive definitions: identity

- Consider the following rules of identity (Martin-Löf 1971):

$$\frac{}{t = t} =_I \qquad \frac{t = s \quad C(x/y, x/z)}{C(t/y, s/z)} =_E$$

- The usual rule of indiscernibility of identicals

$$\frac{t = s \quad A(t/z)}{A(s/z)}$$

can be obtained by taking $C(x/y, x/z)$ as $A(x/y) \rightarrow A(x/z)$ and considering then the following derivation:

$$\frac{t = s \quad \frac{\frac{A(x/z)}{A(x/y) \rightarrow A(x/z)} \rightarrow_I (1)}{A(t/y) \rightarrow A(s/z)} =_E \quad A(t/z)}{A(s/z)} \rightarrow_E$$

Inductive definitions: identity

- ▶ The rules of identity satisfy harmony:

$$\frac{\frac{}{t = t} =_I \quad \frac{\mathcal{D}(x/y, x/z) \quad C(x/y, x/z)}{C(t/y, t/z)} =_E}{C(t/y, t/z)} \quad \text{reduces to} \quad \frac{\mathcal{D}(t/y, t/z) \quad C(t/y, t/z)}{C(t/y, t/z)}$$

- ▶ And they also satisfy stability:

Inductive definitions: identity

- ▶ The rules of identity satisfy harmony:

$$\frac{\frac{\overline{t = t} =_I \quad \mathcal{D}(x/y, x/z)}{C(x/y, x/z)}}{C(t/y, t/z)} =_E \quad \text{reduces to} \quad \frac{\mathcal{D}(t/y, t/z)}{C(t/y, t/z)}$$

- ▶ And they also satisfy stability:

Inductive definitions: identity

- ▶ The rules of identity satisfy harmony:

$$\frac{\frac{\overline{t = t} =_I \quad \mathcal{D}(x/y, x/z)}{C(x/y, x/z)} =_E}{C(t/y, t/z)} \quad \text{reduces to} \quad \frac{\mathcal{D}(t/y, t/z)}{C(t/y, t/z)}$$

- ▶ And they also satisfy stability:

Inductive definitions: identity

- The rules of identity satisfy harmony:

$$\frac{\frac{\overline{t = t} =_I \quad \mathcal{D}(x/y, x/z)}{C(x/y, x/z)}}{C(t/y, t/z)} =_E \quad \text{reduces to} \quad \frac{\mathcal{D}(t/y, t/z)}{C(t/y, t/z)}$$

- And they also satisfy stability:

$$\frac{\mathcal{D} \quad [t = s]}{\mathcal{D}'(t/y, s/z)} \quad \text{expands to} \quad \frac{\mathcal{D} \quad \frac{\overline{[x = x]} =_I \quad \mathcal{D}'(x/y, x/z)}{t = s} \quad C(x/y, x/z)}{C(t/y, s/z)} =_E$$

Inductive definitions: identity

- ▶ The rules of identity satisfy harmony:

$$\frac{\frac{\overline{t = t} =_I \quad \mathcal{D}(x/y, x/z)}{C(x/y, x/z)}}{C(t/y, t/z)} =_E \quad \text{reduces to} \quad \frac{\mathcal{D}(t/y, t/z)}{C(t/y, t/z)}$$

- ▶ And they also satisfy stability:

$$\frac{\mathcal{D}}{t = s} \quad \text{expands to} \quad \frac{\mathcal{D} \quad \frac{t = s \quad \overline{x = x} =_I}{t = s} =_E}{t = s}$$

Inductive definitions: identity

- The rules of identity satisfy harmony:

$$\frac{\frac{\overline{t = t} =_I \quad \mathcal{D}(x/y, x/z)}{C(x/y, x/z)}}{C(t/y, t/z)} =_E \quad \text{reduces to} \quad \frac{\mathcal{D}(t/y, t/z)}{C(t/y, t/z)}$$

- And they also satisfy stability:

$$\frac{\mathcal{D}}{t = s} \quad \text{expands to} \quad \frac{\frac{\mathcal{D}}{t = s} \quad \frac{}{x/y = x/z} =_I}{t/y = s/z} =_E$$

Where

$$\left. \begin{array}{l} \mathcal{D}'(t/y, s/z) \\ C(t/y, s/z) \end{array} \right\} \text{ is } t/y = s/z \text{ and } \left. \begin{array}{l} \mathcal{D}'(x/y, x/z) \\ C(x/y, x/z) \end{array} \right\} \text{ is } x/y = x/z$$

Inductive definitions: identity

- ▶ The rules of identity satisfy harmony:

$$\frac{\frac{\overline{t = t} =_I \quad \mathcal{D}(x/y, x/z)}{C(x/y, x/z)} \quad \text{reduces to} \quad \mathcal{D}(t/y, t/z)}{C(t/y, t/z)} =_E \quad C(t/y, t/z)$$

- ▶ And they also satisfy stability:

$$\frac{\mathcal{D}}{t = s} \quad \text{expands to} \quad \frac{\mathcal{D} \quad \frac{\overline{t = s} \quad \overline{x/y = x/z} =_I}{x/y = x/z}}{t/y = s/z} =_E$$

Where

$$\left. \begin{array}{l} \mathcal{D}'(t/y, s/z) \\ C(t/y, s/z) \end{array} \right\} \text{ is } t/y = s/z \text{ and } \left. \begin{array}{l} \mathcal{D}'(x/y, x/z) \\ C(x/y, x/z) \end{array} \right\} \text{ is } x/y = x/z$$

Hence, deducibility of identicals becomes a particular case of generalized expansion.

Table of contents

Part I. Harmony and Stability

Part II. Stability and Identity

Equivalence relations and the identification of proofs

- ▶ The reflexive, symmetric, and transitive closure of
 - ▶ the *reduction* relation corresponding to harmony,
 - ▶ the *standard expansion* relation corresponding to deducibility of identicals

yields two equivalence relations on derivations: \equiv_β and \equiv_η .

- ▶ Now, if we want to take proofs as the fundamental entities on which founding a semantics (as it is the idea of proof-theoretic semantics), then we have to say what proofs are, that is, we have to assign them a genuine ontological status.
- ▶ A first crucial step in this direction is to possess a criterion of identification for proofs, in agreement with the Quinean epistemological principle that there is no entity without identity.
- ▶ An idea could be to use the \equiv_β and \equiv_η in order to establish this criterion: equivalent derivations correspond to – or better, represent – the same proof.

Equivalence relations and the identification of proofs

- ▶ The reflexive, symmetric, and transitive closure of
 - ▶ the *reduction* relation corresponding to harmony,
 - ▶ the *standard expansion* relation corresponding to deducibility of identicals

yields two equivalence relations on derivations: \equiv_β and \equiv_η .

- ▶ Now, if we want to take proofs as the fundamental entities on which founding a semantics (as it is the idea of proof-theoretic semantics), then we have to say what proofs are, that is, we have to assign them a genuine ontological status.
- ▶ A first crucial step in this direction is to possess a criterion of identification for proofs, in agreement with the Quinean epistemological principle that there is no entity without identity.
- ▶ An idea could be to use the \equiv_β and \equiv_η in order to establish this criterion: equivalent derivations correspond to – or better, represent – the same proof.

Equivalence relations and the identification of proofs

- ▶ The reflexive, symmetric, and transitive closure of
 - ▶ the *reduction* relation corresponding to harmony,
 - ▶ the *standard expansion* relation corresponding to deducibility of identicals

yields two equivalence relations on derivations: \equiv_β and \equiv_η .

- ▶ Now, if we want to take proofs as the fundamental entities on which founding a semantics (as it is the idea of proof-theoretic semantics), then we have to say what proofs are, that is, we have to assign them a genuine ontological status.
- ▶ A first crucial step in this direction is to possess a criterion of identification for proofs, in agreement with the Quinean epistemological principle that there is no entity without identity.
- ▶ An idea could be to use the \equiv_β and \equiv_η in order to establish this criterion: equivalent derivations correspond to – or better, represent – the same proof.

Equivalence relations and the identification of proofs

- ▶ The reflexive, symmetric, and transitive closure of
 - ▶ the *reduction* relation corresponding to harmony,
 - ▶ the *standard expansion* relation corresponding to deducibility of identicals

yields two equivalence relations on derivations: \equiv_β and \equiv_η .

- ▶ Now, if we want to take proofs as the fundamental entities on which founding a semantics (as it is the idea of proof-theoretic semantics), then we have to say what proofs are, that is, we have to assign them a genuine ontological status.
- ▶ A first crucial step in this direction is to possess a criterion of identification for proofs, in agreement with the Quinean epistemological principle that there is no entity without identity.
- ▶ An idea could be to use the \equiv_β and \equiv_η in order to establish this criterion: equivalent derivations correspond to – or better, represent – the same proof.

Equivalence relations and the identification of proofs

- ▶ The reflexive, symmetric, and transitive closure of
 - ▶ the *reduction* relation corresponding to harmony,
 - ▶ the *standard expansion* relation corresponding to deducibility of identicals

yields two equivalence relations on derivations: \equiv_β and \equiv_η .

- ▶ Now, if we want to take proofs as the fundamental entities on which founding a semantics (as it is the idea of proof-theoretic semantics), then we have to say what proofs are, that is, we have to assign them a genuine ontological status.
- ▶ A first crucial step in this direction is to possess a criterion of identification for proofs, in agreement with the Quinean epistemological principle that there is no entity without identity.
- ▶ An idea could be to use the \equiv_β and \equiv_η in order to establish this criterion: equivalent derivations correspond to – or better, represent – the same proof.

Equivalence relations and the identification of proofs

- ▶ The reflexive, symmetric, and transitive closure of
 - ▶ the *reduction* relation corresponding to harmony,
 - ▶ the *standard expansion* relation corresponding to deducibility of identicals

yields two equivalence relations on derivations: \equiv_β and \equiv_η .

- ▶ Now, if we want to take proofs as the fundamental entities on which founding a semantics (as it is the idea of proof-theoretic semantics), then we have to say what proofs are, that is, we have to assign them a genuine ontological status.
- ▶ A first crucial step in this direction is to possess a criterion of identification for proofs, in agreement with the Quinean epistemological principle that there is no entity without identity.
- ▶ An idea could be to use the \equiv_β and \equiv_η in order to establish this criterion: equivalent derivations correspond to – or better, represent – the same proof.

Equivalence relations and the identification of proofs

- ▶ The reflexive, symmetric, and transitive closure of
 - ▶ the *reduction* relation corresponding to harmony,
 - ▶ the *standard expansion* relation corresponding to deducibility of identicals

yields two equivalence relations on derivations: \equiv_β and \equiv_η .

- ▶ Now, if we want to take proofs as the fundamental entities on which founding a semantics (as it is the idea of proof-theoretic semantics), then we have to say what proofs are, that is, we have to assign them a genuine ontological status.
- ▶ A first crucial step in this direction is to possess a criterion of identification for proofs, in agreement with the Quinean epistemological principle that there is no entity without identity.
- ▶ An idea could be to use the \equiv_β and \equiv_η in order to establish this criterion: equivalent derivations correspond to – or better, represent – the same proof.

Equivalence relations and the identification of proofs

- ▶ The reflexive, symmetric, and transitive closure of
 - ▶ the *reduction* relation corresponding to harmony,
 - ▶ the *standard expansion* relation corresponding to deducibility of identicals

yields two equivalence relations on derivations: \equiv_β and \equiv_η .

- ▶ Now, if we want to take proofs as the fundamental entities on which founding a semantics (as it is the idea of proof-theoretic semantics), then we have to say what proofs are, that is, we have to assign them a genuine ontological status.
- ▶ A first crucial step in this direction is to possess a criterion of identification for proofs, in agreement with the Quinean epistemological principle that there is no entity without identity.
- ▶ An idea could be to use the \equiv_β and \equiv_η in order to establish this criterion: equivalent derivations correspond to – or better, represent – the same proof.

Equivalence relations and the identification of proofs

- In an intuitionistic setting we have that:

If \mathcal{D} is a closed and β -normal (i.e. peak-free) derivation of a closed sentence, then the last rule of \mathcal{D} is an introduction rule.

- This fact guarantees the respect of the canonicity condition: β -normal derivations *directly* represent BHK proofs.
- Non- β -normal derivations *indirectly* represent BHK proofs: modulo the \equiv_β these derivations can be associated to β -normal derivations and thus to canonical proofs.
- The Church-Rosser theorem for β -reduction warrants the following criterion for proofs identity (cf. Prawitz 1971: 257):

If \mathcal{D}_1 and \mathcal{D}_2 are β -normal, $\mathcal{D}_1 \equiv_\beta \mathcal{D}_2$ iff $\mathcal{D}_1 \cong \mathcal{D}_2$

(where \cong indicates the identity between proofs).

Equivalence relations and the identification of proofs

- ▶ In an intuitionistic setting we have that:

If \mathcal{D} is a closed and β -normal (i.e. peak-free) derivation of a closed sentence, then the last rule of \mathcal{D} is an introduction rule.

- ▶ This fact guarantees the respect of the canonicity condition: β -normal derivations *directly* represent BHK proofs.
- ▶ Non- β -normal derivations *indirectly* represent BHK proofs: modulo the \equiv_β these derivations can be associated to β -normal derivations and thus to canonical proofs.
- ▶ The Church-Rosser theorem for β -reduction warrants the following criterion for proofs identity (cf. Prawitz 1971: 257):

If \mathcal{D}_1 and \mathcal{D}_2 are β -normal, $\mathcal{D}_1 \equiv_\beta \mathcal{D}_2$ iff $\mathcal{D}_1 \cong \mathcal{D}_2$

(where \cong indicates the identity between proofs).

Equivalence relations and the identification of proofs

- In an intuitionistic setting we have that:

If \mathcal{D} is a closed and β -normal (i.e. peak-free) derivation of a closed sentence, then the last rule of \mathcal{D} is an introduction rule.

- This fact guarantees the respect of the canonicity condition:
 β -normal derivations *directly* represent BHK proofs.
- Non- β -normal derivations *indirectly* represent BHK proofs:
modulo the \equiv_β these derivations can be associated to β -normal derivations and thus to canonical proofs.
- The Church-Rosser theorem for β -reduction warrants the following criterion for proofs identity (cf. Prawitz 1971: 257):

If \mathcal{D}_1 and \mathcal{D}_2 are β -normal, $\mathcal{D}_1 \equiv_\beta \mathcal{D}_2$ iff $\mathcal{D}_1 \cong \mathcal{D}_2$

(where \cong indicates the identity between proofs).

Equivalence relations and the identification of proofs

- ▶ In an intuitionistic setting we have that:

If \mathcal{D} is a closed and β -normal (i.e. peak-free) derivation of a closed sentence, then the last rule of \mathcal{D} is an introduction rule.

- ▶ This fact guarantees the respect of the canonicity condition:
 β -normal derivations *directly* represent BHK proofs.
- ▶ Non- β -normal derivations *indirectly* represent BHK proofs:
modulo the \equiv_β these derivations can be associated to β -normal derivations and thus to canonical proofs.
- ▶ The Church-Rosser theorem for β -reduction warrants the following criterion for proofs identity (cf. Prawitz 1971: 257):

If \mathcal{D}_1 and \mathcal{D}_2 are β -normal, $\mathcal{D}_1 \equiv_\beta \mathcal{D}_2$ iff $\mathcal{D}_1 \cong \mathcal{D}_2$

(where \cong indicates the identity between proofs).

Equivalence relations and the identification of proofs

- In an intuitionistic setting we have that:

If \mathcal{D} is a closed and β -normal (i.e. peak-free) derivation of a closed sentence, then the last rule of \mathcal{D} is an introduction rule.

- This fact guarantees the respect of the canonicity condition:
 β -normal derivations *directly* represent BHK proofs.
- Non- β -normal derivations *indirectly* represent BHK proofs:
modulo the \equiv_β these derivations can be associated to β -normal derivations and thus to canonical proofs.
- The Church-Rosser theorem for β -reduction warrants the following criterion for proofs identity (cf. Prawitz 1971: 257):

If \mathcal{D}_1 and \mathcal{D}_2 are β -normal, $\mathcal{D}_1 \equiv_\beta \mathcal{D}_2$ iff $\mathcal{D}_1 \cong \mathcal{D}_2$

(where \cong indicates the identity between proofs).

Equivalence relations and the identification of proofs

- ▶ In an intuitionistic setting we have that:

If \mathcal{D} is a closed and β -normal (i.e. peak-free) derivation of a closed sentence, then the last rule of \mathcal{D} is an introduction rule.

- ▶ This fact guarantees the respect of the canonicity condition:
 β -normal derivations *directly* represent BHK proofs.
- ▶ Non- β -normal derivations *indirectly* represent BHK proofs:
modulo the \equiv_β these derivations can be associated to β -normal derivations and thus to canonical proofs.

- ▶ The Church-Rosser theorem for β -reduction warrants the following criterion for proofs identity (cf. Prawitz 1971: 257):

If \mathcal{D}_1 and \mathcal{D}_2 are β -normal, $\mathcal{D}_1 \equiv_\beta \mathcal{D}_2$ iff $\mathcal{D}_1 \cong \mathcal{D}_2$

(where \cong indicates the identity between proofs).

Equivalence relations and the identification of proofs

- ▶ In an intuitionistic setting we have that:

If \mathcal{D} is a closed and β -normal (i.e. peak-free) derivation of a closed sentence, then the last rule of \mathcal{D} is an introduction rule.

- ▶ This fact guarantees the respect of the canonicity condition:
 β -normal derivations *directly* represent BHK proofs.
- ▶ Non- β -normal derivations *indirectly* represent BHK proofs:
modulo the \equiv_β these derivations can be associated to β -normal derivations and thus to canonical proofs.
- ▶ The Church-Rosser theorem for β -reduction warrants the following criterion for proofs identity (cf. Prawitz 1971: 257):

If \mathcal{D}_1 and \mathcal{D}_2 are β -normal, $\mathcal{D}_1 \equiv_\beta \mathcal{D}_2$ iff $\mathcal{D}_1 \cong \mathcal{D}_2$

(where \cong indicates the identity between proofs).

Equivalence relations and the identification of proofs

- In an intuitionistic setting we have that:

If \mathcal{D} is a closed and β -normal (i.e. peak-free) derivation of a closed sentence, then the last rule of \mathcal{D} is an introduction rule.

- This fact guarantees the respect of the canonicity condition: β -normal derivations *directly* represent BHK proofs.
- Non- β -normal derivations *indirectly* represent BHK proofs: modulo the \equiv_β these derivations can be associated to β -normal derivations and thus to canonical proofs.
- The Church-Rosser theorem for β -reduction warrants the following criterion for proofs identity (cf. Prawitz 1971: 257):

If \mathcal{D}_1 and \mathcal{D}_2 are β -normal, $\mathcal{D}_1 \equiv_\beta \mathcal{D}_2$ iff $\mathcal{D}_1 \cong \mathcal{D}_2$

(where \cong indicates the identity between proofs).

Equivalence relations and the identification of proofs

- In the $\{\rightarrow, \wedge\}$, the previous criteria of proofs identity can be extended to $\beta\eta$ -reduction:

If \mathcal{D}_1 and \mathcal{D}_2 are $\beta\eta$ -normal, $\mathcal{D}_1 \equiv_{\beta\eta} \mathcal{D}_2$ iff $\mathcal{D}_1 \cong \mathcal{D}_2$

- In particular, this extension does not yield the identification of all the derivations of a certain sentence. For example, the two following derivations cannot be identified:

$$\frac{\frac{A \xrightarrow{1} A}{A \wedge B \rightarrow (A \rightarrow A)} \rightarrow_I}{(A \rightarrow A) \rightarrow (A \wedge B \rightarrow (A \rightarrow A))} \rightarrow_I (1)$$
$$\frac{\frac{\frac{A \wedge B}{A} \wedge_{E_1} \rightarrow_I}{A \rightarrow A} \rightarrow_I}{A \wedge B \rightarrow (A \rightarrow A)} \rightarrow_I (1)}{(A \rightarrow A) \rightarrow (A \wedge B \rightarrow (A \rightarrow A))} \rightarrow_I$$

- However, we have seen that stability seems to be preferable to simple deducibility of identicals, within a proof-theoretic semantics setting. What happens if we consider a generalized form of η , “induced” by stability? And what happens if we go outside the fragment $\{\rightarrow, \wedge\}$?

Equivalence relations and the identification of proofs

- In the $\{\rightarrow, \wedge\}$, the previous criteria of proofs identity can be extended to $\beta\eta$ -reduction:

If \mathcal{D}_1 and \mathcal{D}_2 are $\beta\eta$ -normal, $\mathcal{D}_1 \equiv_{\beta\eta} \mathcal{D}_2$ iff $\mathcal{D}_1 \cong \mathcal{D}_2$

- In particular, this extension does not yield the identification of all the derivations of a certain sentence. For example, the two following derivations cannot be identified:

$$\frac{\frac{A \xrightarrow{1} A}{A \wedge B \rightarrow (A \rightarrow A)} \rightarrow_I}{(A \rightarrow A) \rightarrow (A \wedge B \rightarrow (A \rightarrow A))} \rightarrow_I (1)$$
$$\frac{\frac{\frac{A \wedge B}{A} \wedge_{E_1}}{A \rightarrow A} \rightarrow_I}{A \wedge B \rightarrow (A \rightarrow A)} \rightarrow_I (1)}{(A \rightarrow A) \rightarrow (A \wedge B \rightarrow (A \rightarrow A))} \rightarrow_I$$

- However, we have seen that stability seems to be preferable to simple deducibility of identicals, within a proof-theoretic semantics setting. What happens if we consider a generalized form of η , “induced” by stability? And what happens if we go outside the fragment $\{\rightarrow, \wedge\}$?

Equivalence relations and the identification of proofs

- In the $\{\rightarrow, \wedge\}$, the previous criteria of proofs identity can be extended to $\beta\eta$ -reduction:

If \mathcal{D}_1 and \mathcal{D}_2 are $\beta\eta$ -normal, $\mathcal{D}_1 \equiv_{\beta\eta} \mathcal{D}_2$ iff $\mathcal{D}_1 \cong \mathcal{D}_2$

- In particular, this extension does not yield the identification of all the derivations of a certain sentence. For example, the two following derivations cannot be identified:

$$\frac{\frac{A \xrightarrow{1} A}{A \wedge B \rightarrow (A \rightarrow A)} \rightarrow_I}{(A \rightarrow A) \rightarrow (A \wedge B \rightarrow (A \rightarrow A))} \rightarrow_I (1)$$
$$\frac{\frac{\frac{A \wedge B}{A} \wedge_{E_1} \rightarrow_I}{A \rightarrow A} \rightarrow_I (1)}{A \wedge B \rightarrow (A \rightarrow A)} \rightarrow_I (1)$$
$$\frac{(A \rightarrow A) \rightarrow (A \wedge B \rightarrow (A \rightarrow A))}{(A \rightarrow A) \rightarrow (A \wedge B \rightarrow (A \rightarrow A))} \rightarrow_I$$

- However, we have seen that stability seems to be preferable to simple deducibility of identicals, within a proof-theoretic semantics setting. What happens if we consider a generalized form of η , “induced” by stability? And what happens if we go outside the fragment $\{\rightarrow, \wedge\}$?

Equivalence relations and the identification of proofs

- In the $\{\rightarrow, \wedge\}$, the previous criteria of proofs identity can be extended to $\beta\eta$ -reduction:

If \mathcal{D}_1 and \mathcal{D}_2 are $\beta\eta$ -normal, $\mathcal{D}_1 \equiv_{\beta\eta} \mathcal{D}_2$ iff $\mathcal{D}_1 \cong \mathcal{D}_2$

- In particular, this extension does not yield the identification of all the derivations of a certain sentence. For example, the two following derivations cannot be identified:

$$\frac{\frac{A \xrightarrow{1} A}{A \wedge B \rightarrow (A \rightarrow A)} \rightarrow_I}{(A \rightarrow A) \rightarrow (A \wedge B \rightarrow (A \rightarrow A))} \rightarrow_I (1)$$
$$\frac{\frac{\frac{A \wedge B}{A} \wedge_{E_1} \rightarrow_I}{A \rightarrow A} \rightarrow_I (1)}{A \wedge B \rightarrow (A \rightarrow A)} \rightarrow_I (1)$$
$$\frac{(A \rightarrow A) \rightarrow (A \wedge B \rightarrow (A \rightarrow A))}{(A \rightarrow A) \rightarrow (A \wedge B \rightarrow (A \rightarrow A))} \rightarrow_I$$

- However, we have seen that stability seems to be preferable to simple deducibility of identicals, within a proof-theoretic semantics setting. What happens if we consider a generalized form of η , “induced” by stability? And what happens if we go outside the fragment $\{\rightarrow, \wedge\}$?

Equivalence relations and the identification of proofs

- In the $\{\rightarrow, \wedge\}$, the previous criteria of proofs identity can be extended to $\beta\eta$ -reduction:

If \mathcal{D}_1 and \mathcal{D}_2 are $\beta\eta$ -normal, $\mathcal{D}_1 \equiv_{\beta\eta} \mathcal{D}_2$ iff $\mathcal{D}_1 \cong \mathcal{D}_2$

- In particular, this extension does not yield the identification of all the derivations of a certain sentence. For example, the two following derivations cannot be identified:

$$\frac{\frac{A \xrightarrow{1} A}{A \wedge B \rightarrow (A \rightarrow A)} \rightarrow_I}{(A \rightarrow A) \rightarrow (A \wedge B \rightarrow (A \rightarrow A))} \rightarrow_I (1)$$
$$\frac{\frac{\frac{A \wedge B}{A} \wedge_{E1} \rightarrow_I}{A \rightarrow A} \rightarrow_I (1)}{(A \rightarrow A) \rightarrow (A \wedge B \rightarrow (A \rightarrow A))} \rightarrow_I$$

- However, we have seen that stability seems to be preferable to simple deducibility of identicals, within a proof-theoretic semantics setting. What happens if we consider a generalized form of η , “induced” by stability? And what happens if we go outside the fragment $\{\rightarrow, \wedge\}$?

Equivalence relations and the identification of proofs

- In the $\{\rightarrow, \wedge\}$, the previous criteria of proofs identity can be extended to $\beta\eta$ -reduction:

If \mathcal{D}_1 and \mathcal{D}_2 are $\beta\eta$ -normal, $\mathcal{D}_1 \equiv_{\beta\eta} \mathcal{D}_2$ iff $\mathcal{D}_1 \cong \mathcal{D}_2$

- In particular, this extension does not yield the identification of all the derivations of a certain sentence. For example, the two following derivations cannot be identified:

$$\frac{\frac{A \xrightarrow{1} A}{A \wedge B \rightarrow (A \rightarrow A)} \rightarrow_I}{(A \rightarrow A) \rightarrow (A \wedge B \rightarrow (A \rightarrow A))} \rightarrow_I (1)$$
$$\frac{\frac{\frac{A \wedge B \xrightarrow{1} A}{A \rightarrow A} \wedge_{E1}}{A \wedge B \rightarrow (A \rightarrow A)} \rightarrow_I}{(A \rightarrow A) \rightarrow (A \wedge B \rightarrow (A \rightarrow A))} \rightarrow_I (1)$$

- However, we have seen that stability seems to be preferable to simple deducibility of identicals, within a proof-theoretic semantics setting. What happens if we consider a generalized form of η , “induced” by stability? And what happens if we go outside the fragment $\{\rightarrow, \wedge\}$?

Equivalence relations and the identification of proofs

- In the $\{\rightarrow, \wedge\}$, the previous criteria of proofs identity can be extended to $\beta\eta$ -reduction:

If \mathcal{D}_1 and \mathcal{D}_2 are $\beta\eta$ -normal, $\mathcal{D}_1 \equiv_{\beta\eta} \mathcal{D}_2$ iff $\mathcal{D}_1 \cong \mathcal{D}_2$

- In particular, this extension does not yield the identification of all the derivations of a certain sentence. For example, the two following derivations cannot be identified:

$$\frac{\frac{A \xrightarrow{1} A}{A \wedge B \rightarrow (A \rightarrow A)} \rightarrow_I}{(A \rightarrow A) \rightarrow (A \wedge B \rightarrow (A \rightarrow A))} \rightarrow_I (1)$$
$$\frac{\frac{\frac{A \wedge B}{A} \wedge_{E_1} \rightarrow_I}{A \rightarrow A} \rightarrow_I}{A \wedge B \rightarrow (A \rightarrow A)} \rightarrow_I (1)}{(A \rightarrow A) \rightarrow (A \wedge B \rightarrow (A \rightarrow A))} \rightarrow_I$$

- However, we have seen that stability seems to be preferable to simple deducibility of identicals, within a proof-theoretic semantics setting. What happens if we consider a generalized form of η , “induced” by stability? And what happens if we go outside the fragment $\{\rightarrow, \wedge\}$?

Equivalence relations and the identification of proofs

- In the $\{\rightarrow, \wedge\}$, the previous criteria of proofs identity can be extended to $\beta\eta$ -reduction:

If \mathcal{D}_1 and \mathcal{D}_2 are $\beta\eta$ -normal, $\mathcal{D}_1 \equiv_{\beta\eta} \mathcal{D}_2$ iff $\mathcal{D}_1 \cong \mathcal{D}_2$

- In particular, this extension does not yield the identification of all the derivations of a certain sentence. For example, the two following derivations cannot be identified:

$$\frac{\frac{A \xrightarrow{1} A}{A \wedge B \rightarrow (A \rightarrow A)} \rightarrow_I}{(A \rightarrow A) \rightarrow (A \wedge B \rightarrow (A \rightarrow A))} \rightarrow_I (1)$$
$$\frac{\frac{\frac{A \wedge B}{A} \wedge_{E1}}{A \rightarrow A} \rightarrow_I}{A \wedge B \rightarrow (A \rightarrow A)} \rightarrow_I (1)}{(A \rightarrow A) \rightarrow (A \wedge B \rightarrow (A \rightarrow A))} \rightarrow_I$$

- However, we have seen that stability seems to be preferable to simple deducibility of identicals, within a proof-theoretic semantics setting. What happens if we consider a generalized form of η , “induced” by stability? And what happens if we go outside the fragment $\{\rightarrow, \wedge\}$?

Collapse on negative sentences*

$$\begin{array}{c}
 \mathcal{D}_1 \quad \frac{\frac{A}{\perp}}{\neg A} \xrightarrow{1} \frac{A \Rightarrow \perp}{\rightarrow_E} (1) \\
 \frac{\perp}{\neg B} \rightarrow_I \\
 \hline
 \frac{\perp}{\neg A} \rightarrow_I (4)
 \end{array}
 \qquad
 \begin{array}{c}
 \mathcal{D}_2 \quad \frac{\frac{A}{\perp}}{\neg A} \xrightarrow{2} \frac{A \Rightarrow \perp}{\rightarrow_E} (2) \\
 \frac{\perp}{B} \perp_E \\
 \frac{\perp}{B \Rightarrow \perp} \xrightarrow{3} \rightarrow_E (3)
 \end{array}$$

*Thanks to a hint of Luiz Carlos Pereira.

Collapse on negative sentences*

$$\begin{array}{c}
 \mathcal{D}_1 \quad \frac{\frac{\neg A}{\perp} \quad \frac{A}{\perp}}{\neg B} \rightarrow_I \quad \frac{A \Rightarrow \perp}{\rightarrow_E (1)} \\
 \hline
 \frac{\perp}{\neg A} \rightarrow_I (4)
 \end{array}
 \qquad
 \begin{array}{c}
 \mathcal{D}_2 \quad \frac{\frac{\neg A}{\perp} \quad \frac{A}{\perp}}{\perp} \rightarrow_E (2) \quad \frac{A \Rightarrow \perp}{\rightarrow_E (2)} \\
 \hline
 \frac{\perp}{B} \rightarrow_E (3)
 \end{array}$$

$\rightsquigarrow \beta$
(\rightarrow)

$$\begin{array}{c}
 \mathcal{D}_1 \quad \frac{\frac{\neg A}{\perp} \quad \frac{A}{\perp}}{\neg A} \rightarrow_I (2) \quad \frac{A \Rightarrow \perp}{\rightarrow_E (1)} \\
 \hline
 \frac{\perp}{\neg A} \rightarrow_I (2)
 \end{array}$$

*Thanks to a hint of Luiz Carlos Pereira.

Collapse on negative sentences*

$$\begin{array}{c}
 \mathcal{D}_1 \quad \frac{\frac{\neg A}{\perp} \quad \frac{A}{\perp}}{\neg B} \rightarrow_I \quad \frac{A \Rightarrow \perp}{\rightarrow_E (1)} \quad \frac{\frac{\neg A}{\perp} \quad \frac{A}{\perp}}{A \Rightarrow \perp} \rightarrow_E (2) \\
 \hline
 \frac{\perp}{\neg A} \rightarrow_I (4)
 \end{array}$$

\rightsquigarrow_β
(\rightarrow)

$$\begin{array}{c}
 \mathcal{D}_1 \quad \frac{\frac{\neg A}{\perp} \quad \frac{A}{\perp}}{\neg A} \rightarrow_I (2) \quad \frac{A \Rightarrow \perp}{\rightarrow_E (1)}
 \end{array}$$

\rightsquigarrow_η
(\rightarrow)_{Sym}

\mathcal{D}_1
 $\neg A$

*Thanks to a hint of Luiz Carlos Pereira.

Collapse on negative sentences*

$$\begin{array}{c}
 \mathcal{D}_1 \quad \frac{\frac{\frac{\neg A}{\perp} \xrightarrow{1} \perp}{\neg B} \rightarrow_I}{\neg A} \xrightarrow{4} \perp \xrightarrow{2} \perp \xrightarrow{1} \perp \xrightarrow{E} (1) \\
 \mathcal{D}_2 \quad \frac{\frac{\frac{\neg A}{\perp} \xrightarrow{1} \perp}{B} \xrightarrow{3} \perp \xrightarrow{2} \perp \xrightarrow{1} \perp \xrightarrow{E} (2)}{\neg A} \xrightarrow{4} \perp \xrightarrow{2} \perp \xrightarrow{1} \perp \xrightarrow{E} (3) \\
 \hline
 \frac{\perp}{\neg A} \rightarrow_I (4)
 \end{array}
 \quad \rightsquigarrow_{\eta} (\perp)
 \quad
 \begin{array}{c}
 \mathcal{D}_2 \quad \frac{\frac{\frac{\neg A}{\perp} \xrightarrow{1} \perp}{\neg A} \xrightarrow{2} \perp \xrightarrow{1} \perp \xrightarrow{E} (1)}{\neg A} \xrightarrow{2} \perp \xrightarrow{1} \perp \xrightarrow{E} (2)
 \end{array}$$

$$\rightsquigarrow_{\beta} (\rightarrow)$$

$$\begin{array}{c}
 \mathcal{D}_1 \quad \frac{\frac{\frac{\neg A}{\perp} \xrightarrow{1} \perp}{\neg A} \xrightarrow{2} \perp \xrightarrow{1} \perp \xrightarrow{E} (1)}{\neg A} \xrightarrow{2} \perp \xrightarrow{1} \perp \xrightarrow{E} (2)
 \end{array}$$

$$\rightsquigarrow_{\eta} (\rightarrow)_{Sym}$$

$$\begin{array}{c}
 \mathcal{D}_1 \\
 \neg A
 \end{array}$$

*Thanks to a hint of Luiz Carlos Pereira.

Collapse on negative sentences*

$$\begin{array}{c}
 \mathcal{D}_1 \quad \frac{\frac{4}{A} \quad \frac{\perp}{\neg A}}{\perp} \xrightarrow{A \Rightarrow \perp} \frac{\perp}{\neg B} \xrightarrow{\rightarrow_I} \frac{\perp}{\neg A} \xrightarrow{\rightarrow_I} \frac{\perp}{\neg A} \quad (4) \\
 \mathcal{D}_2 \quad \frac{\frac{4}{A} \quad \frac{\perp}{\neg A}}{\perp} \xrightarrow{A \Rightarrow \perp} \frac{\perp}{B} \xrightarrow{\rightarrow_E} \frac{\perp}{B} \xrightarrow{\rightarrow_E} \frac{\perp}{B} \quad (3) \\
 \sim\!\!\sim\eta \quad (\perp)
 \end{array}$$

$$\begin{array}{c}
 \mathcal{D}_2 \quad \frac{\frac{2}{A} \quad \frac{\perp}{\neg A}}{\perp} \xrightarrow{A \Rightarrow \perp} \frac{\perp}{\neg A} \xrightarrow{\rightarrow_I} \frac{\perp}{\neg A} \quad (2) \\
 \sim\!\!\sim\eta \quad (\perp)_{Sym}
 \end{array}$$

$$\begin{array}{c}
 \mathcal{D}_1 \quad \frac{\frac{2}{A} \quad \frac{\perp}{\neg A}}{\perp} \xrightarrow{A \Rightarrow \perp} \frac{\perp}{\neg A} \xrightarrow{\rightarrow_I} \frac{\perp}{\neg A} \quad (2) \\
 \sim\!\!\sim\eta \quad (\rightarrow)_{Sym}
 \end{array}$$

$$\begin{array}{c}
 \mathcal{D}_2 \quad \frac{\frac{2}{A} \quad \frac{\perp}{\neg A}}{\perp} \xrightarrow{A \Rightarrow \perp} \frac{\perp}{\neg A} \xrightarrow{\rightarrow_I} \frac{\perp}{\neg A} \quad (2) \\
 \mathcal{D}_1 \\
 \neg A
 \end{array}$$

*Thanks to a hint of Luiz Carlos Pereira.

Collapse on negative sentences*

$$\begin{array}{c}
 \mathcal{D}_1 \quad \frac{\frac{4}{A} \quad \frac{\perp}{\perp}}{\neg A} \xrightarrow{1} \frac{A \Rightarrow \perp}{\rightarrow_E (1)} \\
 \frac{\perp}{\neg B} \rightarrow_I \\
 \hline
 \frac{\perp}{\neg A} \rightarrow_I (4)
 \end{array}
 \quad
 \begin{array}{c}
 \mathcal{D}_2 \quad \frac{\frac{4}{A} \quad \frac{\perp}{\perp}}{\neg A} \xrightarrow{2} \frac{A \Rightarrow \perp}{\rightarrow_E (2)} \\
 \frac{\perp}{B} \xrightarrow{1} \frac{\perp}{\rightarrow_E (1)} \\
 \frac{\perp}{B} \xrightarrow{3} \frac{\perp}{\rightarrow_E (3)} \\
 \hline
 \frac{\perp}{\neg A} \rightarrow_I (4)
 \end{array}
 \quad
 \begin{array}{c}
 \rightsquigarrow_{\eta} \\
 (\perp)
 \end{array}
 \quad
 \begin{array}{c}
 \mathcal{D}_2 \quad \frac{\frac{2}{A} \quad \frac{\perp}{\perp}}{\neg A} \xrightarrow{1} \frac{A \Rightarrow \perp}{\rightarrow_E (1)} \\
 \frac{\perp}{\neg A} \rightarrow_I (2)
 \end{array}$$

$$\begin{array}{c}
 \rightsquigarrow_{\beta} \\
 (\rightarrow)
 \end{array}
 \quad
 \begin{array}{c}
 \rightsquigarrow_{\eta} \\
 (\perp)_{Sym}
 \end{array}$$

$$\begin{array}{c}
 \mathcal{D}_1 \quad \frac{\frac{2}{A} \quad \frac{\perp}{\perp}}{\neg A} \xrightarrow{1} \frac{A \Rightarrow \perp}{\rightarrow_E (1)} \\
 \frac{\perp}{\neg A} \rightarrow_I (2)
 \end{array}
 \quad
 \begin{array}{c}
 \mathcal{D}_2 \quad \frac{\frac{2}{A} \quad \frac{\perp}{\perp}}{\neg A} \xrightarrow{1} \frac{A \Rightarrow \perp}{\rightarrow_E (1)} \\
 \frac{\perp}{\neg A} \rightarrow_I (2)
 \end{array}$$

$$\begin{array}{c}
 \rightsquigarrow_{\eta} \\
 (\rightarrow)_{Sym}
 \end{array}
 \quad
 \begin{array}{c}
 \rightsquigarrow_{\eta} \\
 (\rightarrow)_{Sym}
 \end{array}$$

$$\begin{array}{c}
 \mathcal{D}_1 \\
 \neg A
 \end{array}
 \quad
 \begin{array}{c}
 \mathcal{D}_2 \\
 \neg A
 \end{array}$$

*Thanks to a hint of Luiz Carlos Pereira.

Collapse on negative sentences*

$$\begin{array}{c}
 \mathcal{D}_1 \quad \frac{\frac{4}{A} \quad \frac{\perp}{\perp} \quad A \Rightarrow \perp \quad (1)}{\frac{\perp}{\neg A} \rightarrow_I} \\
 \frac{\perp}{\neg B} \rightarrow_I \\
 \hline
 \frac{\perp}{\neg A} \rightarrow_I \quad (4)
 \end{array}
 \quad
 \begin{array}{c}
 \mathcal{D}_2 \quad \frac{4}{A} \quad \frac{\perp}{\perp} \quad A \Rightarrow \perp \quad (2)}{\frac{\perp}{\neg A} \rightarrow_I} \\
 \frac{\perp}{B} \perp_E \quad (3)}{\frac{\perp}{\neg A} \rightarrow_I} \\
 \hline
 \frac{\perp}{\neg A} \rightarrow_I \quad (4)
 \end{array}
 \quad
 \equiv_{\eta}
 \quad
 \begin{array}{c}
 \mathcal{D}_2 \quad \frac{2}{A} \quad \frac{\perp}{\perp} \quad A \Rightarrow \perp \quad (1)}{\frac{\perp}{\neg A} \rightarrow_I} \\
 \frac{\perp}{\neg A} \rightarrow_I \quad (2)
 \end{array}$$

$$\equiv_{\beta}$$

$$\begin{array}{c}
 \mathcal{D}_1 \quad \frac{2}{A} \quad \frac{\perp}{\perp} \quad A \Rightarrow \perp \quad (1)}{\frac{\perp}{\neg A} \rightarrow_I} \\
 \frac{\perp}{\neg A} \rightarrow_I \quad (2)
 \end{array}
 \quad
 \equiv_{\eta}
 \quad
 \begin{array}{c}
 \mathcal{D}_2 \quad \frac{2}{A} \quad \frac{\perp}{\perp} \quad A \Rightarrow \perp \quad (1)}{\frac{\perp}{\neg A} \rightarrow_I} \\
 \frac{\perp}{\neg A} \rightarrow_I \quad (2)
 \end{array}$$

$$\equiv_{\eta}$$

$$\begin{array}{c}
 \mathcal{D}_1 \\
 \neg A
 \end{array}
 \quad
 \begin{array}{c}
 \mathcal{D}_2 \\
 \neg A
 \end{array}$$

*Thanks to a hint of Luiz Carlos Pereira.

Collapse on negative sentences*

$$\begin{array}{ccc}
 \begin{array}{c}
 \mathcal{D}_1 \quad \frac{4}{A} \\
 \frac{\neg A \quad \frac{\perp}{\perp} \quad A \Rightarrow \frac{1}{\perp}}{\frac{\perp}{\neg B} \rightarrow_I} \rightarrow_E (1) \\
 \hline
 \frac{\perp}{\neg A} \rightarrow_I (4)
 \end{array}
 &
 \begin{array}{c}
 \mathcal{D}_2 \quad \frac{4}{A} \\
 \frac{\neg A \quad \frac{\perp}{\perp} \quad A \Rightarrow \frac{2}{\perp}}{\frac{\perp}{B} \rightarrow_E (2)} \\
 \hline
 \frac{\perp}{B} \rightarrow_E (3)
 \end{array}
 &
 \equiv_{\eta}
 \begin{array}{c}
 \mathcal{D}_2 \quad \frac{2}{A} \\
 \frac{\neg A \quad \frac{\perp}{\perp} \quad A \Rightarrow \frac{1}{\perp}}{\frac{\perp}{\neg A} \rightarrow_I (2)} \rightarrow_E (1)
 \end{array} \\
 \equiv_{\beta} & & \equiv_{\eta} \\
 \begin{array}{c}
 \mathcal{D}_1 \quad \frac{2}{A} \\
 \frac{\neg A \quad \frac{\perp}{\perp} \quad A \Rightarrow \frac{1}{\perp}}{\frac{\perp}{\neg A} \rightarrow_I (2)} \rightarrow_E (1)
 \end{array}
 &
 &
 \begin{array}{c}
 \mathcal{D}_2 \quad \frac{2}{A} \\
 \frac{\neg A \quad \frac{\perp}{\perp} \quad A \Rightarrow \frac{1}{\perp}}{\frac{\perp}{\neg A} \rightarrow_I (2)} \rightarrow_E (1)
 \end{array} \\
 \equiv_{\eta} & & \equiv_{\eta} \\
 \mathcal{D}_1 \quad \neg A & \equiv_{\beta\eta} & \mathcal{D}_2 \quad \neg A
 \end{array}$$

*Thanks to a hint of Luiz Carlos Pereira.

Collapse on negative sentences*

$$\begin{array}{ccc}
 \begin{array}{c}
 \mathcal{D}_1 \quad \frac{4}{A} \\
 \frac{\neg A \quad \frac{\perp}{\perp} \quad A \Rightarrow \perp \quad (1)}{\frac{\perp}{\neg B} \rightarrow_I} \\
 \hline
 \frac{\perp}{\neg A} \rightarrow_I \quad (4)
 \end{array}
 &
 \begin{array}{c}
 \mathcal{D}_2 \quad \frac{4}{A} \\
 \frac{\neg A \quad \frac{\perp}{\perp} \quad A \Rightarrow \perp \quad (2)}{\frac{\perp}{B} \perp_E} \\
 \frac{\perp}{\perp} \quad B \Rightarrow \perp \quad (3) \\
 \hline
 \frac{\perp}{\neg A} \rightarrow_I \quad (4)
 \end{array}
 &
 \equiv_{\eta}
 \begin{array}{c}
 \mathcal{D}_2 \quad \frac{2}{A} \\
 \frac{\neg A \quad \frac{\perp}{\perp} \quad A \Rightarrow \perp \quad (1)}{\frac{\perp}{\neg A} \rightarrow_I \quad (2)}
 \end{array} \\
 \\
 \equiv_{\beta}
 &
 &
 \equiv_{\eta} \\
 \begin{array}{c}
 \mathcal{D}_1 \quad \frac{2}{A} \\
 \frac{\neg A \quad \frac{\perp}{\perp} \quad A \Rightarrow \perp \quad (1)}{\frac{\perp}{\neg A} \rightarrow_I \quad (2)}
 \end{array}
 &
 &
 \begin{array}{c}
 \mathcal{D}_2 \quad \frac{2}{A} \\
 \frac{\neg A \quad \frac{\perp}{\perp} \quad A \Rightarrow \perp \quad (1)}{\frac{\perp}{\neg A} \rightarrow_I \quad (2)}
 \end{array} \\
 \\
 \equiv_{\eta}
 &
 &
 \equiv_{\eta} \\
 \mathcal{D}_1 \quad \neg A
 &
 \equiv_{\beta\eta}
 &
 \mathcal{D}_2 \quad \neg A
 \end{array}$$

Any two derivations \mathcal{D}_1 and \mathcal{D}_2 of $\neg A$ are $\beta\eta$ -equivalent, and can thus be identified.

*Thanks to a hint of Luiz Carlos Pereira.

Collapse on identity sentences (1)

$$\frac{\mathcal{D} \quad \frac{t = t \quad \overline{x/w = x/w} =_I}{t/w = t/w} =_E}{t/w = t/w}$$

Collapse on identity sentences (1)

$$\frac{\mathcal{D} \quad \frac{t = t \quad \overline{x/w = x/w}}{=_I}}{t/w = t/w} =_E$$

$$\begin{array}{c} \rightsquigarrow_{\eta} \\ (=)_{Sym} \end{array}$$

$$\frac{\mathcal{D}}{t = t}$$

Collapse on identity sentences (1)

$$\frac{\mathcal{D} \quad \frac{t = t \quad \overline{x/w = x/w}}{x/w = x/w} =_I}{t/w = t/w} =_E$$

$$\begin{array}{c} \rightsquigarrow_{\eta} \\ (=)_{Sym} \end{array}$$

$$\frac{\mathcal{D}}{t = t}$$

$$\left. \begin{array}{l} \mathcal{D}'(t/w, t/w) \\ C(t/w, t/w) \end{array} \right\} \text{ is } t/w = t/w$$

$$\left. \begin{array}{l} \mathcal{D}'(x/w, x/w) \\ C(x/w, x/w) \end{array} \right\} \text{ is } x/w = x/w$$

Collapse on identity sentences (1)

$$\frac{\mathcal{D} \quad \frac{t = t \quad \overline{x/w = x/w} =_I}{t/w = t/w} =_E}{\quad} \quad \overset{\rightsquigarrow_{\eta}}{(\Rightarrow)_{Sym}} \quad \overline{t/w = t/w} =_I$$

$$\overset{\rightsquigarrow_{\eta}}{(\Rightarrow)_{Sym}}$$

$$\frac{\mathcal{D}}{t = t}$$

$$\left. \begin{array}{l} \mathcal{D}'(t/w, t/w) \\ C(t/w, t/w) \end{array} \right\} \text{ is } t/w = t/w$$

$$\left. \begin{array}{l} \mathcal{D}'(x/w, x/w) \\ C(x/w, x/w) \end{array} \right\} \text{ is } x/w = x/w$$

Collapse on identity sentences (1)

$$\frac{\mathcal{D} \quad \frac{t = t \quad \overline{x/w = x/w} =_I}{t/w = t/w} =_E}{\overline{t/w = t/w} =_I} \quad \begin{matrix} \rightsquigarrow_{\eta} \\ (=)_{Sym} \end{matrix}$$

$$\begin{matrix} \rightsquigarrow_{\eta} \\ (=)_{Sym} \end{matrix}$$

$$\frac{\mathcal{D} \quad t = t}{t = t}$$

$$\left. \begin{matrix} \mathcal{D}'(t/w, t/w) \\ C(t/w, t/w) \end{matrix} \right\} \text{ is } \overline{t/w = t/w} =_I$$

$$\left. \begin{matrix} \mathcal{D}'(x/w, x/w) \\ C(x/w, x/w) \end{matrix} \right\} \text{ is } \overline{x/w = x/w} =_I$$

$$\left. \begin{matrix} \mathcal{D}'(t/w, t/w) \\ C(t/w, t/w) \end{matrix} \right\} \text{ is } t/w = t/w$$

$$\left. \begin{matrix} \mathcal{D}'(x/w, x/w) \\ C(x/w, x/w) \end{matrix} \right\} \text{ is } x/w = x/w$$

Collapse on identity sentences (1)

$$\frac{\mathcal{D} \quad \frac{t = t \quad \overline{x/w = x/w} =_I}{t/w = t/w} =_E}{t/w = t/w} \equiv_{\eta} \overline{t/w = t/w} =_I$$

\equiv_{η}

$$\frac{\mathcal{D}}{t = t}$$

Collapse on identity sentences (1)

$$\frac{\mathcal{D} \quad \frac{t = t \quad \frac{x/w = x/w}{=_I}}{=_E}}{t/w = t/w} \equiv_{\eta} \frac{}{t/w = t/w} =_I$$
$$\equiv_{\eta}$$
$$\frac{\mathcal{D}}{t = t}$$

- This means that any derivation \mathcal{D} of $t = t$ can be identified with the derivation obtained just by an application of $=_I$, i.e. by a reflexivity proof.

Collapse on identity sentences (1)

$$\frac{\mathcal{D} \quad t = t \quad \frac{x/w = x/w}{=_I}}{t/w = t/w} =_E \quad \equiv_\eta \quad \frac{}{t/w = t/w} =_I$$

\equiv_η

$$\frac{\mathcal{D}}{t = t}$$

- ▶ This means that any derivation \mathcal{D} of $t = t$ can be identified with the derivation obtained just by an application of $=_I$, i.e. by a reflexivity proof.
- ▶ At first glance, this could seem to be an odd situation.

Collapse on identity sentences (1)

$$\frac{\mathcal{D} \quad t = t \quad \frac{x/w = x/w}{t/w = t/w} =_E}{t/w = t/w} =_I \quad \equiv_{\eta} \quad \frac{x/w = x/w}{t/w = t/w} =_I$$
$$\equiv_{\eta}$$
$$\frac{\mathcal{D} \quad t = t}{t = t}$$

- ▶ This means that any derivation \mathcal{D} of $t = t$ can be identified with the derivation obtained just by an application of $=_I$, i.e. by a reflexivity proof.
- ▶ At first glance, this could seem to be an odd situation. However, it can become acceptable if we think of it in the following way:

Collapse on identity sentences (1)

$$\begin{array}{c}
 \mathcal{D} \\
 \frac{t = t \quad \frac{}{x/w = x/w} =_I}{t/w = t/w} =_E \quad \equiv_{\eta} \quad \frac{}{t/w = t/w} =_I \\
 \\
 \equiv_{\eta} \\
 \mathcal{D} \\
 t = t
 \end{array}$$

- ▶ This means that any derivation \mathcal{D} of $t = t$ can be identified with the derivation obtained just by an application of $=_I$, i.e. by a reflexivity proof.
- ▶ At first glance, this could seem to be an odd situation. However, it can become acceptable if we think of it in the following way: there is only one canonical proof of $t = t$ – its reflexivity proof – and all the other proofs of $t = t$ correspond to this canonical proof.

Collapse on identity sentences (1)

$$\frac{\mathcal{D} \quad t = t \quad \frac{x/w = x/w}{t/w = t/w} =_E}{t/w = t/w} =_I \quad \equiv_{\eta} \quad \frac{x/w = x/w}{t/w = t/w} =_I$$

$$\equiv_{\eta}$$

$$\frac{\mathcal{D} \quad t = t}{t = t}$$

- ▶ This means that any derivation \mathcal{D} of $t = t$ can be identified with the derivation obtained just by an application of $=_I$, i.e. by a reflexivity proof.
- ▶ At first glance, this could seem to be an odd situation. However, it can become acceptable if we think of it in the following way: there is only one canonical proof of $t = t$ – its reflexivity proof – and all the other proofs of $t = t$ correspond to this canonical proof.
- ▶ This is an acceptable formulation since it says that all the proofs of a *trivial* identity sentence like $t = t$ are in fact themselves trivial since they are simple reflexivity proofs.

Collapse on identity sentences (2)

- Suppose now to consider a richer syntactical setting allowing one to transform a term t into another term s – and *vice versa* – by mean of some set of rewriting rules.

Collapse on identity sentences (2)

- Suppose now to consider a richer syntactical setting allowing one to transform a term t into another term s – and *vice versa* – by mean of some set of rewriting rules.

$$\frac{\mathcal{D} \quad \frac{}{x/y = x/z} =_I}{t/y = s/z} =_E$$

Collapse on identity sentences (2)

- Suppose now to consider a richer syntactical setting allowing one to transform a term t into another term s – and *vice versa* – by mean of some set of rewriting rules.

$$\frac{\mathcal{D} \quad \frac{}{x/y = x/z} =_I}{t/y = s/z} =_E$$

$$\begin{array}{c} \rightsquigarrow_{\eta} \\ (=)_{Sym} \end{array}$$

$$\frac{}{t = s} \mathcal{D}$$

Collapse on identity sentences (2)

- Suppose now to consider a richer syntactical setting allowing one to transform a term t into another term s – and *vice versa* – by mean of some set of rewriting rules.

$$\frac{\mathcal{D} \quad \frac{t = s \quad x/y = x/z}{x/y = x/z} =_I}{t/y = s/z} =_E$$

$$\begin{array}{c} \rightsquigarrow_{\eta} \\ (=)_{Sym} \end{array}$$

$$\frac{\mathcal{D}}{t = s}$$

$$\left. \begin{array}{l} \mathcal{D}'(t/y, s/z) \\ C(t/y, s/z) \end{array} \right\} \text{ is } t/y = s/z$$

$$\left. \begin{array}{l} \mathcal{D}'(x/y, x/z) \\ C(x/y, x/z) \end{array} \right\} \text{ is } x/y = x/z$$

Collapse on identity sentences (2)

- Suppose now to consider a richer syntactical setting allowing one to transform a term t into another term s – and *vice versa* – by mean of some set of rewriting rules.

$$\frac{\mathcal{D} \quad \frac{t = s \quad x/y = x/z}{t/y = s/z} =_E}{t/y = s/z} =_I \quad \overset{\rightsquigarrow_{\eta}}{(\Rightarrow)_{Sym}} \quad \frac{}{t/y = s/z} =_I$$

$$\overset{\rightsquigarrow_{\eta}}{(\Rightarrow)_{Sym}}$$

$$\frac{\mathcal{D}}{t = s}$$

$$\left. \begin{array}{l} \mathcal{D}'(t/y, s/z) \\ C(t/y, s/z) \end{array} \right\} \text{ is } t/y = s/z$$

$$\left. \begin{array}{l} \mathcal{D}'(x/y, x/z) \\ C(x/y, x/z) \end{array} \right\} \text{ is } x/y = x/z$$

Collapse on identity sentences (2)

- Suppose now to consider a richer syntactical setting allowing one to transform a term t into another term s – and *vice versa* – by mean of some set of rewriting rules.

$$\frac{\mathcal{D} \quad \frac{t = s \quad x/y = x/z}{t/y = s/z} =_E}{\quad} \quad \begin{array}{c} \rightsquigarrow_{\eta} \\ (=)_{Sym} \end{array} \quad \frac{\quad}{t/y = s/z} =_I$$

$$\begin{array}{c} \rightsquigarrow_{\eta} \\ (=)_{Sym} \end{array}$$

t rewrites on s or s rewrites on t

$$\mathcal{D} \quad t = s$$

$$\left. \begin{array}{l} \mathcal{D}'(t/y, s/z) \\ C(t/y, s/z) \end{array} \right\} \text{ is } \frac{\quad}{t/y = s/z} =_I$$

$$\left. \begin{array}{l} \mathcal{D}'(t/y, s/z) \\ C(t/y, s/z) \end{array} \right\} \text{ is } t/y = s/z$$

$$\left. \begin{array}{l} \mathcal{D}'(x/y, x/z) \\ C(x/y, x/z) \end{array} \right\} \text{ is } \frac{\quad}{x/y = x/z} =_I$$

$$\left. \begin{array}{l} \mathcal{D}'(x/y, x/z) \\ C(x/y, x/z) \end{array} \right\} \text{ is } x/y = x/z$$

Collapse on identity sentences (2)

- Suppose now to consider a richer syntactical setting allowing one to transform a term t into another term s – and *vice versa* – by mean of some set of rewriting rules.

$$\frac{\mathcal{D} \quad \frac{}{x/y = x/z} =_I}{t/y = s/z} =_E \quad \quad \quad \begin{array}{c} \rightsquigarrow_{\eta} \\ (=)_{Sym} \end{array} \quad \frac{}{t/y = t/z} =_I$$

$$\begin{array}{c} \rightsquigarrow_{\eta} \\ (=)_{Sym} \end{array}$$

t rewrites on s or s rewrites on t

$$\mathcal{D} \\ t = s$$

$$\left. \begin{array}{l} \mathcal{D}'(t/y, t/z) \\ C(t/y, t/z) \end{array} \right\} \text{ is } \frac{}{t/y = t/z} =_I$$

$$\left. \begin{array}{l} \mathcal{D}'(t/y, s/z) \\ C(t/y, s/z) \end{array} \right\} \text{ is } t/y = s/z$$

$$\left. \begin{array}{l} \mathcal{D}'(x/y, x/z) \\ C(x/y, x/z) \end{array} \right\} \text{ is } \frac{}{x/y = x/z} =_I$$

$$\left. \begin{array}{l} \mathcal{D}'(x/y, x/z) \\ C(x/y, x/z) \end{array} \right\} \text{ is } x/y = x/z$$

Collapse on identity sentences (2)

- Suppose now to consider a richer syntactical setting allowing one to transform a term t into another term s – and *vice versa* – by mean of some set of rewriting rules.

$$\frac{\mathcal{D} \quad \frac{t = s \quad \overline{x/y = x/z} =_I}{t/y = s/z} =_E}{\quad} \quad \begin{array}{c} \rightsquigarrow_{\eta} \\ (=)_{Sym} \end{array} \quad \overline{t/y = t/z} =_I$$

$$\begin{array}{c} \rightsquigarrow_{\eta} \\ (=)_{Sym} \end{array}$$

t rewrites on s or s rewrites on t

$$\mathcal{D} \\ t = s$$

$$\left. \begin{array}{l} \mathcal{D}'(s/y, s/z) \\ C(s/y, s/z) \end{array} \right\} \text{ is } \overline{s/y = s/z} =_I$$

$$\left. \begin{array}{l} \mathcal{D}'(t/y, s/z) \\ C(t/y, s/z) \end{array} \right\} \text{ is } t/y = s/z$$

$$\left. \begin{array}{l} \mathcal{D}'(x/y, x/z) \\ C(x/y, x/z) \end{array} \right\} \text{ is } \overline{x/y = x/z} =_I$$

$$\left. \begin{array}{l} \mathcal{D}'(x/y, x/z) \\ C(x/y, x/z) \end{array} \right\} \text{ is } x/y = x/z$$

Collapse on identity sentences (2)

- Suppose now to consider a richer syntactical setting allowing one to transform a term t into another term s – and *vice versa* – by mean of some set of rewriting rules.

$$\frac{\mathcal{D} \quad \frac{t = s \quad \overline{x/y = x/z} =_I}{t/y = s/z} =_E}{t/y = s/z} \equiv_\eta \frac{\overline{t/y = s/z} =_I}{t/y = s/z}$$

\equiv_η

t rewrites on s or s rewrites on t

\mathcal{D}

$t = s$

Collapse on identity sentences (2)

- Suppose now to consider a richer syntactical setting allowing one to transform a term t into another term s – and *vice versa* – by mean of some set of rewriting rules.

$$\begin{array}{c}
 \mathcal{D} \\
 \frac{t = s \quad \frac{}{x/y = x/z} =_I}{t/y = s/z} =_E \quad \equiv_\eta \quad \frac{}{t/y = s/z} =_I \\
 \\
 \equiv_\eta \quad \quad \quad t \text{ rewrites on } s \text{ or } s \text{ rewrites on } t \\
 \\
 \mathcal{D} \\
 t = s
 \end{array}$$

- This situation becomes particularly problematic if, for every t and s , and every derivation \mathcal{D} of $t = s$, it is always possible to make that t rewrite on s , and *vice versa*.

Collapse on identity sentences (2)

- Suppose now to consider a richer syntactical setting allowing one to transform a term t into another term s – and *vice versa* – by mean of some set of rewriting rules.

$$\begin{array}{c}
 \mathcal{D} \\
 \frac{t = s \quad \frac{}{x/y = x/z} =_I}{t/y = s/z} =_E \quad \equiv_\eta \quad \frac{}{t/y = s/z} =_I \\
 \\
 \equiv_\eta \quad \quad \quad t \text{ rewrites on } s \text{ or } s \text{ rewrites on } t \\
 \\
 \mathcal{D} \\
 t = s
 \end{array}$$

- This situation becomes particularly problematic if, for every t and s , and every derivation \mathcal{D} of $t = s$, it is always possible to make that t rewrite on s , and *vice versa*.
- In this way, every proof of a every identity sentence would be identified with a reflexivity proof.

Identity sentences in Martin-Löf's ITT

- ▶ This situation can be better explained by considering Martin-Löf's intuitionistic type theory (1984).
- ▶ In Martin-Löf's type theory proofs are denoted – or better, *coded* – by λ -terms, i.e. computable functions.

In this way one can produce *judgements* like:

- ▶ $t : A$, which says that t codes a proof of the proposition A , i.e. t is a proof-term of type A ;
- ▶ $t \equiv s : A$, which says that (the codes of A) t and s are equivalent with respect to certain transformations (viz. rewrite operations); since t and s are λ -terms, the most fundamental of these transformations is the β -reduction (t and s are thus computationally equivalent codes).
- ▶ The rules for identity are the following:*

*Note that in Martin-Löf's type theory identity is not absolute, but relative to a type A .

Identity sentences in Martin-Löf's ITT

- ▶ This situation can be better explained by considering Martin-Löf's intuitionistic type theory (1984).
- ▶ In Martin-Löf's type theory proofs are denoted – or better, *coded* – by λ -terms, i.e. computable functions.

In this way one can produce *judgements* like:

- ▶ $t : A$, which says that t codes a proof of the proposition A , i.e. t is a proof-term of type A ;
- ▶ $t \equiv s : A$, which says that (the codes of A) t and s are equivalent with respect to certain transformations (viz. rewrite operations); since t and s are λ -terms, the most fundamental of these transformations is the β -reduction (t and s are thus computationally equivalent codes).
- ▶ The rules for identity are the following:*

*Note that in Martin-Löf's type theory identity is not absolute, but relative to a type A .

Identity sentences in Martin-Löf's ITT

- ▶ This situation can be better explained by considering Martin-Löf's intuitionistic type theory (1984).
- ▶ In Martin-Löf's type theory proofs are denoted – or better, *coded* – by λ -terms, i.e. computable functions.

In this way one can produce *judgements* like:

- ▶ $t : A$, which says that t codes a proof of the proposition A , i.e. t is a proof-term of type A ;
- ▶ $t \equiv s : A$, which says that (the codes of A) t and s are equivalent with respect to certain transformations (viz. rewrite operations); since t and s are λ -terms, the most fundamental of these transformations is the β -reduction (t and s are thus computationally equivalent codes).
- ▶ The rules for identity are the following:*

*Note that in Martin-Löf's type theory identity is not absolute, but relative to a type A .

Identity sentences in Martin-Löf's ITT

- ▶ This situation can be better explained by considering Martin-Löf's intuitionistic type theory (1984).
- ▶ In Martin-Löf's type theory proofs are denoted – or better, *coded* – by λ -terms, i.e. computable functions.

In this way one can produce *judgements* like:

- ▶ $t : A$, which says that t codes a proof of the proposition A , i.e. t is a proof-term of type A ;
- ▶ $t \equiv s : A$, which says that (the codes of A) t and s are equivalent with respect to certain transformations (viz. rewrite operations); since t and s are λ -terms, the most fundamental of these transformations is the β -reduction (t and s are thus computationally equivalent codes).
- ▶ The rules for identity are the following:*

*Note that in Martin-Löf's type theory identity is not absolute, but relative to a type A .

Identity sentences in Martin-Löf's ITT

- ▶ This situation can be better explained by considering Martin-Löf's intuitionistic type theory (1984).
- ▶ In Martin-Löf's type theory proofs are denoted – or better, *coded* – by λ -terms, i.e. computable functions.

In this way one can produce *judgements* like:

- ▶ $t : A$, which says that t codes a proof of the proposition A , i.e. t is a proof-term of type A ;
- ▶ $t \equiv s : A$, which says that (the codes of A) t and s are equivalent with respect to certain transformations (viz. rewrite operations); since t and s are λ -terms, the most fundamental of these transformations is the β -reduction (t and s are thus computationally equivalent codes).
- ▶ The rules for identity are the following:*

*Note that in Martin-Löf's type theory identity is not absolute, but relative to a type A .

Identity sentences in Martin-Löf's ITT

- ▶ This situation can be better explained by considering Martin-Löf's intuitionistic type theory (1984).
- ▶ In Martin-Löf's type theory proofs are denoted – or better, *coded* – by λ -terms, i.e. computable functions.

In this way one can produce *judgements* like:

- ▶ $t : A$, which says that t codes a proof of the proposition A , i.e. t is a proof-term of type A ;
- ▶ $t \equiv s : A$, which says that (the codes of A) t and s are equivalent with respect to certain transformations (viz. rewrite operations); since t and s are λ -terms, the most fundamental of these transformations is the β -reduction (t and s are thus computationally equivalent codes).
- ▶ The rules for identity are the following:*

*Note that in Martin-Löf's type theory identity is not absolute, but relative to a type A .

Identity sentences in Martin-Löf's ITT

- ▶ This situation can be better explained by considering Martin-Löf's intuitionistic type theory (1984).
- ▶ In Martin-Löf's type theory proofs are denoted – or better, *coded* – by λ -terms, i.e. computable functions.

In this way one can produce *judgements* like:

- ▶ $t : A$, which says that t codes a proof of the proposition A , i.e. t is a proof-term of type A ;
- ▶ $t \equiv s : A$, which says that (the codes of A) t and s are equivalent with respect to certain transformations (viz. rewrite operations); since t and s are λ -terms, the most fundamental of these transformations is the β -reduction (t and s are thus computationally equivalent codes).
- ▶ The rules for identity are the following:*

*Note that in Martin-Löf's type theory identity is not absolute, but relative to a type A .

Identity sentences in Martin-Löf's ITT

- ▶ This situation can be better explained by considering Martin-Löf's intuitionistic type theory (1984).
- ▶ In Martin-Löf's type theory proofs are denoted – or better, *coded* – by λ -terms, i.e. computable functions.

In this way one can produce *judgements* like:

- ▶ $t : A$, which says that t codes a proof of the proposition A , i.e. t is a proof-term of type A ;
- ▶ $t \equiv s : A$, which says that (the codes of A) t and s are equivalent with respect to certain transformations (viz. rewrite operations); since t and s are λ -terms, the most fundamental of these transformations is the β -reduction (t and s are thus computationally equivalent codes).
- ▶ The rules for identity are the following:*

*Note that in Martin-Löf's type theory identity is not absolute, but relative to a type A .

Identity sentences in Martin-Löf's ITT

- ▶ This situation can be better explained by considering Martin-Löf's intuitionistic type theory (1984).
- ▶ In Martin-Löf's type theory proofs are denoted – or better, *coded* – by λ -terms, i.e. computable functions.

In this way one can produce *judgements* like:

- ▶ $t : A$, which says that t codes a proof of the proposition A , i.e. t is a proof-term of type A ;
 - ▶ $t \equiv s : A$, which says that (the codes of A) t and s are equivalent with respect to certain transformations (viz. rewrite operations); since t and s are λ -terms, the most fundamental of these transformations is the β -reduction (t and s are thus computationally equivalent codes).
- ▶ The rules for identity are the following:*

*Note that in Martin-Löf's type theory identity is not absolute, but relative to a type A .

Identity sentences in Martin-Löf's ITT

- ▶ This situation can be better explained by considering Martin-Löf's intuitionistic type theory (1984).
- ▶ In Martin-Löf's type theory proofs are denoted – or better, *coded* – by λ -terms, i.e. computable functions.

In this way one can produce *judgements* like:

- ▶ $t : A$, which says that t codes a proof of the proposition A , i.e. t is a proof-term of type A ;
- ▶ $t \equiv s : A$, which says that (the codes of A) t and s are equivalent with respect to certain transformations (viz. rewrite operations); since t and s are λ -terms, the most fundamental of these transformations is the β -reduction (t and s are thus computationally equivalent codes).
- ▶ The rules for identity are the following:*

*Note that in Martin-Löf's type theory identity is not absolute, but relative to a type A .

Identity sentences in Martin-Löf's ITT

- ▶ This situation can be better explained by considering Martin-Löf's intuitionistic type theory (1984).
- ▶ In Martin-Löf's type theory proofs are denoted – or better, *coded* – by λ -terms, i.e. computable functions.

In this way one can produce *judgements* like:

- ▶ $t : A$, which says that t codes a proof of the proposition A , i.e. t is a proof-term of type A ;
- ▶ $t \equiv s : A$, which says that (the codes of A) t and s are equivalent with respect to certain transformations (viz. rewrite operations); since t and s are λ -terms, the most fundamental of these transformations is the β -reduction (t and s are thus computationally equivalent codes).
- ▶ The rules for identity are the following:*

$$\frac{t \equiv s : A}{\mathbf{r}(t) : t =_A s} =_I \qquad \frac{p : t =_A s}{t \equiv s : A} =_E$$

*Note that in Martin-Löf's type theory identity is not absolute, but relative to a type A .

Identity sentences in Martin-Löf's ITT

- ▶ This situation can be better explained by considering Martin-Löf's intuitionistic type theory (1984).
- ▶ In Martin-Löf's type theory proofs are denoted – or better, *coded* – by λ -terms, i.e. computable functions.

In this way one can produce *judgements* like:

- ▶ $t : A$, which says that t codes a proof of the proposition A , i.e. t is a proof-term of type A ;
- ▶ $t \equiv s : A$, which says that (the codes of A) t and s are equivalent with respect to certain transformations (viz. rewrite operations); since t and s are λ -terms, the most fundamental of these transformations is the β -reduction (t and s are thus computationally equivalent codes).
- ▶ The rules for identity are the following:*

$$\frac{t \equiv s : A}{\mathbf{r}(t) : t =_A s} =_I \qquad \frac{p : t =_A s}{t \equiv s : A} =_E$$

where \mathbf{r} represents a reflexivity proof.

*Note that in Martin-Löf's type theory identity is not absolute, but relative to a type A .

Identity sentences in Martin-Löf's ITT

- ▶ This situation can be better explained by considering Martin-Löf's intuitionistic type theory (1984).
- ▶ In Martin-Löf's type theory proofs are denoted – or better, *coded* – by λ -terms, i.e. computable functions.

In this way one can produce *judgements* like:

- ▶ $t : A$, which says that t codes a proof of the proposition A , i.e. t is a proof-term of type A ;
- ▶ $t \equiv s : A$, which says that (the codes of A) t and s are equivalent with respect to certain transformations (viz. rewrite operations); since t and s are λ -terms, the most fundamental of these transformations is the β -reduction (t and s are thus computationally equivalent codes).
- ▶ The rules for identity are the following:*

$$\frac{t \equiv s : A}{\mathbf{r}(s) : t =_A s} =_I \qquad \frac{p : t =_A s}{t \equiv s : A} =_E$$

where \mathbf{r} represents a reflexivity proof.

*Note that in Martin-Löf's type theory identity is not absolute, but relative to a type A .

Extensional Martin-Löf's ITT

- ▶ These rules allow one to pass from an equivalence by rewriting (\equiv) to a propositional identity ($=_A$), and *vice versa*.
- ▶ What is problematic is the *vice versa* direction: the rule $=_E$ makes one to loose the proof structure coded by the term p .
- ▶ This becomes more clear by considering that

Extensional Martin-Löf's ITT

- ▶ These rules allow one to pass from an equivalence by rewriting (\equiv) to a propositional identity ($=_A$), and *vice versa*.
- ▶ What is problematic is the *vice versa* direction: the rule $=_E$ makes one to loose the proof structure coded by the term p .
- ▶ This becomes more clear by considering that

Extensional Martin-Löf's ITT

- ▶ These rules allow one to pass from an equivalence by rewriting (\equiv) to a propositional identity ($=_A$), and *vice versa*.
- ▶ What is problematic is the *vice versa* direction: the rule $=_E$ makes one to loose the proof structure coded by the term p .
- ▶ This becomes more clear by considering that

Extensional Martin-Löf's ITT

- ▶ These rules allow one to pass from an equivalence by rewriting (\equiv) to a propositional identity ($=_A$), and *vice versa*.
- ▶ What is problematic is the *vice versa* direction: the rule $=_E$ makes one to loose the proof structure coded by the term p .
- ▶ This becomes more clear by considering that

$$\begin{array}{c} \mathcal{D} \\ p : t =_A s \end{array} \quad \text{can be (simply) expanded to} \quad \frac{\frac{\mathcal{D} \quad p : t =_A s}{t \equiv s : A} =_E}{\mathbf{r}(t) : t =_A s} =_I$$

Extensional Martin-Löf's ITT

- ▶ These rules allow one to pass from an equivalence by rewriting (\equiv) to a propositional identity ($=_A$), and *vice versa*.
- ▶ What is problematic is the *vice versa* direction: the rule $=_E$ makes one to loose the proof structure coded by the term p .
- ▶ This becomes more clear by considering that

$$\begin{array}{c} \mathcal{D} \\ p : t =_A s \end{array} \quad \text{can be (simply) expanded to} \quad \frac{\frac{\mathcal{D} \quad p : t =_A s}{t \equiv s : A} =_E}{\mathbf{r}(s) : t =_A s} =_I$$

Extensional Martin-Löf's ITT

- ▶ These rules allow one to pass from an equivalence by rewriting (\equiv) to a propositional identity ($=_A$), and *vice versa*.
- ▶ What is problematic is the *vice versa* direction: the rule $=_E$ makes one to loose the proof structure coded by the term p .
- ▶ This becomes more clear by considering that

$$\begin{array}{c} \mathcal{D} \\ p : t =_A s \end{array} \quad \text{can be (simply) expanded to} \quad \frac{\frac{\mathcal{D} \quad p : t =_A s}{t \equiv s : A} =_E}{\mathbf{r}(s) : t =_A s} =_I$$

- ▶ This means that if we close \equiv by (the) η -equivalence (induced by this expansion) we have that $p \equiv \mathbf{r}(t) \equiv \mathbf{r}(s)$.

Extensional Martin-Löf's ITT

- ▶ These rules allow one to pass from an equivalence by rewriting (\equiv) to a propositional identity ($=_A$), and *vice versa*.
- ▶ What is problematic is the *vice versa* direction: the rule $=_E$ makes one to loose the proof structure coded by the term p .
- ▶ This becomes more clear by considering that

$$\begin{array}{c} \mathcal{D} \\ p : t =_A s \end{array} \quad \text{can be (simply) expanded to} \quad \frac{\frac{p : t =_A s}{t \equiv s : A} =_E}{\mathbf{r}(s) : t =_A s} =_I$$

- ▶ This means that if we close \equiv by (the) η -equivalence (induced by this expansion) we have that $p \equiv \mathbf{r}(t) \equiv \mathbf{r}(s)$. Thus, every proof of $t =_A s$ collapses on a reflexivity proof \mathbf{r} , no matter that it is on t or on s .

Extensional Martin-Löf's ITT

- ▶ These rules allow one to pass from an equivalence by rewriting (\equiv) to a propositional identity ($=_A$), and *vice versa*.
- ▶ What is problematic is the *vice versa* direction: the rule $=_E$ makes one to loose the proof structure coded by the term p .
- ▶ This becomes more clear by considering that

$$\begin{array}{c} \mathcal{D} \\ p : t =_A s \end{array} \quad \text{can be (simply) expanded to} \quad \frac{\frac{\mathcal{D} \quad p : t =_A s}{t \equiv s : A} =_E}{\mathbf{r}(s) : t =_A s} =_I$$

- ▶ This means that if we close \equiv by (the) η -equivalence (induced by this expansion) we have that $p \equiv \mathbf{r}(t) \equiv \mathbf{r}(s)$. Thus, every proof of $t =_A s$ collapses on a reflexivity proof \mathbf{r} , no matter that it is on t or on s .
- ▶ We then have that $\llbracket t =_A s \rrbracket = \begin{cases} \mathfrak{P}_{t=_A s} & \text{if } t =_A s \text{ is provable} \\ \emptyset & \text{otherwise} \end{cases}$

Extensional Martin-Löf's ITT

- ▶ These rules allow one to pass from an equivalence by rewriting (\equiv) to a propositional identity ($=_A$), and *vice versa*.
- ▶ What is problematic is the *vice versa* direction: the rule $=_E$ makes one to loose the proof structure coded by the term p .
- ▶ This becomes more clear by considering that

$$\begin{array}{c} \mathcal{D} \\ p : t =_A s \end{array} \quad \text{can be (simply) expanded to} \quad \frac{\frac{\mathcal{D} \quad p : t =_A s}{t \equiv s : A} =_E}{\mathbf{r}(s) : t =_A s} =_I$$

- ▶ This means that if we close \equiv by (the) η -equivalence (induced by this expansion) we have that $p \equiv \mathbf{r}(t) \equiv \mathbf{r}(s)$. Thus, every proof of $t =_A s$ collapses on a reflexivity proof \mathbf{r} , no matter that it is on t or on s .
- ▶ We then have that $\llbracket t =_A s \rrbracket = \begin{cases} \{\mathbf{r}(t)\} & \text{if } t =_A s \text{ is provable} \\ \emptyset & \text{otherwise} \end{cases}$

Extensional Martin-Löf's ITT

- ▶ These rules allow one to pass from an equivalence (\equiv) to a propositional identity ($=_A$), and *vice versa*.
- ▶ What is problematic is the *vice versa* direction: the rule $=_E$ makes one to loose the proof structure coded by the term p .
- ▶ This becomes more clear by considering that

$$\begin{array}{c} \mathcal{D} \\ p : t =_A s \end{array} \quad \text{can be (simply) expanded to} \quad \frac{\frac{\mathcal{D} \quad p : t =_A s}{t \equiv s : A} =_E}{\mathbf{r}(s) : t =_A s} =_I$$

- ▶ This means that if we close \equiv by (the) η -equivalence (induced by this expansion) we have that $p \equiv \mathbf{r}(t) \equiv \mathbf{r}(s)$. Thus, every proof of $t =_A s$ collapses on a reflexivity proof \mathbf{r} , no matter that it is on t or on s .
- ▶ We then have that $\llbracket t =_A s \rrbracket = \begin{cases} \{\mathbf{r}(s)\} & \text{if } t =_A s \text{ is provable} \\ \emptyset & \text{otherwise} \end{cases}$

Extensional Martin-Löf's ITT

- ▶ These rules allow one to pass from an equivalence by rewriting (\equiv) to a propositional identity ($=_A$), and *vice versa*.
- ▶ What is problematic is the *vice versa* direction: the rule $=_E$ makes one to loose the proof structure coded by the term p .
- ▶ This becomes more clear by considering that

$$\begin{array}{c} \mathcal{D} \\ p : t =_A s \end{array} \quad \text{can be (simply) expanded to} \quad \frac{\frac{\mathcal{D} \quad p : t =_A s}{t \equiv s : A} =_E}{\mathbf{r}(s) : t =_A s} =_I$$

- ▶ This means that if we close \equiv by (the) η -equivalence (induced by this expansion) we have that $p \equiv \mathbf{r}(t) \equiv \mathbf{r}(s)$. Thus, every proof of $t =_A s$ collapses on a reflexivity proof \mathbf{r} , no matter that it is on t or on s .
- ▶ We then have that $\llbracket t =_A s \rrbracket = \begin{cases} 1 & \text{if } t =_A s \text{ is provable} \\ 0 & \text{otherwise} \end{cases}$

Extensional Martin-Löf's ITT

- ▶ These rules allow one to pass from an equivalence by rewriting (\equiv) to a propositional identity ($=_A$), and *vice versa*.
- ▶ What is problematic is the *vice versa* direction: the rule $=_E$ makes one to loose the proof structure coded by the term p .
- ▶ This becomes more clear by considering that

$$\begin{array}{c} \mathcal{D} \\ p : t =_A s \end{array} \quad \text{can be (simply) expanded to} \quad \frac{\frac{\mathcal{D} \quad p : t =_A s}{t \equiv s : A} =_E}{\mathbf{r}(s) : t =_A s} =_I$$

- ▶ This means that if we close \equiv by (the) η -equivalence (induced by this expansion) we have that $p \equiv \mathbf{r}(t) \equiv \mathbf{r}(s)$. Thus, every proof of $t =_A s$ collapses on a reflexivity proof \mathbf{r} , no matter that it is on t or on s .
- ▶ We then have that $\llbracket t =_A s \rrbracket = \begin{cases} \mathbf{T} & \text{if } t =_A s \text{ is provable} \\ \mathbf{F} & \text{otherwise} \end{cases}$

Extensional Martin-Löf's ITT

- ▶ These rules allow one to pass from an equivalence by rewriting (\equiv) to a propositional identity ($=_A$), and *vice versa*.
- ▶ What is problematic is the *vice versa* direction: the rule $=_E$ makes one to loose the proof structure coded by the term p .
- ▶ This becomes more clear by considering that

$$p : t =_A s \quad \text{can be (simply) expanded to} \quad \frac{\mathcal{D} \quad \frac{p : t =_A s}{t \equiv s : A} =_E}{\mathbf{r}(s) : t =_A s} =_I$$

- ▶ This means that if we close \equiv by (the) η -equivalence (induced by this expansion) we have that $p \equiv \mathbf{r}(t) \equiv \mathbf{r}(s)$. Thus, every proof of $t =_A s$ collapses on a reflexivity proof \mathbf{r} , no matter that it is on t or on s .
- ▶ We then have that $\llbracket t =_A s \rrbracket = \begin{cases} \mathbf{T} & \text{if } t =_A s \text{ is provable} \\ \mathbf{F} & \text{otherwise} \end{cases}$
- ▶ This corresponds to an *extensional* version of Martin-Löf's intuitionistic type theory.

Extensional Martin-Löf's ITT

- ▶ Where does Martin-Löf's $=_E$ rule come from?

Extensional Martin-Löf's ITT

- ▶ Where does Martin-Löf's $=_E$ rule come from?
- ▶ The (generalized) η -equivalence

$$\begin{array}{c} \mathcal{D} \\ [t = s] \\ \mathcal{D}'(t/y, s/z) \\ C(t/y, s/z) \end{array} \quad \equiv_{\eta} \quad \frac{\begin{array}{c} \mathcal{D} \quad \mathcal{D}'(x/y, x/z) \\ t = s \quad C(x/y, x/z) \end{array}}{C(t/y, s/z)} \quad \begin{array}{c} \overline{[x = x]} =_I \\ =_E \end{array}$$

Extensional Martin-Löf's ITT

- ▶ Where does Martin-Löf's $=_E$ rule come from?
- ▶ The (generalized) η -equivalence

$$\frac{\mathcal{D} \quad [t = s] \quad \mathcal{D}'(t/y, s/z)}{C(t/y, s/z)} \equiv_{\eta} \frac{\mathcal{D} \quad \frac{\overline{[x = x]} =_I \quad \mathcal{D}'(x/y, x/z)}{C(x/y, x/z)}}{C(t/y, s/z)} =_E$$

can be internalized at the level of proof-terms by the following η_{Conv} rule:

Extensional Martin-Löf's ITT

- Where does Martin-Löf's $=_E$ rule come from?
- The (generalized) η -equivalence

$$\frac{\mathcal{D} \quad [t = s]}{\mathcal{D}'(t/y, s/z)} \equiv_{\eta} \frac{\mathcal{D} \quad \overline{[x = x]} =_I \quad \mathcal{D}'(x/y, x/z)}{t = s \quad \frac{C(x/y, x/z)}{C(t/y, s/z)} =_E} \equiv_E$$

can be internalized at the level of proof-terms by the following η_{Conv} rule:

$$\frac{[x : A], [y : A, z : A, u : y =_A z] \quad d : t =_A s \quad d'(x/y, x/z, \mathbf{r}(x)/u) : C(x/y, x/z)}{J(t, s, d, \hat{x} : A.d'(x/y, x/z, \mathbf{r}(x)/u)) \equiv d'(t/y, s/z, d/u) : C(t/y, s/z)} \eta_{Conv}$$

where d corresponds to \mathcal{D} , and d' corresponds to \mathcal{D}' .

Extensional Martin-Löf's ITT

- Where does Martin-Löf's $=_E$ rule come from?
- The (generalized) η -equivalence

$$\frac{\mathcal{D} \quad [t = s] \quad \mathcal{D}'(t, s) \quad C(t, s)}{\equiv_{\eta}} \quad \frac{\overline{[x = x]} =_I \quad \mathcal{D}'(x, x) \quad C(x, x)}{t = s \quad C(t, s)} =_E$$

can be internalized at the level of proof-terms by the following η_{Conv} rule:

$$\frac{[x : A] \quad d : t =_A s \quad d'(x, x, \mathbf{r}(x)) : C(x, x)}{J(t, s, d, \hat{x} : A.d'(x, x, \mathbf{r}(x))) \equiv d'(t, s, d) : C(t, s)} \eta_{Conv}$$

where d corresponds to \mathcal{D} , and d' corresponds to \mathcal{D}' .

Extensional Martin-Löf's ITT

- ▶ Martin-Löf's $=_E$ rule can be derived via the η_{Conv} . Consider the following derivation:

Extensional Martin-Löf's ITT

- ▶ Martin-Löf's $=_E$ rule can be derived via the η_{Conv} . Consider the following derivation:

Extensional Martin-Löf's ITT

- Martin-Löf's $=_E$ rule can be derived via the η_{Conv} . Consider the following derivation:

$$\frac{
 \frac{
 \frac{d : t =_A s \quad [x/y : A]^1}{J(t, s, d, \hat{x} : A.x/y) \equiv t/y : A} \eta_{Conv} (1)
 }{t/y \equiv J(t, s, d, \hat{x} : A.x/y) : A} Sym
 \quad
 \frac{
 \frac{d : t =_A s \quad [x/z : A]^2}{J(t, s, d, \hat{x} : A.x/z) \equiv s/z : A} \eta_{Conv} (2)
 }{t/y \equiv s/z : A} Trans
 }{t/y \equiv s/z : A}$$

Extensional Martin-Löf's ITT

- Martin-Löf's $=_E$ rule can be derived via the η_{Conv} . Consider the following derivation:

$$\frac{
 \frac{
 \frac{d : t =_A s \quad [x : A]^1}{J(t, s, d, \hat{x} : A.x) \equiv t : A} \eta_{Conv} (1)
 }{t \equiv J(t, s, d, \hat{x} : A.x) : A} Sym
 \quad
 \frac{
 \frac{d : t =_A s \quad [x : A]^2}{J(t, s, d, \hat{x} : A.x) \equiv s : A} \eta_{Conv} (2)
 }{t \equiv s : A} Trans
 }{t \equiv s : A}$$

Extensional Martin-Löf's ITT

- ▶ Martin-Löf's $=_E$ rule can be derived via the η_{Conv} . Consider the following derivation:

$$\frac{\frac{\frac{d : t =_A s \quad [x : A]^1}{J(t, s, d, \hat{x} : A.x) \equiv t : A} \eta_{Conv} (1)}{t \equiv J(t, s, d, \hat{x} : A.x) : A} Sym \quad \frac{\frac{d : t =_A s \quad [x : A]^2}{J(t, s, d, \hat{x} : A.x) \equiv s : A} \eta_{Conv} (2)}{t \equiv s : A} Trans$$

- ▶ As shown by Streicher (1993) these two rules are in fact inter-derivable.

Extensional Martin-Löf's ITT

- ▶ Martin-Löf's $=_E$ rule can be derived via the η_{Conv} . Consider the following derivation:

$$\frac{\frac{\frac{d : t =_A s \quad [x : A]^1}{J(t, s, d, \hat{x} : A.x) \equiv t : A} \eta_{Conv} (1)}{t \equiv J(t, s, d, \hat{x} : A.x) : A} Sym \quad \frac{\frac{d : t =_A s \quad [x : A]^2}{J(t, s, d, \hat{x} : A.x) \equiv s : A} \eta_{Conv} (2)}{t \equiv s : A} Trans$$

- ▶ As shown by Streicher (1993) these two rules are in fact inter-derivable.
- ▶ The generalized expansion seems then to be the main responsible of the collapsing phenomena.

Possible solutions (1)

- ▶ Which solution should we adopt in order to avoid collapses?
- ▶ Drop stability, and work with harmony and deducibility of identical, i.e. work modulo $\equiv_{\beta\eta}$, where \equiv_{η} is induced by simple expansion and not the generalized one.

Problem 1: certain expressions would be governed by inference rules which could be considered as ambiguous definitions of these expressions, since different sets of elimination rules could be associated to the same set of introduction rules (cf. the problem of quantum disjunction).

Possible solutions (1)

- ▶ Which solution should we adopt in order to avoid collapses?
- ▶ Drop stability, and work with harmony and deducibility of identical, i.e. work modulo $\equiv_{\beta\eta}$, where \equiv_{η} is induced by simple expansion and not the generalized one.

Problem 1: certain expressions would be governed by inference rules which could be considered as ambiguous definitions of these expressions, since different sets of elimination rules could be associated to the same set of introduction rules (cf. the problem of quantum disjunction).

Possible solutions (1)

- ▶ Which solution should we adopt in order to avoid collapses?
- ▶ Drop stability, and work with harmony and deducibility of identical, i.e. work modulo $\equiv_{\beta\eta}$, where \equiv_{η} is induced by simple expansion and not the generalized one.

Problem 1: certain expressions would be governed by inference rules which could be considered as ambiguous definitions of these expressions, since different sets of elimination rules could be associated to the same set of introduction rules (cf. the problem of quantum disjunction).

Possible solutions (1)

- ▶ Which solution should we adopt in order to avoid collapses?
- ▶ Drop stability, and work with harmony and deducibility of identical, i.e. work modulo $\equiv_{\beta\eta}$, where \equiv_{η} is induced by simple expansion and not the generalized one.

Problem 1: certain expressions would be governed by inference rules which could be considered as ambiguous definitions of these expressions, since different sets of elimination rules could be associated to the same set of introduction rules (cf. the problem of quantum disjunction).

Possible solutions (1)

- ▶ Which solution should we adopt in order to avoid collapses?
- ▶ Drop stability, and work with harmony and deducibility of identical, i.e. work modulo $\equiv_{\beta\eta}$, where \equiv_{η} is induced by simple expansion and not the generalized one.

Problem 1: certain expressions would be governed by inference rules which could be considered as ambiguous definitions of these expressions, since different sets of elimination rules could be associated to the same set of introduction rules (cf. the problem of quantum disjunction).

Possible solutions (1)

Problem 2: consider the following derivations

Possible solutions (1)

Problem 2: consider the following derivations

$$\mathcal{D}_1 = \frac{\frac{N^1x}{Nx \rightarrow Nx} \rightarrow_I (1)}{\forall x(Nx \rightarrow Nx) \forall_I} \quad \text{and} \quad \frac{\frac{N^2x}{N0} N_{I_1} \quad \frac{N^1y}{Ny'} N_{I_2}}{\frac{Nx}{Nx \rightarrow Nx} \rightarrow (2)} N_E(1) = \mathcal{D}_2$$
$$\frac{\frac{Nx}{Nx \rightarrow Nx} \rightarrow (2)}{\forall x(Nx \rightarrow Nx) \forall_I}$$

Possible solutions (1)

Problem 2: consider the following derivations

$$\mathcal{D}_1 = \frac{\frac{N_x}{N_x \rightarrow N_x} \rightarrow_I (1)}{\forall x(N_x \rightarrow N_x)} \forall_I \not\equiv_{\beta} \frac{\frac{N_x}{N0} N_{I_1} \quad \frac{N_y}{N_{y'}} N_{I_2}}{\frac{N_x}{N_x \rightarrow N_x} \rightarrow (2)} N_E(1) = \mathcal{D}_2$$

Possible solutions (1)

Problem 2: consider the following derivations

$$\mathcal{D}_1 = \frac{\frac{N^1x}{Nx \rightarrow Nx} \rightarrow_I (1)}{\forall x(Nx \rightarrow Nx)} \forall_I \quad \text{simply expands to} \quad \frac{\frac{N^2x}{N0} N_{I_1} \quad \frac{N^1y}{Ny'} N_{I_2}}{\frac{Nx}{Nx \rightarrow Nx} \rightarrow (2)} N_E(1) = \mathcal{D}_2$$
$$\frac{\frac{Nx}{Nx \rightarrow Nx} \rightarrow (2)}{\forall x(Nx \rightarrow Nx)} \forall_I$$

Possible solutions (1)

Problem 2: consider the following derivations

$$\mathcal{D}_1 = \frac{\frac{N_x}{\forall x(Nx \rightarrow Nx)} \rightarrow_I (1)}{\forall x(Nx \rightarrow Nx)} \equiv_{\eta} \frac{\frac{N_x}{\forall x(Nx \rightarrow Nx)} \rightarrow_I (1)}{\frac{N_x}{\forall x(Nx \rightarrow Nx)} \rightarrow_I (1)} \equiv_{\eta} \mathcal{D}_2$$

- Working modulo \equiv_{η} means that \mathcal{D}_1 and \mathcal{D}_2 can be identified.

Possible solutions (1)

Problem 2: consider the following derivations

$$\mathcal{D}_1 = \frac{\frac{N^1x}{Nx \rightarrow Nx} \rightarrow_I (1)}{\forall x(Nx \rightarrow Nx) \forall_I} \equiv_{\eta} \frac{\frac{N^2x}{N0} N_{I_1} \quad \frac{N^1y}{Ny'} N_{I_2}}{\frac{Nx}{Nx \rightarrow Nx} \rightarrow (2)} N_E(1) = \mathcal{D}_2$$
$$\frac{\frac{Nx}{Nx \rightarrow Nx} \rightarrow (2)}{\forall x(Nx \rightarrow Nx) \forall_I}$$

- ▶ Working modulo \equiv_{η} means that \mathcal{D}_1 and \mathcal{D}_2 can be identified.
- ▶ Thanks to the Curry-Howard correspondance, derivations correspond to computable functions.

Possible solutions (1)

Problem 2: consider the following derivations

$$\begin{array}{c}
 \frac{u : Nx}{\lambda u. u : Nx \rightarrow Nx} \rightarrow_I (1) \\
 \frac{\lambda u. u : Nx \rightarrow Nx}{\lambda u. u : \forall x (Nx \rightarrow Nx)} \forall_I
 \end{array}
 \equiv_{\eta}
 \begin{array}{c}
 \frac{u : Nx \quad \frac{\frac{v : Ny}{S(v) : Ny'} N_{I_2}}{O : N0} N_{I_1}}{\text{It}(u, O, \hat{v}.S(v)) : Nx} N_E(1) \\
 \frac{\text{It}(u, O, \hat{v}.S(v)) : Nx}{\lambda u. \text{It}(u, O, \hat{v}.S(v)) : Nx \rightarrow Nx} \rightarrow (2) \\
 \frac{\lambda u. \text{It}(u, O, \hat{v}.S(v)) : Nx \rightarrow Nx}{\lambda u. \text{It}(u, O, \hat{v}.S(v)) : \forall x (Nx \rightarrow Nx)} \forall_I
 \end{array}$$

- ▶ Working modulo \equiv_{η} means that \mathcal{D}_1 and \mathcal{D}_2 can be identified.
- ▶ Thanks to the Curry-Howard correspondance, derivations correspond to computable functions.

Possible solutions (1)

Problem 2: consider the following derivations

$$\begin{array}{c}
 \frac{u : Nx}{\lambda u. u : Nx \rightarrow Nx} \rightarrow_I (1) \\
 \frac{\lambda u. u : Nx \rightarrow Nx}{\lambda u. u : \forall x (Nx \rightarrow Nx)} \forall_I
 \end{array}
 \equiv_{\eta}
 \begin{array}{c}
 \frac{u : Nx \quad \frac{\frac{v : Ny}{S(v) : Ny'} N_{I_2}}{O : N0} N_{I_1}}{\text{It}(u, O, \hat{v}.S(v)) : Nx} N_E(1) \\
 \frac{\text{It}(u, O, \hat{v}.S(v)) : Nx}{\lambda u. \text{It}(u, O, \hat{v}.S(v)) : Nx \rightarrow Nx} \rightarrow (2) \\
 \frac{\lambda u. \text{It}(u, O, \hat{v}.S(v)) : Nx \rightarrow Nx}{\lambda u. \text{It}(u, O, \hat{v}.S(v)) : \forall x (Nx \rightarrow Nx)} \forall_I
 \end{array}$$

- ▶ Working modulo \equiv_{η} means that \mathcal{D}_1 and \mathcal{D}_2 can be identified.
- ▶ Thanks to the Curry-Howard correspondance, derivations correspond to computable functions. Thus, the functions $\lambda u. u$ and $\lambda u. \text{It}(u, O, \hat{v}.S(v))$ would be identified.

Possible solutions (1)

Problem 2: consider the following derivations

$$\begin{array}{c}
 \frac{u : {}^1Nx}{\lambda u. u : Nx \rightarrow Nx} \rightarrow_I (1) \\
 \frac{\lambda u. u : Nx \rightarrow Nx}{\lambda u. u : \forall x(Nx \rightarrow Nx)} \forall_I
 \end{array}
 \equiv_{\eta}
 \begin{array}{c}
 \frac{u : {}^2Nx \quad \frac{\overline{\mathbf{O} : N0} \quad N_{I_1} \quad \frac{v : {}^1Ny}{\mathbf{S}(v) : Ny'} N_{I_2}}{\mathbf{It}(u, \mathbf{O}, \hat{v}. \mathbf{S}(v)) : Nx} N_{E(1)} \\
 \frac{\lambda u. \mathbf{It}(u, \mathbf{O}, \hat{v}. \mathbf{S}(v)) : Nx \rightarrow Nx}{\lambda u. \mathbf{It}(u, \mathbf{O}, \hat{v}. \mathbf{S}(v)) : \forall x(Nx \rightarrow Nx)} \rightarrow (2) \quad \forall_I
 \end{array}$$

- ▶ Working modulo \equiv_{η} means that \mathcal{D}_1 and \mathcal{D}_2 can be identified.
- ▶ Thanks to the Curry-Howard correspondance, derivations correspond to computable functions. Thus, the functions $\lambda u. u$ and $\lambda u. \mathbf{It}(u, \mathbf{O}, \hat{v}. \mathbf{S}(v))$ would be identified.
- ▶ The problem is that even if these two functions give the same outputs in presence of the same inputs, they compute in different ways.

Possible solutions (1)

Problem 2: consider the following derivations

$$\begin{array}{c}
 \frac{u : {}^1Nx}{\lambda u. u : Nx \rightarrow Nx} \rightarrow_I (1) \\
 \hline
 \lambda u. u : \forall x (Nx \rightarrow Nx) \quad \forall_I
 \end{array}
 \equiv_{\eta}
 \begin{array}{c}
 \frac{u : {}^2Nx \quad \frac{\frac{\frac{v : {}^1Ny}{S(v) : Ny'} N_{I_2}}{O : N0} N_{I_1}}{It(u, O, \hat{v}.S(v)) : Nx} N_{I_2}}{\lambda u. It(u, O, \hat{v}.S(v)) : Nx \rightarrow Nx} \rightarrow (2) \\
 \hline
 \lambda u. It(u, O, \hat{v}.S(v)) : \forall x (Nx \rightarrow Nx) \quad \forall_I
 \end{array}$$

- ▶ Working modulo \equiv_{η} means that \mathcal{D}_1 and \mathcal{D}_2 can be identified.
- ▶ Thanks to the Curry-Howard correspondance, derivations correspond to computable functions. Thus, the functions $\lambda u. u$ and $\lambda u. It(u, O, \hat{v}.S(v))$ would be identified.
- ▶ The problem is that even if these two functions give the same outputs in presence of the same inputs, they compute in different ways. E.g. their time complexity is different: constant vs. linear.

Possible solutions (1)

Problem 2: consider the following derivations

$$\begin{array}{c}
 \frac{u : {}^1Nx}{\lambda u. u : Nx \rightarrow Nx} \rightarrow_I (1) \\
 \frac{\lambda u. u : Nx \rightarrow Nx}{\lambda u. u : \forall x(Nx \rightarrow Nx)} \forall_I
 \end{array}
 \equiv_{\eta}
 \begin{array}{c}
 \frac{u : {}^2Nx \quad \frac{\frac{\frac{}{\mathbf{O} : N0} N_{I_1}}{\mathbf{S}(v) : Ny'} N_{I_2}}{\mathbf{It}(u, \mathbf{O}, \hat{v}. \mathbf{S}(v)) : Nx} N_{E(1)}}{\lambda u. \mathbf{It}(u, \mathbf{O}, \hat{v}. \mathbf{S}(v)) : Nx \rightarrow Nx} \rightarrow (2) \\
 \frac{\lambda u. \mathbf{It}(u, \mathbf{O}, \hat{v}. \mathbf{S}(v)) : Nx \rightarrow Nx}{\lambda u. \mathbf{It}(u, \mathbf{O}, \hat{v}. \mathbf{S}(v)) : \forall x(Nx \rightarrow Nx)} \forall_I
 \end{array}$$

- ▶ Working modulo \equiv_{η} means that \mathcal{D}_1 and \mathcal{D}_2 can be identified.
- ▶ Thanks to the Curry-Howard correspondance, derivations correspond to computable functions. Thus, the functions $\lambda u. u$ and $\lambda u. \mathbf{It}(u, \mathbf{O}, \hat{v}. \mathbf{S}(v))$ would be identified.
- ▶ The problem is that even if these two functions give the same outputs in presence of the same inputs, they compute in different ways. E.g. their time complexity is different: constant vs. linear.
- ▶ These two functions are thus intensionnaly different, since they correspond to different algorithms.

Possible solutions (1)

Problem 2: consider the following derivations

$$\begin{array}{c}
 \frac{u : {}^1Nx}{\lambda u. u : Nx \rightarrow Nx} \rightarrow_I (1) \\
 \frac{\lambda u. u : Nx \rightarrow Nx}{\lambda u. u : \forall x(Nx \rightarrow Nx)} \forall_I
 \end{array}
 \equiv_{\eta}
 \begin{array}{c}
 \frac{u : {}^2Nx \quad \frac{\frac{\frac{v : {}^1Ny}{S(v) : Ny'} N_{I_2}}{O : N0} N_{I_1}}{It(u, O, \hat{v}.S(v)) : Nx} N_{E(1)}}{\lambda u. It(u, O, \hat{v}.S(v)) : Nx \rightarrow Nx} \rightarrow (2) \\
 \frac{\lambda u. It(u, O, \hat{v}.S(v)) : Nx \rightarrow Nx}{\lambda u. It(u, O, \hat{v}.S(v)) : \forall x(Nx \rightarrow Nx)} \forall_I
 \end{array}$$

- ▶ Working modulo \equiv_{η} means that \mathcal{D}_1 and \mathcal{D}_2 can be identified.
- ▶ Thanks to the Curry-Howard correspondance, derivations correspond to computable functions. Thus, the functions $\lambda u. u$ and $\lambda u. It(u, O, \hat{v}.S(v))$ would be identified.
- ▶ The problem is that even if these two functions give the same outputs in presence of the same inputs, they compute in different ways. E.g. their time complexity is different: constant vs. linear.
- ▶ These two functions are thus intensionnaly different, since they correspond to different algorithms. They should then be kept distinct.

Possible solutions (2): Propositional case

- ▶ In order to work within a genuine intensional account, it seems then natural to ban any kind of expansion, and consider only \equiv_β .
- ▶ This solution is not so restrictive as it could seem. Consider the following situation where \mathcal{D} is a closed derivation:

$$\begin{array}{ccc}
 \frac{\mathcal{D} \quad \frac{\frac{A}{B} \xrightarrow{m} A \Rightarrow B}{A \rightarrow B} \xrightarrow{n} A \Rightarrow B}{A \rightarrow B} \xrightarrow{(m)} \rightsquigarrow \beta & \frac{\frac{A}{B} \xrightarrow{l} A \Rightarrow B}{A \rightarrow B} \xrightarrow{(l)} \rightsquigarrow \beta & \frac{\frac{A}{B} \xrightarrow{m} A \Rightarrow B}{A \rightarrow B} \xrightarrow{(m)} \rightsquigarrow \beta \\
 \parallel & \parallel & \parallel \\
 \mathcal{D}_1 & \mathcal{D}_2 & \mathcal{D}_3
 \end{array}$$

- ▶ \mathcal{D}_3 is obtained from \mathcal{D}_2 by β -reduction. But, \mathcal{D}_2 also corresponds to the simple η -expansion of \mathcal{D}_3 (cf. Martin-Löf 1975; Tranchini, Pistone & Petrolo 2016).
- ▶ Thus, with β -reduction it is possible to recover some instances of η -expansion which do not lead to any collapse.

Possible solutions (2): Propositional case

- In order to work within a genuine intensional account, it seems then natural to ban any kind of expansion, and consider only \equiv_β .
- This solution is not so restrictive as it could seem. Consider the following situation where \mathcal{D} is a closed derivation:

$$\begin{array}{ccc}
 \frac{\mathcal{D} \quad \frac{\frac{A}{B} \xrightarrow{m} A \Rightarrow B}{A \rightarrow B} \xrightarrow{n} B}{A \rightarrow B} \xrightarrow{I(m)} \rightsquigarrow \beta & \frac{\frac{A}{B} \xrightarrow{l} A \Rightarrow B}{A \rightarrow B} \xrightarrow{I(l)} \rightsquigarrow \beta & \frac{\frac{A}{B} \xrightarrow{m} A \Rightarrow B}{A \rightarrow B} \xrightarrow{I(m)} \rightsquigarrow \beta \\
 \parallel & \parallel & \parallel \\
 \mathcal{D}_1 & \mathcal{D}_2 & \mathcal{D}_3
 \end{array}$$

- \mathcal{D}_3 is obtained from \mathcal{D}_2 by β -reduction. But, \mathcal{D}_2 also corresponds to the simple η -expansion of \mathcal{D}_3 (cf. Martin-Löf 1975; Tranchini, Pistone & Petrolo 2016).
- Thus, with β -reduction it is possible to recover some instances of η -expansion which do not lead to any collapse.

Possible solutions (2): Propositional case

- In order to work within a genuine intensional account, it seems then natural to ban any kind of expansion, and consider only \equiv_β .
- This solution is not so restrictive as it could seem. Consider the following situation where \mathcal{D} is a closed derivation:

$$\begin{array}{ccc}
 \frac{\mathcal{D} \quad \frac{\frac{A}{B} \xRightarrow{m} A \Rightarrow B}{A \rightarrow B} \xrightarrow{I(m)} \sim\sim\beta}{A \rightarrow B} \xrightarrow{E(n)} & \frac{\frac{\frac{A}{B} \xRightarrow{l} A \Rightarrow B}{A \rightarrow B} \xrightarrow{I(l)} \sim\sim\beta}{A \rightarrow B} \xrightarrow{E(n)} & \frac{\frac{\frac{A}{B} \xRightarrow{l} A \Rightarrow B}{A \rightarrow B} \xrightarrow{I(l)}}{A \rightarrow B} \xrightarrow{E(n)} \\
 \parallel & \parallel & \parallel \\
 \mathcal{D}_1 & \mathcal{D}_2 & \mathcal{D}_3
 \end{array}$$

- D_3 is obtained from D_2 by β -reduction.
But, D_2 also corresponds to the simple η -expansion of \mathcal{D}_3 (cf. Martin-Löf 1975; Tranchini, Pistone & Petrolo 2016).
- Thus, with β -reduction it is possible to recover some instances of η -expansion which do not lead to any collapse.

Possible solutions (2): Propositional case

- ▶ In order to work within a genuine intensional account, it seems then natural to ban any kind of expansion, and consider only \equiv_β .
- ▶ This solution is not so restrictive as it could seem. Consider the following situation where \mathcal{D} is a closed derivation:

$$\begin{array}{ccc}
 \frac{\mathcal{D} \quad \frac{\frac{m}{A} \quad A \Rightarrow B}{B} \rightarrow_I (m)}{A \rightarrow B \quad A \rightarrow B} \rightarrow_E (n) & \rightsquigarrow_\beta & \frac{\frac{l}{A} \quad \mathcal{D}^*}{B} \rightarrow_I (l) \quad \frac{\frac{m}{A} \quad A \Rightarrow B}{B} \rightarrow_I (m)}{A \rightarrow B \quad A \rightarrow B} \rightarrow_E (n) \\
 \parallel & & \parallel \\
 \mathcal{D}_1 & & \mathcal{D}_2
 \end{array}$$

$$\begin{array}{ccc}
 & & \frac{l}{A} \quad \mathcal{D}^* \\
 & & B \\
 & & \rightarrow_I (l)
 \end{array}$$

$$\begin{array}{ccc}
 & & \parallel \\
 & & \mathcal{D}_3
 \end{array}$$

- ▶ \mathcal{D}_3 is obtained from \mathcal{D}_2 by β -reduction.
But, \mathcal{D}_2 also corresponds to the simple η -expansion of \mathcal{D}_3 (cf. Martin-Löf 1975; Tranchini, Pistone & Petrolo 2016).
- ▶ Thus, with β -reduction it is possible to recover some instances of η -expansion which do not lead to any collapse.

Possible solutions (2): Propositional case

- ▶ In order to work within a genuine intensional account, it seems then natural to ban any kind of expansion, and consider only \equiv_β .
- ▶ This solution is not so restrictive as it could seem. Consider the following situation where \mathcal{D} is a closed derivation:

$$\begin{array}{ccc}
 \frac{\mathcal{D} \quad \frac{\frac{m}{A} \quad A \Rightarrow B}{B} \rightarrow_I (m)}{A \rightarrow B \quad A \rightarrow B} \rightarrow_E (n) & \rightsquigarrow_\beta & \frac{\frac{l}{A} \quad \mathcal{D}^* \quad B}{A \rightarrow B} \rightarrow_I (l) \quad \frac{\frac{m}{A} \quad A \Rightarrow B}{B} \rightarrow_I (m)}{A \rightarrow B \quad A \rightarrow B} \rightarrow_E (n) \\
 \parallel & & \parallel \\
 \mathcal{D}_1 & & \mathcal{D}_2
 \end{array}$$

$$\begin{array}{ccc}
 & & \frac{l}{A} \quad \mathcal{D}^* \quad B}{A \rightarrow B} \rightarrow_I (l) \\
 & & \parallel \\
 & & \mathcal{D}_3
 \end{array}$$

- ▶ \mathcal{D}_3 is obtained from \mathcal{D}_2 by β -reduction.
But, \mathcal{D}_2 also corresponds to the simple η -expansion of \mathcal{D}_3 (cf. Martin-Löf 1975; Tranchini, Pistone & Petrolo 2016).
- ▶ Thus, with β -reduction it is possible to recover some instances of η -expansion which do not lead to any collapse.

Possible solutions (2): Propositional case

- In order to work within a genuine intensional account, it seems then natural to ban any kind of expansion, and consider only \equiv_β .
- This solution is not so restrictive as it could seem. Consider the following situation where \mathcal{D} is a closed derivation:

$$\begin{array}{ccc}
 \frac{\mathcal{D} \quad \frac{\frac{m}{A} \quad A \Rightarrow B}{B} \rightarrow_I (m)}{A \rightarrow B \quad A \rightarrow B} \rightarrow_E (n) & \rightsquigarrow_\beta & \frac{\frac{l}{A} \quad \mathcal{D}^*}{B} \rightarrow_I (l) \quad \frac{\frac{m}{A} \quad A \Rightarrow B}{B} \rightarrow_I (m)}{A \rightarrow B \quad A \rightarrow B} \rightarrow_E (n) \\
 \parallel & & \parallel \\
 \mathcal{D}_1 & & \mathcal{D}_2
 \end{array}$$

$$\begin{array}{ccc}
 & & \frac{l}{A} \quad \mathcal{D}^* \\
 & & B \\
 & & \rightarrow_I (l)
 \end{array}$$

$$\begin{array}{ccc}
 & & \parallel \\
 & & \mathcal{D}_3
 \end{array}$$

- D_3 is obtained from D_2 by β -reduction.
But, D_2 also corresponds to the simple η -expansion of \mathcal{D}_3 (cf. Martin-Löf 1975; Tranchini, Pistone & Petrolo 2016).
- Thus, with β -reduction it is possible to recover some instances of η -expansion which do not lead to any collapse.

Possible solutions (2): Propositional case

- In order to work within a genuine intensional account, it seems then natural to ban any kind of expansion, and consider only \equiv_β .
- This solution is not so restrictive as it could seem. Consider the following situation where \mathcal{D} is a closed derivation:

$$\begin{array}{ccc}
 \frac{\mathcal{D} \quad \frac{\frac{A}{B} \xRightarrow{m} A \Rightarrow B}{A \rightarrow B} \xrightarrow{I (m)} \rightsquigarrow_\beta \quad \frac{A}{B} \xRightarrow{m} A \Rightarrow B}{A \rightarrow B} \xrightarrow{E (n)} \rightsquigarrow_\beta}{A \rightarrow B} \parallel \mathcal{D}_1 & \frac{\frac{l}{A} \mathcal{D}^* \quad \frac{A}{B} \xRightarrow{m} A \Rightarrow B}{A \rightarrow B} \xrightarrow{I (l)} \rightsquigarrow_\beta \quad \frac{A}{B} \xRightarrow{m} A \Rightarrow B}{A \rightarrow B} \xrightarrow{E (n)} \rightsquigarrow_\beta}{A \rightarrow B} \parallel \mathcal{D}_2 & \frac{l}{A} \mathcal{D}^* \quad \frac{A}{B} \xRightarrow{m} A \Rightarrow B}{A \rightarrow B} \xrightarrow{I (l)} \rightsquigarrow_\beta \quad \frac{A}{B} \xRightarrow{m} A \Rightarrow B}{A \rightarrow B} \xrightarrow{E (n)} \rightsquigarrow_\beta}{A \rightarrow B} \parallel \mathcal{D}_3
 \end{array}$$

- D_3 is obtained from D_2 by β -reduction.
But, D_2 also corresponds to the simple η -expansion of \mathcal{D}_3 (cf. Martin-Löf 1975; Tranchini, Pistone & Petrolo 2016).
- Thus, with β -reduction it is possible to recover some instances of η -expansion which do not lead to any collapse.

Possible solutions (2): Propositional case

- In order to work within a genuine intensional account, it seems then natural to ban any kind of expansion, and consider only \equiv_β .
- This solution is not so restrictive as it could seem. Consider the following situation where \mathcal{D} is a closed derivation:

$$\begin{array}{ccc}
 \frac{\mathcal{D} \quad \frac{\frac{m}{A} \quad A \Rightarrow B}{B} \rightarrow_I (m)}{A \rightarrow B \quad A \rightarrow B} \rightarrow_E (n) & \rightsquigarrow_\beta & \frac{\frac{l}{A} \quad \frac{m}{B} \quad A \Rightarrow B}{B} \rightarrow_I (l) \quad \frac{A \rightarrow B}{A \rightarrow B} \rightarrow_E (n)}{A \rightarrow B} \rightarrow_I (l) \\
 \parallel & & \parallel \\
 \mathcal{D}_1 & & \mathcal{D}_2
 \end{array}$$

$$\begin{array}{ccc}
 \frac{\frac{l}{A} \quad \frac{m}{B} \quad A \Rightarrow B}{B} \rightarrow_I (l) & \rightsquigarrow_\beta & \frac{l}{A} \quad \frac{B}{B} \rightarrow_I (l)}{A \rightarrow B} \rightarrow_I (l) \\
 \parallel & & \parallel \\
 \mathcal{D}_2 & & \mathcal{D}_3
 \end{array}$$

- D_3 is obtained from D_2 by β -reduction.
But, D_2 also corresponds to the simple η -expansion of \mathcal{D}_3 (cf. Martin-Löf 1975; Tranchini, Pistone & Petrolo 2016).
- Thus, with β -reduction it is possible to recover some instances of η -expansion which do not lead to any collapse.
- This imposes a hierarchical order on the principles used in proof-theoretic semantics: deducibility of identicals (and possibly stability) has not to be taken as an independent principle, but has to be induced by harmony.

Possible solutions (2): Propositional case

- In order to work within a genuine intensional account, it seems then natural to ban any kind of expansion, and consider only \equiv_β .
- This solution is not so restrictive as it could seem. Consider the following situation where \mathcal{D} is a closed derivation:

$$\begin{array}{ccc}
 \frac{\mathcal{D} \quad \frac{\frac{m}{A} \quad A \Rightarrow B}{B} \rightarrow_I (m)}{A \rightarrow B \quad A \rightarrow B} \rightarrow_E (n) & \rightsquigarrow_\beta & \frac{\frac{l}{A} \quad B}{A \rightarrow B} \rightarrow_I (l) \\
 \parallel & & \parallel \\
 \mathcal{D}_1 & & \mathcal{D}_2
 \end{array}$$

$$\begin{array}{ccc}
 \frac{\frac{m}{A} \quad A \Rightarrow B}{B} \rightarrow_I (m) & \rightsquigarrow_\beta & \frac{l}{A} \quad B \\
 \frac{A \rightarrow B \quad A \rightarrow B}{A \rightarrow B} \rightarrow_E (n) & & \frac{B}{A \rightarrow B} \rightarrow_I (l) \\
 \parallel & & \parallel \\
 \mathcal{D}_2 & & \mathcal{D}_3
 \end{array}$$

- \mathcal{D}_3 is obtained from \mathcal{D}_2 by β -reduction.
But, \mathcal{D}_2 also corresponds to the simple η -expansion of \mathcal{D}_3 (cf. Martin-Löf 1975; Tranchini, Pistone & Petrolo 2016).
- Thus, with β -reduction it is possible to recover some instances of η -expansion which do not lead to any collapse.
- In other words, the no less conditions have to be induced after having established the no more conditions.

Possible solutions (2): Identity case

- ▶ Not imposing η -equivalence from the beginning corresponds – in the framework of Martin-Löf's ITT – to drop the η_{Conv} rule allowing one to conclude:

$$J(t, s, d, \hat{x} : A.d'(x, x, \mathbf{r}(x))) \equiv d'(t, s, d) : C(t, s)$$

from a proof d of $t =_A s$ and a proof $d'(x, x, \mathbf{r}(x))$ of $C(x, x)$.

- ▶ The collapse of all proofs of $t =_A s$, for all t and s , is thus avoided.
- ▶ However, even without the η_{Conv} , it is still possible to have a *propositional version* of the η relation, that is, to find a p such that

$$p : J(t, s, d, \hat{x} : A.d'(x, x, \mathbf{r}(x))) =_{C(t, s)} d'(t, s, d)$$

(see Hofmann & Streicher 1998).

Possible solutions (2): Identity case

- ▶ Not imposing η -equivalence from the beginning corresponds – in the framework of Martin-Löf's ITT – to drop the η_{Conv} rule allowing one to conclude:

$$J(t, s, d, \hat{x} : A.d'(x, x, \mathbf{r}(x))) \equiv d'(t, s, d) : C(t, s)$$

from a proof d of $t =_A s$ and a proof $d'(x, x, \mathbf{r}(x))$ of $C(x, x)$.

- ▶ The collapse of all proofs of $t =_A s$, for all t and s , is thus avoided.
- ▶ However, even without the η_{Conv} , it is still possible to have a *propositional version* of the η relation, that is, to find a p such that

$$p : J(t, s, d, \hat{x} : A.d'(x, x, \mathbf{r}(x))) =_{C(t,s)} d'(t, s, d)$$

(see Hofmann & Streicher 1998).

Possible solutions (2): Identity case

- ▶ Not imposing η -equivalence from the beginning corresponds – in the framework of Martin-Löf's ITT – to drop the η_{Conv} rule allowing one to conclude:

$$J(t, s, d, \hat{x} : A.d'(x, x, \mathbf{r}(x))) \equiv d'(t, s, d) : C(t, s)$$

from a proof d of $t =_A s$ and a proof $d'(x, x, \mathbf{r}(x))$ of $C(x, x)$.

- ▶ The collapse of all proofs of $t =_A s$, for all t and s , is thus avoided.
- ▶ However, even without the η_{Conv} , it is still possible to have a *propositional version* of the η relation, that is, to find a p such that

$$p : J(t, s, d, \hat{x} : A.d'(x, x, \mathbf{r}(x))) =_{C(t, s)} d'(t, s, d)$$

(see Hofmann & Streicher 1998).

Possible solutions (2): Identity case

- ▶ Not imposing η -equivalence from the beginning corresponds – in the framework of Martin-Löf's ITT – to drop the η_{Conv} rule allowing one to conclude:

$$J(t, s, d, \hat{x} : A.d'(x, x, \mathbf{r}(x))) \equiv d'(t, s, d) : C(t, s)$$

from a proof d of $t =_A s$ and a proof $d'(x, x, \mathbf{r}(x))$ of $C(x, x)$.

- ▶ The collapse of all proofs of $t =_A s$, for all t and s , is thus avoided.
- ▶ However, even without the η_{Conv} , it is still possible to have a *propositional version* of the η relation, that is, to find a p such that

$$p : J(t, s, d, \hat{x} : A.d'(x, x, \mathbf{r}(x))) =_{C(t, s)} d'(t, s, d)$$

(see Hofmann & Streicher 1998).

Possible solutions (2): Identity case

- ▶ Not imposing η -equivalence from the beginning corresponds – in the framework of Martin-Löf's ITT – to drop the η_{Conv} rule allowing one to conclude:

$$J(t, s, d, \hat{x} : A.d'(x, x, \mathbf{r}(x))) \equiv d'(t, s, d) : C(t, s)$$

from a proof d of $t =_A s$ and a proof $d'(x, x, \mathbf{r}(x))$ of $C(x, x)$.

- ▶ The collapse of all proofs of $t =_A s$, for all t and s , is thus avoided.
- ▶ However, even without the η_{Conv} , it is still possible to have a *propositional version* of the η relation, that is, to find a p such that

$$p : J(t, s, d, \hat{x} : A.d'(x, x, \mathbf{r}(x))) =_{C(t,s)} d'(t, s, d)$$

(see Hofmann & Streicher 1998).

Intensionality and extensionality again

- ▶ Similarly, we have that the judgment

$$\lambda u.u \equiv \lambda u.\text{It}(u, \text{O}, \hat{v}.S(v)) : \forall x(Nx \rightarrow Nx)$$

is not assertable, since without simple expansion there is no way to rewrite $\lambda u.u$ on $\lambda u.\text{It}(u, \text{O}, \hat{v}.S(v))$, and *vice versa* (as we said, the two terms are not β -equivalent).

- ▶ On the contrary, the judgment

$$p : \lambda u.u =_{\forall x(Nx \rightarrow Nx)} \lambda u.\text{It}(u, \text{O}, \hat{v}.S(v))$$

is assertable, for a certain p .

- ▶ This means that even if we do not have a procedure for transforming $\lambda u.u$ into $\lambda u.\text{It}(u, \text{O}, \hat{v}.S(v))$, and *vice versa*, we can however *say* that they are identical.
- ▶ The reason is that $\lambda u.u$ and $\lambda u.\text{It}(u, \text{O}, \hat{v}.S(v))$ correspond to different algorithms, but they are denotationally equivalent.
- ▶ Thus, while the relation \equiv is intensional, the identity predicate $=_A$ (for a certain A) is extensional.

Intensionality and extensionality again

- ▶ Similarly, we have that the judgment

$$\lambda u.u \equiv \lambda u.\text{It}(u, \text{O}, \hat{v}.\text{S}(v)) : \forall x(Nx \rightarrow Nx)$$

is not assertable, since without simple expansion there is no way to rewrite $\lambda u.u$ on $\lambda u.\text{It}(u, \text{O}, \hat{v}.\text{S}(v))$, and *vice versa* (as we said, the two terms are not β -equivalent).

- ▶ On the contrary, the judgment

$$p : \lambda u.u =_{\forall x(Nx \rightarrow Nx)} \lambda u.\text{It}(u, \text{O}, \hat{v}.\text{S}(v))$$

is assertable, for a certain p .

- ▶ This means that even if we do not have a procedure for transforming $\lambda u.u$ into $\lambda u.\text{It}(u, \text{O}, \hat{v}.\text{S}(v))$, and *vice versa*, we can however *say* that they are identical.
- ▶ The reason is that $\lambda u.u$ and $\lambda u.\text{It}(u, \text{O}, \hat{v}.\text{S}(v))$ correspond to different algorithms, but they are denotationally equivalent.
- ▶ Thus, while the relation \equiv is intensional, the identity predicate $=_A$ (for a certain A) is extensional.

Intensionality and extensionality again

- ▶ Similarly, we have that the judgment

$$\lambda u.u \equiv \lambda u.\text{It}(u, \text{O}, \hat{v}.\text{S}(v)) : \forall x(Nx \rightarrow Nx)$$

is not assertable, since without simple expansion there is no way to rewrite $\lambda u.u$ on $\lambda u.\text{It}(u, \text{O}, \hat{v}.\text{S}(v))$, and *vice versa* (as we said, the two terms are not β -equivalent).

- ▶ On the contrary, the judgment

$$p : \lambda u.u =_{\forall x(Nx \rightarrow Nx)} \lambda u.\text{It}(u, \text{O}, \hat{v}.\text{S}(v))$$

is assertable, for a certain p .

- ▶ This means that even if we do not have a procedure for transforming $\lambda u.u$ into $\lambda u.\text{It}(u, \text{O}, \hat{v}.\text{S}(v))$, and *vice versa*, we can however *say* that they are identical.
- ▶ The reason is that $\lambda u.u$ and $\lambda u.\text{It}(u, \text{O}, \hat{v}.\text{S}(v))$ correspond to different algorithms, but they are denotationally equivalent.
- ▶ Thus, while the relation \equiv is intensional, the identity predicate $=_A$ (for a certain A) is extensional.

Intensionality and extensionality again

- ▶ Similarly, we have that the judgment

$$\lambda u.u \equiv \lambda u.\text{It}(u, \text{O}, \hat{v}.\text{S}(v)) : \forall x(Nx \rightarrow Nx)$$

is not assertable, since without simple expansion there is no way to rewrite $\lambda u.u$ on $\lambda u.\text{It}(u, \text{O}, \hat{v}.\text{S}(v))$, and *vice versa* (as we said, the two terms are not β -equivalent).

- ▶ On the contrary, the judgment

$$p : \lambda u.u =_{\forall x(Nx \rightarrow Nx)} \lambda u.\text{It}(u, \text{O}, \hat{v}.\text{S}(v))$$

is assertable, for a certain p .

- ▶ This means that even if we do not have a procedure for transforming $\lambda u.u$ into $\lambda u.\text{It}(u, \text{O}, \hat{v}.\text{S}(v))$, and *vice versa*, we can however *say* that they are identical.
- ▶ The reason is that $\lambda u.u$ and $\lambda u.\text{It}(u, \text{O}, \hat{v}.\text{S}(v))$ correspond to different algorithms, but they are denotationally equivalent.
- ▶ Thus, while the relation \equiv is intensional, the identity predicate $=_A$ (for a certain A) is extensional.

Intensionality and extensionality again

- ▶ Similarly, we have that the judgment

$$\lambda u.u \equiv \lambda u.\text{It}(u, \text{O}, \hat{v}.\text{S}(v)) : \forall x(Nx \rightarrow Nx)$$

is not assertable, since without simple expansion there is no way to rewrite $\lambda u.u$ on $\lambda u.\text{It}(u, \text{O}, \hat{v}.\text{S}(v))$, and *vice versa* (as we said, the two terms are not β -equivalent).

- ▶ On the contrary, the judgment

$$p : \lambda u.u =_{\forall x(Nx \rightarrow Nx)} \lambda u.\text{It}(u, \text{O}, \hat{v}.\text{S}(v))$$

is assertable, for a certain p .

- ▶ This means that even if we do not have a procedure for transforming $\lambda u.u$ into $\lambda u.\text{It}(u, \text{O}, \hat{v}.\text{S}(v))$, and *vice versa*, we can however *say* that they are identical.
- ▶ The reason is that $\lambda u.u$ and $\lambda u.\text{It}(u, \text{O}, \hat{v}.\text{S}(v))$ correspond to different algorithms, but they are denotationally equivalent.
- ▶ Thus, while the relation \equiv is intensional, the identity predicate $=_A$ (for a certain A) is extensional.

The beginning of homotopy type theory

- ▶ More precisely, Hoffman & Streicher (1998) set up a groupoid interpretation of type theory in which they showed that there exists an equivalent relation \simeq on the set of terms of type $t =_A s$ such that

$f \simeq g$ iff there exists a term p such that $p : f =_{(t=_A s)} g$.

- ▶ In terms of spaces, f and g correspond to paths from the point a to the point b in the space A and p corresponds to a homotopy relative to their endpoints from f to g .
- ▶ These ideas have been one of the starting points of the project of homotopy type theory.

The beginning of homotopy type theory

- ▶ More precisely, Hoffman & Streicher (1998) set up a groupoid interpretation of type theory in which they showed that there exists an equivalent relation \simeq on the set of terms of type $t =_A s$ such that

$f \simeq g$ iff there exists a term p such that $p : f =_{(t=_A s)} g$.

- ▶ In terms of spaces, f and g correspond to paths from the point a to the point b in the space A and p corresponds to a homotopy relative to their endpoints from f to g .
- ▶ These ideas have been one of the starting points of the project of homotopy type theory.

The beginning of homotopy type theory

- ▶ More precisely, Hoffman & Streicher (1998) set up a groupoid interpretation of type theory in which they showed that there exists an equivalent relation \simeq on the set of terms of type $t =_A s$ such that

$f \simeq g$ iff there exists a term p such that $p : f =_{(t=_A s)} g$.

- ▶ In terms of spaces, f and g correspond to paths from the point a to the point b in the space A and p corresponds to a homotopy relative to their endpoints from f to g .
- ▶ These ideas have been one of the starting points of the project of homotopy type theory.

The beginning of homotopy type theory

- More precisely, Hoffman & Streicher (1998) set up a groupoid interpretation of type theory in which they showed that there exists an equivalent relation \simeq on the set of terms of type $t =_A s$ such that

$f \simeq g$ iff there exists a term p such that $p : f =_{(t=_A s)} g$.

- In terms of spaces, f and g correspond to paths from the point a to the point b in the space A and p corresponds to a homotopy relative to their endpoints from f to g .
- These ideas have been one of the starting points of the project of homotopy type theory.

References (1)



Dummett, M. (1991). *The Logical Basis of Metaphysics*. London: Duckworth.



Francez, N. & Dyckhoff, R. (2012). A note on harmony. *Journal of Philosophical Logic*, 41: 613–628.



Hacking, I. (1979). What is logic?. *The Journal of Philosophy*, 76: 285–319.



Hofmann, M. & Streicher, T. (1998). The groupoid interpretation of type theory. In G. Sambin and J. Smith (eds.), *Twenty-Five Years of Constructive Type Theory*, 83–111. Oxford: OUP.



Jacinto, B. & Read, S. (2016) General-elimination stability. *Studia Logica*, Online First.

References (2)



Martin-Löf, P. (1971). Hauptsatz for the intuitionistic theory of iterated inductive definitions. In J.E. Fenstad (ed.), *Proceedings of the Second Scandinavian Logic Symposium*, 179–216. Amsterdam: North-Holland.



Martin-Löf, P. (1975). About models for intuitionistic type theories and the notion of definitional equality. In S. Kanger (ed.), *Proceedings of the Third Scandinavian Logic Symposium*, 81–109. Amsterdam: North-Holland.



Martin-Löf, P. (1984). *Intuitionistic Type Theory*. Naples: Bibliopolis.



Naibo, A. & Petrolo, M. (2015). Are uniqueness and deducibility of identicals the same?. *Theoria*, 81: 143–181.










Negri, S. & von Plato, J. (2001). *Structural Proof Theory*. Cambridge: CUP.



Pfenning, F. & Davies, R. (2001). A judgmental reconstruction of modal logic. *Mathematical Structures in Computer Science*, 11: 511–540.

References (3)

-  Prawitz, D. (1965). *Natural Deduction: A proof-theoretical study*. Stockholm: Almqvist & Wiksell.
-  Prawitz, D. (1971). Ideas and results in proof theory. In J.E. Fenstad (ed.), *Proceedings of the Second Scandinavian Logic Symposium*, 235–307. Amsterdam: North-Holland.
-  Prior, A.N. (1960). The runabout inference-ticket. *Analysis*, 21 (2): 38–39.
-  Schroeder-Heister, P. (1984). A natural extension of natural deduction. *The Journal of Symbolic Logic*, 49: 1284–1300.
-  Streicher, T. (1993). *Investigations into Intensional Type Theory*. Habilitationsschrift, Ludwig-Maximilians-Universität München.
-  Tranchini, L. (2016). Proof-theoretic harmony: towards an intensional account. *Synthese*. Online First.
-  Tranchini, L., Pistone, P. & Petrolo, M. (2016). The naturality of natural deduction. Manuscript (arXiv:1607.06603).