



Searches for New Physics With Tau Leptons at the CMS Experiment

George Uttley

Imperial College London
Department of Physics

A thesis submitted to Imperial College London
for the degree of Doctor of Philosophy

The copyright of this thesis rests with the author and is made available under a Creative Commons Attribution Non-Commercial No Derivatives licence. Researchers are free to copy, distribute or transmit the thesis on the condition that they attribute it, that they do not use it for commercial purposes and that they do not alter, transform or build upon it. For any reuse or redistribution, researchers must make clear to others the licence terms of this work.

Abstract

The Standard Model of particle physics is currently the best model of the fundamental particles and their interactions. However, there are still significant theoretical issues and experimental tensions with the model. The theoretical issues include the hierarchy problem which forecasts the breakdown of the Standard Model when looking at the size of corrections needed to calculate the mass of the newest found member of the theory, the Higgs boson particle. The current experimental tensions include the B-anomalies and the g-2 measurement. These results, although they do not yet sit at the required 5σ deviation for a discovery, offer the most prominent leads into where new physics may be hiding. Looking for signatures of theoretical explanations of these anomalies offers excellent search options for new fundamental particles. This thesis describes the search for new physics that can explain both the theoretical problems and experimental tensions. This is done using tau leptons seen during Run-2 of the Large Hadron Collider (LHC) at the Compact Muon Solenoid experiment. The Beyond Standard Model theories searched for range from Supersymmetry, leptoquarks, to type-X two Higgs doublet models. Each theory is separately studied and an analysis is tailored to find its most sensitive signature. In the process of optimisation, data-driven background modelling is improved to aid the reliability of the results. Two local excesses of size $\approx 3\sigma$ for a resonance produced via gluon fusion at masses 100 GeV and 1.2 TeV is observed, otherwise good agreement was observed between the Standard Model background prediction and data. Leading constraints are placed on scenarios of the Minimal Supersymmetric Standard Model phase space and limits that shrink the allowed explanation to the B anomalies are set. The type X two Higgs doublet model as an explanation to the g-2 anomaly is completed excluded by this thesis.

Declaration

I declare that the work in this thesis is mine. Figures and results taken from other sources are indicated by a reference in the text or figure caption. Figures labelled "CMS" are sourced from CMS publications. Figures labelled "CMS Supplementary" have been made public as supplementary material accompanying a publication. All figures with these labels, including those made by myself, also include the relevant reference in the caption. The analyses presented in this document were developed in collaboration with other members of the CMS experiment. Chapters 1–3 do not contain original work by myself, but are intended to give an description of the theoretical motivation, and apparatus and methods used to generate and collect the data, that are critical to the work presented in following chapters. The high-mass additional Higgs boson search, described in Chapter 4, and the origin of the background modelling methods was pre-established in Ref. [1]. The work described in this chapter was done in collaboration with the other members of the CMS $H \rightarrow \tau\tau$ group. For the additional Higgs boson component of this chapter, I contributed to the low-mass optimisation, derivation and parametrisation of fake factors, evaluation of uncertainties and the final statistical fits to data. The vector leptoquark interpretation of the results was solely my work. The results in Chapter 4.4 were made public in Ref. [2]. Chapter 5 contains work performed in collaboration with the Imperial College London $H \rightarrow \tau\tau$ group. I was responsible for the full work flow of this analysis including signal simulation, background modelling, optimisation, uncertainty modelling, interpretations and the final statistical fits to data. This analysis is currently not published, but a publication is planned in the near future. Chapter 6 includes interpretations of the results in Chapters 4 and 5, that are entirely my personal work.

George Uttley

Acknowledgements

I would like to thank Imperial College London High Energy Physics group and STFC for giving me the opportunity to conduct this research. Thank you to my supervisor, David Colling, for all the support and guidance, in particular the aid I received whilst working through the pandemic. I also owe a massive thank you to Daniel Winterbottom, for all of the help and advice, as well as the patience he showed with me throughout my PhD. I am also grateful to the remaining members of the $H \rightarrow \tau\tau$ group for the all of interesting discussions and the continual motivation whilst writing this thesis. Thank you to my friends and family, who have continually supported me throughout my life and are the reason I was able to complete this work. In particular, thank you to Emmy for her unwavering care and encouragement throughout this journey.

George Uttley

Contents

1 Theory and Motivation	21
1.1 The Standard Model of Particle Physics	22
1.1.1 Fundamental Particles and Interactions	22
1.1.2 Higgs Sector	25
1.2 Extended Higgs Sector	27
1.3 Theoretical Problems and Potential Solutions	29
1.3.1 Hierarchy Problem	29
1.4 Experimental Tensions and Potential Solutions	31
1.4.1 B Anomalies	31
1.4.2 g-2 Anomaly	32
2 The LHC and CMS Experiment	35
2.1 The LHC	35
2.2 The CMS Detector	38
2.2.1 Tracker	38
2.2.2 Electromagnetic calorimeter	40
2.2.3 Hadron calorimeter	41
2.2.4 Muon system	43
2.2.5 Triggering and Computing	45
3 Object Reconstruction	48
3.1 Tracks and vertices	48
3.2 Particle flow	49
3.3 Muons	52
3.4 Electrons	55
3.5 Jets	56
3.6 b jets	58
3.7 Missing transverse energy	58

3.8 Taus	60
4 Searches for New Physics in $\tau^+\tau^-$ Final States	67
4.1 Signal Modelling	68
4.1.1 Additional Higgs Bosons	68
4.1.2 Vector Leptoquarks	71
4.2 Event Selection	73
4.2.1 Trigger Requirements	75
4.2.2 Offline Requirements	75
4.3 Search Optimisation	77
4.3.1 High Mass Optimisation	78
4.3.2 Low Mass Optimisation	79
4.3.3 Standard Model Higgs Boson Optimisation	79
4.4 Background Modelling Overview	81
4.5 QCD Estimation in the $e\mu$ Channel	82
4.6 Embedding Method	83
4.7 Fake Factor Method	85
4.7.1 Determination Regions	86
4.7.2 Parametrisation	88
4.7.3 Corrections	92
4.7.4 Applying Fake Factors	94
4.8 MC Corrections	96
4.9 Uncertainty Model	97
4.10 Signal Extraction	101
4.11 Postfit Plots	104
4.12 Model Independent Results	105
4.12.1 Limits	105
4.12.2 Significance and Compatibility	110
4.12.3 2D Likelihood Scans	112
4.13 Model Dependent Limits	112
5 Search for New Physics in $\tau^+\tau^-\tau^+\tau^-$ Final States	117
5.1 Signal Modelling	118
5.2 Event Selection	120
5.2.1 Trigger Requirements	122
5.2.2 Offline Requirements	123

5.3	Search Optimisation	124
5.4	Background Modelling Overview	125
5.5	ZZ Modelling	126
5.6	Machine Learning Fake Factor Method	126
5.6.1	BDT Reweighting	128
5.6.2	Fitting Regions	128
5.6.3	Variables Used	130
5.6.4	Machine Learning Subtraction Method	130
5.6.5	Fitting	132
5.6.6	Applying Fake Factors	136
5.7	Uncertainty Model	137
5.8	Signal Extraction	139
5.8.1	Postfit Plots	139
5.9	Model Independent Results	140
5.9.1	Limits	140
5.9.2	Compatibility	143
5.10	Model Dependent Limits	143
6	Conclusion	148
6.1	Global Interpretations of Results	148
6.2	Summary	152

List of Figures

1.1	Diagram of the fundamental particles the constitute the SM. Also displayed are the fermion generation shown in Roman numerals, the particles measured mass, charge, spin and colours available for the strong interaction. This is taken and adjusted from Reference [3]	23
1.2	One-loop corrections to the Higgs boson's mass by a fermion f (a) and a scalar S (b).	29
1.3	Examples of one-loop Bar-Zee feynman diagrams for the contribution from ϕ and A in (a) and H^\pm in (b).	33
1.4	Examples of two-loop Bar-Zee feynman diagrams for the contribution from ϕ and A in (a) and H^\pm in (b). Note that other fundamental particles can also contribute to the loop.	33
2.1	A schematic diagram of the CERN [4].	36
2.2	The total integrated luminosity for proton-proton collisions collected by the Compact Muon Solenoid (CMS) experiment between 2010 and May 2023 [5].	37
2.3	A perspective view of the CMS detector [6].	39
2.4	Schematic of the CMS tracker (pre-pixel upgrade) in the r - z plane, showing the position of the pixel detector as well as the TIB, TID, TOB, and TEC strip detectors. The lines represent detector modules and the double lines represent back-to-back modules [6].	39
2.5	A schematic of the CMS ECAL detector [6].	42
2.6	A schematic of the CMS HCAL showing the arrangement of the HB, HE, HO, and HF calorimeters [6].	42
2.7	A schematic of the L1 trigger workflow for object reconstruction [7].	46

3.1	The p_T resolution for muon identification using the muon system (black), the inner tracker (blue), and the full system (red). This is shown for the pseduorapidity regions $0 < \eta < 0.8$ (a) and $1.2 < \eta < 2.4$ (b) [6].	53
3.2	Performance of the electron BDT using $Z \rightarrow ee$ enriched events from data (dots), background-enriched events from data (triangles), and $Z \rightarrow ee$ MC events (purple solid histogram). This is shown for the performance in the barrel (a) and the endcaps (b) for a BDT that does not utilise trigger information during training [8].	56
3.3	Mis-identification probability 2σ bands against efficiency for b tagging using the DeepJet algorithm, for AK4jets with $p_T > 30$ GeV AND $ \eta < 2.5$ utilising MC with parameters from the 2017 era of data taking. Two bands are displayed, one for the identification of a b quark rather than light quarks and gluons (blue) and one for the identification of a b quark rather than a c quark (red) [9].	59
3.4	The distance between the hadronic tau and electron or photon η (a) and ϕ (b) with respect the the electron or photon p_T . The binned values (points) and the fitted functions f and g (red line), that encapsulates 95% of all electron and photons are shown in both cases [10].	62
3.5	The architecture of the DeepTau neural network, comprising of three sets of input variables: inner cells, outer cells, and high-level features. These sets are processed separately through subnetworks and their outputs are concatenated. Five fully connected layers process the concatenated output to calculate the probabilities for a candidate to be a hadronic tau, electron, muon, or jet. The high-level input subnetwork consists of three fully connected layers, taking 47 inputs and yielding 57 outputs. Complex subnetworks process the features of inner and outer cells separately, with fully connected layers followed by concatenation and additional fully connected layers. Convolutional layers progressively reduce the grid size for both inner and outer cells. For inner cells, there are 5 convolutional layers, while for outer cells, there are 10 convolutional layers. The number of trainable parameters (TP) for the different subnetworks are also provided [11].	64

3.6 Efficiency comparisons of electrons (a), muons (b) and jets (c) passing the DeepTau identification discriminators versus hadronic tau efficiency. (a) uses $Z \rightarrow ee$, (b) uses $Z \rightarrow \mu\mu$ and (c) uses W+jets MC, with only objects with $p_T < 100$ GeV used. Working points are indicated by full circles and previous algorithms described in Reference [12] are shown in blue triangles and green squares [11].	66
4.1 Diagram (a) shows the production of neutral Higgs bosons from gluon fusion. The dominant loop contributions to this diagrams are from top-only, bottom-only and top-bottom interference. Diagrams (b) and (c) show production in association with b quarks.	68
4.2 p_T density distributions of the A (top) and H (bottom) boson, with contributions to the gluon fusion loop displayed individually and summed. These are shown for $\tan\beta$ values of 10 (left) and 30 (right) where $m_A = 500$ GeV in the MSSM M_h^{125} scenario.	70
4.3 Feynman diagrams showing the contribution from U_1 vector leptoquarks to the final state with a pair of oppositely charged τ leptons. q represents any SM quark. Diagram (a) shows t-channel, (b) pair and (c) single production of a vector leptoquark.	72
4.4 The generator level $m_{\tau\tau}$ density distributions of the t-channel vector leptoquark signal and the interference with Drell-Yan. This is shown in the VLQ BM 1 scenario for a leptoquark of mass 1 TeV for coupling strengths of $g_U = 3$ (a) and $g_U = 5$ (b).	74
4.5 Diagram of inputs to the D_ζ variable [2].	77
4.6 Overview of the categories used for the extraction of the signal in the high mass optimisation procedure.	80
4.7 Overview of the categories used for the extraction of the signal in the low mass optimisation procedure.	80
4.8 Schematic of the embedding method to model genuine di- τ backgrounds from di-muon events in data [13].	84
4.9 Closure plot showing the di-muon mass on the dedicated embedding validation samples.	85
4.10 Schematic of the regions used for fake factor derivation.	87

4.11 A 2D heat map of the F_F determined from $t\bar{t}$ MC for the full run-2 dataset in the combined $e\tau_h$ and $\mu\tau_h$ channels. This is shown with respect to the $\tau_h p_T$ and the p_T of the jet matched to the τ_h . The ratio of jet to $\tau_h p_T$ categorisation used is shown split by the dashed lines.	89
4.12 Flow chart of the algorithm used to determine where binned values are taken instead of the fit. The blue box represents the input, the green diamonds represent the decisions and the red boxes represent the outputs.	90
4.13 F_F fits in $\tau_h\tau_h$ channel for the QCD $N_{\text{pre b jets}} = 0$ category with 2018 data. The three jet p_T to $\tau_h p_T$ categories are shown.	91
4.14 F_F fits in $\mu\tau_h$ channel for the QCD and W + Jets $N_{\text{pre b jets}} = 0$ category with 2018 data. The three jet p_T to $\tau_h p_T$ categories are shown for each process.	91
4.15 Diagram showing how fake MET arises from mismodelling jet constituents in τ_h identification.	92
4.16 Determination Region closure correction fits with 2018 data. (a) is the correction parametrised by ΔR in events with $N_{\text{b jets}} = 0$ in the $\tau_h\tau_h$ channel. (b) and (c) show the correction for the $\mu\tau_h$ channel parametrised by the specific correction variables defined in Equation 4.6 and 4.7 for QCD and W + jets processes respectively.	93
4.17 Determination Region to Application Region closure correction fits with 2018 data. (a) is the correction moving from same sign to opposite sign τ leptons the parametrised by leading $\tau_h p_T$ in events with $N_{\text{b-jets}} = 0$ in the $\tau_h\tau_h$ channel. (b) and (c) show the correction for the $\mu\tau_h$ channel moving from same sign to opposite sign τ leptons and high m_T to low m_T both parametrised by the the muon p_T for QCD and W + jets processes respectively.	94
4.18 The expected Application Region fractions of the processes in the $\mu\tau_h$ channel. (a) and (b) show the no b tag Tight- m_T and Loose- m_T categories and (c) and (d) show the b tag Tight- m_T and Loose- m_T categories respectively.	95

4.19 Distributions of $m_{\tau\tau}$ in the no b tag second highest (left) and highest (right) p_T category for the $\tau_h\tau_h$ (top), the combined $e\tau_h$ and $\mu\tau_h$ (middle) and the $e\mu$ (bottom) channels. The solid histograms show the stacked background predictions after a signal plus background fit to the data. The best fit gluon fusion signal for $m_\phi = 100$ GeV is shown by the red line. Also shown by a blue dashed line on the bottom pad is the ratio of the background predictions for the background only fit to the signal plus background fit	106
4.20 Distributions of m_T^{tot} in the $\tau_h\tau_h$ no b tag (a) and b tag (b) categories, the combined $e\tau_h$ and $\mu\tau_h$ no b tag (c) and b tag (d) Tight- m_T categories and the $e\mu$ no b tag (e) and b tag (f) Medium- D_ζ categories. The solid histograms show the stacked background predictions after a background only fit to the data. The best fit gluon fusion signal for $m_\phi = 1.2$ TeV is shown by the red line, b associated production and U_1 signals are also shown for illustrative purposes.	107
4.21 Expected (dashed line) and observed (solid line and dots) 95% CL upper limits on the product of the cross sections and branching fraction for the decay into τ leptons for (a) gg ϕ and (b) bb ϕ production in a mass range of $60 \leq m_\phi \leq 3500$ GeV. The dark green and bright yellow bands indicate the central 68% and 95% intervals for the expected exclusion limit.	108
4.22 Comparison of the expected 95% CL upper limits on the product of the cross sections and branching fraction for the decay into τ leptons for (a) gg ϕ and (b) bb ϕ production, split by the $\tau\tau$ decay products fit individually.	109
4.23 Comparison of the expected 95% CL upper limits on the product of the cross sections and branching fraction for the decay into τ leptons for (a) gg ϕ and (b) bb ϕ production, split by the CMS result detailed in this thesis and the ATLAS result from Reference [14].	109
4.24 Local p -value and significance of a gg ϕ (left) and bb ϕ (right) signal as a function of m_ϕ	110
4.25 Compatibility plots of the 100 GeV excess split into analysis channels (a) and categories (b). In each case the fitted signal strength is decoupled in the bin shown on the plot.	111
4.26 Compatibility plots of the 1.2 TeV excess split by analysis channels. In each case the fitted signal strength is decoupled in each channel.	111

4.27 Maximum likelihood scans, including 68% and 95% CL contours obtained from the signal likelihood for the model-independent search. The scans are shown for selected values of m_ϕ between 60 GeV and 3.5 TeV.	113
4.28 Expected and observed 95% CL exclusion contours in the MSSM M_h^{125} (a) and $M_{h,EFT}^{125}$ (b) scenarios. The exclusion limit only on background expectation is shown as a dashed black line, the dark and bright grey bands show the 68% and 95% intervals of the expected exclusion and the observed exclusion contour is shown by the blue area. The parameter space where m_h deviates by more than ± 3 GeV from the observed SM Higgs boson mass is shown by a red hatched area.	115
4.29 Expected and observed 95% CL upper limits on g_U in the VLQ BM 1 (a) and 2 (b) scenarios, in a mass range of $1 < m_U < 5$ TeV. The exclusion limit only on background expectation is shown as a dashed black line, the dark and bright grey bands show the 68% and 95% intervals of the expected exclusion and the observed exclusion contour is shown by the blue area. The 95% confidence interval for the preferred region from the global fit presented in Reference [15] is also shown by the green shaded area.	116
5.1 Diagram of production of two additional neutral Higgs bosons from an off-shell Z boson and their decays to tau leptons.	118
5.2 Generator level distributions of the visible mass densities for A (a) and ϕ (b) for the signal process.	119
5.3 Calculated production cross sections for the $Z^* \rightarrow \phi A$ process, varying the masses of ϕ and A.	120
5.4 Calculated branching fractions of A (a) and ϕ (b) decaying to a pair of τ leptons for various mass scenarios.	121
5.5 Calculated branching fractions in the $\cos(\beta - \alpha)$ - $\tan \beta$ phase space for ϕ of mass 200 GeV decaying to a pair of τ leptons, in the scenario where $m_A = 160$ GeV.	121
5.6 Overview of the categories used for the extraction of the signal in the $Z^* \rightarrow \phi A \rightarrow 4\tau$	124
5.7 Distributions in the $\mu\mu\mu\mu$ channel are shown for the variables total transverse mass, m_T^{tot} , (a) and the total mass, $m_{\mu\mu\mu\mu}$ (b). The solid histograms show the stacked background predictions.	127

5.8 Comparison of histograms produced via the BDT subtraction method to histogram subtraction. Also shown is the histogram produced when no subtraction is performed. The uncertainty bands contain statistical uncertainties and uncertainties derived on the non-closure of the method. The is shown for four of the fitted variable: τ_h - p_T , the ratio of τ_h - p_T to jet- p_T , the τ_h HPS decay mode and the τ_h - η	133
5.9 Average fake factors (reweights) calculated by the BDT reweighting method shown individually in regions A, B and D as defined in Figure 4.10. The is shown for four of the fitted variable: τ_h - p_T , the ratio of τ_h - p_T to jet- p_T , the τ_h HPS decay mode and the τ_h - η	134
5.10 Comparison of histograms produced using the fake factors applied to the fitted fail region compared to fitted pass region. The uncertainty bands contain statistical uncertainties and uncertainties derived on the non-closure of the method. The is shown for three of the fitted variable: τ_h - p_T , the ratio of τ_h - p_T to jet- p_T , the τ_h HPS decay mode and the τ_h - η . The χ^2 divided by number of degrees of freedom between the two histograms is also shown.	135
5.11 Distributions of m_T^{tot} in the high (a), medium (b) and low (c) statistic categories. As the bin sizes vary drastically between channels, the bin sizes are kept constant and the bin intervals are shown on x axis. The high statistic categories consistent of only the $\tau_h\tau_h\tau_h$ channel, the medium statistic categories include the $ee\tau_h\tau_h$ and $\mu\mu\tau_h\tau_h$ 0S Leptons categories, and the low statistic categories show the $e\tau_h\tau_h\tau_h$, $\mu\tau_h\tau_h\tau_h$, $\tau_h\tau_h\tau_h\tau_h$, $ee\tau_h\tau_h$ SS Leptons , $\mu\mu\tau_h\tau_h$ SS Leptons , $e\mu\tau_h\tau_h$ SS Leptons and $e\mu\tau_h\tau_h$ 0S Leptons channels and categories. The solid histograms show the stacked background predictions after a background-only fit to the data. The $m_\phi = 200$ GeV and $m_A = 160$ GeV signal scaled to 0.01 pb is also shown by a blue line for illustrative purposes.	141
5.12 Expected (dashed line) and observed (solid line and dots) 95% CL upper limits on the product of the cross sections and branching fractions for the decay of both additional Higgs bosons into τ leptons. Different m_A hypotheses are scaled by different orders of magnitudes (written on plot) in order to make mass points distinguishable. The dark green and bright yellow bands indicate the central 68% and 95% intervals for the expected exclusion limit.	142

5.13 Comparison of the expected 95% CL upper limits on the product of the cross sections and branching fractions for the decay into τ leptons, split by the $\tau\tau\tau\tau$ decay products fit individually.	144
5.14 Compatibility plots for the $m_A = 100$ GeV and $m_\phi = 100$ GeV mass scenario in analysis decay channels. In each case the fitted signal strength is decoupled in the bin shown on the plot. The combined best fit value and its $1sigma$ variation are shown by the blue line and band respectively. The black dashed line indicates a signal strength of zero.	144
5.15 Expected and observed 95% CL exclusion contours on the m_A - $\tan\beta$ phase space in the type X 2HDM alignment scenario for m_ϕ scenarios of 100 GeV (a) and 200 GeV (b). The exclusion limit only on background expectation is shown as a dashed black line, the dark and bright grey bands show the 68% and 95% intervals of the expected exclusion and the observed exclusion contour is shown by the blue area.	146
5.16 Expected and observed 95% CL exclusion contours on the $\cos(\beta - \alpha)$ - $\tan\beta$ phase space in the type X 2HDM alignment scenario with m_ϕ equal to 200 GeV and m_A scenarios of 100 GeV (a) and 160 GeV (b). The exclusion limit only on background expectation is shown as a dashed black line, the dark and bright grey bands show the 68% and 95% intervals of the expected exclusion and the observed exclusion contour is shown by the blue area.	147
6.1 Expected and observed 95% CL exclusion contours on the m_A - $\tan\beta$ phase space in the type X 2HDM alignment scenario for m_ϕ scenarios of 100 GeV (a) and 200 GeV (b). The exclusion limit only on background expectation is shown as a dashed black line, the dark and bright grey bands show the 68% and 95% intervals of the expected exclusion and the observed exclusion contour is shown by the blue area. The limit obtained by HIGGSTOOLS is shown in the red contour.	150

List of Tables

1.1	Table showing which Higgs doublet different fermion groups different types of 2HDMs couple to. By convention the u quark is chosen to couple to Φ_2 .	28
1.2	Table showing the couplings of fermion groups to additional neutral Higgs bosons in different types of 2HDMs. These are dependent on the mixing angles α and β . t_x , s_x and c_x represent $\tan x$, $\sin x$ and $\cos x$ respectively.	28
1.3	Best fit values and uncertainties for mixing matrix parameters as given in Reference [15].	32
1.4	Regions of interest for g-2 anomaly with respect to the type X 2HDM in the normal and inverted alignment scenarios as suggested in Reference [16].	34
3.1	Electron effective areas used for the ρ -corrected isolation computation [8].	57
3.2	Measured branching fractions for the tau lepton. h represents a charged hadron either a pion or a kaon [17].	61
3.3	Target efficiencies of the DeepTau working points with respect to electrons, muons and jets.	65
4.1	Branching fractions of the decays of two τ leptons.	74
4.2	Lower trigger light lepton thresholds p_T in GeV for the $e\tau_h$ and $\mu\tau_h$ channels.	75
4.3	pdfs used for nuisance parameters.	101
5.1	Branching fractions of four τ leptons, where e and μ represent the leptonic decay of the τ and τ_h represent the hadronic decay of the τ .	122
5.2	Vetoos set on the number of leptons for each channel.	123

5.3 Regions modelled by the fake factor method when using the leading τ_h (τ_h^1), the sub-leading τ_h (τ_h^2) and both, to model the jet $\rightarrow \tau_h$ background and whether that specific object is a genuine τ lepton (τ) or a jet (j). Also shown is a combination of three regions to fully model all possible combinations.	136
---	-----

Chapter 1

Theory and Motivation

The Standard Model ([SM](#)) of particle physics is our current best theory to describe the fundamental particles and their interactions. The [SM](#) describes the strong force, as well as unifying the weak and electromagnetic forces. The latter is partly done through the Brout-Englet-Higgs ([BEH](#)) mechanism for spontaneous symmetry breaking, that allows many fundamental particles to obtain masses. It also predicts a new scalar boson, named the Higgs boson. In 2012, the ATLAS [18] and CMS [19] collaborations discovered a Higgs boson-like particle when colliding protons at high energy at the Large Hadron Collider ([LHC](#)) and further measurements of the particle’s properties have been consistent with the [SM](#) prediction. This discovery experimentally completed the [SM](#) particle constituents.

However, the [SM](#) is not without theoretical problems and experimental tensions. Firstly, the hierarchy problem describes the issue of lightness of the observed Higgs boson mass and the “unnatural” balancing of inputs needed to explain the theorised loop corrections to the predicted mass. These are orders of magnitude larger than the observed mass. A solution to this problem is supersymmetry ([SUSY](#)), but no experimental evidence for this theory has yet been found that separates it from the [SM](#). Secondly, results from B physics [20–27] and the g–2 experiment [28] have shown deviations from the [SM](#) predictions. Although not the statistical significance for a discovery, they offer intriguing hints at potential Beyond Standard Model ([BSM](#)) physics. [BSM](#) particles, produced from extended Higgs sectors or otherwise, have been theorised to explain these deviations. This chapter will explain the [SM](#) and the Higgs sector theory, as well as detailing the [BSM](#) extensions that can help resolve the theoretical problems and experimental tensions.

One caveat to the B physics results presented in this thesis, is the exclusion of the updated R_K and R_{K^*} measurements from the LHCb collaboration [29]. This is due to the interpretation of the B anomalies, described in this chapter and used in Chapter 4, being concluded prior to the release of Reference [29]. The phenomenological affect of Reference [29], on the best fit of new physics to the B anomalies, is yet to be fully determined.

1.1 The Standard Model of Particle Physics

1.1.1 Fundamental Particles and Interactions

The **SM** is a set of fundamental particles, as shown in Figure 1.1, and rules that govern the interactions between particles. The interactions between these particles are able to model the strong, weak and electromagnetic force, unifying the latter two into one electroweak interaction. The **SM** consists of 6 quarks, 3 charged leptons and 3 neutrinos, which are grouped into fermions because of their shared half integer spin. Each of these particles has an anti-partner with opposite quantum numbers but the same mass. The **SM** also consists of a number of bosons with integer spin, that describe the fundamental forces of nature: the strong, weak, and electromagnetic forces. The gluon is the mediator of the strong force, the W and Z bosons mediate the weak force, and the photon mediates the electromagnetic force.

The **SM** is a renormalisable quantum field theory that is built on the principle of local gauge invariance. The $SU(3)_C \otimes SU(2)_L \otimes U(1)_Y$ is the gauge symmetry group of the **SM**. This means that the Lagrangian, that governs the particles interactions, is invariant under such a transformation. $SU(2)_L \otimes U(1)_Y$ is the symmetry of the electroweak unification and $SU(3)_C$ is the symmetry of the theory for strong force, name Quantum Chromodynamics (**QCD**).

The quantum number associated with the **QCD** $SU(3)_C$ symmetry is the colour charge C. Quarks and gluons carry a colour charge and so interact with the strong force. One characteristic feature of **QCD** is confinement, which requires neutral colour charges and so quarks must be observed in a bound state, named hadrons. Another key property is asymptotic freedom, which weakens the interaction strength at higher energy or very short distances [30, 31].




Figure 1.1: Diagram of the fundamental particles the constitute the SM. Also displayed are the fermion generation shown in Roman numerals, the particles measured mass, charge, spin and colours available for the strong interaction. This is taken and adjusted from Reference [3]

Electroweak unification was initially proposed by Glashow [32], Weinberg [33] and Salam [34] to combine the theories for the weak and electromagnetic forces into one. It is built on the premise of the Dirac equation. The Dirac Lagrangian for a massless field, ψ , is shown below.

$$\mathcal{L}_{\text{Dirac}} = i\bar{\psi}\gamma^\mu\partial_\mu\psi \quad (1.1)$$

The $SU(2)_L$ transformation operates on the weak isospin, I , only for the left-handed spinors and the $U(1)_Y$ transformation operates on the weak hypercharge, $Y = 2(Q - I_3)$, where Q is the charge of the fundamental particles and I_3 is the third component of the weak isospin. The reason for the distinction of the $SU(2)$ symmetry to act only on left-handed spinors is from the observed chirality of fermions under the weak interaction. By invoking gauge invariance of the Dirac Lagrangian under such a transformation, four associated gauge fields are present, $\mathbf{W}_\mu = (W_\mu^1, W_\mu^2, W_\mu^3)$ and B_μ .

$$\begin{aligned} \mathcal{L}_{\text{Electroweak}} = & \bar{\psi}_L \left(\partial_\mu + \frac{i}{2}g\mathbf{W}_\mu \cdot \boldsymbol{\sigma} + g'YB_\mu \right) \psi_L \\ & + \bar{\psi}_R \left(\partial_\mu + g'YB_\mu \right) \psi_R - \frac{1}{4}\mathbf{W}_{\mu\nu} \cdot \mathbf{W}^{\mu\nu} - \frac{1}{4}B_{\mu\nu}B^{\mu\nu}, \end{aligned} \quad (1.2)$$

where the partial derivatives have been replaced with the covariant derivative and the added field tensors $\mathbf{W}_{\mu\nu}$ and $B_{\mu\nu}$ are defined as,

$$\mathbf{W}_{\mu\nu} = \partial_\mu\mathbf{W}_\nu - \partial_\nu\mathbf{W}_\mu - ig[\mathbf{W}_\mu, \mathbf{W}_\nu] \quad (1.3)$$

$$B_{\mu\nu} = \partial_\mu B_\nu - \partial_\nu B_\mu. \quad (1.4)$$

These fields can be rotated into the physical fields for the photon, Z and W boson with the following transformations,

$$\begin{aligned} W_\mu^\pm &= \frac{1}{\sqrt{2}}(W_\mu^1 \mp W_\mu^2), \\ Z_\mu &= W_\mu^3 \cos\theta_w - B_\mu \sin\theta_w, \\ A_\mu &= W_\mu^3 \sin\theta_w - B_\mu \cos\theta_w, \end{aligned} \quad (1.5)$$

where θ_w represents the weak mixing angle and is defined such that

$$\sin\theta_w = \frac{g'}{\sqrt{g^2 + g'^2}}, \quad \cos\theta_w = \frac{g}{\sqrt{g^2 + g'^2}}. \quad (1.6)$$

The W and Z bosons were discovered by the UA1 and UA2 collaborations in 1983 [35, 36], which confirmed the predictions made by electroweak unification. However, the W and Z bosons were measured to have a non-zero mass and Lagrangian mass terms of the form $\frac{1}{2}m_Z^2Z_\mu Z^\mu$ or $m_W^2W_\mu^-W^{+\mu}$ cannot be included as they are not invariant under the $SU(2)_L \otimes U(1)_Y$ gauge symmetry. The same issue also exists in massive fermions, as the fermion mass term's left- and right-handed chiral states, $m\bar{\psi}\psi = m(\bar{\psi}_R\psi_L + \bar{\psi}_L\psi_R)$, transform differently and therefore does not remain invariant under the gauge transformation. To resolve this, the BEH mechanism for spontaneous symmetry breaking was theorised.

1.1.2 Higgs Sector

The BEH mechanism was proposed in the 1960s by Englert and Brout [37], Higgs [38–40] and Guralnik, Hagen and Kibble [41, 42]. It works on the principles of spontaneous symmetry breaking of the $SU(2)_L \otimes U(1)_Y$ gauge symmetry. It does this by introducing a gauge invariant field that has a non-zero vacuum expectation value (VEV). The formalism of this is shown below, starting from a complex scalar doublet, Φ .

$$\Phi = \begin{pmatrix} \phi^+ \\ \phi^0 \end{pmatrix}, \quad (1.7)$$

where ϕ^+ and ϕ^0 are complex functions. The Lagrangian and the potential, chosen to fulfil the conditions stated above, then takes the form,

$$\mathcal{L}_{\text{Complex Scalar}} = (\partial_\mu \Phi)^\dagger (\partial^\mu \Phi) - V(\Phi) \quad (1.8)$$

with,

$$V(\phi) = \mu^2 \Phi^\dagger \Phi + \lambda (\Phi^\dagger \Phi)^2, \quad (1.9)$$

where μ^2 and λ are two real parameters, λ is required to be positive for the vacuum to be stable. If μ^2 is negative, Φ will have a non-zero VEV and the field will be able to spontaneously break the gauge symmetry. In the vacuum state, the field must satisfy the criteria,

$$\Phi^\dagger \Phi = -\frac{\mu^2}{2\lambda}. \quad (1.10)$$

This potential fulfils the criteria of a non-zero **VEV** as Φ must be non-zero. Choosing a gauge, to remove the massless scalar bosons predicted by Goldstone's theorem [43], by taking ϕ^+ to be zero and ϕ to be real. The ground state of Φ is found as,

$$\langle 0 | \Phi | 0 \rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ \nu \end{pmatrix}, \quad (1.11)$$

where $\nu^2 = \mu^2/\lambda$ and ν is the non-zero **VEV**. The complex doublet can then be written as an expansion around the minimum of the potential,

$$\Phi = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ \nu + h(x) \end{pmatrix} \quad (1.12)$$

Applying the covariant derivative for the electroweak gauge symmetry to produce the Lagrangian for the field, Φ , yields

$$\begin{aligned} \mathcal{L}_{\text{Higgs}} = & \frac{1}{2}(\partial_\mu h)(\partial^\mu h) + \frac{1}{8}g^2(W_\mu^1 + iW_\mu^2)(W^{1\mu} - iW^{2\mu})(\nu + h)^2 \\ & + \frac{1}{8}(gW_\mu^3 - g'B_\mu)(gW^{3\mu} - g'B^\mu)(\nu + h)^2 \\ & + \mu^2(\nu + h)^2 + \lambda(\nu + h)^4. \end{aligned} \quad (1.13)$$

Mass terms for W_μ^1 and W_μ^2 are now present and hence the mass of the W boson is $m_W = \frac{1}{2}g\nu$. Once again the remaining states need to be rotated to give the photon and the Z boson fields. Setting up the non-diagonal mass matrix for the fields W_μ^3 and B_μ , and calculating the eigenvalues to find the diagonal basis, masses of two fields are found, one equal to 0 and the other equal to $\frac{1}{2}\nu\sqrt{g^2 + g'^2}$ which represent the photon and Z boson mass respectively. The transformation to the photon and Z fields as parametrised in Equation 1.5 utilises,

$$\tan \theta_W = \frac{g'}{g}. \quad (1.14)$$

A mass term for the Higgs boson also arises from the potential of the **BEH** mechanism with $m_h = \sqrt{-2\mu^2}$. The same mechanism can also be used to add mass terms for quarks and charged leptons. Gauge invariant couplings to the Higgs bosons and fermion mass terms can be generated from the **BEH** mechanism applied to a Lagrangian term of the form,

$$\mathcal{L}_{\text{Yukawa}} = \lambda_f (\bar{\psi}_L \Phi \psi_R + \bar{\psi}_R \Phi \psi_L). \quad (1.15)$$

Upon spontaneous symmetry breaking, λ_f becomes proportional to the mass of the fermion, and hence the couplings are stronger between heavier fermions and the Higgs field.

1.2 Extended Higgs Sector

There is no theoretical limitation to have only one Higgs doublet in the theory. Therefore, a natural extension to the **SM** Higgs sector is the two Higgs doublet model (**2HDM**). The Lagrangian for such a theory is shown below.

$$\mathcal{L}_{\text{2HDM}} = (D_\mu \Phi_1)^\dagger (D_\mu \Phi_1) + (D_\mu \Phi_2)^\dagger (D_\mu \Phi_2) - V_{\text{2HDM}}(\Phi_1, \Phi_2) \quad (1.16)$$

where,

$$\begin{aligned} V_{\text{2HDM}} = & m_{11}^2 |\Phi_1|^2 + m_{22}^2 |\Phi_2|^2 + (m_{12}^2 \Phi_1^\dagger \Phi_2 + \text{h.c.}) \\ & + \frac{\lambda_1}{2} |\Phi_1|^4 + \frac{\lambda_2}{2} |\Phi_2|^4 + \lambda_3 |\Phi_1|^2 |\Phi_2|^2 + \lambda_4 |\Phi_1^\dagger \Phi_2| \\ & + \frac{1}{2} \left[\lambda_5 (\Phi_1^\dagger \Phi_2)^2 + \lambda_6 |\Phi_1|^2 \Phi_1^\dagger \Phi_2 + \lambda_7 |\Phi_2|^2 \Phi_1^\dagger \Phi_2 + \text{h.c.} \right], \end{aligned} \quad (1.17)$$

where m_{ij} represents the terms in the mass matrix and λ_i parametrises the self couplings of the Higgs sector. After the **BEH** mechanism is applied, **2HDMs** predict 5 Higgs bosons; 1 lighter and 1 heavier charge-parity (**CP**)-even (h and H), 1 **CP**-odd (A) and 2 charged particles (H^\pm). The Lagrangian for the Yukawa interactions with the Higgs bosons in such a theory are,

$$\begin{aligned} \mathcal{L}_{\text{yukawa}}^{\text{2HDM}} = & - \sum_{f=u,d,l} \left(\frac{m_f}{\nu} g_h^f \bar{f} f h + \frac{m_f}{\nu} g_H^f \bar{f} f H - i \frac{m_f}{\nu} g_A^f \bar{f} \gamma_5 f A \right) \\ & - \left[\frac{\sqrt{2} V_{ud}}{\nu} \bar{u} (m_u g_A^u P_L + m_d g_A^d P_R) d H^+ + \frac{\sqrt{2} m_l g_A^l}{\nu} \bar{\nu}_L l_R H^+ + \text{h.c.} \right], \end{aligned} \quad (1.18)$$

where u , d , l and ν represent up-like quark, down-like quark, charged lepton and neutrino fields, where the subscript L and R are the left- and right-handed projection performed via the projection operators, P . m_f are the fermion masses, ν is the vacuum expectation value of the **SM** Higgs doublet, and g are the couplings (relative to the **SM** couplings) of fermion fields to Higgs fields, h , H , A and H^+ . There are four main types of 2HDM that are dependent on which Higgs doublet couples to which group of fermions, named type I, II, X (lepton-specific) and Y (flipped). The

couplings of the fermion groups to the Higgs doublets are shown in Table 1.1. By convention Φ_2 is chosen to couple to up-like quarks.

	Type I	Type II	Type X	Type Y
u	Φ_2	Φ_2	Φ_2	Φ_2
d	Φ_2	Φ_1	Φ_2	Φ_1
l	Φ_2	Φ_1	Φ_1	Φ_2

Table 1.1: Table showing which Higgs doublet different fermion groups different types of 2HDMs couple to. By convention the u quark is chosen to couple to Φ_2 .

The type of **2HDM** determines the formulae for the couplings, g , that are functions on two parameters: the **CP**-even (α) and **CP**-odd (β) mixing angles of the mass matrices. These relative couplings are shown in Table 1.2.

	Type I	Type II	Type X	Type Y
g_h^u	c_α/s_β	c_α/s_β	c_α/s_β	c_α/s_β
g_h^d	c_α/s_β	$-s_\alpha/c_\beta$	c_α/s_β	$-s_\alpha/c_\beta$
g_h^l	c_α/s_β	$-s_\alpha/c_\beta$	$-s_\alpha/c_\beta$	c_α/s_β
g_H^u	s_α/s_β	s_α/s_β	s_α/s_β	s_α/s_β
g_H^d	s_α/s_β	c_α/c_β	s_α/s_β	c_α/c_β
g_H^l	s_α/s_β	c_α/c_β	c_α/c_β	s_α/s_β
g_A^u	$1/t_\beta$	$1/t_\beta$	$1/t_\beta$	$1/t_\beta$
g_A^d	$1/t_\beta$	t_β	$-1/t_\beta$	t_β
g_A^l	$1/t_\beta$	t_β	t_β	$-1/t_\beta$

Table 1.2: Table showing the couplings of fermion groups to additional neutral Higgs bosons in different types of 2HDMs. These are dependent on the mixing angles α and β . t_x , s_x and c_x represent $\tan x$, $\sin x$ and $\cos x$ respectively.

To match the observed Higgs boson measurements to a **CP**-even boson predicted by a **2HDM**, a linear combination of these two states are taken.

$$h_{\text{obs}} = \sin(\beta - \alpha)h + \cos(\beta - \alpha)H \quad (1.19)$$

Assuming a non-degeneracy of the observed Higgs boson mass, two possible alignment limits are acquired: the normal scenario where $h_{\text{obs}} = h$ and $\cos(\beta - \alpha) = 0$ and the inverted scenario where $h_{\text{obs}} = H$ and $\sin(\beta - \alpha) = 0$. In the normal scenario the

values of the coupling ratios from Table 1.2 are: $\cos \alpha / \sin \beta = 1$, $\sin \alpha / \cos \beta = -1$, $\sin \alpha / \sin \beta = -1 / \tan \beta$ and $\cos \alpha / \cos \beta = \tan \beta$. Whilst in the inverted scenario the ratios become: $\cos \alpha / \sin \beta = 1 / \tan \beta$, $\sin \alpha / \cos \beta = \tan \beta$, $\sin \alpha / \sin \beta = 1$ and $\cos \alpha / \cos \beta = 1$.

In the physical basis the 2HDM depends on the following parameters:

$$m_h, m_H, m_A, m_{H^\pm}, \tan \beta, \cos(\beta - \alpha), m_{12}^2, \lambda_6, \lambda_7 \quad (1.20)$$

It is common to require a \mathbb{Z}_2 symmetry to a 2HDM in order to maintain naturalness and avoid quadratic divergences, as well as suppressing flavour changing neutral currents. In this case the basis of parameters is shrunk, as $\lambda_6 = \lambda_7 = 0$.

1.3 Theoretical Problems and Potential Solutions

1.3.1 Hierarchy Problem

The current best measurement for the SM Higgs boson's mass from the CMS experiment is 125.38 ± 0.14 GeV [44] in natural units, which are used throughout this thesis. The lightness is a concern when considering “naturalness” and physics beyond the SM. The Higgs boson has loop corrections to calculations of its mass. Known heavy fermions already provide significant contributions (orders of magnitude greater than the observed mass) to the Higgs boson's calculated mass. On top of this, treating the SM as an effective field theory, new physics would be expected in the unexplored regions between the weak scale and the Planck scale. But if a new fermion were to be found, f , or a heavy scalar, S , in such a range, the Higgs boson would be subject to even greater changes in its predicted mass. The Feynman diagrams for the mass corrections for a fermion and a scalar are shown in Figure 1.2.




Figure 1.2: One-loop corrections to the Higgs boson's mass by a fermion f (a) and a scalar S (b).

Figure 1.2a shows a representation of the fermionic correction to the Higgs boson's

mass. The Lagrangian term for the coupling of the Higgs field to fermions is $-\lambda_f h \bar{f} f$. Hence, it can be determined that the correction to the mass is,

$$\Delta m_h^2 = -\frac{\lambda_f}{16\pi^2} \left[\Lambda_{UV}^2 - 2m_f^2 \ln \left(\frac{\Lambda_{UV}}{m_f} \right) \right] + \dots, \quad (1.21)$$

where Λ_{UV} is the ultraviolet momentum cut off [45]. Beyond this, our effective field theory would be expected to break down and for new physics to be found.

Figure 1.2b illustrates a correction to the Higgs mass by a scalar particle. The coupling of a scalar to the Higgs field is represented by the Lagrangian term $-\lambda_S |h|^2 |S|^2$. The Higgs boson's mass correction for such a term is derived to be,

$$\Delta m_h^2 = \frac{\lambda_S^2}{16\pi^2} \left[\Lambda_{UV}^2 - 2m_S^2 \ln \left(\frac{\Lambda_{UV}}{m_S} \right) \right] + \dots \quad (1.22)$$

Equation 1.21 and 1.22 show that if the mass of the scalar is equivalent to that of the fermion and $\lambda_f = \lambda_S^2$, then the Higgs boson's mass corrections cancel. This offers a solution to the hierarchy problem, by introducing a new symmetry that extends the **SM**. The symmetry relates fermions and bosons and is known as **SUSY**. It states that fermions and bosons exist in groups called supermultiplets. Each supermultiplet contains fermion and boson states which are superpartners of one another. On-shell each supermultiplet must have an equivalent number of fermionic and bosonic degrees of freedom. In order for this to also hold off-shell, an auxiliary field is added to balance the number of degrees of freedom.

If **SUSY** is an unbroken theory, then it would be expected for the superpartners to have the same mass as the **SM** particles. This has not been seen experimentally, therefore **SUSY** must be a broken theory in the vacuum state. Soft **SUSY** breaking can be introduced through the addition of **SUSY** violating Lagrangian term $\mathcal{L}_{\text{soft}}$, where

$$\mathcal{L} = \mathcal{L}_{\text{SUSY}} + \mathcal{L}_{\text{soft}}. \quad (1.23)$$

$\mathcal{L}_{\text{soft}}$ contains only mass terms and coupling parameters. Defining m_{soft} as the largest mass scale involved in the soft Lagrangian, m_{soft} also then defines the mass splitting between the **SM** and supersymmetric particles. If the mass splitting becomes significant, the hierarchy problem would be reintroduced as corrections to the Higgs boson's mass would again become large.

The Minimal Supersymmetric Standard Model ([MSSM](#)) is the simplest implementation of [SUSY](#). It introduces sets of new particles named squarks, sleptons, gauginos and Higgsinos as superpartners to quarks, leptons, gauge bosons and the Higgs bosons. Also added are neutralinos and charginos that are combinations of gauginos and Higgsinos. Additional contributions to the particle content comes from the Higgs sector. The Higgs sector of the [MSSM](#) is required to be extended to two Higgs doublets in order to maintain the gauge symmetries and to cancel quantum mechanical inconsistencies [45]. In particular a type II [2HDM](#) is needed to ensure natural Yukawa couplings, minimal flavour violation and tree-level mass relations, that all cannot be achieved with a different extended Higgs sector [45]. At tree-level, the [MSSM](#) Higgs sector is only dependent on m_A and $\tan\beta$. At high-order accuracies, benchmark scenarios are needed to set the remaining free parameters.

1.4 Experimental Tensions and Potential Solutions

1.4.1 B Anomalies

Measurements from the B physics experiments such as LHCb [20,24,26,27], BaBar [21, 22] and Belle [23, 25], testing lepton flavour conservation, have found deviations away from the [SM](#) expectation of lepton universality. The differences are observed in both neutral current ($b \rightarrow sl^+l^-$) and charged current ($b \rightarrow c\tau\nu$) transitions. These B anomalies have prompted the idea for a short range lepton flavour violating interaction. This interaction is theorised to be mediated by a new “leptoquark” particle [46, 47]. In an attempt to fit a model that offers a combined explanation of these results, it was found that a U_1 vector leptoquark was the only leptoquark that could offer a simultaneous explanation of all anomalous results [15]. Such a leptoquark would couple to fermions by the Lagrangian shown below.

$$\mathcal{L}_U = \frac{g_U}{\sqrt{2}} U^\mu [\beta_L^{i\alpha} (\bar{q}_L^i \gamma_\mu l_L^\alpha) + \beta_R^{i\alpha} (\bar{d}_R^i \gamma_\mu e_R^\alpha)] + \text{h.c.} \quad (1.24)$$

where g_U is the coupling scaling parameter and β_L and β_R are the left and right-handed mixing matrices,

$$\beta_L = \begin{pmatrix} 0 & 0 & 0 \\ 0 & \beta_L^{s\mu} & \beta_L^{s\tau} \\ 0 & \beta_L^{b\mu} & 1 \end{pmatrix}, \quad \beta_R = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & \beta_R^{b\tau} \end{pmatrix}. \quad (1.25)$$

The coupling g_U is defined such that $\beta_L^{b\tau} = 1$, and the negligible matrix elements for the fit are set to 0. The fit to B anomalies performed in Reference [15], found the best fit values for each left-handed mixing matrix parameter based on two scenarios for $\beta_R^{b\tau}$, namely $\beta_R^{b\tau} = 0$ and $\beta_R^{b\tau} = -1$. These are named VLQ BM 1 and VLQ BM 2 respectively. These represent no and maximal right-handed contributions. The best fit results to the matrix parameters are shown in Table 1.3.

Scenario	$\beta_R^{b\tau}$	$\beta_L^{b\mu}$	$\beta_L^{s\tau}$	$\beta_L^{s\mu}$
VLQ BM 1	0	$-0.15^{+0.13}_{-0.11}$	$0.19^{+0.06}_{-0.09}$	$0.014^{+0.01}_{-0.01}$
VLQ BM 2	-1	$-0.14^{+0.12}_{-0.11}$	$0.19^{+0.05}_{-0.08}$	$0.03^{+0.01}_{-0.02}$

Table 1.3: Best fit values and uncertainties for mixing matrix parameters as given in Reference [15].

The fit also provides a 1σ and 2σ bound on allowed values for the ratio of the vector leptoquark mass m_U , to the coupling constant g_U , and these are,

$$\begin{aligned} \text{VLQ BM 1, } 1\sigma &: 0.70 < g_U/m_U \text{ (1/TeV)} < 1.09, \\ \text{VLQ BM 1, } 2\sigma &: 0.57 < g_U/m_U \text{ (1/TeV)} < 1.38, \\ \text{VLQ BM 2, } 1\sigma &: 0.49 < g_U/m_U \text{ (1/TeV)} < 0.67, \\ \text{VLQ BM 2, } 2\sigma &: 0.39 < g_U/m_U \text{ (1/TeV)} < 1.25. \end{aligned} \quad (1.26)$$

1.4.2 g-2 Anomaly

The measurement of the muon anomalous magnetic moment from the Fermilab National Accelerator Laboratory muon g-2 experiment [28], combined with earlier results from the Brookhaven National Laboratory E821 measurement [48], finds the difference of a_μ between experiment and SM prediction to be,

$$\Delta a_\mu^{\text{obs}} = a_\mu^{\text{exp}} - a_\mu^{\text{SM}} = (251 \pm 59) \times 10^{-11}, \quad (1.27)$$

where $a_\mu = (g - 2)_\mu / 2$. This is a 4.2σ deviation away from the SM expectation. One potential solution to this deviation is a 2HDM. This contributes in two ways to Δa_μ , one-loop and two-loop Bar-Zee contributions [49, 50], that can be mediated by any of the additional Higgs bosons.

The one-loop contribution to Δa_μ is positive for ϕ and negative from A and H^\pm , however the more significant contribution comes from two-loop Bar-Zee diagrams




Figure 1.3: Examples of one-loop Bar-Zee feynman diagrams for the contribution from ϕ and A in (a) and H^\pm in (b).




Figure 1.4: Examples of two-loop Bar-Zee feynman diagrams for the contribution from ϕ and A in (a) and H^\pm in (b). Note that other fundamental particles can also contribute to the loop.

with heavy fermions in the loop which gives a positive shift to Δa_μ . In order to fulfil the requirements for the anomaly, an enhanced couplings between additional Higgs bosons to muons is needed. This gives two options: a type II or a type X [2HDM](#), both at large values of $\tan\beta$. The type II [2HDM](#), that also offers enhanced couplings to down-like quarks, is heavily constrained by Large Electron-Positron ([LEP](#)), Tevatron and [LHC](#) searches and an available region of phase space to explain the anomaly is not easily found. However, the type X [2HDM](#), with only lepton coupling enhancements at high $\tan\beta$, is relatively unconstrained due to suppressed quark initiated production modes of additional Higgs bosons.

Reference [16] puts an upper limit on the $m_A \lesssim 200$ GeV, that is also required for an explanation of $\Delta a_\mu^{\text{obs}}$. Further constraints are placed on the phase space from theoretical stabilities, electroweak precision measurements and collider bounds. The

final available phase space for a type X 2HDM in the alignment scenario to explain the g-2 anomaly is shown in Table 1.4.

Scenario	$\tan \beta$	m_A (GeV)	m_ϕ (GeV)	m_{H^\pm} (GeV)
Normal	≥ 90	[62.5,145]	[130,245]	[95,285]
Inverted	≥ 120	[70,105]	[100,120]	[95,185]

Table 1.4: Regions of interest for g-2 anomaly with respect to the type X 2HDM in the normal and inverted alignment scenarios as suggested in Reference [16].

Chapter 2

The LHC and CMS Experiment

This chapter will explain the apparatus used to collect the datasets that are utilised for the physics analysed described in Chapters 4 and 5. This is split into two parts. Firstly, an overview of the LHC and a description of how proton-proton collisions at a center-of-mass energy (\sqrt{s}) of 13 TeV are achieved. Secondly, is an explanation of the CMS detector, including a look at the individual sub-detectors, that are crucial to the reconstruction of particles originating from the proton-proton collisions.

2.1 The LHC

The LHC [51], located at the European Organization for Nuclear Research (CERN) just outside Geneva, is a synchrotron measuring 27 km in circumference, installed in the tunnel previously used by the LEP accelerator [52]. Upon design, its primary purpose was to provide collisions between proton beams, generating center-of-mass energies of up to 14 TeV and an instantaneous luminosity of approximately $10^{34} \text{ cm}^{-2}\text{s}^{-1}$. Fig 2.1 illustrates the arrangement of the LHC and the CERN accelerator complex. Protons are supplied to the LHC through a sequence of accelerators: Linac4 (Linac2 pre-2020), Proton Synchrotron Booster (PSB), Proton Synchrotron (PS), and Super Proton Synchrotron (SPS), that successively raise the energies of the protons to 50 MeV, 1.4 GeV, 25 GeV, and 450 GeV respectively. When the final energy has been achieved, the SPS injects bunched protons into the LHC's beam pipes as two counter-rotating beams. Each bunch containing over 10^{11} protons, with each one separated by 25 ns and each beam consists of 2808 bunches. The protons are accelerated to collision energy using eight 400 MHz radio frequency (RF) cavities and kept in a circular trajectory with 1232 niobium-titanium superconducting dipole magnets. These magnets must be maintained at their op-




Figure 2.1: A schematic diagram of the CERN [4].

erating temperature of 1.9 K to generate magnetic fields up to 8.4 T, requiring the use of superfluid helium. The bunches are collided at four intersections point surrounded by the ALICE [53], ATLAS [54], CMS [6] and LHCb [55] detectors, at a collision rate of 40 MHz.

The rate of events for a process produced in **LHC** collisions can be expressed as,

$$R_{\text{event}} = L_{\text{inst}} \sigma(\sqrt{s}), \quad (2.1)$$

where σ represents the cross section of the process and is a function of the center-of-mass energy, and L_{inst} denotes the **LHC** machine instantaneous luminosity, that depends only on the beam parameters and can be calculated by,

$$L_{\text{inst}} = \frac{n_b N_b^2 f_{\text{rev}} \gamma_r}{4\pi \epsilon_n \beta^*} F. \quad (2.2)$$

where n_b is the number of bunches per beam, N_b is the number of particles per bunch, f_{rev} is the revolution frequency, γ_r is the relativistic gamma factor, ϵ_n is the normalized transverse beam emittance, β^* is the beta-function at the collision point, and F is a reduction factor which accounts for the crossing angle of the beams at the collision point. One disadvantage of an increase to the instantaneous luminosity is the increase of pileup (PU), defined as the number of additional inelastic proton-proton collisions per bunch crossing.




Figure 2.2: The total integrated luminosity for proton-proton collisions collected by the CMS experiment between 2010 and May 2023 [5].

The LHC began its first physics collisions in 2010, initially colliding beams at $\sqrt{s} = 7$ TeV, which was increased to 8 TeV throughout 2012. Following this data-taking period, known as Run 1, the LHC underwent a two-year Long Shutdown 1 (LS1) to undergo upgrades in preparation for an increase in \sqrt{s} . In 2015, the LHC was restarted, initiating its Run 2 phase of data collection at $\sqrt{s} = 13$ TeV, which lasted until the end of 2018. During Run 2, the LHC achieved and surpassed its original design luminosity by obtaining a record peak luminosity of $2.1 \times 10^{34} \text{ cm}^{-2}\text{s}^{-1}$ in 2018. Subsequently, the LHC performed a second update period, the Long Shutdown 2 (LS2), lasting for approximately three years, where-after the data collection for Run 3 began at $\sqrt{s} = 13.6$ TeV. Fig 2.2 illustrates the total integrated luminosity.

ity of proton-proton collisions delivered to the CMS detector at the time of writing. The data used for the analyses described in this thesis correspond to the full Run 2 dataset collected during the 2016–2018 data-taking periods at 13 TeV. Only data recorded by CMS, where all sub-detectors were functioning correctly, are certified for use in physics analyses. This equates to 36.3 fb^{-1} , 41.5 fb^{-1} and 59.7 fb^{-1} of data collected in 2016, 2017 and 2018 respectively.

2.2 The CMS Detector

The CMS detector was engineered to fulfil the rigorous demands of the LHC physics program. Its primary objective is to achieve sensitivity to the Higgs boson and novel phenomena at the TeV energy scale. Weighing 12,500 tonnes and measuring 21.6 m in length with a diameter of 14.6 m, the CMS detector uses an array of sub-detectors encircling the central beam axis. The layout of the CMS detector is shown in Figure 2.3. A superconducting solenoid, operating at a magnetic field strength of 3.8 T, which surrounds the inner tracker, electromagnetic calorimeter, and hadronic calorimeter. Situated outside the solenoid within the iron return yoke are gaseous muon detectors, positioned to accurately measure muons. The CMS detector adopts a coordinate system centred at the collision point, with the y-axis oriented vertically, the x-axis directed radially inward toward the LHC center, and the z-axis aligned with the beam direction. Transverse energy (E_T) and transverse momentum (p_T) are defined in the x-y plane, and the azimuthal angle (ϕ) and polar angle (θ) are measured relative to the x-axis and z-axis, respectively. r is used as the radial distance in the x-y plane. Pseudorapidity (η), defined as $\eta = -\ln[\tan(\theta/2)]$, is used due to its gauge invariance, and distances in the ϕ - η plane are characterised by the metric $\Delta R = \sqrt{\Delta\phi^2 + \Delta\eta^2}$.

2.2.1 Tracker

Closest to the interaction point in the CMS experiment is the tracker [6, 56, 57], which is essential for precise measurements of charged particle trajectories and the determination of primary and secondary vertices. To meet the requirements of high granularity and fast response for the large number of particles generated in each bunch crossing, as well as being radiation-hard to deal with the particle flux, a silicon tracking detector is utilised. The tracker consists of a pixel detector and a




Figure 2.3: A perspective view of the CMS detector [6].

silicon strip detector as shown in Figure 2.4.




Figure 2.4: Schematic of the CMS tracker (pre-pixel upgrade) in the r - z plane, showing the position of the pixel detector as well as the TIB, TID, TOB, and TEC strip detectors. The lines represent detector modules and the double lines represent back-to-back modules [6].

The pixel detector covers the pseudorapidity range $|\eta| < 2.5$ and is composed of

three cylindrical pixel modules and two disk pixel modules. It contains 66 million silicon pixels, each with dimensions of $100 \times 150 \mu\text{m}^2$. This configuration provides a spatial resolution of $15\text{--}20 \mu\text{m}$ in both the $r\text{-}\phi$ plane and the z direction, enabling 3-dimensional vertex reconstruction. The original tracker was built to handle an instantaneous luminosity of $10^{34} \text{ cm}^2\text{s}^{-1}$ and an average PU of 25. To handle higher luminosities and increased event PU, the pixel detector was upgraded in 2016/2017 to handle approximately double the instantaneous luminosity and PU [57]. The upgraded detector consists of four barrel module layers and three endcap disks, resulting in 124 million pixels.

Surrounding the pixel detector is the silicon strip detector, which is divided into four subsystems: the tracker inner barrel (**TIB**), tracker inner disks (**TID**), tracker outer barrel (**TOB**), and tracker endcaps (**TEC**). The **TIB** and **TID** provide four layers of silicon strip detectors in the barrel region and three disks at each end, extending up to a radius of 55 cm. The **TOB** consists of six barrel layers extending up to an outer radius of 116 cm, while the **TEC** comprises nine disks covering a range of $|z|$ from 124 cm to 282 cm. The silicon strips have various thickness and widths, providing multiple measurements of the $r\text{-}\phi$ position with resolutions ranging from 23–35 μm in the **TIB** to 35–53 μm in the **TOB**.

In addition to the main components, the tracker includes back-to-back mounted micro-strip detector modules to provide additional measurements of the z coordinate in specific regions. The overall tracker layout ensures the presence of at least three hits in the pixel detector (at least four hits for the upgraded detector) and at least nine hits in the silicon-strip tracker, with a minimum of four 2-dimensional measurements among them.

2.2.2 Electromagnetic calorimeter

The CMS electromagnetic calorimeter (**ECAL**) is a hermetic homogeneous calorimeter designed to detect high energy electrons and photons [6, 58]. In total, 75,848 lead tungstate (PbWO_4) crystals are sorted in a barrel and endcap configuration. PbWO_4 was chosen as the crystal material due to its radiation-hardness, high density, short radiation length, and small Molière radius (the average radius containing on average 90% of a shower’s total energy deposit), enabling the construction of a compact and finely granular **ECAL**. Scintillation light produced by showering electrons and pho-

tons in the crystals is converted into an electrical signal by photodetectors such as avalanche photodiodes in the barrel and vacuum phototriodes in the endcaps. The scintillation decay time of the crystals matches the 25 ns [LHC](#) bunch crossing time, ensuring that a significant portion of the light is emitted between bunch crossings.

The [ECAL](#) is placed outside of the tracker but within the magnet bore and covers the pseudorapidity range $|\eta| < 3.0$. It comprises of an ECAL barrel ([EB](#)) covering $|\eta| < 1.479$ and an ECAL endcap ([EE](#)) covering $1.479 < |\eta| < 3.0$. A diagram of this is shown in Figure 2.5. The barrel region consists of 61,200 crystals with 360-fold granularity in ϕ and 170-fold granularity in η . The crystals are tapered with a front face area of 0.0174×0.0174 in $\eta\phi$ ($22 \times 22 \text{ mm}^2$) and a length of 230 mm. The endcaps house 7,324 crystals arranged in a rectangular x - y grid, each with a front face area of $28.62 \times 28.62 \text{ mm}^2$ and a length of 220 mm.

The [ECAL](#) system also includes preshower detectors placed in front of each endcap to identify photons from neutral pion decays and improve electron identification and position resolution. These detectors consist of lead radiators to initiate showering and silicon-strip sensors to measure the deposited energy. The [ECAL](#) energy resolution is parametrised as a function of the incident particle energy E , with terms for the stochastic S , noise N , and constant C contributions.

$$\left(\frac{\sigma}{E}\right)^2 = \left(\frac{S}{\sqrt{E}}\right)^2 + \left(\frac{N}{E}\right) + C^2 \quad (2.3)$$

The stochastic term captures fluctuations in lateral shower containment and photon yield, the noise term accounts for electronics noise and pileup, and the constant term arises from non-uniformity of the longitudinal response and calibration errors. Measurements using electron beams have determined the values of $S = 0.028 \text{ GeV}^{1/2}$, $N = 0.12 \text{ GeV}$, and $C = 0.003$ for the [ECAL](#) energy resolution.

2.2.3 Hadron calorimeter

The [CMS](#) detector includes the hadronic calorimeter ([HCAL](#)), which plays a crucial role in measuring the energies of strongly interacting particles such as hadronic jets and missing energy signals. The [HCAL](#) is located outside of the [ECAL](#) and covers the pseudorapidity region $|\eta| < 5.2$. It consists of four sub-detectors: the hadron barrel ([HB](#)), hadron endcap ([HE](#)), hadron outer ([HO](#)), and hadron forward ([HF](#)) calorimeters, arranged as shown in Figure 2.6.




Figure 2.5: A schematic of the CMS ECAL detector [6].




Figure 2.6: A schematic of the CMS HCAL showing the arrangement of the HB, HE, HO, and HF calorimeters [6].

The **HB** and **HE** calorimeters are positioned within the magnet bore, covering the pseudorapidity regions $|\eta| < 1.3$ and $1.3 < |\eta| < 3.0$, respectively. They are constructed using alternating layers of brass absorber plates and plastic scintillator tiles. Brass is chosen as the absorber material due to its short nuclear interaction length and non-magnetic properties. The **HB** provides 5.82 to 10.6 interaction lengths of absorber material, while the **HE** provides approximately 10 interaction lengths. The plastic scintillator tiles collect scintillation light emitted by charged particles in the hadronic showers, which is then read out using hybrid photodiodes.

To extend the **HCAL**'s containment capability in the central detector region ($|\eta| < 1.3$), the **HO** calorimeter is utilised. It is placed outside of the solenoid coil and utilizes plastic scintillator tiles as the active material. The **HO** enhances the thickness of the **HCAL** to a minimum of 11.8 interaction lengths, improving the measurement of late starting or highly penetrating showers.

In the forward regions ($|\eta| > 3.0$), the **HF** detector extends the pseudorapidity coverage of the **HCAL** up to $|\eta| < 5.2$. The **HF** is subjected to a high flux of incoming particles, and therefore, requires extremely radiation-hard materials. Quartz fibers embedded in steel absorbers are employed as the active material. Charged particles in the showers generate Cherenkov light in the quartz fibers, which is detected by photomultiplier tubes.

The energy resolution of the **HCAL** can be parametrised as a function of the incident particle energy E , using the formula,

$$\left(\frac{\sigma}{E}\right)^2 = \left(\frac{S}{\sqrt{E}}\right)^2 + C^2 \quad (2.4)$$

where the stochastic term S is measured to be $0.943 \text{ GeV}^{1/2}$ and the constant term C is measured to be 0.084. During the **LS1** period, upgrades were performed on the photomultiplier tubes of the **HF**.

2.2.4 Muon system

The muon system in the **CMS** detector plays a crucial role in the accurate identification and measurement of muons, which are essential for various physics analyses. The system is designed with three primary functions: efficient muon identification, precise momentum measurement, and triggering capability.

Located outside the solenoid coil and **HO** calorimeter, the muon system is strategically positioned between the iron plates forming the flux return yoke. This arrangement takes advantage of the high field solenoidal magnet and the yoke’s structure to enable the desired functions. The muon system covers a wide pseudorapidity range, specifically $|\eta| < 2.4$. It consists of three types of gaseous detectors: drift tube (**DT**) chambers, cathode strip chambers (**CSC**), and resistive plate chambers (**RPC**).

In the region of $|\eta| < 1.2$, the muon system employs **DT** chambers, which are organised into four stations. These include inner and outer stations positioned inside and outside the magnet return yoke, as well as two stations inter-spaced between the iron layers of the yoke. The **DT** chambers are composed of rectangular drift cells filled with a mixture of argon and carbon dioxide gas. Each cell features an anode wire running along its length. The cells are arranged into superlayers, each containing four layers of cells. The **DT** chambers in the three innermost muon stations consist of three superlayers, whereas the outer station contains only two superlayers. The spatial resolution of the **DT** chambers is measured to be 77–123 μm for the ϕ coordinate measurement and 133–393 μm for the z coordinate measurement.

In the region of $0.9 < |\eta| < 2.4$, the **CSC**s are employed. These chambers are trapezoidal-shaped multiwire proportional chambers with six layers of gas. The gas mixture used includes argon, carbon dioxide, and tetrafluoromethane. The **CSC** modules are organized into four stations positioned between the endcap layers of the magnet’s return yoke. They consist of layers of cathode strips and anode wires, allowing for measurements of the muon’s ϕ and r coordinates, respectively. The spatial resolution per chamber of the **CSC**s ranges from 45 μm to 143 μm .

To enhance the system’s capabilities in the $|\eta| < 1.6$ region, **RPC**s are used. These parallel-plate detectors feature a double-gap module design with anode and cathode plates separated by a gas gap. The **RPC**s are embedded within the barrel and endcap iron yokes. In the barrel, six layers of **RPC** chambers are present, with two layers in each of the two innermost muon stations and one layer in each of the two outer stations. In the endcap, the **RPC**s consist of four layers positioned at either side of the three iron disks. While the **RPC**s exhibit a lower spatial resolution compared to the **DT**s and **CSC**s (0.78–1.38 cm per chamber), they offer a remarkably fast response time, making them suitable for dedicated muon triggering.

To achieve precise muon momentum resolution, information from the tracker hits is combined with the measurements obtained from the muon systems. The standalone muon system alone achieves a momentum resolution of approximately 9% for muons with transverse momentum (p_T) up to 200 GeV. However, for muons with a p_T of 1 TeV, the momentum resolution varies between 15% and 40% depending on the muon’s pseudorapidity.

2.2.5 Triggering and Computing

The [LHC](#) collides protons at a rate of 40 MHz during its operation. However, it is impractical to read out and store every event due to the high data volume involved. To address this challenge, the [CMS](#) detector employs a dedicated trigger system that selectively chooses the most interesting events, reducing the recorded rate to around 1 kHz.

The [CMS](#) trigger system consists of two stages: the Level-1 ([L1](#)) trigger and the high-level trigger ([HLT](#)). The [L1](#) trigger, implemented with custom built programmable electronics, operates as the first stage and reduces the event rate to approximately 100 kHz. During the upgrade in 2015/2016, the [L1](#) system was enhanced to accommodate the increased instantaneous luminosity and pileup conditions. Upgrades included improved muon momentum resolution, electron/photon isolation, hadronic tau identification and isolation, jet finding algorithms with pileup subtraction, and enhanced trigger menu capabilities.

The [L1](#) trigger utilises a time multiplexed architecture, where energy deposits recorded in the calorimeters and hits from the muon detectors are processed. In the calorimeter trigger, energy deposits from the HCAL and ECAL are passed through two layers (Layer 1 and Layer 2), enabling object identification and the selection of the best candidates based on their p_T . Simultaneously, hits from the muon detectors are combined in the muon trigger’s tracking-finder and sorting/merging layers, producing a sorted list of muon candidates for the entire detector. The global trigger then integrates the information from both calorimeter and muon triggers to make a decision on event selection within a maximum storage time of 3.2 μ s. A diagrammatic representation of this is shown in Figure 2.7.




Figure 2.7: A schematic of the L1 trigger workflow for object reconstruction [7].

The second stage of the trigger system is the [HLT](#), which is a software based trigger running on a multi-processor farm with thousands of CPU cores. The [HLT](#) utilises information from the full detector, including the tracker, calorimeters, and muon systems. By closely following the algorithms and selections used offline, the [HLT](#) ensures good momentum resolution and high identification efficiencies for selected objects. From the initial event rate of approximately 100 kHz received from the [L1](#) trigger, the [HLT](#) further reduces the output rate to about 1 kHz.

The events selected by the [HLT](#) produce a significant amount of data, approximately $\mathcal{O}(10 \text{ pb})$ per year. To process, store, and facilitate easy access for analysts worldwide, [CMS](#) collaborates with other [LHC](#) experiments through the Worldwide LHC Computing Grid ([WLCG](#)). The [WLCG](#) combines computing resources from research institutions globally, organized into three tiers. Tier-0, comprising the CERN Data Center and the Wigner Research Centre in Budapest, performs full data reconstruction and stores a copy of the data on tape. The data is then distributed to Tier-1 centers, which handle the storage of raw, reconstructed, and simulated data, as well as the distribution to Tier-2 sites. Tier-2 sites, located at universities

and research institutes worldwide, provide computing resources for data production, reconstruction, and analysis by researchers.

Chapter 3

Object Reconstruction

This chapter will overview the methods used to reconstruct the objects in each event, that are critical for the searches described in Chapters 4 and 5. The topics discussed here are the reconstruction of tracks and vertices, the particle flow algorithm, the calculation of the missing transverse energy ([MET](#)), the measurements of jets and the tagging of their flavours, and the identification of electrons, muons, and tau leptons.

3.1 Tracks and vertices

The Combinatorial Track Finder ([CTF](#)) algorithm [59] is employed by [CMS](#) to reconstruct particle tracks. This algorithm consists of four steps to estimate trajectory parameters and uncertainties. Initially, track seeds are formed using hits in the first few layers of the tracker. These seeds provide initial estimates of the trajectory. Next, the track seeds are extrapolated using a Kalman filter [60], searching for additional hits along the expected trajectory in successive detector layers. The trajectory parameters are continuously updated as hits are added to the track candidate. The extrapolation process continues until the final detector layer is reached.

Afterwards, a Kalman filter and smoother are used to fit the final trajectory iteratively. Spurious hits are identified and removed after each iteration until no more spurious hits are found. This fitting process provides the most accurate estimates of the trajectory parameters. Subsequently, tracks must pass a set of quality criteria, and any failing tracks are discarded.

The [CTF](#) algorithm performs six iterations, gradually reconstructing more challeng-

ing tracks and recovering missed tracks from previous iterations. Associated hits are removed after each iteration, reducing combinatorial complexity and facilitating the search for complicated tracks in later iterations. Tracking efficiencies for muons and pions with transverse momentum (p_T) greater than 500 MeV have been measured to be over 99.3% and 98.5%, respectively [61].

Once all tracks are reconstructed, the positions of interaction vertices, including the primary vertex and additional vertices from PU, are determined. Vertex reconstruction begins by selecting tracks consistent with being promptly produced near the beamspot. Tracks are then clustered based on the z coordinate of their closest approach to the beamspot, using a deterministic annealing algorithm [62]. Candidate vertices are retained if at least two of their associated tracks are incompatible with other vertices. The candidate vertices are fitted using an adaptive vertex fitter [63] to determine the best estimate of their three dimensional positions. The primary interaction vertex is determined as the vertex with the highest summed physics-object p_T , where physics objects are jets clustered from tracks associated with the candidate vertex using the anti- k_T algorithm [63]. The efficiency of vertex reconstruction has been measured, with values exceeding 98.7% for vertices containing at least two tracks and exceeding 99.9% for vertices with four or more tracks [61].

3.2 Particle flow

The particle flow (PF) algorithm [64–66] is used for the reconstruction of long lived particles in an event, such as electrons, photons, muons, and hadrons. All the sub-detectors of CMS are used to achieve accurate identification and measurements of particle energies and directions. The final output of the algorithm is a list of particles and their four vectors, that can then be used to construct more complicated objects such as jets, hadronically decaying taus, and the MET.

The tracking and calorimeter clustering information are the initial inputs to the PF algorithm. The tracking information includes the high resolution momentum and direction of charged hadrons calculated by an iterative-tracking strategy [67], that is not achievable with the calorimeters. The calorimeter clustering is used to achieve multiple objectives, including detecting and measuring the energy and direction of neutral particles, separating neutral particles from charged hadrons, reconstructing electrons and Bremsstrahlung photons, and aiding in the energy measurement

of charged hadrons with inaccurate track parameters. The clustering algorithm involves identifying “cluster seeds”, growing “topological clusters”, and generating “particle flow clusters” based on energy thresholds and cell-cluster distances in each sub-detector, except for the **HF**, where each cell forms a cluster, and it explained in more detail below.

The clustering algorithm used in the calorimeters can be split into three stages. Firstly, “cluster seeds” are identified as local maxima of energy in calorimeter cells above a specified threshold. Secondly, “topological clusters” are formed by combining cells that have at least one side in common with an existing cluster cell and exceed a given energy threshold. These thresholds corresponding to the electronics noise are 80 MeV in the **ECAL** barrel, 300 MeV in the **ECAL** end caps and 800 MeV in the **HCAL**. The number of seeds dictates how many “particle flow clusters” are formed from a topological cluster. Utilising the granularity of the calorimeter, an iterative process is employed to determine cluster energies and positions by distributing the energy of each cell among all particle flow clusters based on the distance between the cell and the cluster.

Muon reconstruction involves the matching of tracks from the inner tracker to tracks in the muon system [68,69]. Initially, tracks are independently reconstructed in each system. Standalone muon tracks are reconstructed in the muon system using seed groups of **CSC** or **DT** segments. These seeds provide an initial trajectory estimate, which is extended using a Kalman filter (**KF**) to incorporate hits from **DT**, **CSC**, and **RPC** sub-detectors. In the tracker muon reconstruction, tracks in the inner tracker with p_T greater than 0.5 GeV and total momentum greater than 5 GeV are extrapolated to the muon system. A track qualifies as a tracker muon if it is matched to at least one muon segment based on their positions in a local x-y coordinate frame. The global muon reconstruction starts with standalone muon tracks and matches them to tracks in the inner detector. If a match is found, a combined fit using the **KF** is performed with hits from both systems.

Electrons reconstruction initially works by finding matching tracks with **ECAL** energy clusters [8]. The path of electrons towards the **ECAL** involves traversing a considerable amount of material, which constitutes the tracker. The thickness of this material varies depending on the η region, ranging from approximately 0.4 to 2 radiation lengths [8]. Within this material, electrons emit bremsstrahlung photons,

resulting in a significant energy loss. At regions with maximal material thickness, electrons lose an average of 86% of their energy, predominantly spreading out in the ϕ direction due to magnetic field induced bending [8]. To accurately measure the initial electron energy, the energy from all emitted photons is measured through the supercluster (**SC**) algorithms. These algorithms initiate clustering with seed crystals, expand the clusters based on energy thresholds, and merge them into the **SC**.

For reconstructing electron tracks, the standard **CTF** algorithm is not suitable due to the substantial radiative losses in the tracker material. These losses lead to reduced hit collection efficiency and inaccurate trajectory parameter estimation. To address this, a dedicated tracking procedure is employed, which identifies seeds consisting of two or three hits in the tracker using two complementary methods: **ECAL**-based seeding and tracker-based seeding. **ECAL**-based seeding uses SC energy and position to estimate the trajectory, while tracker-based seeding starts with tracks reconstructed using the **CTF** algorithm. The selected seeds are then extrapolated and smoothed the Gaussian sum filter (**GSF**) instead of the **KF**. The **GSF**, with a Gaussian mixture to model bremsstrahlung energy loss, provides a more accurate estimation of the trajectory parameters.

PF elements are connected by a link algorithm that evaluates the quality of the connection between each pair of elements by defining a distance. Elements linked directly or indirectly form “blocks” that serve as inputs for the particle reconstruction and identification algorithm. The granularity of the **CMS** sub-detectors ensures that blocks typically contain one to three elements, allowing the algorithm to perform independently of event complexity. The link between a charged-particle track and a calorimeter cluster is established by extrapolating the track to different depths in the **ECAL** and **HCAL**, and the track is linked to a cluster if the extrapolated position lies within the cluster boundaries. Similarly, links between calorimeter clusters are established based on their positions in the more granular calorimeter. A link between a charged particle track in the tracker and a muon track in the muon system is established through a global fit, with the link distance determined by the χ^2 value of the fit.

The **PF** algorithm operates on each block by following a specific set of steps. Initially, global muons within the block are converted into “**PF** muons” if their combined momentum aligns with that determined solely from the tracker. Afterwards, elec-

tron reconstruction and identification take place, where tracks are pre-identified as potential electrons based on their characteristics. Identified electrons become “[PF](#) electrons”, and their corresponding tracks and [ECAL](#) clusters are removed.

Next, tighter quality criteria are applied to the remaining tracks, ensuring that the measured p_T uncertainty is smaller than the expected calorimetric energy resolution for charged hadrons. Some tracks are rejected based on these criteria, but the energy from genuine tracks is still used in the reconstruction process. The remaining elements in the block can give rise to charged hadrons, photons, neutral hadrons, or additional muons.

To detect neutral particles, tracks are linked to [ECAL](#) and [HCAL](#) clusters by comparing momenta and energy measurements. Multiple tracks can be linked to the same [HCAL](#) cluster, while the closest cluster is retained when a track is linked to multiple [HCAL](#) clusters. Similar considerations apply to [ECAL](#) clusters, where ordering based on distance is used to preserve appropriate links.

In cases where the calibrated calorimetric energy is significantly lower than the total track momentum, a search is conducted for additional muons and fake tracks. Tracks are progressively removed based on p_T uncertainty until a certain threshold is reached. The impact of this procedure on the overall track selection is minimal.

Finally, the remaining tracks in the block are identified as “[PF](#) charged hadrons”, with momentum derived directly from the track measurements. If the calibrated energy of linked [ECAL](#) and [HCAL](#) clusters exceeds the associated charged particle momentum significantly, “[PF](#) photons” and possibly “[PF](#) neutral hadrons” are created. The precedence is given to photons over neutral hadrons due to their higher contribution in jet energy. Unclassified [ECAL](#) and [HCAL](#) clusters give rise to “[PF](#) photons” and “[PF](#) neutral hadrons” using the calibration procedure applied to [HCAL](#) clusters.

3.3 Muons

Three muon identification techniques are described by the [PF](#) algorithm; tracker, global and [PF](#) muons. Tracker muons and global muons differ in their requirements for muon segments. Subsequently, tracker muon reconstruction efficiency if larger




Figure 3.1: The p_T resolution for muon identification using the muon system (black), the inner tracker (blue), and the full system (red). This is shown for the psuedorapidity regions $0 < \eta < 0.8$ (a) and $1.2 < \eta < 2.4$ [6].

than global reconstruction efficiency, when dealing with muons with a momentum below 5 GeV. This is because many low momentum muons are likely to reach only up to the first muon station. However, global muon reconstruction enhances momentum resolution for muons with a p_T exceeding 200 GeV, whilst below this the inner tracker information is dominant to the resolution, as shown in Figure 3.1.

The **Medium** muon identification [69] is used for the analyses described in subsequent chapters. The requirements for this identification are:

- i) The muon is reconstructed by the tracker or global muon reconstruction algorithm.
- ii) The impact parameters, d_{xy} and d_z , defined as the distance in the x-y and z plane of the closest approach to the primary vertex (**PV**), between the muon track and the primary vertex are restricted as $d_{xy} < 0.045$ cm and $d_z < 0.2$ cm to ensure the muon is associated with the primary vertex.

- iii) At least 80% of the tracker hits have to be valid.

In addition, either of the following two sets of criteria must be satisfied:

- i) The muon is reconstructed by the global muon reconstruction algorithm.
- ii) The χ^2 divided by the number of degrees of freedom of the global track fit is smaller than 3.
- iii) The χ^2 of the tracker standalone position match [70] is smaller than 12.
- iv) The χ^2 of the track kink finder [69] is less than 20.
- v) The muon segment compatibility is > 0.303 .

or,

- i) The muon segment compatibility is > 0.451

The efficiency of the **Medium** muon identification is approximately 90%, with a misidentification probability of approximately 0.1%.

To reduce the contamination from muons originating from heavy-flavoured quark decays, muons are required to be isolated from hadronic activity in the detector. The isolation is based on the p_T of the photon (γ), neutral (h^0) and charged hadron (h^\pm if originating from the **PV** and h^\pm , **PU** if not originating from **PV**) **PF** candidates within a cone size of $\Delta R < 0.4$ of the selected muon (μ). A relative combined isolation variable is defined as:

$$I_{\text{rel}} = \frac{1}{p_T^\mu} \left(\sum p_T^{h^\pm} + \max \left(0, \sum p_T^{h^0} + \sum p_T^\gamma - \Delta\beta \sum p_T^{h^\pm, \text{PU}} \right) \right), \quad (3.1)$$

where the sums loop through all objects with the cone and $\Delta\beta$ is the energy estimate of neutral particles due to pileup, which is taken to be 0.5. This represents an estimated fraction of transverse momenta between the other contributions within the cone that originate from the interaction at the **PV**, and the muon itself. The muon isolation can then be used to separate muons, with a low value of I_{rel} , from other hadronic objects, at higher values of I_{rel} .

3.4 Electrons

In addition to the **PF** algorithm to reconstruct electrons, they are required to pass an identification variable based on a boosted decision tree (**BDT**) discriminator to distinguish genuine electrons from backgrounds like misidentified jets, electrons from photon conversions, and electrons produced in heavy-flavour decays [8]. The following variables are used as input to the **BDT**:

- i) Shower shape variables.
- ii) The fraction of energy deposited in the **HCAL**.
- iii) The fraction of energy lost through bremsstrahlung.
- iv) Track quality variables.
- v) The variable $1/E_{SC} - 1/p$, where E_{SC} is the **SC** energy and p is the momentum of the track at the point of closest approach to the vertex.
- vi) Variables comparing the geometrical matching between the track and the **SC**.

The **BDT** was trained on a Z/γ^* monte carlo (**MC**) sample generated with MADGRAPH, in 3 η bins for electrons with $p_T > 10$ GeV. The performance of the **BDT** is shown in Figure 3.2. This thesis uses the version of the identification **BDT** without the isolation included in the training, instead using an additional cut on the electron isolation. This allows for easily accessible sideband regions (orthogonal regions to the signal) used for estimation of background estimation.

The electron are required to pass the 90% efficiency working point of the **BDT**, which corresponds to a misidentification probability of 2% [8]. Finally electrons are also subject to the same impact parameter requirements as the muons: the impact parameters d_{xy} and d_z between the electron track and the primary vertex are restricted as $d_{xy} < 0.045$ cm and $d_z < 0.2$ cm to ensure the electron is associated with the primary vertex.

Similarly to muons, an isolation variable is used to further separate electrons from other backgrounds. Different to for muons, the isolation used is based on photon, neutral and charged hadron **PF** candidates within a smaller cone size of $\Delta R < 0.3$ of the selected electron. A different method to estimate the neutral particles due




Figure 3.2: Performance of the electron BDT using $Z \rightarrow ee$ enriched events from data (dots), background-enriched events from data (triangles), and $Z \rightarrow ee$ MC events (purple solid histogram). This is shown for the performance in the barrel (a) and the endcaps (b) for a BDT that does not utilise trigger information during training [8].

to PU is used, namely the rho-effective-area method. The pileup in this method is estimated as $PU = \rho \cdot A_{\text{eff}}$, where ρ is the event-specific average pile-up energy density per unit area in the ϕ - η plane and the A_{eff} is the effective area specific to the given type of isolation. The rho-effective-area subtracted relative combined isolation variable is defined as

$$I_{\text{rel}} = \frac{1}{p_T^e} \left(\sum p_T^{h^\pm} + \max \left(0, \sum p_T^{h^0} + \sum p_T^\gamma - \rho A_{\text{eff}} \right) \right), \quad (3.2)$$

where A_{eff} is measured in bins of η as listed in Table 3.1.

3.5 Jets

Quarks and gluons undergo fragmentation and hadronisation, resulting in collimated sprays of hadrons called jets [71]. The process of combining particles into jets is accomplished through the anti- k_T jet clustering algorithm [72] as implemented in FastJet [73]. This algorithm relies two distance parameters: d_{ij} , representing the

η range	A_{eff}
$0.0 \leq \eta < 1.0$	0.1440
$1.0 \leq \eta < 1.479$	0.1562
$1.479 \leq \eta < 2.0$	0.1032
$2.0 \leq \eta < 2.2$	0.0859
$2.2 \leq \eta < 2.3$	0.1116
$2.3 \leq \eta < 2.4$	0.1321
$2.4 \leq \eta < 5.0$	0.1654

Table 3.1: Electron effective areas used for the ρ -corrected isolation computation [8].

distance between objects i and j , and d_{iB} , representing the distance between object i and the beam. These objects can be individual particles or clusters of particles. The distance parameters are defined as,

$$d_{ij} = \min(p_{T,i}^{-2}, p_{T,j}^{-2}) \frac{\Delta R_{ij}^2}{R^2}, \quad (3.3)$$

$$d_{iB} = p_{T,i}^{-2},$$

where ΔR_{ij} is the separation in δR of objects i and j , and R is the radius parameter controlling the typical size of the jets. The clustering process unfolds as follows:

- i) Compute the minimum of d_{ij} and d_{iB} for all objects.
- ii) If the minimum corresponds to d_{ij} , merge objects i and j into a single entity.
- iii) If the minimum corresponds to d_{iB} , set object i as a jet and remove it from the object list.
- iv) Repeat steps (i) to (iii) until no objects remain.

Chapters 4 and 5 use jets clustered with a radius parameter of $R = 0.4$, often called AK4 jets. The impact of PU on jet energy and substructure is addressed by using the charged hadron subtraction (**CHS**) technique [74]. **CHS** excludes charged hadrons that are not associated with the primary vertex during jet clustering. To reject misidentified jets a discriminator evaluated on variables like energy fractions carried by different types of **PF** candidates, as well as charged and neutral particle multiplicities is used [74]. For data collected in 2017, the region with $2.65 < |\eta| < 3.139$

experiences substantial **ECAL** noise, leading to a notable increase in the number of jets. To veto this, jets with $p_T < 50$ GeV within this η range are removed and a **PU** jet identification discriminant [75]. For τ selections in the later chapters, to exclude selected jets from electron, muon or hadronic tau candidates, the jets are required to be separated from the selected τ candidate by $\Delta R > 0.5$.

3.6 b jets

For determining whether a jet is initiated by a b quark, the **DeepJet** algorithm [76, 77] is used. This heavy-flavour jet identifier, uses properties of the jet to distinguish jets originating from b quarks, c quarks, the remainder of the lighter quarks grouped, and gluons. Core to the b tagging in this algorithm is the reconstruction of a secondary vertex (**SV**) due to the lifetime of b quarks, allowing for displaced tracks. The **DeepJet** deep neural network (**DNN**) uses a combination of high-level variables such as the **SV**, as well as utilising low-level variable of jet constituents to separate between initiators. The **DNN** uses a mixture of convolutional and dense layers to perform this categorisation. The performance of the algorithm is shown in Figure 3.3.

For b jet selection, jets are required to have $p_T > 20$ GeV and $|\eta| < 2.4$ (2.5) in 2016 (2017 and 2018), and are considered b tagged if their discriminator value is larger than some threshold that represents a mis-identification rate of 1%. This corresponds to a b tagging efficiency around 80%.

3.7 Missing transverse energy

The **MET** is used as a feature to understand particles passing through the **CMS**, such as neutrinos in the **SM**, and weakly interacting particles from hypothetical **BSM** extensions. This information cannot be pulled from the detector itself but instead the absence of detection can be used to infer the presence of such an object. The **MET** is nominally calculated as the negative vector sum of all transverse momenta of the **PF** candidates in the collision event,

$$\vec{E}_T^{\text{miss}} = - \sum_i \vec{p}_T^i. \quad (3.4)$$

However, this raw **PFMET** is inaccurate due to the p_T thresholds of the calorimeters, the non-linearity of the calorimeter response and due to reconstruction inefficiencies.




Figure 3.3: Mis-identification probability 2σ bands against efficiency for b tagging using the DeepJet algorithm, for AK4jets with $p_T > 30$ GeV AND $|\eta| < 2.5$ utilising MC with parameters from the 2017 era of data taking. Two bands are displayed, one for the identification of a b quark rather than light quarks and gluons (blue) and one for the identification of a b quark rather than a c quark (red) [9].

cies [78]. This is fixed with jet energy corrections, determined for jets with $p_T > 15$ GeV and less than 90% for their energies deposited in the **ECAL** for each of the problems mentioned, and then the corrected **MET** is calculated by,

$$\vec{E}_T^{\text{miss,corr}} = \vec{E}_T^{\text{miss}} - \sum_{\text{jets}} (\vec{p}_{T,\text{jet}}^{\text{corr}} - \vec{p}_{T,\text{jet}}^{\text{raw}}), \quad (3.5)$$

where “corr” and “raw” are the corrected and uncorrected values of the jet momenta, respectively.

Further corrections are required to account for **PU** effects, and the PUPPI algorithm is used for this purpose [79]. The algorithm is designed to address the impact of **PU** on observables involving clustered hadrons, such as jets, missing transverse momentum, and lepton isolation. It achieves this by combining information about the local particle distribution, event **PU** properties, and tracking data. PUPPI operates at the level of individual particle candidates, before any clustering is performed. It assigns a weight to each particle, ranging from 0 to 1, based on the information from surrounding particles. A weight of 1 is given to particles believed to originate from the **PV**. These per-particle weights are then used to rescale the four-momenta of the particles, effectively correcting for the impact of **PU** and so altering the raw calculation of the **MET** to,

$$\vec{E}_T^{\text{miss}} = - \sum_i w_i \cdot \vec{p}_T^i. \quad (3.6)$$

3.8 Taus

Fundamental to this thesis is the identification of tau particles. The tau lepton is measured to have a mean lifetime of 2.9×10^{-13} s. This short lifetimes means that the tau lepton is not directly observable in the **CMS** detector. In order to detect these particles, it is important to understand how the tau decays. Due to the heavy nature of the particle, it does not only decay leptonically, but unlike the muon, it can also decay hadronically. A list of prominent decays of the tau lepton are shown in the Table 3.2. These decays can be split into three groups: the 17.8% of taus that decay to an electron (e), the 17.4% that decay into a muon (μ), and hadronic tau decays (τ_h) that make up the final 64.8% of tau decays. The leptonic decays of the tau can be accounted for by the identification of electrons and muons as discussed

in the previous subsection.

Decay Mode	Branching Fraction
Leptonic Decay (e, μ)	35.2%
$e^- \bar{\nu}_e \nu_\tau$	17.8%
$\mu^- \bar{\nu}_\mu \nu_\tau$	17.4%
Hadronic Decay (τ_h)	64.8%
$h^- \pi^0 \nu_\tau$	25.9%
$h^- \nu_\tau$	11.5%
$h^- h^+ h^+ \nu_\tau$	9.8%
$h^- 2\pi^0 \nu_\tau$	9.3%
$h^- h^+ h^- \pi^0 \nu_\tau$	4.8%
other	3.2%

Table 3.2: Measured branching fractions for the tau lepton. h represents a charged hadron either a pion or a kaon [17].

A two step process is used to identify hadronic taus. The hadron-plus-strips ([HPS](#)) algorithm is used to initially identify hadronic taus based on jets produced by the anti- k_T algorithm with a distance parameter of $\Delta R = 0.4$. To capture the energy deposits left by π^0 candidates in the ECAL, the photon and electron constituents of the jet responsible for seeding the tau reconstruction are assembled into strips. All electrons or photons used are required to have $p_T > 0.5$ GeV. The initial iteration of the [HPS](#) algorithm used a fixed strip size of $\Delta\eta \times \Delta\phi$ equal to 0.05×0.20 . However, this technique was updated to a dynamical strip size to account for the multiple scatterings of $e^+ e^-$ products from a pi^0 decay, falling outside of the fixed window. A reliance of the hadronic tau p_T spectrum of the required strip size was observed and so the following iterative algorithm was proposed to resolve this issue.

- i) The highest p_T electron or photon (not previously grouped into a strip) is used to initiate a new strip.
- ii) The second highest p_T electron or photon deposition within,

$$\Delta\eta = f(p_T^{e/\gamma}) + f(p_T^{\text{strip}}), \quad \Delta\phi = g(p_T^{e/\gamma}) + g(p_T^{\text{strip}}) \quad (3.7)$$

of the strip is then combined with the strip. These functions are determined from a fit to hadronic tau MC, so that 95% of electrons and photons are con-

tained in a single strip. These fits are shown in Figure 3.4 and correspond to,

$$f(p_T) = 0.20 p_T^{-0.66}, \quad g(p_T) = 0.35 p_T^{-0.71} \quad (3.8)$$

where the p_T is in units of GeV. These functions have lower and upper caps of 0.05 to 0.3 for $\Delta\phi$ and 0.05 to 0.15 for $\Delta\eta$.

- iii) Recalculate the strip position using a weight average of the average p_T of all the electron and photon strip constituents.

$$\eta_{\text{strip}} = \frac{1}{p_T^{\text{strip}}} \sum p_T^{e/\gamma} \eta_{e/\gamma}, \quad \phi_{\text{strip}} = \frac{1}{p_T^{\text{strip}}} \sum p_T^{e/\gamma} \phi_{e/\gamma} \quad (3.9)$$

- iv) Repeat steps ii and iii until no other electron or photon candidate fulfilling condition ii is found.




Figure 3.4: The distance between the hadronic tau and electron or photon η (a) and ϕ (b) with respect the the electron or photon p_T . The binned values (points) and the fitted functions f and g (red line), that encapsulates 95% of all electron and photons are shown in both cases [10].

Charged hadrons (prongs) are also required to have $p_T > 0.5$ GeV and originate from the [PV](#), with a loose transverse impact parameter of $d_{xy} < 0.1$ cm. Further constraints are placed on the reconstructed masses of the specific resonances produced by a hadronic tau decay, if produced by the decay. In particular, the visible mass positions and widths of the grouped charged hadrons and strips are optimised to match the ρ (770 MeV) and a_1 (1260 MeV) decays. This is performed to maximise the fraction of

hadronic tau efficiency to the probability of misidentification from jets. The visible decay products of each decay shown in Table 3.2 are reconstructed by:

- i) $h^-\pi^0$: One charged hadron candidate and no strips.
- ii) h^- : One charged hadron candidate and one strip with mass $0.3 < m_\tau < 1.3\sqrt{p_T/100}$ GeV. The mass is required to be between 1.3 and 4.2 GeV.
- iii) $\pi^-\pi^-\pi^+$: Three charged hadron candidates with mass $0.8 < m_\tau < 1.5$ GeV. The tracks are required to originate within $\Delta z < 0.4$ cm of the same vertex.
- iv) $h^-2\pi^0$: One charged hadron candidate and two strips. The τ_h mass should be $0.4 < m_\tau < 1.2\sqrt{p_T/100}$ GeV. The mass is required to be between 1.2 and 4.0 GeV.
- v) $\pi^-\pi^-\pi^+\pi^0$: Three charged hadron candidates and one strip.

The remaining hadronic decays of the tau leptons are not included in this thesis. The decay mode (**DM**) of the hadronic tau that is reconstructed by a **HPS** is quantified by the following formula relating the number of charged hadron candidates N_C and the number of strips N_N .

$$\text{DM} = 5(N_C - 1) + N_N \quad (3.10)$$

The second step of the identification, comes from a multiclass **DNN**-based algorithm named **DeepTau**, that seeks to discriminate hadronic tau decays from electron, muons, and most importantly quark or gluon jets, that can be misidentified as hadronic tau decays [11]. It uses a **DNN** architecture that consists of multiple interconnected layers of nodes, that attempts to learn whether the input object is an hadronic tau decay, an electron, a muon or a jet. The algorithm takes inputs from reconstructed particles surrounding the **HPS** hadronic tau candidate, including information about energies, momenta, and spatial positions. Convolutional layers are used to efficiently process these inputs by dividing them into smaller regions in η - ϕ space, which allows the algorithm to extract local patterns and features. It also incorporates high-level features of the hadronic tau candidate calculated from the **HPS** algorithm, such as the four-momentum, charge, **DM**, isolation variables used in previous multivariate analysis (**MVA**) [12], impact parameters, η and ϕ strip information, as well as event level information such as variables related to the **PU**. The architecture of algorithm is shown in Figure 3.5.




Figure 3.5: The architecture of the DeepTau neural network, comprising of three sets of input variables: inner cells, outer cells, and high-level features. These sets are processed separately through subnetworks and their outputs are concatenated. Five fully connected layers process the concatenated output to calculate the probabilities for a candidate to be a hadronic tau, electron, muon, or jet. The high-level input subnetwork consists of three fully connected layers, taking 47 inputs and yielding 57 outputs. Complex subnetworks process the features of inner and outer cells separately, with fully connected layers followed by concatenation and additional fully connected layers. Convolutional layers progressively reduce the grid size for both inner and outer cells. For inner cells, there are 5 convolutional layers, while for outer cells, there are 10 convolutional layers. The number of trainable parameters (TP) for the different subnetworks are also provided [11].

The DeepTau DNN is trained using a large dataset, incorporating examples of hadronic tau decays and background processes of electrons, muons and jets. The output of the DNN is 4 scores, that represent the probability that the object is a hadronic tau (y_τ), an electron (y_e), a muon (y_μ) or a jet (y_{jet}). From these raw scores, additional scores are calculated for the probability that an object is a hadronic tau rather than an electron, muon or jet. These are defined as,

$$D_i^{\text{score}} = \frac{y_\tau}{y_i + y_\tau}, \quad i \in (e, \mu, \text{jet}). \quad (3.11)$$

From this, working points (WP)s are defined to match specific efficiencies of hadronic tau identification with respect to each of the other 3 objects, named D_e^{WP} , D_μ^{WP} and $D_{\text{jet}}^{\text{WP}}$. The target efficiencies for different WP are shown in Table 3.3.

WP	VVTight	VTight	Tight	Medium	Loose	VLoose	VVLoose	VVVLoose
D_e^{WP}	60%	70%	80%	90%	95%	98%	99%	99.5%
D_μ^{WP}	-	-	99.5%	99.8%	99.9%	99.95%	-	-
$D_{\text{jet}}^{\text{WP}}$	40%	50%	60%	70%	80%	90%	95%	98%

Table 3.3: Target efficiencies of the DeepTau working points with respect to electrons, muons and jets.

Due to the lack of muons misidentified as hadronic taus, high efficiencies can be required for D_μ^{WP} . This is also the reason for why the target efficiencies for D_e^{WP} are higher than for $D_{\text{jet}}^{\text{WP}}$, where the large challenge jets misidentified as hadronic taus is present. The misidentification probabilities for the different hadronic tau efficiencies are shown in Figure 3.6. Also shown in these plots, is the performance comparison to an older MVA algorithm for electrons and jets, as well as a cut based algorithm for muons, both described in Reference [12]. Significant improvements with respect to the previous algorithm is observed, which has a results in a significant improvement in analyses utilising tau leptons at the CMS experiment. The different D_e^{WP} , D_μ^{WP} and $D_{\text{jet}}^{\text{WP}}$ can be used to optimise the hadronic tau identification for analyses attempting to separate tau enriched signals from backgrounds of misidentified objects. Two examples of analyses that utilise this are described in Chapter 4 and 5.




Figure 3.6: Efficiency comparisons of electrons (a), muons (b) and jets (c) passing the DeepTau identification discriminators versus hadronic tau efficiency. (a) uses $Z \rightarrow ee$, (b) uses $Z \rightarrow \mu\mu$ and (c) uses $W + \text{jets}$ MC, with only objects with $p_T < 100 \text{ GeV}$ used. Working points are indicated by full circles and previous algorithms described in Reference [12] are shown in blue triangles and green squares [11].

Chapter 4

Searches for New Physics in $\tau^+\tau^-$ Final States

The $\tau^+\tau^-$ final states are a powerful tool to search for new physics at collider experiments. As the heaviest lepton, τ particles are sensitive to resonant production of new neutral particles where the couplings have mass hierarchy. They are also sensitive to non-resonant effects from new physics mediators. This chapter details the searches for two such areas of new physics: additional Higgs bosons and vector leptoquarks. These searches are split up into three sections:

- i) A model independent search for single narrow spin-0 resonance (ϕ), produced via gluon fusion ($gg\phi$) or in association with a b quark ($bb\phi$). The **SM** Higgs boson is treated as a background and the Yukawa couplings of the spin-0 resonance that contribute to the gluon fusion loop are set to **SM** values.
- ii) A search for the **MSSM** Higgs sector in a number of benchmark scenarios. The benchmark scenarios were proposed in References [80–82] and described fully in Reference [83]. The M_h^{125} and $M_{h,\text{EFT}}^{125}$ scenarios are shown in this chapter. The production of the observed Higgs boson particle at 125 GeV is also used to constrain the available phase space.
- iii) A search for the t-channel exchange of a U_1 vector leptoquark. Two scenarios are taken, motivated by the best fit to the B anomalies as detailed in Section 1.4.1.

These searches are performed with the full run-2 dataset (138 fb^{-1}) collected by the **CMS** experiment. The search for additional Higgs bosons had previously been performed with data collected in 2016 (39 fb^{-1}) and results were consistent with the **SM** background prediction [1].

4.1 Signal Modelling

4.1.1 Additional Higgs Bosons

Extended Higgs sectors, such as that of the [MSSM](#), can be probed by direct searches for the additional bosons and further precise measurements of the [SM](#) Higgs boson. This search for an extended Higgs sector is motivated by a Type II [2HDM](#), such as the [MSSM](#). In these models $\tan\beta$ enhances couplings of additional Higgs bosons to down-type quarks and leptons, whilst up-type quark couplings are suppressed. This divides the dominant production modes of the Higgs boson into two categories: Gluon fusion and production in association with a bottom quark. Examples of the Feynman diagrams for such processes are shown in Figure 4.1, where in this chapter ϕ is used to represent the h , H , and A bosons.




Figure 4.1: Diagram (a) shows the production of neutral Higgs bosons from gluon fusion. The dominant loop contributions to this diagram are from top-only, bottom-only and top-bottom interference. Diagrams (b) and (c) show production in association with b quarks.

With the $\tan\beta$ enhancement, the decays of additional Higgs bosons to τ leptons and b quarks are most dominant. τ leptons are identified with a higher purity than b quarks at the [CMS](#) detector. It is also easier to separate $\tau^+\tau^-$ from the large [QCD](#) multijet background produced from the high energy proton-proton collisions. This was confirmed with the 2016 dataset and although no deviations were observed, the

strongest limits on the **MSSM** phase space was placed by the $\tau^+\tau^-$ final states [1, 84].

For this analysis, signal templates for the production of additional Higgs bosons over a mass range of 60 GeV to 3.5 TeV are generated. Gluon fusion is simulated at next-to-leading-order (**NLO**) precision using the **2HDM** implementation of PowHEG 2.0 [85–88]. The kinematic properties are highly dependent on the contributions to the loop, and these are dependent on the specific signal model. To account for the t quark only, b quark only, and tb-interference loop contributions at the **NLO** plus parton shower prediction, weights are calculated from the p_T spectra to split these contributions up. Once individual templates have been determined for each contribution to the loop, the **2HDM** samples can be scaled to the **MSSM** scenario prediction with the following formula,

$$\frac{d\sigma_{\text{MSSM}}}{dp_T} = \left(\frac{Y_{t,\text{MSSM}}}{Y_{t,\text{2HDM}}} \right)^2 \frac{d\sigma_{\text{2HDM}}^t(Q_t)}{dp_T} + \left(\frac{Y_{b,\text{MSSM}}}{Y_{b,\text{2HDM}}} \right)^2 \frac{d\sigma_{\text{2HDM}}^b(Q_b)}{dp_T} + \left(\frac{Y_{t,\text{MSSM}}}{Y_{t,\text{2HDM}}} \frac{Y_{b,\text{MSSM}}}{Y_{b,\text{2HDM}}} \right) \left\{ \frac{d\sigma_{\text{2HDM}}^{t+b}(Q_{tb})}{dp_T} - \frac{d\sigma_{\text{2HDM}}^t(Q_{tb})}{dp_T} - \frac{d\sigma_{\text{2HDM}}^b(Q_{tb})}{dp_T} \right\}, \quad (4.1)$$

where Q_i are resummation scales that depend on the mass of the additional Higgs boson, and σ_j^i and Y_j^i are the determined cross sections and Yukawa couplings of the relevant theory j . Further contributions from any supersymmetric partners were checked and account for less than a few percent and so are neglected. The p_T reweighting is done separately for the scalar and pseudoscalar additional Higgs bosons, as the p_T distributions can differ. The benchmark scenarios provide the relative Yukawa couplings (to calculate the cross sections) and branching fractions of the **MSSM** Higgs bosons. An example of the distributions for gluon fusion production, in the **MSSM** M_h^{125} scenario with $m_A = 500$ GeV whilst varying $\tan\beta$, is shown in Figure 4.2. The distributions peak at a higher p_T for the top quark loop, therefore at smaller $\tan\beta$, where the top quark contribution is dominant, an additional Higgs boson would be more boosted.

Production in association with bottom quarks is simulated at **NLO** precision using the corresponding PowHEG 2.0 implementation in the four-flavour scheme [85–88]. All additional Higgs boson signal generation is performed using the parton distribution function (PDF) NNPDF3.1 [89, 90]. τ lepton decay, parton showering and hadronisation are all modelled with the PYTHIA event generator where the PU pro-




Figure 4.2: p_T density distributions of the A (top) and H (bottom) boson, with contributions to the gluon fusion loop displayed individually and summed. These are shown for $\tan\beta$ values of 10 (left) and 30 (right) where $m_A = 500 \text{ GeV}$ in the $MSSM M_h^{125}$ scenario.

file is matched to data [91, 92]. All events generated are passed through a **GEANT4**-based [93] simulation of the CMS detector and reconstructed in the same way as data.

The model independent and model dependent additional neutral Higgs boson searches explained in this chapter, utilise the same signal templates generated. The differences lie in scaling of the gluon fusion cross section and loop contributions (set to the Yukawa couplings for the model independent interpretation), and the b associated production cross section, as well as taking into account the branching fractions in the model dependent interpretation. The model dependent search for the **MSSM** also looks to find differences from the observed **SM** Higgs boson and the predicted **MSSM SM-like** Higgs boson. In each **MSSM** benchmark scenario, an uncertainty of ± 3 GeV is given on the prediction for the **SM** Higgs boson mass. This uncertainty is to reflect the contribution from any unknown higher-order corrections. The value of the mass is allowed to vary within this window, however the Yukawa couplings are rescaled the observed mass.

4.1.2 Vector Leptoquarks

The best fit to the B anomalies in the available phase space for vector leptoquarks yielded large b quark and τ lepton couplings to the U_1 particle. The possible production modes of a $\tau^+\tau^-$ final state are shown in Figure 4.3.




Figure 4.3: Feynman diagrams showing the contribution from U_1 vector leptoquarks to the final state with a pair of oppositely charged τ leptons. q represents any SM quark. Diagram (a) shows t-channel, (b) pair and (c) single production of a vector leptoquark.

Pair and single production of a vector leptoquark are dependent on its strong coupling, which is highly model dependent. For large mass, m_U , the probability of producing an on-shell U_1 singlet or pair is heavily suppressed due to the momentum of the initial partons. These production processes are not discussed further in this chapter. A search for single and pair production at the CMS experiment was performed and no statistically significant derivation was observed [94]. Single production places the loosest constraints on the available phase space, out of the processes mentioned. Pair production puts a lower limit on the leptoquark mass, as the process is approximately independent of g_U . However, this limit is heavily dependent on the value chosen for the strong coupling.

The t-channel process contain two vertices with a U_1 vector leptoquark, a quark and a τ lepton, and hence the cross section will scale with g_U^4 . From the best fit to B anomalies, the vertex is dominated by the b quark and hence the initial state will be mostly from $b\bar{b}$, with sub-dominant contributions from $b\bar{s}$, $s\bar{b}$ and $s\bar{s}$. Although there are no additional b quarks in the final state in the leading-order (LO) process, initial state radiation can lead to additional b quarks in the final state. In this search the two scenarios discussed in Section 1.4.1 are considered. The only non negligible

freely floating parameter in each fit, for $\tau^+\tau^-$ final states in the m_U - g_U phase space is the $\beta_L^{s\tau}$ parameter. This is set to the best fit value.

The signal process of the U_1 t-channel exchange is simulated in the five flavour scheme ([FS](#)) at [LO](#) precision using the MADGRAPH5_aMC@NLO v2.6.5 event generator [95]. Events are generated with one or fewer outgoing partons from the matrix element and the MLM prescription [96] is then used for matching, with a scale set to 40 GeV. Negligible dependence of the U_1 decay width is observed, for simulation this is chosen to approximately match the value predicted by the B anomaly fit. Samples with a mass between 1 and 5 TeV are generated.

The interference between the U_1 signal and $Z/\gamma^* \rightarrow \tau\tau$ production was checked and a large destructive affect was observed, with magnitude dependent on g_U . To account for this, separate samples are produced for this interference, generated in the same way as the t-channel exchange. The interference samples are then generated with extra statistics at large di- τ mass values, in order to have a sufficient number of events in the regions of interest for the signal. The cross section of these interference samples scale with g_U^2 . Examples of the generator level di- τ mass distributions are shown in [Figure 4.4](#).

The t-channel signal produces a broad distribution in $m_{\tau\tau}$ due to its non-resonant nature. The interference is mostly a destructive effect (except for at small $m_{\tau\tau}$), with the yield becoming less negative at higher $m_{\tau\tau}$. The interference peaks negatively between 100 and 200 GeV and in this region the combined yield can be negative. Due to the difference in scaling of the two effects, at small g_U the interference is more dominant than the signal and hence the yield of the combined result is reduced.

4.2 Event Selection

The possible decays of two τ leptons and their branching fractions, where the τ decay is grouped into three categories; e , μ and τ_h as defined in [Section 3](#), are shown in [Table 4.1](#). For this search the four largest branching fraction channels are used: $\tau_h\tau_h$, $e\tau_h$, $\mu\tau_h$ and $e\mu$. This accounts for approximately 94% of di- τ events. The two same lepton channels are neglected due to small branching ratio and the dominating $Z \rightarrow ee$ and $Z \rightarrow \mu\mu$ backgrounds.




Figure 4.4: The generator level $m_{\tau\tau}$ density distributions of the t-channel vector leptoquark signal and the interference with Drell-Yan. This is shown in the VLQ BM 1 scenario for a leptoquark of mass 1 TeV for coupling strengths of $g_U = 3$ (a) and $g_U = 5$ (b).

Channel	Branching Fraction
$\tau_h\tau_h$	42.0%
$e\tau_h$	23.1%
$\mu\tau_h$	22.6%
$e\mu$	6.2%
ee	3.2%
$\mu\mu$	3.0%

Table 4.1: Branching fractions of the decays of two τ leptons.

4.2.1 Trigger Requirements

In these four final state pairs, a number of different online trigger requirements are needed. In the $\tau_h\tau_h$ channel, two possible triggers are available: the double- τ_h and single- τ_h triggers. The single- τ_h trigger has a high p_T threshold at 120 (180) GeV for events recorded in 2016 (2017-2018), whilst the double- τ_h has a p_T threshold at 40 GeV. Therefore, the double- τ_h trigger is used unaccompanied when the τ_h has p_T below the single- τ_h threshold, and the union of single- τ_h and double- τ_h triggered events are taken above the threshold.

In the $e\tau_h$ and $\mu\tau_h$ channels, there are three possible triggers available: the single- e/μ , single- τ_h and the $e/\mu-\tau_h$ cross-trigger. The p_T thresholds of light lepton in these triggers are shown in Table 4.2. The cross-trigger is used for events where the light lepton has p_T between the thresholds for the cross-trigger and single- e/μ . The light lepton used in the cross-trigger is required to be in the central barrel of the detector within $|\eta| < 2.1$, and the τ_h triggered on is required to have $p_T > 30$ GeV. Above these light lepton cross-trigger p_T thresholds, the single- e/μ trigger is used. At τ_h p_T above the single- τ_h thresholds, the single- τ_h trigger is used in combination with the single- e/μ trigger.

Year/ Trigger	$e-\tau_h$ cross-trigger	single- e	$\mu-\tau_h$ cross-trigger	single- μ
2016	23	26	20	23
2017	25	28	20	25
2018	25	33	21	25

Table 4.2: Lower trigger light lepton thresholds p_T in GeV for the $e\tau_h$ and $\mu\tau_h$ channels.

In the $e\mu$ channel, there are three possible triggers available: the single- e , single- μ and the $e-\mu$ cross-trigger. However, only the cross-trigger is used in this analysis, due to the larger efficiencies of correctly selecting light leptons. The e and μ are required to have $p_T > 15$ GeV and $|\eta| < 2.4$.

4.2.2 Offline Requirements

All offline selections stated are in addition the object selection discussed in Chapter 3. In this analysis, τ_h candidates are required to pass the DeepTau Medium $D_{\text{jet}}^{\text{WP}}$,

as described in Section 3.8. The D_e^{WP} and D_μ^{WP} choices are dependent on the decay channel. The **VVLoose**, **Tight**, **VVLoose** D_e^{WP} and the **VLoose**, **VLoose** and **Tight** D_μ^{WP} are used in the $\tau_h\tau_h$, $e\tau_h$ and $\mu\tau_h$ channels respectively. The tighter working point for the same light lepton discrimination as tagged in the event is used to remove light leptons misidentified as τ_h from the $Z \rightarrow ll$ process. The light lepton isolation requirement is $I_{\text{rel}} < 0.15$ for both electrons and muons, except for in the $e\mu$ channel where the muon is required to have $I_{\text{rel}} < 0.2$.

The selected τ lepton decay candidates are required to have opposite charge and to be separated by more than $\Delta R > 0.5$ in all channels except $e\mu$ where $\Delta R > 0.3$. In events where the numbers of an object in the event is greater than the required number of objects in the $\tau\tau$ decay channel, the objects are sorted by the maximum $D_{\text{jet}}^{\text{score}}$ if a τ_h candidate or minimum I_{rel} if a light lepton candidate and the leading objects are chosen. In order to maintain orthogonality between channels, events with additional light leptons passing looser selections than the nominal requirements, are rejected from the selection. The looser selections vetoed, help to suppress the $Z \rightarrow ll$ background process further, as well as keeping the channels orthogonal.

In the $e\tau_h$ and $\mu\tau_h$ channels, a cut is placed on the transverse mass between the light lepton \vec{p}_T and the missing \vec{p}_T at 70 GeV, where the transverse momentum is defined as,

$$m_T(\vec{p}_T^i, \vec{p}_T^j) = \sqrt{2p_T^i p_T^j (1 - \cos \Delta\phi)}, \quad (4.2)$$

where $\Delta\phi$ to the azimuthal angle between \vec{p}_T^i and \vec{p}_T^j . The variable is used to remove $W + \text{jets}$ background events, where a jet is misidentified as a τ_h , and the **MET** (neutrino) and light lepton from the W decay are aligned and hence the event has a large $m_T(\vec{p}_T^{e/\mu}, \vec{p}_{\text{MET}})$.

In the $e\mu$ channel, an additional cut is placed on the variable D_ζ , which is defined as,

$$D_\zeta = p_\zeta^{\text{miss}} - 0.85 p_\zeta^{\text{vis}}; \quad p_\zeta^{\text{miss}} = \vec{p}_T^{\text{miss}} \cdot \hat{\zeta}; \quad p_\zeta^{\text{vis}} = (\vec{p}_T^e + \vec{p}_T^\mu) \cdot \hat{\zeta} \quad (4.3)$$

where $\vec{p}_T^{e/\mu}$ corresponds to the transverse momentum vector of the electron or muon and $\hat{\zeta}$ to the bisectional direction between the electron and the muon in the transverse plane. A diagram of the inputs is shown Figure 4.5.




Figure 4.5: Diagram of inputs to the D_ζ variable [2].

The linear combination is optimised for genuine di- τ events to peak around $D_\zeta = 0$ GeV. It is motivated by the expectation that in di- τ decays from a resonance, the visible and missing (from τ neutrinos) momenta are roughly aligned and of similar magnitudes. In $W + \text{jets}$ and $t\bar{t}$ events the directions of the visible and missing products are expected to be more randomly distributed and lead to a non-peaking D_ζ . Therefore, only events with $D_\zeta > -35$ GeV are considered for signal events. No b tag events failing this cut are vetoed, and events with a b tag also failing this cut are used for a $t\bar{t}$ control region and discussed further in Section 4.4.

4.3 Search Optimisation

The optimisation of the signal extraction depends on which of the three scenarios, set out at beginning of this section, is being searched for. The components of the optimisation are named the high-mass, low-mass and **SM** Higgs optimisation procedures. For the model independent search (i) the high- or low-mass optimisation procedures are used depending on whether the mass of the resonance is greater or less than 250 GeV. The search for the **MSSM** Higgs sector (ii) uses the high mass or the **SM** Higgs optimisation procedures depending on whether the reconstructed di- τ mass is greater or less than 250 GeV. The presence of more than one neutrino in the event, means that it is not possible to fully reconstruct the di- τ mass directly. Instead, the likelihood-based **SVFit** algorithm is used [97]. Finally, the search for vector leptoquarks (iii) uses only the high-mass optimisation procedure. The procedures are discussed in detail below.

4.3.1 High Mass Optimisation

The high-mass optimisation procedure follows what was done in Reference [1]. Each event is initially split into categories depending on whether it has 0 or ≥ 1 b tagged jets. Firstly, this helps target the additional Higgs boson production modes gluon fusion and b associated production respectively. In particular, $b\bar{b}\phi$ has final state b quarks that if tagged can significantly aid the sensitivity to this signal. Secondly, the production of the dominant initial state b quarks for a t-channel vector leptoquark signal, can lead to additional b jets in the final state. The reduced backgrounds in b tagged events allows for a more sensitive vector leptoquark search in this category.

The $e\tau_h$ and $\mu\tau_h$ channels are further subdivided into categories depending on the transverse mass between the light lepton and missing transverse momentum vectors as defined in Equation 4.2. The corresponding categories are defined as:

- **Tight- m_T :** $m_T(\vec{p}_T^{e/\mu}, \vec{p}_T^{\text{miss}}) < 40$ GeV;
- **Loose- m_T :** $40 \leq m_T(\vec{p}_T^{e/\mu}, \vec{p}_T^{\text{miss}}) < 70$ GeV.

The majority of the signal events fall within the **Tight- m_T** sub-category. The **Loose- m_T** category is used improve the signal acceptance for resonant masses of $m_\phi > 700$ GeV.

The $e\mu$ channel, is subdivided into three signal categories based of the cuts on the variable D_ζ as defined in Equation 4.3. The three categories are defined as:

- **Low- D_ζ :** $-35 \leq D_\zeta < -10$ GeV;
- **Medium- D_ζ :** $-10 \leq D_\zeta[\text{GeV}] < 30$ GeV;
- **High- D_ζ :** $D_\zeta[\text{GeV}] \geq 30$ GeV.

By design, the majority of signal events are located in the **Medium- D_ζ** sub-category. The **Low-** and **High- D_ζ** categories are used to catch the tail of the signal distributions. A schematic of all high mass optimisation categories are shown in Figure 4.6.

Once all category divisions have been applied, events are drawn in histograms based on a discriminating variable. The discriminating variable used in this analysis is m_T^{tot} and is defined below.

$$m_T^{\text{tot}} = \sqrt{m_T(\vec{p}_T^{\tau_1}, \vec{p}_T^{\text{miss}})^2 + m_T(\vec{p}_T^{\tau_2}, \vec{p}_T^{\text{miss}})^2 + m_T(\vec{p}_T^{\tau_1}, \vec{p}_T^{\tau_2})^2}, \quad (4.4)$$

where τ_1 and τ_2 refer to the visible products of the two τ lepton decays. This variable provides excellent discriminating power between higher mass resonant signals compared to other non-peaking backgrounds, whilst still maintaining some separation between signal masses. It is also excellent at separating the high-mass non-resonant di- τ signatures from backgrounds, where a di- τ mass does not represent the mass of a resonance. This is due to the use of the transverse momenta and MET in the variable definition. For a t-channel signal where the mediator has high mass, no significant mass separation is expected in any variable.

4.3.2 Low Mass Optimisation

The low-mass optimisation procedure loosely follows the high-mass procedure with a few key difference. Categories that are only sensitive to high-mass signals are dropped. This includes the `Low- D_ζ` and `Loose- m_T` categories. Each no b tag subcategory is further divided into four bins of reconstructed di- τ visible p_T with bin edges: 0, 50, 100, 200 and ∞ . This is not done in the b tag subcategories due to the lack of statistics in this region. A schematic of the categories used in the low mass optimisation procedure is shown in Figure 4.7. The final difference with the high-mass optimisation procedure is the discriminator used. In the low-mass optimisation procedure the reconstructed di- τ mass is used. This helps to separate signal events from the Z boson peak in this region.

4.3.3 Standard Model Higgs Boson Optimisation

Finally, the `SM` Higgs optimisation procedure is taken from the [CMS SM](#) $H \rightarrow \tau\tau$ analysis and is detailed in Reference [98]. This was previously used for simplified template cross section measurements. This uses a neural network (`NN`) based categorisation to obtain the most precise estimates from data of the `SM` Higgs produced via gluon fusion, vector boson fusion or vector boson associated production. The `NN` based analysis introduces 26 categories, 8 of which are optimised to pull out the Higgs boson signal. Although the `NN` is trained specifically to target events with an `SM`-like Higgs boson, signal events with differing masses can also enter the `NN` categories.

	No b tag			b tag		
e μ	Low- D_ζ	Medium- D_ζ	High- D_ζ	Low- D_ζ	Medium- D_ζ	High- D_ζ
e τ_h	Loose- m_T	Tight- m_T		Loose- m_T	Tight- m_T	
$\mu\tau_h$	Loose- m_T	Tight- m_T		Loose- m_T	Tight- m_T	
$\tau_h\tau_h$						
$t\bar{t}(e\mu)$						
	Signal region (SR)			$D_\zeta < -35 \text{ GeV}$		
	Control region					

Figure 4.6: Overview of the categories used for the extraction of the signal in the high mass optimisation procedure [2].

	No b tag			b tag		
e μ	Medium- D_ζ $p_T^{\tau\tau} < 50 \text{ GeV}$	High- D_ζ $p_T^{\tau\tau} < 50 \text{ GeV}$		Medium- D_ζ	High- D_ζ	
	$50 < p_T^{\tau\tau} < 100 \text{ GeV}$	$50 < p_T^{\tau\tau} < 100 \text{ GeV}$				
	$100 < p_T^{\tau\tau} < 200 \text{ GeV}$	$100 < p_T^{\tau\tau} < 200 \text{ GeV}$				
	$p_T^{\tau\tau} > 200 \text{ GeV}$	$p_T^{\tau\tau} > 200 \text{ GeV}$				
e τ_h	Tight- m_T $p_T^{\tau\tau} < 50 \text{ GeV}$			Tight- m_T $p_T^{\tau\tau} < 50 \text{ GeV}$		
	$50 < p_T^{\tau\tau} < 100 \text{ GeV}$					
	$100 < p_T^{\tau\tau} < 200 \text{ GeV}$					
	$p_T^{\tau\tau} > 200 \text{ GeV}$					
$\mu\tau_h$	Tight- m_T $p_T^{\tau\tau} < 50 \text{ GeV}$			Tight- m_T $p_T^{\tau\tau} < 50 \text{ GeV}$		
	$50 < p_T^{\tau\tau} < 100 \text{ GeV}$					
	$100 < p_T^{\tau\tau} < 200 \text{ GeV}$					
	$p_T^{\tau\tau} > 200 \text{ GeV}$					
$\tau_h\tau_h$						
$t\bar{t}(e\mu)$						
	Signal region (SR)			$D_\zeta < -35 \text{ GeV}$		
	Control region					

Figure 4.7: Overview of the categories used for the extraction of the signal in the low mass optimisation procedure [2].

4.4 Background Modelling Overview

The relevant backgrounds for this analysis include Drell-Yan, $t\bar{t}$, W+jets, QCD, di-boson, single-top, and electroweak W and Z bosons production. These are split into a five categories:

- i) Events containing only genuine τ leptons.
- ii) Events with a jet misidentified as a τ_h ($\text{jet} \rightarrow \tau_h$) in the $e\tau_h$, $\mu\tau_h$ or $\tau_h\tau_h$ channels.
- iii) Events with jets faking both light leptons ($\text{jet} \rightarrow l$) in the $e\mu$ channel.
- iv) Events from $t\bar{t}$ with a prompt light lepton (e or μ not from a τ decay) and the other object (if there are not two prompt light lepton) is from a genuine τ lepton.
- v) Other events. This is a small contribution and so it is grouped.
 - Non $t\bar{t}$ events satisfying (iv).
 - Events with a light lepton misidentified as a τ_h and no jet misidentified as a τ_h .
 - Events with a jet misidentified as a light lepton and the other object is from genuine τ leptons in the $e\tau_h$, $\mu\tau_h$ or $\tau_h\tau_h$ channels.
 - Events with a muon misidentified as an electron in the $e\mu$ channel.
 - Events with one jet misidentified as a light lepton and the other object from a prompt light lepton in the $e\mu$ channel.

Backgrounds from (i) consists of largely $Z/\gamma^* \rightarrow \tau\tau$ events but there are also smaller contributions from other processes. This background is modelled by a data-simulation hybrid method called the “embedding method” [13] and this is described in detail in Section 4.6. Group (ii) is dominated by QCD, W + jets and $t\bar{t}$ events with a $\text{jet} \rightarrow \tau_h$ misidentification. This is modelled from data by the fake factor method (F_F) [99, 100] and is explained in Section 4.7. Group (iii) is modelled from data to describe the QCD multijet contribution to the background in the $e\mu$ channel. The method to obtain this background is described in Section 4.5. The data driven background estimations for (i), (ii) and (iii) contribute $\approx 98\%$ of all expected background events in the $\tau_h\tau_h$ channel, $\approx 90\%$ in $e\tau_h$ and $\mu\tau_h$ channels and $\approx 50\%$ in the $e\mu$ channel. The final groups, (iv) and (v), are modelled with MC simulations. The $t\bar{t}$ process from (iv) is separated because of its large contribution to the phase

space where a b jet is required.

In 2016 (2017–2018), the $W + \text{jets}$ and $Z \rightarrow ll$ processes are simulated using the MADGRAPH5_aMC@NLO 2.2.2 (2.4.2) event generator at LO [95]. Supplementary samples are generated with up to four outgoing partons in the hard interaction to increase the number of simulated events in regions of high signal purity. For diboson production, MADGRAPH5_aMC@NLO is used at NLO precision [95], and the FxFx [96] (MLM [101]) prescription is used to match the NLO (LO) matrix element calculation with the parton shower model. Samples for $t\bar{t}$ [102] and (t-channel) single top quark production [103] are generated at NLO precision using PowHeg 2.0 [85–88], and for single top quark production in association with a W boson (tW channel) [104], PowHeg version 1.0 at NLO precision is used. When compared with data, $W + \text{jets}$, $Z \rightarrow ll$, $t\bar{t}$, and single top quark events in the tW channel are normalised to their cross sections at next-to-next-to-leading-order (NNLO) precision [105–107], while single top quark (t-channel) and di-boson events are normalized to their cross sections at NLO precision or higher [107–109].

4.5 QCD Estimation in the $e\mu$ Channel

The QCD model in the $e\mu$ channel, that attempts to model events where two jets are misidentified as an electron muon pair, is taken from data with a same sign pair with a transfer factor (F_T). The transfer factor determines differences from the same sign to opposite sign region and is calculated from a sideband region with an anti-isolated muon ($0.2 < I_{\text{rel}} < 0.5$). F_T is initially parametrised by the ΔR between the electron and muon, and the number of jets in the event, however additional dependencies on the electron and muon p_T enter via a correction.

Good agreement is observed in events with no b jets, when applying F_T onto same sign events compared to opposite sign events where both regions have an anti-isolated muon. However, in events with b jets an additional correction is needed. This is determined to be ≈ 0.75 (differs very slightly between data taking years). As this correction is large, it is validated by switching the light lepton anti-isolation, so that the election is required to have $0.15 < I_{\text{rel}} < 0.5$. Also, events where both light leptons are anti-isolated are looked at. The correction for b tagged events is equivalent in all three regions, and a global average of the three is taken for the final correction.

To understand the physical reason for the large difference in no b tag and b tag events in same sign and opposite pairs, studies were performed on simulated samples. It was observed that the electron-muon pair is usually produced from pairs of heavy quarks, $pp \rightarrow b\bar{b}$ ($c\bar{c}$). If the two jets are initiated from the heavy quarks there is a large bias towards opposite sign jets due to the opposite signs of the quark anti-quark pair. However, if one of the heavy quarks is tagged as a b jet, another object has to be the jet initiator (a radiated gluon for example) and there is therefore no charge preference in the pair. As F_T is originally fit inclusively in numbers of b jets and the 0 bin is dominant, the correction over predicts the opposite sign to same sign ratio and so a large correction is needed as observed.

4.6 Embedding Method

The background for genuine di- τ lepton pairs is modelled via the embedding method [13]. This is a hybrid method that utilises both data and MC techniques to produce high statistic samples, where the bulk of the event comes from data. This minimises both the chance of MC fluctuations and the size of the uncertainties. The background is dominated by $Z \rightarrow \tau\tau$ decay however there will be smaller contributions from $t\bar{t}$ and di-boson processes.

The algorithm first selects $\mu\mu$ events from data. The selection is chosen to naturally target the pure $Z \rightarrow \mu\mu$ region but still be loose enough to catch events from other processes, so not to introduce a bias on the Z boson mass. Events are required to pass the double- μ trigger with minimum requirements on the invariant mass of the two muons ($m_{\mu\mu}$) and the p_T of the leading and trailing muons. Also required at the trigger level is a loose association of the track to the PV and a loose isolation in the tracker. Offline objects matched to the trigger muons, are then required to have standard d_z and η selections and originate from a global muon-track, as defined in Section 3. The muon pair are required to have opposite charge and have $m_{\mu\mu} > 20$ GeV. The fraction of processes within this selection is tested with MC background samples and a QCD model from same sign muon pairs with an extrapolation factor. Approximately 97% of selected events are expected to come from $Z/\gamma^* \rightarrow \mu\mu$ events with smaller contributions from $Z/\gamma^* \rightarrow \tau\tau$ ($\tau \rightarrow \mu$), di-boson, $t\bar{t}$ and QCD. The di-boson and $t\bar{t}$ relative contributions are greater at higher $m_{\mu\mu}$ and in events with tagged b jets whilst the QCD contribution is largest at lower $m_{\mu\mu}$. The events selected are biased by detector acceptances. Therefore, corrections on the recon-

struction and identification efficiencies are performed in muon η and p_T using the “tag-and-probe” method, as described in Reference [110].

Next, all energy deposits in the detector from the selected muons are removed. This involves removing the hits on global-muon track in the tracker, hits in the muons systems and clusters in the calorimeters that intercept the muon trajectory. Once completed, the selected muons and their kinematic properties are replaced with a simulated τ lepton. To account for the difference in mass between the muon and τ , the muons are boosted into the center-of-mass frame of the di-muon system and then this four-vector is taken for the τ but boosted back into the laboratory frame. The event simulation is performed from the PV. The τ lepton decay is then simulated with PYTHIA [91, 92] and separate samples are produced for different $\tau\tau$ decay channels. Only the decay of the τ leptons are then processed through the detector simulation and the remainder of the $\mu\mu$ event is added back. A schematic of the process is shown in Figure 4.8.




Figure 4.8: Schematic of the embedding method to model genuine di- τ backgrounds from di-muon events in data [13].

The embedding method is validated on dedicated samples, where the muons from

data are replaced by simulated muons instead of τ leptons. A plot of the agreement from these dedicated samples with data is shown in Figure 4.9.




Figure 4.9: Closure plot showing the di-muon mass on the dedicated embedding validation samples.

4.7 Fake Factor Method

Backgrounds in which a jet is misidentified as a τ_h can be difficult to model using MC due to the poor description of the jet $\rightarrow \tau_h$ fake rate in simulation. In addition, the small probability of a jet being misidentified as a τ_h necessitates the production of high statistics MC samples at a significant computational expense. These shortcomings motivate the use of data-driven estimates for these processes. One such procedure is the F_F method [99, 100].

The F_F method utilises regions in the data to model the jet $\rightarrow \tau_h$ background. Firstly, the determination regions are defined, which are jet $\rightarrow \tau_h$ enriched control regions orthogonal to the signal region. They are then purified by subtracting off any non jet $\rightarrow \tau_h$ events with MC. It is used to calculate F_F by taking the ratio of number of jet $\rightarrow \tau_h$ events that pass the nominal τ_h ID requirement ($N(\text{Nominal})$),

to the number of jet fake events that fail the nominal τ_h identification but pass a looser alternative τ_h identification requirement ($N(\text{Alternative} \&\& \text{!Nominal})$), as shown in Equation 4.5.

$$F_F = \frac{N(\text{Nominal})}{N(\text{Alternative} \&\& \text{!Nominal})}. \quad (4.5)$$

In the remaining text this numerator and denominator are referred to as the pass and fail regions. The derivation of this ratio is done differentially with respect to key parameters that differ in the two regions. Once F_F have been derived it is common to calculate corrections in other sideband regions (a region orthogonal to the signal region) and combine F_F measured from different processes. Finally, the F_F are applied to the application region. This is defined as the signal region but with the criteria that the jet $\rightarrow \tau_h$ events fail the nominal τ_h ID but pass the looser alternative τ_h identification requirement. This now models the background from jet $\rightarrow \tau_h$ events in signal region.

The following Sections 4.7.1–4.7.4 detail the complexities of how this method is applied to this analysis. For these searches the nominal τ_h ID used is the **Medium** $D_{\text{jet}}^{\text{WP}}$ and the alternative τ_h ID used is the **VVLoose** $D_{\text{jet}}^{\text{WP}}$.

4.7.1 Determination Regions

The F_F are measured separately in each year of data taking period (2016, 2017, 2018), in each channel containing a τ_h ($e\tau_h$, $\mu\tau_h$, $\tau_h\tau_h$) and in enriched regions of dominant processes that contribute jet $\rightarrow \tau_h$ events. In the $e\tau_h$ and $\mu\tau_h$ channels F_F are measured for three processes: **QCD**, W + jets and $t\bar{t}$. In the $\tau_h\tau_h$ channel F_F are measured only for the dominant **QCD** process. The **QCD** process is assumed to produce two jets misidentified as τ_h candidates, and so the F_F is chosen to be calculated from leading p_T τ_h candidate only. Section 4.7.4 discusses how a single jet misidentified as a τ_h candidate events in the $\tau_h\tau_h$ channel are modelled.

Each separate measurement region is split into three sideband regions based on two cuts that surround the signal region. These regions are named the **Determination Region (A)**, **Alternative Determination Region (B)** and **Correction Region (D)** and are schematically shown in Figure 4.10.

Region A is used to measure and fit the F_F . Region B is an alternative region used




Figure 4.10: Schematic of the regions used for fake factor derivation.

to measure and fit the F_F to account for the difference in fake factors between A and C. These alternative F_F are applied to the fail τ_h identification region in D and corrections are calculated comparing it to the pass region in D. The total F_F per measurement region is calculated as the product of the F_F derived in region A and the correction calculated from region B to D.

The selection for x_C , x_D , y_C and y_A , as defined in Figure 4.10, in each separate measurement region are shown below. These are chosen to balance the number of events and the purity of each background in the region.

i) $\tau_h\tau_h$ QCD

y_C : The τ_h candidates are required to have the opposite sign.

y_A : The τ_h candidates are required to have the same sign.

x_C : The subleading τ_h passes the Medium $D_{\text{jet}}^{\text{WP}}$.

x_D : The subleading τ_h fails the VVLoose $D_{\text{jet}}^{\text{WP}}$ but passes the VVVLoose $D_{\text{jet}}^{\text{WP}}$.

ii) $e\tau_h$ and $\mu\tau_h$ QCD

y_C : The e/μ and τ_h candidates are required to have the opposite sign.

y_A : The e/μ and τ_h candidates are required to have the same sign and the e/μ to have $I_{\text{rel}} > 0.05$.

x_C : The e/μ candidate is required to have $I_{\text{rel}} < 0.15$.

x_D : The e/μ candidate is required to have $0.25 < I_{\text{rel}} < 0.5$.

iii) $e\tau_h$ and $\mu\tau_h$ W + Jets

y_C : The m_T between the e/μ and the MET < 70 GeV.

y_A : The m_T between the e/μ and the $\text{MET} > 70$ GeV and no b jets in the event.

x_C : Data.

x_D : W + Jets MC.

iv) $e\tau_h$ and $\mu\tau_h$ $t\bar{t}$

y_C : Data.

y_A : MC ($t\bar{t}$ in B and W + Jets D).

x_C : $m_T < 70$ GeV.

x_D : $m_T > 70$ GeV and no b jets.

In the $\mu\tau_h$ and $e\tau_h$ channels, W + jets jet $\rightarrow \tau_h$ events are in general the most significant and QCD contributes a smaller fraction. $t\bar{t}$ inclusively is small but becomes more significant when searching for events with a b jet. The additional $I_{\text{rel}} > 0.05$ requirement in these channels for QCD is to reduce processes producing genuine leptons and the $N_{\text{b-jets}} = 0$ requirement for W + Jets is to reduce $t\bar{t}$ contamination. It is not possible to define a Determination Region that is sufficiently pure in $t\bar{t}$ events to make a reasonable measurement of $F_F^{t\bar{t}}$ from data. Therefore $F_F^{t\bar{t}}$ are derived from MC. A comparison of the $F_F^{\text{W+jets}}$ measured in data and MC shows only $\sim 10\text{--}20\%$ differences in the fake rates in data and MC. This observation coupled with the fact that the $t\bar{t}$ contribution is small compared to the other processes means that any bias introduced by using $F_F^{t\bar{t}}$ measured in MC is small compared to the uncertainties on the F_F , discussed in Section 4.9. In many cases, selections on the regions B and D are applied which are tighter than required, in order to purify the regions as much as possible, while maintaining enough statistics to perform the fit.

4.7.2 Parametrisation

The raw F_F^i take into account dependencies on N_{jets} via an analysis category tailed variable $N_{\text{pre b jets}}$, the p_T of the τ_h candidate ($p_T^{\tau_h}$) and the p_T of the jet that seeds the HPS algorithm to the τ_h (p_T^{jet}). $N_{\text{pre b jets}}$ is the number of jets in the event with $|\eta| < 2.4$ and $p_T > 20$ GeV, which is the η and p_T selection for a b tagged jet. It is defined to map the dependence of F_F^i on the number of jets and describe the categorising variable $N_{\text{b jets}}$ well. Although not local to the τ_h , it helps control other dependencies on the constituents of the event. It is the number of jets in the event

with $|\eta| < 2.4$ and $p_T > 20$ GeV. These are the same η and p_T thresholds required for a b jet. The data is split into two bins of $N_{\text{pre b jets}}$, equal to 0 and greater than 0. It is then further split by the ratio of p_T^{jet} to $p_T^{\tau_h}$. An example of the dependence of these two transverse momentums on the fake factor is shown in Figure 4.11.




Figure 4.11: A 2D heat map of the F_F determined from $t\bar{t}$ MC for the full run-2 dataset in the combined $e\tau_h$ and $\mu\tau_h$ channels. This is shown with respect to the τ_h p_T and the p_T of the jet matched to the τ_h . The ratio of jet to τ_h p_T categorisation used is shown split by the dashed lines.

It is motivated by the observation that the F_F are largest when the p_T^{jet} and $p_T^{\tau_h}$ are closest. The physical motivation for this is when they are close, the τ_h candidate is likely to be isolated from any other hadronic activity and so more likely to be identified as a τ . However, when p_T^{jet} is larger than $p_T^{\tau_h}$, the candidate is likely surrounded by other hadronic activity and so more likely to be a jet fake. When p_T^{jet} is less than $p_T^{\tau_h}$, charge pions are likely not close enough to the PV to be clustered into the jet and so the event is less likely to be classified as a hadronic tau. This leads to the F_F dependence as seen in Figure 4.11.

For all divisions of the phase space, dependence on the $p_T^{\tau_h}$ is fit using the superposition of a Landau function and a constant in the low- p_T region. The fake factors are seen to rise sharply at high- p_T . This increase happens in either the bin $140 <$

$p_T^{\tau_h} < 200$ GeV or $p_T^{\tau_h} > 200$ GeV. To map this effect, binned values are taken based on the algorithm shown in Figure 4.12 and the fit is used below the minimum bin.




Figure 4.12: Flow chart of the algorithm used to determine where binned values are taken instead of the fit. The blue box represents the input, the green diamonds represent the decisions and the red boxes represent the outputs.

The fits are flattened at $p_T^{\tau_h}$ values where there is no significant downwards shift or at the final bin to avoid extrapolating past the fitted range. F_F fits with respect to $p_T^{\tau_h}$ are shown in Figures 4.13-4.14. The F_F are highest in the lowest $p_T^{\text{jet}}/p_T^{\tau_h}$ bin and lowest in the highest $p_T^{\text{jet}}/p_T^{\tau_h}$ bin, as expected. Otherwise the F_F fall with p_T in each category until the thresholds used for the high p_T binning algorithm.




Figure 4.13: F_F fits in $\tau_h\tau_h$ channel for the QCD $N_{\text{pre b jets}} = 0$ category with 2018 data. The three jet p_T to $\tau_h p_T$ categories are shown.




Figure 4.14: F_F fits in $\mu\tau_h$ channel for the QCD and W + Jets $N_{\text{pre b jets}} = 0$ category with 2018 data. The three jet p_T to τ_h p_T categories are shown for each process.

4.7.3 Corrections

In the $\tau_h\tau_h$ channel, the measured F_F^{QCD} are then corrected to account for non-closures in other variable in the **Determination Region**. The only significance non-closures are observed for E_T^{miss} related variables and are largest for events with $N_{\text{pre b jets}} = 0$. Closure corrections are performed for the variable ΔR between the τ_h candidates in bins of $N_{\text{b jets}}$. In the $\mu\tau_h$ and $e\tau_h$ channels, the measured F_F^{QCD} and $F_F^{\text{W+jets}}$ are corrected for non-closures observed in the E_T^{miss} variables and $p_T^{e/\mu}$ distributions. A study was performed to determine the nature of these non-closures and it was found that the cause was due to fake **MET** arising from mismeasurement of the energies of particles in a jet. A diagram of this effect is shown in Figure 4.15.




Figure 4.15: Diagram showing how fake MET arises from mismodelling jet constituents in τ_h identification.

To correct for this effect, the **QCD** fake factors are corrected as a function of C_{QCD} , where C_{QCD} is defined as,

$$C_{\text{QCD}} = \frac{E_T^{\text{miss}} \cos \Delta\phi(\vec{p}_T^{\text{miss}}, \vec{p}_T^{\tau_h})}{p_T^{\tau_h}}. \quad (4.6)$$

where $\Delta\phi(\vec{p}_T^{\text{miss}}, \vec{p}_T^{\tau_h})$ is the separation in the azimuthal angle between the the missing \vec{p}_T^{miss} and $\vec{p}_T^{\tau_h}$. The numerator quantifies the missing transverse momentum in the direction of the τ_h candidate. Once divided by the τ_h p_T , C_{QCD} is a measure of the fraction of missing to visible τ_h transverse momentum aligned with the τ_h . For $\text{W} + \text{jets}$ and $t\bar{t}$ the situation is slightly different due to the presence of genuine missing energy from neutrinos. In this case, the correction variable is modified to approximately subtract the genuine **MET** from the total. This approximation assumes the neutrino is back-to-back and balanced with the light lepton (which is

exactly true for W bosons produce at rest in the transverse direction). The equation then becomes,

$$C_W = \frac{(E_T^{\text{miss}} + p_T^{e/\mu}) \cos \Delta\phi(\vec{p}_T^{\text{miss}} + \vec{p}_T^{e/\mu}, \vec{p}_T^{\tau_h})}{p_T^{\tau_h}}. \quad (4.7)$$

When either correction variable is non 0, a larger quantity of fake MET is expected in the event. In these regions a large correction is needed due to the mis-measured jet energy spectrum shifting the τ_h candidate isolation and so shifting the τ identification scores. Examples of these closure corrections are shown in Figure 4.16

After the Determination Region is modelled well for all variables of interest, extrapolation corrections from the fake factors derived in B applied to region D are calculated. In the $\tau_h\tau_h$ the correction is parametrised by the p_T of the leading τ_h candidate, in the $e\tau_h$ and $\mu\tau_h$ channels it is parametrised by the p_T of the light lepton. Where statistics allow, these corrections are calculated in the high-mass optimisation procedure categories. Examples of the extrapolation corrections are shown in Figure 4.17.




Figure 4.16: Determination Region closure correction fits with 2018 data. (a) is the correction parametrised by ΔR in events with N_b jets = 0 in the $\tau_h\tau_h$ channel. (b) and (c) show the correction for the $\mu\tau_h$ channel parametrised by the specific correction variables defined in Equation 4.6 and 4.7 for QCD and W + jets processes respectively.




Figure 4.17: Determination Region to Application Region closure correction fits with 2018 data. (a) is the correction moving from same sign to opposite sign τ leptons the parametrised by leading τ_h p_T in events with $N_{b\text{-jets}} = 0$ in the $\tau_h\tau_h$ channel. (b) and (c) show the correction for the $\mu\tau_h$ channel moving from same sign to opposite sign τ leptons and high m_T to low m_T both parametrised by the muon p_T for QCD and $W + \text{jets}$ processes respectively.

4.7.4 Applying Fake Factors

In the $e\tau_h$ and $\mu\tau_h$ channels the F_F^i measured for the different processes are combined into an overall factor, F_F , using

$$F_F = \sum_i f_i \cdot F_F^i, \quad (4.8)$$

where the factor f_i is defined as

$$f_i = \frac{N_{\text{AR}}^i}{\sum_j N_{\text{AR}}^j}, \quad (4.9)$$

which is the fraction of events with a jet $\rightarrow \tau_h$ originating from process i over the total number of jet $\rightarrow \tau_h$ events for all processes in the Application Region. These fraction of events are estimated with MC, with a QCD model extrapolated from same sign τ pairs, and examples are shown in Figure 4.18. It is observed that $W + \text{jets}$ is the dominating process in this region, however, there are affects from QCD at low m_T and from $t\bar{t}$ in the b tagged categories. These fractions are then multiplied by the relevant corrected fake factor and applied to the fail region in C, where this region is purified by subtracting any non jet $\rightarrow \tau_h$ events with MC.

The $\tau_h\tau_h$ channel has two τ_h candidates that a jet can be misidentified as. For this analysis, the F_F are only applied the leading τ_h candidate failing the τ_h identification




Figure 4.18: The expected Application Region fractions of the processes in the $\mu\tau_h$ channel. (a) and (b) show the no b tag Tight- m_T and Loose- m_T categories and (c) and (d) show the b tag Tight- m_T and Loose- m_T categories respectively.

in C. This models all events where the leading τ_h candidate is a jet $\rightarrow\tau_h$. However, this leaves a small fraction of events, where the leading candidate is a genuine τ and the sub-leading candidate is a jet $\rightarrow\tau_h$. This contribution (mostly from W + jets) is added back with MC.

4.8 MC Corrections

The corrections stated in this section apply both to simulated and embedding samples, as the τ decay is simulated, however they are derived separately. For electrons and muons, corrections are applied to triggers, tracking efficiencies, and the identification and isolation requirements. Using the tag-and-probe method, they are obtained in bins of p_T and η of the corresponding lepton, with $Z \rightarrow ee$ and $Z \rightarrow \mu\mu$ events. These corrections are generally no more than a few percent. The electron energy scale is adjusted to the scale measured in data using the Z boson mass peak in $Z \rightarrow ee$ events.

Similarly, corrections are derived for the efficiency of triggering and identification efficiency of τ_h candidates. In the $e\tau_h$ and $\mu\tau_h$ channels, trigger efficiency corrections are obtained from the ratio of fits to data versus simulated samples, for the trigger efficiency as a function of p_T . For the $\tau_h\tau_h$ channel, this is instead done using the binned values of the τ_h decay modes. The identification efficiency corrections are derived as a function of the p_T of the τ_h candidate. Corrections to the energy scale of the τ_h candidates and of electrons misidentified as τ_h candidates are obtained from likelihood scans of discriminating observables, such as the reconstructed τ_h candidate mass. For muons misidentified as τ_h candidates, the energy scale correction is negligible.

The magnitude and resolution of the missing transverse energy needs correcting in embedding events, to take into account the incomplete removal of energy deposits from muons replaced by simulated τ decays during the embedding procedure. These corrections are derived by comparing p_T^{miss} in embedded events to fully simulated events.

In fully simulated events, a specific trigger inefficiency caused by a shift in the timing of the inputs of the ECAL L1 trigger in the region at $|\eta| > 2.5$ during the 2016 and 2017 data taking [111], is needed to be corrected for. This effect is named

“prefiring”. This resulted in a loss of efficiency for events containing an electron or jet with p_T larger than approximately 50 or 100 GeV, in 2016 and 2017 respectively. Corresponding corrections are derived from data and applied to the simulation.

Corrections to the energy of jets are calculated in bins of the jet p_T and η . These range from subpercent levels in the central part of the detector to a few percent in the forward region. The energy resolution of the simulated jets is also tuned to match that of the data. A correction is applied to the missing transverse momentum, based on differences in the estimated hadronic recoil between data and simulation. An MC to data correction for a b jet passing the selection criteria is also determined. These corrections can alter the number of b jets in a simulated event.

Any differences from simulated events to data, where an electron or muon is reconstructed as a τ_h , are corrected from the pure $Z \rightarrow ee$ and $Z \rightarrow \mu\mu$ regions. Similarly, a correction is applied to account for residual differences in the $\mu \rightarrow e$ misidentification rate between data and simulation.

Further MC to data corrections are applied to the di-lepton mass and p_T spectra in simulated $Z \rightarrow ll$ events. These are derived from $Z \rightarrow \mu\mu$ events. Additionally, all simulated $t\bar{t}$ events are weighted to match the top quark p_T distribution observed in data [112].

4.9 Uncertainty Model

The statistical uncertainties are taken into account by the Barlow-Beeston method, described in References [113, 114]. The systematic model is split into uncertainties based on the online and offline reconstruction of objects and the background and signal modelling. An uncertainty is correlated across channels when it represents a shift on the reconstruction of an object and decorrelated otherwise. It is decorrelated across era of data taking when the shift is derived independently by era. The embedded samples use the same uncertainty scheme as MC but 50% are correlated and 50% uncorrelated with MC uncertainties, because of the shared real data in the measurement.

Hadronic Taus

Uncertainties on the τ_h triggers are obtained from the fitted scale factors used to derive the corrections for the τ_h trigger efficiencies. The legs of the double- τ_h and $e/\mu\text{-}\tau_h$ cross triggers in different decay mode bins are treated as uncorrelated. For the single- τ_h trigger leg, due to limited statistics it is not possible to determine scale factors and uncertainties split by decay mode and therefore a single uncertainty common to all decay modes is applied. The double- τ_h trigger uncertainties are further split in the p_T regions < 100 GeV and > 100 GeV to allow the fit more freedom to adjust the high p_T regions relative to the low p_T regions. Uncertainties are also applied on the energy scale of the τ_h candidates. These uncertainties range between 0.2 and 1.1%. Finally, an uncertainty on the ID efficiency is placed as a function of p_T in the $e\tau_h$ and $\mu\tau_h$ channels, and of the τ_h decay mode in the $\tau_h\tau_h$ channels. These vary between 3-9% and is uncorrelated in each variable bin it is derived in. To account for the different anti-lepton discriminator working points, an uncertainty of 3% per τ_h is applied and treated as uncorrelated between the channels where different $D_{\text{WP}}^{e/\mu}$ are used.

Light leptons

The uncertainty on the trigger efficiencies amounts to 2% per lepton in the $e\tau_h$, $\mu\tau_h$ and $e\mu$ channels. They are basically normalisation uncertainties but implemented as shape uncertainties as they only touch the events triggered by the corresponding cross trigger or single lepton triggers. Uncertainties are also placed on the electron energy scale based off the calibration of ECAL crystals. This information is not reliable for embedding samples and so uncertainties of 0.5-1.25% are placed here. The energy scale variations are negligible and so not included. Another 2% uncertainty is placed on the ID of any electron or muon in the event.

Jets

Jet energy scale and resolution uncertainties arise from a number of sources. These include limited statistical measurements used for calibration, energy measurement changes due to detector ageing, and bias corrections to address differences between simulation and data. Uncertainty ranges are from sub-percent to $\mathcal{O}(10\%)$. Uncertainties are also placed on the tagging of b jets, which vary from 0–3%.

Leptons misidentified as hadronic taus

Uncertainty shifts are applied for the energy scale of leptons faking τ_h parametrised by the p_T of the $e/\mu \rightarrow \tau_h$ fake. The magnitude is 1.0% for muons in all eras and for electrons the uncertainties vary between 0.5 and 6.6 %.

Jets misidentified as hadronic taus

The backgrounds with jets misidentified as τ_h are estimated from data with the fake factor method. There are different sources of uncertainty related to this method. The first uncertainties come from subtracting off other background processes with MC to form the determination region. The subtraction is shifted up and down by 10% to determine new weights. Next, statistical uncertainties on all of the fake factor method fits are accounted for, where the binned values are uncorrelated with the rest of the fit. An uncertainty is also placed on the choice of fit function, the shifts are estimated by comparing the fits to a first order polynomial fit set to constant above 100 GeV. The final systematic variation are then on the extrapolation corrections by applying the corrections twice and not at all. The size of each systematic uncertainty varies from 0–10%, whilst the statistical element from the fits can be larger in the tails of the distributions.

Jets misidentified as light leptons

Background with jets misidentified as electrons or muons from QCD are only considered in the $e\mu$ channel and modelled from data. Uncertainties are placed based on the statistical uncertainties in the determination region which are 2–4% and the extrapolation to the signal region that are $\mathcal{O}(10\%)$.

Muons misidentified as electrons

Background with muons misidentified as electrons are only considered in the $e\mu$ channel. These events are modelled from MC and any generator matched muon identified as an electron are given a 15%–45% uncertainty, that is derived from the initial corrections.

MET

The MET uncertainties is different dependent on process. For all processes that are not $t\bar{t}$ and diboson, the hadronic recoil response and its resolution are varied within

the uncertainties determined during the computation of the recoil corrections. For $t\bar{t}$ and diboson an uncertainty is derived on the energy carried by an unclustered particle [115]. These uncertainties vary between 0–10%.

Background process specific uncertainties

Uncertainties on the $t\bar{t} p_T$ and Z boson $m_{ll}-p_T$ reweighting is placed by applying the correction twice and not at all. An additional uncertainty is placed to cover the $t\bar{t}$ contamination in embedding, where the removed $t\bar{t}$ genuine τ pair is shifted up and down by 10%. Some non-closures are observed in embedded $Z \rightarrow \mu\mu$ control samples. Therefore, these non-closures are taken as an additional shape uncertainty as a function of the Z p_T and $m_{\tau\tau}$. Uncertainties on the normalisation background processes with sizes 4% for $Z \rightarrow ll$ and W + jets production [105], 6% for $t\bar{t}$ production [106, 107], and 5% for diboson and single t quark production [107–109].

Signal process specific uncertainties

For the gg ϕ and bb ϕ processes, the variation of `hdamp` parameter of the PowHEG MC generator as well as the μ_R/μ_F scale variations are used to determine the uncertainties on the p_T spectrum of each contribution at NLO QCD to the Higgs boson production via gluon fusion (top, bottom, top-bottom interference). These are also determined from additional samples produced up to the generator level, and applied as event weights dependent on generator level p_T after the parton shower simulation. For the vector leptoquark signal samples, the PDF and μ_R/μ_F scale variations are applied on an event-by-event basis. These uncertainties are included as shape uncertainties as they may affect the shapes of the m_T^{tot} distribution as well as the predicted signal yields.

Luminosity

(1.2%, 2.3%, 2.5%) normalisation luminosity uncertainty is applied to the (2016, 2017, 2018) templates which originate from MC simulation.

Prefiring

Upper and lower bounds are taken from the efficiency maps and propagated on all MC samples as shape uncertainty for 2016 and 2017. The size of the uncertainty as the weight itself depends on the event topology. In general the uncertainty is at the order of 1%.

4.10 Signal Extraction

A simultaneous binned maximum likelihood fit over all analysis categories is used to extract the results. The likelihood takes the form,

$$\mathcal{L}(\text{data} \mid \mu, \theta) = \prod_i^{N_i} \text{Poisson}\left(n_i \mid \sum_j^{N_j} g_j(\mu_{ij}) \cdot s_{ij}(\theta) + \sum_k^{N_k} b_{ik}(\theta)\right) \cdot p(\hat{\theta} \mid \theta), \quad (4.10)$$

where i loops through all histogram bins and analysis categories. The indices j and k loop over all signal and background processes for the hypothesis being fit. n_i , s_i and b_i are the data observed, signal and background expectation respectively in each bin. θ represents the set of nuisance parameters (corresponding to the systematic uncertainties as detailed in Section 4.9) that parametrise the signal and background modelling. μ are rate parameters and $g(\mu)$ are scaling functions that scale to a signal to a specific hypothesis. The form of the Poisson probabilities are,

$$\text{Poisson}(n \mid x) = \frac{x^n e^{-x}}{n!}. \quad (4.11)$$

Finally, $p(\hat{\theta} \mid \theta)$ represents the probability distribution function (PDF) of each nuisance parameter (θ) with respect to the initial value of the parameter ($\hat{\theta}$).

The PDFs come in two forms, the first is for uncertainties that only affect the normalisation of the process and are modelled by log-normal pdf! (pdf!)s. The second is for uncertainties that affect the shape of the distribution, these are assigned Gaussian PDFs. The $\pm 1\sigma$ shifts for each shape variations are derived and vertical morphing [114] is used to interpolate and extrapolate within and outside the shifts. Both PDFs are dependent on the mean (μ) and standard deviations (σ) and the functional forms are shown in Table 4.3.

Gaussian	Log-normal
$f(x) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{(x-\mu)^2}{2\sigma^2}\right)$	$f(x) = \frac{1}{x\sqrt{2\pi\sigma^2}} \exp\left(-\frac{(\ln x - \mu)^2}{2\sigma^2}\right)$

Table 4.3: pdfs used for nuisance parameters.

The following subsections discuss the results of many such fits. The key fits to understand the results are the background-only fit and the signal-plus-background

fits. The background-only fit is performed with N_j (number of signal processes) set to 0. For all signal-plus-background fits, the fit is done with respect to a single mass hypothesis, however within this mass hypothesis can be a number of signal processes. The model independent resonance search has separate $gg\phi$ and $bb\phi$ signal modes and so two rate parameters $\mu_{gg\phi}$ and $\mu_{bb\phi}$ are needed. When the samples are initially scaled to the cross section times branching ratio ($\sigma \times B(\phi \rightarrow \tau\tau)$) of 1 pb and $g(\mu) = \mu$ for both processes, $\mu_{gg\phi}$ and $\mu_{bb\phi}$ represent the $\sigma \times B(\phi \rightarrow \tau\tau)$ with units of pb. To avoid negative signal strengths, μ will only be taken to be positive. Also used in the following subsections, is a signal-plus-background channel/category compatibility fit. In this fit the signal processes and rate parameters are further split in each channel or category utilising index i in Equation 4.10. This is used to determine the compatibility of the results in different decay channels and analysis categories. In this case, μ is allowed to take negative values to help fully understand the fits to data in each channel or category.

The vector leptoquark search has two signal modes; the t-channel interaction and the interference with Drell-Yan. However, as this is a model-dependent interpretation of these results both these rate parameters scale together. The scaling functions differ between the two processes with $g_{t\text{-channel}}(\mu) = \mu^4$ and $g_{\text{interference}}(\mu) = \mu^2$ to mimic how the cross sections of each process scales. When the initial samples are scaled to cross section at $g_U = 1$, μ corresponds to the coupling g_U .

For the **MSSM** interpretation of the results, there are three Higgs bosons to consider in the signal model (h, H and A) produced via both gluon fusion and in association with b quarks. The $gg\phi$ samples are also split into the separate loop contributors, so the kinematic properties can be properly scaled to **MSSM** prediction, as described in Section 4.1.1. The **SM**-like Higgs boson is considered in the **MSSM** signal model to monitor differences in the observed Higgs boson prediction between the **MSSM** and the **SM**. In each benchmark scenario chosen, the signal prediction depends only on m_A and $\tan\beta$ and the scaling to cross section is shown in Equation 4.1. As the potential scaling functions for **MSSM** interpretations are not necessarily smooth one-to-one mappings, the likelihood is tested for individual points on the m_A - $\tan\beta$ parameter space. At each point, the **MSSM** Higgs bosons are scaled to the theory predicted cross section times branching ratio. To test the **MSSM** hypothesis over the **SM** hypothesis, the single rate parameter μ is used and only allowed to take values of 1 (**MSSM**) and 0 (**SM**) with $g(\mu) = \mu$. As the SM Higgs boson is added to

the background modelling and the [MSSM](#) prediction of the observed Higgs boson is added to the signal model when $\mu = 1$, the [SM](#) Higgs boson prediction must then be subtracted from the signal model.

The confidence intervals in the best fit results are given by the $-2\Delta \ln \mathcal{L}$, where $\Delta \ln \mathcal{L}$ is the difference between $\ln \mathcal{L}$ of the best fit model and the test value of μ . The 68% and 95% confidence level ([CL](#)) regions with two degrees of freedom (as in the model independent resonant search) are determined by $-2\Delta \ln \mathcal{L} = 2.28$ and 5.99 respectively.

Upper limits are placed using the modified frequentest approach [116, 117] with a profile likelihood ratio used for the test statistic, as defined below.

$$q_\mu = -2 \ln \left(\frac{\mathcal{L}(\text{data}|\mu, \hat{\theta}_\mu)}{\mathcal{L}(\text{data}|\hat{\mu}, \hat{\theta}_{\hat{\mu}})} \right), 0 \leq \hat{\mu} \leq \mu, \quad (4.12)$$

where $\hat{\mu}$ and $\hat{\theta}_{\hat{\mu}}$ are the best fit values of μ and θ . $\hat{\theta}_\mu$ are the values of θ that is maximised by the likelihood for a tested value of μ . The bounds on $\hat{\mu}$ are to ensure a positive signal strength with a one-sided confidence interval. The probability of $q_\mu \geq q_\mu^{\text{obs}}$ is,

$$\text{CL}(\mu) = \int_{q_\mu^{\text{obs}}}^{\infty} f(q_\mu | \mu, \theta_\mu^{\text{obs}}), \quad (4.13)$$

where $f(q_\mu | \mu, \theta_\mu^{\text{obs}})$ is the pdf of q_μ . CL_b and CL_{s+b} are then defined by the relevant background-only and signal-plus-background fits. CL_s is defined as the ratio of CL_{s+b} and CL_b and then upper limits are placed at the confidence level of $1 - \text{CL}_s$. The $f(q_\mu | \mu, \theta_\mu^{\text{obs}})$ are determined using the asymptotic approximation [118] and results are cross-checked and deemed consistent with toy MC datasets.

If a deviation from the background expectation is observed, the size of the deviation is quantified by a significance. To test rejection of the background-only hypothesis in favour of the signal-plus-background hypothesis, μ is replaced with 0 in the test statistic. The p -value, p_0 is then,

$$p_0 = \int_{q_0^{\text{obs}}}^{\infty} f(q_0 | 0, \theta_\mu^{\text{obs}}). \quad (4.14)$$

p_0 is uniformly distributed between 0 and 1 for the background-only hypothesis and so the probability and significance of rejecting the background-only hypothesis can be found.

4.11 Postfit Plots

Figures 4.19 and 4.20 show the unblinded distributions in the most sensitive analysis categories. For simplicity, the $e\tau_h$ and $\mu\tau_h$ channels have been combined. Figure 4.19 shows the distributions of the $m_{\tau\tau}$ discriminator in the no b tag low-mass optimisation categories. A signal-plus-background fit for a model-independent gluon fusion resonant mass hypothesis of 100 GeV is shown and the changes in the background modelling when using a background-only fit is displayed in the ratio. Figure 4.20 shows the distributions of the m_T^{tot} discriminator in the high mass optimisation categories. A background-only fit is shown for the stacked background and example signal hypotheses for the model independent 1.2 TeV $gg\phi$ and $bb\phi$ resonances and VLQ (BM 1) 1 TeV mass points are displayed.

In the low-mass optimisation categories, a small excess of events is observed on the Z boson peak in the no b tag categories and reasonable agreement is observed in the b tag categories. The excess of events are distributed in $m_{\tau\tau}$ between 80 and 120 GeV. A signal-plus-background hypothesis is best fit with a 100 GeV $gg\phi$ signal with a cross section times branching ratio of 5.8 pb. In this same fit the $bb\phi$ process is constrained by the b tag categories to give a signal yield of 0. A background-only fit is also performed on the data, it is observed that this can only partly explain the differences observed between background and data. Even after a background-only fit there is still a small excess of data events over the Z boson peak.

In the high-mass optimisation categories, another small excess is observed in high m_T^{tot} bins, particularly in the most sensitive no b tag categories. This excess is best fit by a model independent gluon fusion resonant mass at 1.2 TeV with a cross section times branching ratio of 3.1 fb. There are no considerable differences observed in background modelling between signal-plus-background and background-only fits. This is as the uncertainties in these bins are more statistically dominated and the majority of the systematic uncertainties are constrained in the bulk of the distribution. Good agreement is observed in the rest of the distribution. There is a very small deviation in the b tag categories, but as this can also be explained by a $gg\phi$ signal, the $bb\phi$ signal is heavily constrained and so largely does not contribute to the signal-plus-background fit of the excess. Similar to the $bb\phi$ signal, the VLQ BM 1 signal is constrained by the results in the b tag categories, leading to a small non-zero best fit signal strength, but cannot explain the excess in the no b tag categories.

4.12 Model Independent Results

4.12.1 Limits

95% CL limits are set on the assumption of absence of a signal for the search for a $gg\phi$ or $bb\phi$ resonance and shown in Figure 4.21. In each case, the other process is allowed to float freely in the fit. The excesses observed in the postfit distributions act to weaken the observed limit compared to the expected limit at 100 GeV and 1.2 TeV, as more data was observed than expected. For $gg\phi$ production the expected limits flatten under 100 GeV, due to difficulty of separating signal from the Z boson at this mass. Both sets of limits vary from $\mathcal{O}(10 \text{ pb})$ at 60 GeV to 0.3 fb at 3.5 TeV.

95% CL expected limits are drawn on the fit to each di- τ decay channel individually and are shown in Figure 4.22. This gives a measure of the sensitivity of each channel. In the high-mass optimisation categories, the combined limit is heavily dominated by the $\tau_h\tau_h$ channel. This is mostly driven by branching fraction, as all channels in this mass range have similar signal separation ability. In the high-mass optimisation categories, the combined limit is more a contribution of all channels. In the $\tau_h\tau_h$ channel in this region, the QCD multijet background is the largest fraction of any non $Z \rightarrow \tau\tau$ backgrounds in all channels and so the limit for this channel is weakened and the other channels contribute to the combined limit more. The high double- τ_h trigger p_T thresholds (chosen because of the QCD multijet background) also lowers the signal acceptance in the $\tau_h\tau_h$ channel.

A comparison of the limits are also made with the ATLAS experiment and in particular the results presented in Reference [14]. This ATLAS search looks for the same signal but over a smaller mass range, from 200 GeV to 2.5 TeV. Plots showing the comparison of the expected and observed limits for $gg\phi$ and $bb\phi$ are shown in Figure 4.23. The expected limits from the CMS and ATLAS results are roughly compatible over the shared mass range, except at high mass where the extra statistics from the embedded $Z \rightarrow \tau\tau$ samples compared to MC allow for lower background uncertainties and hence a stronger limit. The ATLAS result observed no excess of events compatible with $gg\phi$ signal at 1.2 TeV, in fact a small deficit was observed. Also, ATLAS observed local excesses at 400 GeV of 2.2σ for $gg\phi$ and 2.7σ for $bb\phi$.




Figure 4.19: Distributions of $m_{\tau\tau}$ in the no b tag second highest (left) and highest (right) p_T category for the $\tau_h\tau_h$ (top), the combined $e\tau_h$ and $\mu\tau_h$ (middle) and the $e\mu$ (bottom) channels. The solid histograms show the stacked background predictions after a signal plus background fit to the data. The best fit gluon fusion signal for $m_\phi = 100$ GeV is shown by the red line. Also shown by a blue dashed line on the bottom pad is the ratio of the background predictions for the background only fit to the signal plus background fit [2].




Figure 4.20: Distributions of m_T^{tot} in the $\tau_h\tau_h$ no b tag (a) and b tag (b) categories, the combined $e\tau_h$ and $\mu\tau_h$ no b tag (c) and b tag (d) Tight- m_T categories and the $e\mu$ no b tag (e) and b tag (f) Medium- D_ζ categories. The solid histograms show the stacked background predictions after a background only fit to the data. The best fit gluon fusion signal for $m_\phi = 1.2$ TeV is shown by the red line, b associated production and U_1 signals are also shown for illustrative purposes [2].




Figure 4.21: Expected (dashed line) and observed (solid line and dots) 95% CL upper limits on the product of the cross sections and branching fraction for the decay into τ leptons for (a) $gg\phi$ and (b) $bb\phi$ production in a mass range of $60 \leq m_\phi \leq 3500$ GeV. The dark green and bright yellow bands indicate the central 68% and 95% intervals for the expected exclusion limit [2].




Figure 4.22: Comparison of the expected 95% CL upper limits on the product of the cross sections and branching fraction for the decay into τ leptons for (a) $gg\phi$ and (b) $bb\phi$ production, split by the $\tau\tau$ decay products fit individually.

None of these excesses are consistent between the ATLAS and CMS results. The ATLAS search does not stretch to the mass of the low mass CMS excess and so cannot be used a cross-check for this.




Figure 4.23: Comparison of the expected 95% CL upper limits on the product of the cross sections and branching fraction for the decay into τ leptons for (a) $gg\phi$ and (b) $bb\phi$ production, split by the CMS result detailed in this thesis and the ATLAS result from Reference [14].

4.12.2 Significance and Compatibility

The p -values and significances at each model independent signal hypothesis are calculated as described in Section 4.10 and shown in Figure 4.24. Identical to the model independent limits, the $gg\phi$ or $bb\phi$ process is allowed to float freely if not the parameter of interest. The excesses for the $gg\phi$ process peak at 100 GeV and 1.2 TeV and quantify to a local (global) significance of 3.1σ (2.7σ) and 2.8σ (2.2σ) respectively. There are also excesses at neighbouring mass points (particularly at high mass), however this is consistent with the mass resolution of the fitted templates for the central values. No deviations beyond 2σ are observed for $bb\phi$ production.




Figure 4.24: Local p -value and significance of a $gg\phi$ (left) and $bb\phi$ (right) signal as a function of m_ϕ [2].

As many different decay channels and categories are used to extract these significances, the signal strength is studied in each channel and category. This is done via compatibility fits as described in Section 4.10, where the signal strength parameter in each channel/category is decoupled. No statistically significant differences are observed in the best fit signal strength in any decay channel or category fit and p -values between each channel or category fit are always above 0.05. Figure 4.25 shows the results of the compatibility fits in the low-mass optimisation categories split by di- τ decay channels and the p_T bins fit. Figure 4.26 shows the compatibility fits in the high-mass optimisation categories split by di- τ decay channels. The low-mass signal strengths are no more dominant in any p_T region than another. In both low-

and high-mass cases, the signal strengths are consistent across di- τ decay channels. There is a small shift in the high mass $e\mu$ categories to a negative signal strength, these categories have little to no sensitivity to this signal in comparison to others and a small deficit is observed in data, resulting in fits for a negative signal strength with a large uncertainty.




Figure 4.25: Compatibility plots of the 100 GeV excess split into analysis channels (a) and categories (b). In each case the fitted signal strength is decoupled in the bin shown on the plot [2].




Figure 4.26: Compatibility plots of the 1.2 TeV excess split by analysis channels. In each case the fitted signal strength is decoupled in each channel [2].

4.12.3 2D Likelihood Scans

As the model-independent search looks for two signal modes at each mass point, the results for both processes happening simultaneously are studied. This is done in the form of two dimensional likelihood scans. The best fit cross section times branching fractions of each process and the 95% and 68% confidence intervals are shown for a number of different mass scenarios in Figure 4.27. The SM prediction in all plots is at (0,0). These results highlight how the excesses at 100 GeV and 1.2 TeV are dominated in the phase space in which $gg\phi$ and not $bb\phi$ signals are allowed. In the 60 GeV example, there are smaller deviations in both $gg\phi$ and $bb\phi$ and the SM background is again over 2σ away. Otherwise, signal strengths are completely compatible with the background expectation.

4.13 Model Dependent Limits

The exclusion contours for two benchmark scenarios of the MSSM, M_h^{125} and $M_{h,EFT}^{125}$, are presented in Figure 4.28. The red hatched regions denote areas where m_h is inconsistent with the observed SM Higgs boson mass within a ± 3 GeV boundary. For low values of $\tan \beta$, higher values of the additional SUSY particle masses, denoted as m_{SUSY} , are needed to explain a mass of approximately 125 GeV for the Higgs boson. In the M_h^{125} scenario, m_{SUSY} is fixed, and the predicted value of m_h is below 122 GeV. In contrast, the $M_{h,EFT}^{125}$ scenario adjusts m_{SUSY} to satisfy the required value of m_h for each point in $(m_A, \tan \beta)$ individually, accounting for the logarithmic corrections associated with the large values of m_{SUSY} using an effective field theory approach. The red hatched region in Figure 4.28 (b) indicates that the required values of m_{SUSY} exceed the GUT scale at very low values of m_A in this scenario. The Higgs boson masses, mixing angle α , and effective Yukawa couplings were calculated using FEYNHIGGS, and branching fractions for the decay into τ leptons and other final states were obtained from a combination of the FEYNHIGGS and HDECAY, following the prescriptions in References [119–121], for the scenarios described in Reference [83].

For the $M_{h,EFT}^{125}$ scenario, the sensitivity sharply drops at $m_A = 2m_t$ due to a drop in the branching fractions for the decay of A and H into τ leptons, when the A and H decays into two on-shell top quarks becomes kinematically accessible. Both scenarios are excluded at 95% CL for $m_A \lesssim 350$ GeV. For $m_A \lesssim 250$ GeV, most of




Figure 4.27: Maximum likelihood scans, including 68% and 95% CL contours obtained from the signal likelihood for the model-independent search. The scans are shown for selected values of m_ϕ between 60 GeV and 3.5 TeV [2].

the ggH/A events do not enter the no b tag categories due to the $m_{\tau\tau} > 250$ GeV requirement. In this parameter space, the sensitivity to the [MSSM](#) is driven by the measurements of the observed Higgs boson, even though H and A still contribute to the categories here. The sensitivity to the H and A enters mainly via the $bb\phi$ signal in the b tag categories, especially for increasing values of $\tan\beta$.

Other [MSSM](#) scenarios are tested and detailed in [2]. One scenario of note is the M_H^{125} scenario, which is the equivalent scenario to the M_h^{125} but with the observed Higgs boson being the heavier [CP](#)-even Higgs boson. Despite the local excess at a resonant mass of 100 GeV, this scenario is entirely excluded by the search. This is mostly due to the sensitivity of the b tag categories to b associated production. The local excess observed at 1.2 TeV is hard to rectify within these [MSSM](#) benchmark scenarios. The lack of any excess in the b tag categories strictly constrains the b associated production cross section times branching fraction. It is not possible within these scenarios to predict the excess of gluon fusion events within the constraints placed on b associated production.

Upper limits of 95% [CL](#) for VLQ BM 1 and 2 are shown Figure 4.29. These are drawn with respect to the leptoquark mass (m_U) and coupling (g_U). The limit on g_U decreases as m_U increases, with values of g_U ranging from 1.3 to 5.2 in VLQ BM 1 and 0.8 to 3.2 in VLQ BM 2. VLQ BM 2 has stronger exclusion limits than VLQ BM 1 due to additional right-handed couplings of the leptoquark with a b quark and a τ lepton. The observed limits fall within the central 95% intervals of the expected limits when no signal is present. The expected limits are also within the 95% confidence interval of the best fit results reported by Reference [15] and described in Section 1.4.1. This indicates that the search is capable of detecting a part of the parameter space that can explain the anomalies observed in b physics and since no significant excess was observed in this search, new constraints are placed on the vector leptoquark phase space.

Similarly to the [MSSM](#) scenarios, the local excess at 1.2 TeV is not consistent with a VLQ BM 1 or 2 vector leptoquark. Again this is due to lack of signal in the b tag categories, where the reduction in backgrounds makes the t-channel signal with initial state radiation the dominant search option.




Figure 4.28: Expected and observed 95% CL exclusion contours in the MSSM M_h^{125} (a) and $M_{h,EFT}^{125}$ (b) scenarios. The exclusion limit only on background expectation is shown as a dashed black line, the dark and bright grey bands show the 68% and 95% intervals of the expected exclusion and the observed exclusion contour is shown by the blue area. The parameter space where m_h deviates by more than ± 3 GeV from the observed SM Higgs boson mass is shown by a red hatched area. [2]




Figure 4.29: Expected and observed 95% CL upper limits on g_U in the VLQ BM 1 (a) and 2 (b) scenarios, in a mass range of $1 < m_U < 5$ TeV. The exclusion limit only on background expectation is shown as a dashed black line, the dark and bright grey bands show the 68% and 95% intervals of the expected exclusion and the observed exclusion contour is shown by the blue area. The 95% confidence interval for the preferred region from the global fit presented in Reference [15] is also shown by the green shaded area [2].

Chapter 5

Search for New Physics in $\tau^+\tau^-\tau^+\tau^-$ Final States

Enhancement from $\tan\beta$ to up or down-like quark couplings to additional Higgs bosons are essential for the majority of searches for extended Higgs sectors, including in the analysis detailed in Chapter 4. However, in some 2HDMs it can be the case that both up and down-like couplings to additional Higgs bosons are suppressed and this parameter space is left relatively untouched by “MSSM-like” searches. In particular, type X 2HDMs where only lepton couplings are enhanced by $\tan\beta$, allow for new physics loop contributions to SM measurements through couplings between leptons and additional Higgs bosons. This is particularly interesting in the context of the g-2 anomaly [28, 48] with reasoning explained in Section 1.4.2. This chapter will detail a search for such an extended Higgs sector, that looks for a production mode not suppressed at high $\tan\beta$, through the process $Z^* \rightarrow \phi A \rightarrow 4\tau$. This search is split up into two sections:

- i) A model independent search for the $Z^* \rightarrow \phi A \rightarrow 4\tau$ process. Both additional particles are required to have narrow width and no assumptions are made on the production cross-section via an off-shell Z boson or the branching fraction of ϕ and A decaying to a pair of tau leptons.
- ii) A search for the type X 2HDM, motivated by the phase space for possible explanations to the g-2 anomaly. The m_A - $\tan\beta$ phase space for scenarios of m_ϕ in the alignment limit is scanned, as well as checks outside of this limit on the $\cos(\beta - \alpha)$ - $\tan\beta$ for specific scenarios of both m_ϕ and m_A .

These searches are performed with the full run-2 dataset (138 fb^{-1}) collected by the CMS experiment.

5.1 Signal Modelling

Any additional Higgs boson produced in the type X [2HDM](#) at high $\tan\beta$ will predominantly decay to tau leptons. To probe the type X [2HDM](#) at high $\tan\beta$, a production process that is not suppressed is required. Reference [16] discusses that the following production modes of two additional Higgs bosons are dominant to produce any of these new particles at high $\tan\beta$:

- i) $pp \rightarrow Z^* \rightarrow \phi A \rightarrow (\tau^-\tau^+)(\tau^-\tau^+)$
- ii) $pp \rightarrow Z^* \rightarrow H^+H^- \rightarrow (\tau^-\nu)(\tau^+\nu)$
- iii) $pp \rightarrow W^{\pm*} \rightarrow H^\pm A \rightarrow (\tau^\pm\nu)(\tau^-\tau^+)$
- iv) $pp \rightarrow W^{\pm*} \rightarrow H^\pm\phi \rightarrow (\tau^\pm\nu)(\tau^-\tau^+)$

As the production cross sections are of the similar magnitudes, the search sensitivities depend on the separation of the signals from background. In general, the more objects you can select in the final state, the smaller the background contributions. This is certainly true in tau enriched final states, where backgrounds can be dominated jets misidentified as hadronic taus and so every extra tau selected reduces this background. In particular, (ii) has the production cross sections [16] far smaller than the observed limit for gluon fusion production of a single resonance shown in Figure 4.21(a) and it is not possible to use tau decay product and [MET](#) alignment to separate the background, so does seem not a viable search option with the run-2 [CMS](#) dataset. The increased background from fewer object selections and looser charge sum selection on (iii) and (iv), makes (i) the golden search channel for a type X [2HDM](#). A Feynman diagram for this process is shown in Figure 5.1.




Figure 5.1: Diagram of production of two additional neutral Higgs bosons from an off-shell Z boson and their decays to tau leptons.

Signal templates for the production of this process with a mass grid for ϕ and A between 100 to 300 and 60 to 160 GeV respectively are generated. These mass ranges are motivated from the results in Table 1.4. The samples are simulated in the five **FS** at **NLO** precision using the **MADGRAPH5_aMC@NLO v2.6.5** event generator [95]. Generation is performed using the parton distribution function **NNPDF3.1** [89, 90], where the τ lepton decay, parton showering and hadronisation are all modelled with the **PYTHIA** event generator with the **PU** profile matched to data [91, 92]. The events are then passed through the **GEANT4**-based [93] simulation of the **CMS** detector and reconstructed in the same way as data. Generator level distributions of di- τ mass distributions from the decay of ϕ and A are shown in Figure 5.2.




Figure 5.2: Generator level distributions of the visible mass densities for A (a) and ϕ (b) for the signal process.

The cross sections in the alignment scenarios are also determined with this procedure and vary from 10 fb (highest m_A and m_ϕ scenario) to 650 fb (lowest m_A and m_ϕ scenario), as shown in Figure 5.3. These are independent of $\tan\beta$, however, out of the alignment scenarios the cross sections for H scales with $\sin^2(\beta - \alpha)$ and h scales with $\cos^2(\beta - \alpha)$.

The branching fractions of ϕ and A to pairs of τ leptons is dependent on both $\tan\beta$ and $\beta - \alpha$. For this analysis, the branching fractions are calculated using **2HDECAY** [122]. In the alignment scenarios, the $A \rightarrow \tau\tau$ branching fractions are approximately 1 above $\tan\beta \approx 2$, where below they sharply drop off and other




Figure 5.3: Calculated production cross sections for the $Z^* \rightarrow \phi A$ process, varying the masses of ϕ and A .

processes such as $A \rightarrow b\bar{b}$ become dominant. This is also true for $\phi \rightarrow \tau\tau$ branching fractions, except in the case where m_ϕ is greater than m_A by more than m_Z , and so the $\phi \rightarrow ZA$ decay becomes kinematically feasible and can dominate at high $\tan\beta$. Examples of this are shown in Figure 5.4. Out of the alignment scenario, the branching fractions of ϕ to tau leptons becomes smaller as the magnitude of the coupling of additional neutral Higgs bosons to taus is reduced, whilst the A branching fractions, like the couplings, are left unchanged. An example of the ϕ branching fractions out of the alignment scenario is shown in Figure 5.5.

5.2 Event Selection

In comparison to Chapter 4, four τ leptons produce a much larger number of possible final states. All variations of e , μ and τ_h final state combinations and their branching fractions are shown in Table 5.1. Just under 90% of the branching ratio goes to decay products containing two or more hadronic taus. These are the main final states that are explored in this analysis. In addition to this, an orthogonal $\tau_h\tau_h\tau_h$ channel is added to target events where reconstruction of the $\tau_h\tau_h\tau_h\tau_h$ channel loses a single τ_h object. This can come about due to low triggering and ID efficiency of τ_h candidates, as well as the high p_T thresholds required for both.




Figure 5.4: Calculated branching fractions of A (a) and ϕ (b) decaying to a pair of τ leptons for various mass scenarios.




Figure 5.5: Calculated branching fractions in the $\cos(\beta - \alpha)$ - $\tan\beta$ phase space for ϕ of mass 200 GeV decaying to a pair of τ leptons, in the scenario where $m_A = 160$ GeV.

In total, the analysis consists of 7 channels.

Channel	Branching Fraction
$e\tau_h\tau_h\tau_h$	19.4%
$\mu\tau_h\tau_h\tau_h$	18.9%
$\tau_h\tau_h\tau_h\tau_h$	17.6%
$e\mu\tau_h\tau_h$	15.6%
$ee\tau_h\tau_h$	8.0%
$\mu\mu\tau_h\tau_h$	7.6%
$ee\mu\tau_h$	4.3%
$e\mu\mu\tau_h$	4.2%
$eee\tau_h$	1.5%
$\mu\mu\mu\tau_h$	1.4%
$eee\mu$	1.4%
$ee\mu\mu$	0.6%
$e\mu\mu\mu$	0.4%
$eeee$	0.1%
$\mu\mu\mu\mu$	0.1%

Table 5.1: Branching fractions of four τ leptons, where e and μ represent the leptonic decay of the τ and τ_h represent the hadronic decay of the τ .

5.2.1 Trigger Requirements

Given that each final state is not exactly triggered on, there is no obvious choice for what triggers to use. A variety of triggers are available for individual and clusters of objects in the final state. The possible triggers for single objects are the single-e and single- μ triggers. This is not the case for the single- τ_h trigger as it has a p_T threshold too high for it to be useful. The possible triggers for clusters of objects are the double-e, double- μ , double- τ_h , e- μ cross, $\mu\tau_h$ cross and e- τ_h cross-triggers. The cross-triggers and double-e/ μ triggers are found to offer little improvement to the signal acceptance and so these events are not included. Any combination of objects in the final state of a channel can be selected by these triggers and the union of events passing each iteration is taken. The trigger p_T and η thresholds for the remaining triggers (single-e/ μ and double- τ_h) are equivalent to what is stated in Section 4.2.1.

5.2.2 Offline Requirements

All offline selections stated are in addition the object selection discussed in Section 3. In this analysis, τ_h candidates are required to pass the **Loose** $D_{\text{jet}}^{\text{WP}}$. The **VVLoose** D_e^{WP} and **VLoose** D_μ^{WP} are used in all decay channels. These working points are chosen to maximise the sensitivity of the analysis and looser cuts are used than in Chapter 4.1.1 to ensure there are enough statistics to account for the stricter selection of more objects in the final states.

On top of the object selection, there are few further selections on the total tau collection. As the two tau pairs from the signals originate from two neutral additional Higgs bosons, the sum of charge of the fully reconstructed objects should be 0. This is applied in the decay channels where there are four objects. In the $\tau_h\tau_h\tau_h\tau_h$ channel, as the assumption is that a hadronic tau has been lost, the absolute value of the sum of the charges of the object is required to be 1. In order to ensure the orthogonality between channels, a number of vetos are needed. Firstly, extra lepton vetos are used in all decay channels, where the absence of additional electrons and muons is required on top of those already part of the selected pair. The kinematic requirements, ID and isolation requirements on these extra leptons are at least as loose as the loosest cuts required for signal electron or muon of the final states. Secondly, an extra τ_h veto is applied to the $\tau_h\tau_h\tau_h\tau_h$ channel, in order to keep it orthogonal to the $\tau_h\tau_h\tau_h\tau_h$ channel. Here any extra τ_h candidate passing the signal selection is vetoed. The constraints set on the number of leptons and τ_h candidates for each channel are shown in Table 5.2.

Channel	e	μ	τ_h
$\tau_h\tau_h\tau_h\tau_h$	0	0	≥ 4
$\tau_h\tau_h\tau_h$	0	0	3
$\mu\tau_h\tau_h\tau_h$	0	1	≥ 3
$e\tau_h\tau_h\tau_h$	1	0	≥ 3
$e\mu\tau_h\tau_h$	1	1	≥ 2
$\mu\mu\tau_h\tau_h$	0	2	≥ 2
$ee\tau_h\tau_h$	2	0	≥ 2

Table 5.2: Veto set on the number of leptons for each channel.

Finally, in channels containing an electron or a muon, a veto on events with one or more b tagged jets is placed. This is done to remove the $t\bar{t}$ background process in a region where little to no signal is expected. This is not done in the fully τ_h channels as the $t\bar{t}$ process is negligible.

5.3 Search Optimisation

Due to the limited statistics in each decay channel, only minimal categorisation of events can be performed and the only divisions of the decay channels happen in final states with two light leptons and two τ_h candidates, where two categories are made. These separate events where the light leptons have the same charge (**SS Leptons**) and where they have opposite charge (**OS Leptons**). This is motivated to separate regions where specific background processes are dominant. In particular, a large portion of the $e\mu\tau_h\tau_h$ and $\mu\mu\tau_h\tau_h$ channel backgrounds come from $Z \rightarrow ll$ (or $Z \rightarrow \tau\tau \rightarrow ll$) with 2 jets misidentified as τ_h candidates, where the two light leptons are of opposite sign. Similarly, the $e\mu\tau_h\tau_h$ channel contains more background events with an opposite sign electron and muon, from a Z decay via taus or from the $t\bar{t}$ process. There is no preference for light lepton charge sums in the signal samples and so more sensitivity to the signal is expected in the **SS Lepton** categories. A summary of the categorisation and b tag selection is shown in Figure 5.6.




Figure 5.6: Overview of the categories used for the extraction of the signal in the $Z^* \rightarrow \phi A \rightarrow 4\tau$.

Similarly to Chapter 4.1.1, events in each 10 analysis channels and categories are drawn out into histogram based on the same discriminating variable, the total transverse mass (m_T^{tot}). However, as there are more objects in the final state, the definition is extended to what is shown below.

$$m_T^{\text{tot}} = \sqrt{\sum_{i=1}^{N_\tau} m_T(\vec{p}_T^{\tau_i}, \vec{p}_T^{\text{miss}})^2 + \sum_{i,j=1; i \neq j}^{N_\tau} m_T(\vec{p}_T^{\tau_i}, \vec{p}_T^{\tau_j})^2}, \quad (5.1)$$

where N_τ is the number of objects in the final state, and τ_i refers to the visible products of the i 'th τ lepton. This variable again provides excellent discriminating power between resonant signals compared to other non-peaking backgrounds, whilst still maintaining some separation between signal masses. The sensitivity of this discriminating variable was tested against many others, including the visible masses of the two bosons which can be separated to $\approx 90\%$ efficiency after selection into the correct pairs. As m_T^{tot} uses information about the whole event, in particular the higher p_T objects and high MET from many τ decays, a greater sensitivity to the signal is observed.

In this analysis, many of the histograms contain very few background events and in order to minimise statistical fluctuations in the background templates, histogram bins are merged based off the fractional statistical uncertainty of each bin. This leads to a number of single and minimally binned channels/categories where background statistics are low but signal sensitivity is high and a few finely binned and high statistic channels/categories able to better classify the signal, if one is observed.

5.4 Background Modelling Overview

The backgrounds are split into a three categories:

- i) Events containing only genuine τ leptons.
- ii) Events with one or more jet misidentified as a τ_h (jet $\rightarrow \tau_h$).
- iii) Events where all τ_h candidates are either correctly identified or a misidentified light lepton.

Background contributions from (i) are mostly from gluon and quark initiated di-Z production and is modelled using MC. The details and validation of this modelling is described in Section 5.5. Background (ii) accounts for a number a different process

with jets misidentified as a τ_h candidate. Examples of this are single-Z, single-W and $t\bar{t}$ productions with additional jets in the events being misidentified. This is modelled with a fake factor method, similar to what is described in Section 4.7, but using machine learning to improve upon method, and discussed further in Section 5.6. The latter contribution is very small in comparison to the others and modelled with MC. Events with jets misidentified as light leptons and no jets misidentified as τ_h candidates have been checked with MC and deemed negligible in all channels.

All MC background samples described in Section 4.4 are modelled in the same manner. In addition, tri-boson samples are used that are generated using MADGRAPH5_aMC@NLO at next-to-LO (NLO) precision [95]. All corrections described in Section 4.8 are applied to MC, with extra corrections are applied on the ZZ \rightarrow 4l process, that are detailed in Section 5.5.

5.5 ZZ Modelling

The di-Z background shapes are modelled using MC. The cross-sections for this analysis are scaled to high order precisions than the signal generation using k factors. For quark initiated di-Z production, PowHEG 2.0 [85–88] is used to determine NNLO/NLO QCD and electroweak corrections as a function of m_{ZZ} , which takes an average value of ≈ 1.2 . Gluon initiated di-Z production corrections from LO to NNLO is derived with HNNLO v2 [123], again as a function of m_{ZZ} . This gives a much larger correction, equal to ≈ 2 . The modelling is validation using a $\mu\mu\mu\mu$ final state and good agreement is observed between data and simulation. Identical object selection is applied as stated in Section 3, except to ensure better statistics within the channel, the I_{rel}^μ cut is loosened to 0.35. The charges of the μ candidates are similarly required to sum to one and no vetos on the number of b jets in the event is applied. Plots of this are shown in Figures 5.7.

5.6 Machine Learning Fake Factor Method

The F_F method, as described in Section 4.7, is an algorithm used to model the jet $\rightarrow \tau_h$ backgrounds. It uses classical reweighting techniques, such as binning the two regions and applying a fit to the ratio in a chosen parametrisation. This method faces difficulties when approached with highly dimensional dependence on




Figure 5.7: Distributions in the $\mu\mu\mu\mu$ channel are shown for the variables total transverse mass, m_T^{tot} , (a) and the total mass, $m_{\mu\mu\mu\mu}$ (b). The solid histograms show the stacked background predictions.

the parametrisation. In this scenario, classical ‘‘bin and fit’’ methods can break down, as for every new parameter included the statistics of each fit is reduced. A compromise must then be reached between variable dependence and fit statistics and is the case for the jet- p_T/τ_h - p_T binning described in Section 4.7.2. It is also the case, that the analysis in Section 4.1.1 uses well checked assumptions on for example, the consistency of τ_h decay modes from determination to signal regions, in order to average across the dependencies on this variable when calculating F_F , however this will not always be the case. Also binned upon in the previous search, is the process dependence of the F_F . The binning used is an admission that the relevant parametrisation of the F_F between process is too complication to model with ‘‘bin and fit’’ methods. It can also be seen that the initial fits, do not do a good job at modelling all variables as corrections are needed.

There is final problem with using a F_F method as described, when there are multiple τ_h candidates in final states. In the $\tau_h\tau_h$ channel of the previous analysis, the F_F were only calculated from the leading- τ_h candidate. This was valid, as the dominant backgrounds had two jet $\rightarrow \tau_h$ candidates rather than one genuine τ_h and one jet $\rightarrow \tau_h$ candidate. However, for this search this assumption is not valid.

All of these reason motivate a more generalised and smarter method to model jet

$\rightarrow \tau_h$ backgrounds, as well as the extra importance on modelling these backgrounds where there are more τ_h candidates in the final state. This is done utilising machine learning and in particular using a BDT for the purpose of multi-dimensional reweighting.

5.6.1 BDT Reweighting

Reference [124] proposes a new method utilising machine learning techniques to solve the issues with dimensionality. It looks to optimise the regions that most need reweighting. One good way to do this is using a decision tree, as with this the data can be split into “leafs” by checking simple conditions. To best choose the regions that need reweighting the algorithm looks to maximise the symmetrised χ^2 .

$$\chi^2 = \sum_{\text{leaf}} \frac{(w_{\text{leaf}, 1} - w_{\text{leaf}, 2})^2}{(w_{\text{leaf}, 1} + w_{\text{leaf}, 2})^2} \quad (5.2)$$

The larger the value of χ^2 the more important reweighting is in this region. The kind of tree is utilised many times in the reweighting algorithm shown below:

- i) Input training dataset with a large number of variables.
- ii) Build a tree as stated above. If not the first loop, use newly determined weights.
- iii) Compute predictions in the leaves $r_{\text{leaf}} = \log \frac{w_{\text{leaf}, 1}}{w_{\text{leaf}, 2}}$. The logarithm is taken so weights in different trees can be summed as usually done in boosting.
- iv) In each leaf MC events by $w = w \times e^{r_{\text{leaf}}}$.

The final two steps are identical to the first approach except for the use of the logarithm for convenience using boosting. The major difference being how the bins used for reweighting are found, and that this step is repeated multiple times.

5.6.2 Fitting Regions

Unlike the standard F_F method, statistics are not a problem using the BDT reweighter when choosing which variables you can use to parametrise the F_F . Therefore, rather than fitting two fake factor regions separately and then a correction from sideband to signal to account for missing parametrisation, regions A,B and D from Figure 4.10

are fit simultaneously to improve statistics. Also, all τ_h candidates are fit simultaneously as separate entries in the dataset and for each τ_h in the event, the remaining τ_h candidates are named the alternative τ_h candidates, rather than the τ_h that is being fit. The sideband variable definitions and cuts, with respect to Figure 4.10, are shown below.

- i) $ee\tau_h\tau_h$, $e\mu\tau_h\tau_h$, $\mu\mu\tau_h\tau_h$, $e\tau_h\tau_h\tau_h$ and $\mu\tau_h\tau_h\tau_h$

y_C : The sum of τ_h candidates is required to be 0.

y_A : The sum of τ_h candidates is required to not be 0.

x_C : All alternative τ_h candidates pass the **Loose** $D_{\text{jet}}^{\text{WP}}$.

x_D : At least one alternative τ_h candidate fails the **Loose** $D_{\text{jet}}^{\text{WP}}$ but has $D_{\text{jet}}^{\text{score}} > 0.1$.

- ii) $\tau_h\tau_h\tau_h$

y_C : The absolute value of sum of τ_h candidates is required to be 1.

y_A : The absolute value of the sum of τ_h candidates is required to not be 1.

x_C : All alternative τ_h candidates pass the **Loose** $D_{\text{jet}}^{\text{WP}}$.

x_D : At least one alternative τ_h candidate fails the **Loose** $D_{\text{jet}}^{\text{WP}}$ but has $D_{\text{jet}}^{\text{score}} > 0.1$.

These selections mimic what is done for the $\tau_h\tau_h$ channel F_F from Section 4.7.1, where the definitions of the alternative sideband variable are extended to more than one other τ_h candidate. The $D_{\text{jet}}^{\text{score}} > 0.1$ selection is used as the alternative tau ID selection for this analysis and is also then extended to the alternative τ_h candidates for this selection. A cut on the score is used instead of a working point as the loosest defined working point does not provide enough statistics to ensure a good fit.

Extrapolating the regions used for fitting to the $\tau_h\tau_h\tau_h\tau_h$ channel, there would be some overlap in the fitting region with the $\tau_h\tau_h\tau_h$ signal region. As this region is expected to be sensitive to signal, this is not used to model jet $\rightarrow \tau_h$ backgrounds in the $\tau_h\tau_h\tau_h\tau_h$ channel. Instead the fit from the $\tau_h\tau_h\tau_h$ channel is used. There are a few variables that are defined differently in the fit between the two channels due to the difference of four to three objects selected, therefore when getting F_F in the $\tau_h\tau_h\tau_h\tau_h$, the lowest $D_{\text{jet}}^{\text{score}}$ unused τ_h candidate is dropped and variables recalculated. This candidate is chosen to be removed to best mimic the initial selection of the τ_h candidates where they are sorted by $D_{\text{jet}}^{\text{score}}$ and the high scoring candidates

are chosen. As shown later in this section, there is no major dependence on the shifted variables and any effect from this removal is covered within the uncertainty model.

5.6.3 Variables Used

The variables used are ones that have been shown to have F_F dependence previously [2, 125], and additional properties of the τ_h candidate and the event. Also added are the variables that take you from A to C and D to C, from Figure 4.10. As all years are fit together, in order to account for any differences in fake fake factor from year to year, this is also added. The final variable added is the p_T ordered ranking of the τ_h candidates in the event. All these variables are shown below.

- i) The [HPS](#) decay mode of the hadronic tau candidate.
- ii) p_T of the hadronic tau candidate.
- iii) The ratio of the p_T of the matched jet to the p_T of the hadronic tau candidate.
- iv) η of the hadronic tau candidate.
- v) The charge of the hadronic tau candidate.
- vi) A boolean of whether the hadronic tau candidate passes a leg of the double tau trigger.
- vii) The total charge of the combined objects.
- viii) The boolean of whether $D_{\text{jet}}^{\text{WP}}$ passes to [Loose](#) WP for the other hadronic tau candidates. These are sorted by p_T .
- ix) Era of data taking.
- x) τ_h p_T ordered event rank.

5.6.4 Machine Learning Subtraction Method

In the standard F_F method, histograms are used to fit the F_F rather than datasets. Using histograms, the small fraction of events which are not jet fakes can easily be subtracted off. To do this, the data histogram is subtracted from by a stacked

MC background produced with generator matching ensuring the event is not a jet $\rightarrow \tau_h$ and this produces a data-**MC** hybrid histogram of predicted jet $\rightarrow \tau_h$ events. However, subtraction is not possible with a full dataset and negative weights do not work with the **BDT** reweighter. Therefore, the only option is to remove like-for-like events in data compared to the non jet $\rightarrow \tau_h$ generator matched MC. An example of this is template matching, which takes an event and can find the closest event in another datasets. But as the fitting dataset is highly dimensional, this requires too much computation. The solution proposed for this is to use a **BDT** to reduce the dimensionality of the datasets effectively the one dimension of an output score of a simple binary classifier.

- i) In each channel, all **MC** in the fitting region is stacked and scaled to cross section (via weighting) and the variables used for reweighting (Section ??) are put into a dataset.
- ii) Events that are jet fakes and not jet fakes are separated into the two classes that will be used for the binary classification.
- iii) To ensure unbiased training, the weights of the two categories are normalised to one another.
- iv) A **BDT** is then trained to separate whether the **MC** is jet $\rightarrow \tau_h$ candidate or not.
- v) The scores of the **BDT** for how likely the entry is not a jet $\rightarrow \tau_h$ candidate is added to the dataset.
- vi) The scores of the non jet $\rightarrow \tau_h$ candidates are drawn into a histogram with a number of bins suitable for the number of statistics and rescaled to cross section to best match what would be observed in data.
- vii) The output score of the **BDT** is added to data events
- viii) Each bin of the **MC** histogram is then looped through:
 - Data events with **BDT** score within the range of the bin are selected.
 - Events within this bin are then randomly sample and removed.
 - This stops when the number of events removed equals to the number of non jet fake events predicted in the **MC** histogram bin.

This method then gives a data-[MC](#) hybrid method to determine a dataset of predicted jet $\rightarrow \tau_h$ candidate and should give near identical results when drawing out histograms in all variables compared to subtracting off [MC](#) non jet $\rightarrow \tau_h$ candidates from a data histogram as used in the original F_F method. It allows for non jet $\rightarrow \tau_h$ -like candidates to be sampled and removed to the correct yield through all variables. It is important to remove events throughout the [MC](#) non jet $\rightarrow \tau_h$ [BDT](#) score histogram as if only the highest score events were removed, these events would come primarily from the tails of the distribution where it is easiest to separate.

An uncertainty is placed on the performance of this algorithm with respect to histogram subtraction. This is calculated by drawing each variable used for the method into a histogram with binning chosen to ensure a sensible number of events in each bin. An uncertainty is then derived from the difference in prediction between the histogram and [BDT](#) subtraction.

To validate this method, example histograms with this uncertainty are shown for the $\mu\tau_h\tau_h\tau_h$ pass region in Figures 5.8, comparing the [BDT](#) subtraction method and histogram subtraction method for a few of the fitted variables.

5.6.5 Fitting

These subtracted datasets are then randomly split 50:50 into a train and test datasets and only the train dataset is fit. The [BDT](#) reweighter has a number of hyperparameters and these are tuned with a scan optimising the Kolmogorov-Smirnov test [126] on the test dataset in each channel separately. Final models are then produced that can optimally model jet $\rightarrow \tau_h$ candidates for all fitted variables. Examples of the fake factor derived in the $\mu\tau_h\tau_h\tau_h$ channel are shown in Figure 5.9.

An uncertainty is placed on the performance of this algorithm. This is again calculated by drawing each variable into a histogram and comparing the histograms from reweighted events with the alternative τ_h ID selections to the events with the nominal τ_h ID in all of the fitted regions simultaneously. Plots of the closure of this method in the $\mu\tau_h\tau_h\tau_h$ accompanied by this uncertainty are shown in Figure 5.10.




Figure 5.8: Comparison of histograms produced via the BDT subtraction method to histogram subtraction. Also shown is the histogram produced when no subtraction is performed. The uncertainty bands contain statistical uncertainties and uncertainties derived on the non-closure of the method. The figure is shown for four of the fitted variables: τ_h - p_T , the ratio of τ_h - p_T to jet- p_T , the τ_h HPS decay mode and the τ_h - η .




Figure 5.9: Average fake factors (reweights) calculated by the BDT reweighting method shown individually in regions A, B and D as defined in Figure 4.10. The is shown for four of the fitted variable: τ_h - p_T , the ratio of τ_h - p_T to jet- p_T , the τ_h HPS decay mode and the τ_h - η .




Figure 5.10: Comparison of histograms produced using the fake factors applied to the fitted fail region compared to fitted pass region. The uncertainty bands contain statistical uncertainties and uncertainties derived on the non-closure of the method. The figure is shown for three of the fitted variable: $\tau_h\text{-}p_T$, the ratio of $\tau_h\text{-}p_T$ to jet- p_T , the τ_h HPS decay mode and the $\tau_h\text{-}\eta$. The χ^2 divided by number of degrees of freedom between the two histograms is also shown.

5.6.6 Applying Fake Factors

Fake factors, F_F^i , have now been calculated for each τ_h candidate, i , in the event and uncertainties determined on each weight. However, it is difficult to generate a full description of any number of jet $\rightarrow \tau_h$ fakes in an event. Taking channels with two τ_h candidates as the simplest example, if the jet $\rightarrow \tau_h$ background is determined purely off the leading τ_h candidate, then events the leading τ_h is genuine and the sub-leading τ_h is a jet, are missed. Similarly, the situation can be flipped if the sub-leading τ_h is chosen to determine the jet $\rightarrow \tau_h$ fake background. A third options can be tried where both τ_h candidates are used to determine the background. However, this will only model events where both τ_h candidates are jets and not where there is a single jet $\rightarrow \tau_h$ fake. It is seen that if the first two attempts at calculating this background from individual candidates are added and the contribution from both τ_h candidates is subtracted, all possible numbers of jet fakes in the event are accounted for.

Region	$\tau_h^1(\tau)\tau_h^2(j)$	$\tau_h^1(j)\tau_h^2(\tau)$	$\tau_h^1(j)\tau_h^2(j)$
R_1 (from τ_h^1)	0	1	1
R_2 (from τ_h^2)	1	0	1
R_{12} (from both)	0	0	1
$R_1 + R_2 - R_{12}$	1	1	1

Table 5.3: Regions modelled by the fake factor method when using the leading τ_h (τ_h^1), the sub-leading τ_h (τ_h^2) and both, to model the jet $\rightarrow \tau_h$ background and whether that specific object is a genuine τ lepton (τ) or a jet (j). Also shown is a combination of three regions to fully model all possible combinations.

This logic is extended to all channels with one exception. The $\tau_h\tau_h\tau_h\tau_h$ channel application region would have overlap with the $\tau_h\tau_h\tau_h$ signal region using this common method, therefore the formula is adjusted to avoid this region. If the overlapped regions are removed from the equation, the scale of the triple and quadruple regions needed to be adjusted to account for this. The caveat to this is that events where there are only one jet $\rightarrow \tau_h$ candidate are not accounted for. As the majority of the background events in this channel comes from [QCD](#) with many jet $\rightarrow \tau_h$ candidates, this contribution is deemed negligible. The formulae for the total jet $\rightarrow \tau_h$ backgrounds in each channel are shown below.

i) $\mu\mu\tau_h\tau_h$, $ee\tau_h\tau_h$ and $e\mu\tau_h\tau_h$

$$R_1 + R_2 - R_{12} \quad (5.3)$$

ii) $\mu\tau_h\tau_h\tau_h$, $e\tau_h\tau_h\tau_h$ and $\tau_h\tau_h\tau_h\tau_h$

$$R_1 + R_2 + R_3 - R_{12} - R_{13} - R_{23} + R_{234} \quad (5.4)$$

iii) $\tau_h\tau_h\tau_h\tau_h$

$$\begin{aligned} & R_{12} + R_{13} + R_{14} + R_{23} + R_{24} + R_{34} \\ & - 2(R_{123} + R_{124} + R_{134} + R_{234}) + 3R_{1234} \end{aligned} \quad (5.5)$$

5.7 Uncertainty Model

The uncertainty model follows the schemes detailed in Section 4.9 for the statistical uncertainties and systematic uncertainties for light leptons, jets, leptons misidentified as hadronic taus, MET, luminosity and prefiring. Updates and additions to the previous uncertainty model are shown below.

Hadronic Taus

An improvement is made to the τ_h identification uncertainty correlation scheme applied to simulated events. The fit used to derive scale factors for the τ_h MC events consists of both statistical and systematic uncertainties. In the previous search, the ID uncertainties were correlated across decay mode and era despite the statistical components of each fit being orthogonal. Therefore for this search the uncertainty scheme contains correlated and decorrelated parts across decay and era. The double- τ_h trigger uncertainties remain unchanged.

Jets misidentified as hadronic taus

The backgrounds with jets misidentified as τ_h are estimated from data with the machine learning (ML) F_F method. There are different sources of uncertainty related to this method. All uncertainties are uncorrelated across decay channels, except in the $\tau_h\tau_h\tau_h\tau_h$ and $\tau_h\tau_h\tau_h$ where the same fit is used, so uncertainties are

correlated.

The initial uncertainties come from removal of the non jet $\rightarrow\tau_h$ backgrounds from the fitting region and this is split into two types. Firstly, an uncertainty is placed on the non-closure of the **BDT** subtraction method, this is done by comparing the distributions in each variable fit using standard histogram subtraction with the **BDT** subtraction method, and the largest shift is taken for each event. This is done separately in the pass and fail τ_h identification regions. The second uncertainty to do with purify the fitting region is motivated by any **MC** mismodelling of the non jet $\rightarrow\tau_h$ objects predicted. This is shifted up and down by 10%, to represent any **MC** mismodelling , and the **BDT** subtraction method and the reweighting is repeated with the differing datasets.

The second source of uncertainties comes from the **BDT** reweighter fit. In a similar way to the subtraction method, an uncertainty is placed on the non closure of the fit, comparing reweighted events in the fail τ_h ID region to the pass τ_h ID region. Further uncertainties are placed on the assumption of the variables used to separate the signal region from the fitting region. The assumption is that the fake factors would be identical no matter what combination of these variables are used. Therefore, the uncertainties are placed by taking the largest shift when changing these variables whilst getting the output to the fit. These are decorrelated in each combination of the separating variables.

Background process specific uncertainties

Specific uncertainties are placed on the di-Z simulated events due to application of k factors. Due to the size of the differences between **NNLO** and lower order predictions for the cross sections, an uncertainty is placed on the size of these yields utilising the k factors derived.

Signal process specific uncertainties

Parton distribution functions, α_s and μ_R/μ_F scale variations are applied on an event-by-event basis to the signal samples. The normalisation of these uncertainties are approximately 6%, 1% and 2% respectively.

5.8 Signal Extraction

The statistical interpretation of the results is done as described in Section 4.10, but with different signal scaling functions, g , and parameters of interest, μ . The two interpretation of the analysis, as stated at the beginning of chapter, have different parameters of interest and scaling functions.

Firstly, the model independent search uses a linear scaling function, $g(\mu) = (\mu)$, and a parameter of interest that represents the cross section of the $Z^* \rightarrow \phi A$ multiplied by the branching fractions of $\phi \rightarrow \tau\tau$ and $A \rightarrow \tau\tau$. Secondly, the interpretation in the type X 2HDM model is done by testing each point in the parameter space, whether that is $m_A \cdot \tan \beta$ for the alignment scenario or $\cos(\beta - \alpha) \cdot \tan \beta$ for scenarios of the remaining parameters. This is done by scaling the samples to the predicted cross sections times branching ratios at that point in the parameter space, and defining a single rate parameter that can only take the values 1 (type X 2HDM) and 0 (SM) with $g(\mu) = \mu$.

5.8.1 Postfit Plots

Figure 5.11 shows the unblinded distributions of the m_T^{tot} discriminator, after a background-only fit to data, in every bin used in the fit. For visualisation, categories with similar event numbers in each bin are displayed on the same plot. A stacked background of events is shown, separated into three groups: events with 1 or more jet $\rightarrow \tau_h$, events where all τ_h candidates are reconstructed correctly, and the remaining events where only light leptons are misidentified as τ_h and not jets. An example signal hypothesis is also shown for a mass hypothesis of $m_A = 160$ GeV and $m_\phi = 200$ GeV scaled to 0.01 pb, which is approximately three times smaller than the predicted cross section for this process.

There are no upward deviations of the number of events observed, that would be consistent with any mass hypotheses for the signal model searched for. The combined results are consistent with a background-only fit to data. Within this combined fit, individual bins and categories such as the $e\mu\tau_h\tau_h$ SS Leptons, have small deficits of events observed in comparison to events expected. The background prediction yield in this category from the F_F method is checked with a comparison to MC for the non-QCD prediction. The background estimations are compatible when the estimation of the fraction of QCD events from the charge inverted region is taken

in account, further validating the F_F background prediction in this category.

5.9 Model Independent Results

5.9.1 Limits

95% CL are set on the cross section for two additional neutral Higgs bosons produced via an off-shell Z boson multiplied by the branching fraction of each additional boson decaying to τ lepton pairs, and are shown in Figure 5.12. In order to show this for all 72 mass hypotheses, the limit on higher masses of the A boson are scaled to larger negative powers of 10, as indicated on the plot. The observed limit falls between the lower 1 and 2σ bands of the expected result. This is consistent with the distributions observed in Figure 5.11, as the small deficits in sensitive channels lead to a strengthening of the limit. The observed limit in comparison to the expected limit is relatively consistent across the signal mass range. This is because of the degeneracy of the signal hypothesis in the sensitive category bins fit. Therefore, the strongest constraints on different mass hypotheses mostly arise from the same bins. The observed limits vary from 20 fb at the lowest mass hypothesis of $m_A = 60$ GeV and $m_\phi = 100$ GeV, to 1.4 fb at the highest mass hypothesis of $m_A = 160$ GeV and $m_\phi = 300$ GeV. It is worth noting that each of the observed limits are well below the predicted cross sections calculated and shown in Figure 5.3, indicating excellent sensitivity to the type X 2HDM in the alignment scenario.

Figure 5.13 shows the effect of individual decay channels, by showing the expected 95% CL limit on fits to each decay channel separately for an $m_A = 100$ GeV scenario. It is observed from this, that the dominant search channels, across the whole m_ϕ range, are the $\tau_h\tau_h\tau_h\tau_h$, $\mu\tau_h\tau_h\tau_h$ and $e\mu\tau_h\tau_h$. The $\tau_h\tau_h\tau_h$ channel can significantly contribute to the combined limit at higher values of m_ϕ , as the distribution of events in m_T^{tot} peaks higher and so can be more easily separated from the jet $\rightarrow\tau_h$ backgrounds. The remaining channels contribute less to the combined limit, as for $e\tau_h\tau_h\tau_h$ more electrons and jets misidentified as τ_h candidates are present due to worse rejection power for the processes that contribute, and for the $ee\tau_h\tau_h$ and $\mu\mu\tau_h\tau_h$ channels the low branching fractions from four τ leptons and difficult background separation where the light leptons have opposite charge, make it difficult to get high signal over background acceptance.




Figure 5.11: Distributions of m_T^{tot} in the high (a), medium (b) and low (c) statistic categories. As the bin sizes vary drastically between channels, the bin sizes are kept constant and the bin intervals are shown on x axis. The high statistic categories consist of only the $\tau_h\tau_h\tau_h$ channel, the medium statistic categories include the $ee\tau_h\tau_h$ and $\mu\mu\tau_h\tau_h$ OS Leptons categories, and the low statistic categories show the $e\tau_h\tau_h\tau_h$, $\mu\tau_h\tau_h\tau_h$, $\tau_h\tau_h\tau_h\tau_h$, $ee\tau_h\tau_h$ SS Leptons, $\mu\mu\tau_h\tau_h$ SS Leptons, $e\mu\tau_h\tau_h$ SS Leptons and $e\mu\tau_h\tau_h$ OS Leptons channels and categories. The solid histograms show the stacked background predictions after a background-only fit to the data. The $m_\phi = 200$ GeV and $m_A = 160$ GeV signal scaled to 0.01 pb is also shown by a blue line for illustrative purposes.




Figure 5.12: Expected (dashed line) and observed (solid line and dots) 95% CL upper limits on the product of the cross sections and branching fractions for the decay of both additional Higgs bosons into τ leptons. Different m_A hypotheses are scaled by different orders of magnitudes (written on plot) in order to make mass points distinguishable. The dark green and bright yellow bands indicate the central 68% and 95% intervals for the expected exclusion limit.

5.9.2 Compatibility

Similarly to in Section 4.12.2, the compatibility of the best fit signal cross section multiplied by branching fraction in each decay channel are determined, and these are shown in Figure 5.14 for a mass hypothesis of $m_\phi = m_A = 100$ GeV for this search. For this fit the signal strength in each channel is allowed to go negative, although unphysical, to show the data effects in each channel. The best fit signal strength of the combined fit is between 1 and 2σ below zero. Four categories fit a signal strength slightly below zero, but the combined fit is mostly dominated by the downward fluctuations of the $e\mu\tau_h\tau_h$ channel. The results in each channel are consistent with the zero value within 2σ . These conclusions are consistent across any mass hypotheses, again due to the degenerate nature of the signal shapes in the fitted bins.

5.10 Model Dependent Limits

The 95% CL exclusion contours, for the type X 2HDM alignment scenario in the m_A - $\tan\beta$ phase space, are shown in Figure 5.15 for two m_ϕ scenarios of 100 and 200 GeV. These exclusion limits are “top-down” limits as the cross sections are unchanged in $\tan\beta$ and the branching fractions to τ leptons are enhanced as $\tan\beta$ increases. The cross section and branching ratios are calculated as described in Section 5.1. In all cases, as previous observed for the model independent interpretation, the observed limit lies between the downwards 1 and 2σ expected bands.

The $m_\phi = 100$ GeV scenario has a moderately flat observed $\tan\beta$ limit between 1.2-1.5, across the m_A range. The very slight weakening of the limit at high values of m_A values is because the $A \rightarrow Zh$ decay becomes more kinematically feasible and so the $A \rightarrow \tau\tau$ branching fraction has to compete with this process. The $A \rightarrow Zh$ decay will become dominant if m_A is raised much beyond the 160 GeV threshold set in this analysis. The $m_\phi = 200$ GeV scenario’s observed limit is weaker at lower values of m_A , ranging $\tan\beta$ values of 10 to 1.6 between $m_A = 60$ and 125 GeV. This weakening at lower values of m_A happens as the $H \rightarrow ZA$ decay becomes more kinematically feasible in this region and so competes with $H \rightarrow \tau\tau$ for the branching fraction. This was not present in the $m_\phi = 100$ GeV scenario as the ϕ boson is not heavy enough. The limit then flattens for the remainder of the m_A phase space shown, as the $H \rightarrow ZA$ and $A \rightarrow Zh$ decays only minimally hinder the $\tau\tau$ branching




Figure 5.13: Comparison of the expected 95% CL upper limits on the product of the cross sections and branching fractions for the decay into τ leptons, split by the $\tau\tau\tau\tau$ decay products fit individually.




Figure 5.14: Compatibility plots for the $m_A = 100$ GeV and $m_\phi = 100$ GeV mass scenario in analysis decay channels. In each case the fitted signal strength is decoupled in the bin shown on the plot. The combined best fit value and its 1σ variation are shown by the blue line and band respectively. The black dashed line indicates a signal strength of zero.

fractions of ϕ and A.

Although not shown here, the limits for intermediate ϕ masses see similar trends at low m_A , where the further the ϕ and A masses are separated the weaker the limit, within this mass range. At higher values of m_ϕ than 200 GeV, it becomes difficult to find stable theories across the m_A range. This is because if the m_ϕ and m_A are too separated, the values of λ_i as shown in Equation 1.17, can become non-perturbative [16]. Within the m_A allowed regions for this region, the limits follow the same trend as seen at lower values of m_ϕ .

95% CL limits are also set outside of the alignment scenario. This is done by varying relevant alignment parameter, $\cos(\beta - \alpha)$ or $\sin(\beta - \alpha)$ depending on m_ϕ , with respect to $\tan\beta$ for the individual mass hypothesis. Two of these are shown in Figure 5.16 for the mass scenario of $m_\phi = 200$ GeV and $m_A = 100$ GeV, as well as $m_\phi = 200$ GeV and $m_A = 160$. The observed limit lies in the equivalent place compared to the expected as seen in all limit setting for this analysis. The alignment limit at these mass points are equivalent to the limit set at $\cos(\beta - \alpha) = 0$. The shapes of the limits represent the region where the loss of cross section out of the alignment scenario and the loss of cross section and branching ratios out of the alignment scenario and at lower $\tan\beta$, is too large that the theory cannot be excluded. The shapes are symmetric in $\cos(\beta - \alpha)$ as no sign dependence is measurable from this process. The widening and narrowing of the limit band in $\tan\beta$ is due to shape of the branching fractions of ϕ , as shown in Figure 5.5. The widest constraint from this search on $|\cos(\beta - \alpha)|$ is at approximately 0.5 at a $\tan\beta$ around 20.




Figure 5.15: Expected and observed 95% CL exclusion contours on the m_A - $\tan\beta$ phase space in the type X 2HDM alignment scenario for m_ϕ scenarios of 100 GeV (a) and 200 GeV (b). The exclusion limit only on background expectation is shown as a dashed black line, the dark and bright grey bands show the 68% and 95% intervals of the expected exclusion and the observed exclusion contour is shown by the blue area.




Figure 5.16: Expected and observed 95% CL exclusion contours on the $\cos(\beta - \alpha)$ - $\tan\beta$ phase space in the type X 2HDM alignment scenario with m_ϕ equal to 200 GeV and m_A scenarios of 100 GeV (a) and 160 GeV (b). The exclusion limit only on background expectation is shown as a dashed black line, the dark and bright grey bands show the 68% and 95% intervals of the expected exclusion and the observed exclusion contour is shown by the blue area.

Chapter 6

Conclusion

6.1 Global Interpretations of Results

The analyses presented in Chapters 4 and 5, although motivated by different physics, are complementary to one another in context of the type X 2HDM. The limits on this phase space from the analysis discussed in Chapter 4, are studied using the HIGGSTOOLS-1 framework [127]. HIGGSTOOLS is a combination of the HIGGS-BOUNDS, HIGGSSIGNALS and HIGGSPREDICTIONS frameworks and these are used for the following purpose:

- HIGGSPREDICTIONS is used to determine theory production cross sections and decay rates.
- HIGGSBOUNDS is used to find direct bounds for searches for new particles.
- HIGGSSIGNALS is used to find the bounds from shifts to the observed Higgs boson's properties.

HIGGSBOUNDS and HIGGSSIGNALS contain a database of results from all key measurements from the LHC, LEP and other colliders. The result from Chapter 4 is included in this database. The widths of additional Higgs bosons and branching fractions calculated with 2HDECAY for Chapter 5 are utilised for the scan of the type X 2HDM parameter space. The cross sections used in each analysis scanned over is scaled to that of the model parameters by the NEUTRALEFFECTIVECOUPLINGS function in HIGGSPREDICTIONS. A 95% CL limit is then tested on each point in the parameter space.

In the alignment scenario, the strongest bounds on the type X **2HDM** at all mass points tested come from the analysis described in Chapter 4. The limits determined for the m_ϕ equal to 100 and 200 GeV scenarios are overlayed onto the limits shown in Figure 5.15 and shown in Figure 6.1.

The previous strongest constraints on the type X **2HDM** alignment limit parameter space comes from the analysis described in Chapter 4. The exclusion limits are at low values of $\tan\beta$ only. As $\tan\beta$ increases, the gluon fusion and b associated production modes are suppressed but the branching ratios of the additional Higgs bosons to τ leptons are enhanced. Therefore, the exclusion limit represents a compromise between suppressed cross sections and enhanced branching ratios. The regions where the product is large enough for that parameter point to be excluded, happens at low $\tan\beta$. In the type X **2HDM**, the b associated production mode is negligible due to no enhancement of couplings to b quarks. The mass hypotheses change the limit due to the non-flat nature of Fig. 4.21(a). The limit in Fig. 6.1(b) is flat as the the strongest constraint comes from the $\phi(H)$ boson, and for an additional neutral **CP**-even Higgs boson $\tan\beta \lesssim 10$ is excluded. However, this is not the case for Figure 6.1(a) where m_ϕ is lighter and the sensitivity is not always driven by the $\phi(h)$ boson. In particular, the limit on a 100 GeV resonance, from Fig. 4.21, is weakest due to the large Drell-Yan background and the local excess observed on top of this peak, and so effects from both additional neutral Higgs bosons are present. Together, the exclusion limits from both analyses yield an almost complete coverage of the type X 2HDM alignment scenario within the mass range searched.

Next the affect of the **BSM** searches and precision measurements of the **SM** Higgs boson on the type X **2HDM** model, outside of the alignment scenario is studied. Bounds are again calculated with HIGGSTOOLS, however the constraints now come from **SM** Higgs measurements as well as **BSM** searches such as described in Chapter 4. The bounds determined, overlayed with the results shown in Figure 5.16, are shown in Figure 6.2.

The limits determined from the Chapter 4, although stringent across values of $\cos(\beta - \alpha)$, are weaker than the constraints from the **SM** Higgs boson when moving outside of the alignment limit. They are nonetheless crucial in setting limits, when very close to or on the alignment limit. The constraints from the **SM** Higgs boson, significantly narrow the region allowed outside of the alignment limit and motivates




Figure 6.1: Expected and observed 95% CL exclusion contours on the m_A - $\tan\beta$ phase space in the type X 2HDM alignment scenario for m_ϕ scenarios of 100 GeV (a) and 200 GeV (b). The exclusion limit only on background expectation is shown as a dashed black line, the dark and bright grey bands show the 68% and 95% intervals of the expected exclusion and the observed exclusion contour is shown by the blue area. The limit obtained by HIGGSTOOLS is shown in the red contour.




Figure 6.2: Expected and observed 95% CL exclusion contours on the $\cos(\beta - \alpha)$ - $\tan\beta$ phase space in the type X 2HDM alignment scenario with m_ϕ equal to 200 GeV and m_A scenarios of 100 GeV (a) and 160 GeV (b). The exclusion limit only on background expectation is shown as a dashed black line, the dark and bright grey bands show the 68% and 95% intervals of the expected exclusion and the observed exclusion contour is shown by the blue area. The limit obtained by HIGGSTOOLS is shown in the red contour.

the continued use of alignment scenarios for extended Higgs sector searches. The combination of both **BSM** and **SM** Higgs boson results, prior to the work performed in Chapter 5, leaves only a small strip of the phase space for new physics to exist. However, the entirety of the phase space for the two mass points is excluded by the combination of the searches detailed in Chapters 4 and 5, as well as precision measurement of the **SM** Higgs boson.

6.2 Summary

This thesis has presented two analyses utilising the full run 2 dataset collected by the **CMS** experiment, from the 13 TeV proton proton collisions at the **LHC**, targeting final states enriched in tau leptons. The motivation for the searches presented in Chapter 4 and 5 are **BSM** theories that attempt to resolve the theoretical issue of the hierarchy problem and the experiment tensions of the B anomalies and the g-2 anomaly. The preceding chapters, act to motivate the new physics models that could potentially resolve these tensions, and the apparatus and methods used for the foundation of data taking and reconstruction required for the searches.

Chapter 4 presents a search for two possible areas of new physics in the di- τ final states. The first of these is a search for additional neutral Higgs bosons, motivated by the type II **2HDM** of the **MSSM**, as a consequence of a solution to the hierarchy problem. This targets two production modes: gluon fusion and production is association with a b quark, with the latter dominant to a search for the **MSSM** at higher values of $\tan\beta$. No deviation is observed for the search for b associated production, that targets event categories that require a minimum of one b jet. This makes it very difficult to coincide an **MSSM** benchmark scenario with any gluon fusion signal. The gluon fusion results yielded two small deviations from the **SM** expectation, peaking at 100 GeV and 1.2 TeV with a local (global) statistical significance of 3.1σ (2.7σ) and 2.8σ (2.2σ) respectively. The two excesses are present in the no b tag categories fit and are compatible across all decay channels and categories fit. Good agreement between data and the background hypothesis is observed in the rest of the mass hypotheses. Limits are set on the cross section of both production modes multiplied by the additional scalars decay to τ leptons and both of these vary from $\mathcal{O}(10 \text{ pb})$ at 60 GeV to 0.3 fb at 3.5 TeV. These results are also interpreted as exclusion limits in **MSSM** benchmark scenarios. In the M_h^{125} scenario, values of $m_A < 500 \text{ GeV}$ are excluded, and in the remaining phase space as well as in the $M_{h,EFT}^{125}$ the strongest

constraints on the phase space are set.

The second area of new physics that is searched for in Chapter 4, is a potential solution to the B anomalies, in vector leptoquarks. The signal model searched for is a non-resonant t-channel interaction producing a di- τ final state, where the initial state is dominated by b quarks. The best fit vector leptoquark is heavily constrained by the b tag categories where no deviation from the SM expectation is observed. It therefore can not be used to explain the deviations observed in the no b tag categories. Limits on vector leptoquark phase space are set and constrain the regions allowed for an explanation to the B anomalies.

Chapter 5 details a search for an extended Higgs sector to explain the g-2 anomaly. This can be done with type X 2HDM at high values of $\tan\beta$. A different signal strategy is needed than in Chapter 4, due to the suppressed coupling between the additional neutral Higgs bosons and quarks in this model. The preferred signal model here is the production of two additional neutral Higgs bosons via an off-shell Z boson. At high values of $\tan\beta$, the branching fractions of the additional neutral Higgs bosons are dominated by di- τ pairs. Therefore, the $\tau\tau\tau\tau$ final states are used to reconstruct this signal. No significant deviation is observed from the background estimation. Limits on the cross section multiplied by branching fractions are set and vary from 20 fb at the lowest mass hypothesis, to 1.4 fb at the highest mass hypothesis. The results are interpreted in terms on the type X 2HDM, and it is found that the constraints from the SM Higgs boson measurements limit the phase space to be very close to the alignment limit. The alignment limit exclusions are found to exclude upwards of $\tan\beta$ approximately equal to 1.5, unless the $H \rightarrow ZA$ becomes kinematically feasible and the limit is weakened. This an exclusion way beyond a possible explanation to the g-2 anomaly. The results from Chapter 4 are complimentary to Chapter 5 in the type X 2HDM alignment limit, as they exclude downwards of $\tan\beta$ approximately equal to 10, and so together exclude the vast majority of the phase space in mass ranges searched.

Bibliography

- [1] CMS Collaboration, *Search for additional neutral MSSM Higgs bosons in the $\tau\tau$ final state in proton-proton collisions at $\sqrt{s} = 13$ TeV*, *JHEP* **09** (2018) 007 [[1803.06553](#)].
- [2] CMS Collaboration, *Searches for additional Higgs bosons and vector leptoquarks in $\tau\tau$ final states in proton-proton collisions at $\sqrt{s} = 13$ TeV*, *CMS Physics Analysis Summary CMS-PAS-HIG-21-001* (2022) .
- [3] D. Galbraith and C. Burgard, “Example: Standard model of physics.” <https://texample.net/tikz/examples/model-physics/>. [Accessed: 25/05/23].
- [4] H. Bartosik and G. Rumolo, *Performance of the LHC injector chain after the upgrade and potential development*, [2203.09202](#).
- [5] CMS Collaboration, “CMS Luminosity public results.” <https://twiki.cern.ch/twiki/bin/view/CMSPublic/LumiPublicResults>. [Accessed: 25/05/23].
- [6] CMS Collaboration, *The CMS Experiment at the CERN LHC*, *JINST* **3** (2008) S08004.
- [7] CMS Collaboration, *CMS Technical Design Report for the Level-1 Trigger Upgrade*, *CMS Technical Design Report CERN-LHCC-2013-011*, *CMS-TDR-12*, *CMS-TDR-012* (2013) .
- [8] CMS Collaboration, *Performance of Electron Reconstruction and Selection with the CMS Detector in Proton-Proton Collisions at $\sqrt{s} = 8$ TeV*, *JINST* **10** (2015) P06005 [[1502.02701](#)].

- [9] CMS Collaboration, “CMS Public Twiki.” <https://twiki.cern.ch/twiki/bin/view/CMSPublic/BTV13TeV2017DeepJet>. [Accessed: 25/05/23].
- [10] CMS Collaboration, *Performance of reconstruction and identification of τ leptons decaying to hadrons and ν_τ in pp collisions at $\sqrt{s} = 13$ TeV*, *JINST* **13** (2018) P10005 [[1809.02816](#)].
- [11] CMS Collaboration, *Identification of hadronic tau lepton decays using a deep neural network*, *JINST* **17** (2022) P07023 [[2201.08458](#)].
- [12] CMS Collaboration, *Performance of reconstruction and identification of τ leptons decaying to hadrons and ν_τ in pp collisions at $\sqrt{s} = 13$ TeV*, *JINST* **13** (2018) P10005 [[1809.02816](#)].
- [13] CMS Collaboration, *An embedding technique to determine $\tau\tau$ backgrounds in proton-proton collision data*, *JINST* **14** (2019) P06032 [[1903.01216](#)].
- [14] ATLAS Collaboration, *Search for heavy Higgs bosons decaying into two tau leptons with the ATLAS detector using pp collisions at $\sqrt{s} = 13$ TeV*, *Phys. Rev. Lett.* **125** (2020) 051801 [[2002.12223](#)].
- [15] C. Cornella, D. A. Faroughy, J. Fuentes-Martin, G. Isidori and M. Neubert, *Reading the footprints of the B-meson flavor anomalies*, *JHEP* **08** (2021) 050 [[2103.16558](#)].
- [16] A. Jueid, J. Kim, S. Lee and J. Song, *Type-X two-Higgs-doublet model in light of the muon $g-2$: Confronting Higgs boson and collider data*, *Phys. Rev. D* **104** (2021) 095008 [[2104.10175](#)].
- [17] R. L. Workman et al., *Review of Particle Physics*, *PTEP* **2022** (2022) 083C01.
- [18] ATLAS Collaboration, *Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC*, *Phys. Lett. B* **716** (2012) 1 [[1207.7214](#)].
- [19] CMS Collaboration, *Observation of a New Boson at a Mass of 125 GeV with the CMS Experiment at the LHC*, *Phys. Lett. B* **716** (2012) 30 [[1207.7235](#)].
- [20] LHCb Collaboration, *Test of lepton universality in beauty-quark decays*, *Nature Phys.* **18** (2022) 277 [[2103.11769](#)].

- [21] BaBar Collaboration, *Evidence for an excess of $B \rightarrow D^{(*)}\tau\nu$ decays*, *J. Phys. Conf. Ser.* **455** (2013) 012039.
- [22] BaBar Collaboration, *Measurement of an excess of $\bar{B} \rightarrow D^{(*)}\tau^-\bar{\nu}_\tau$ decays and implications for charged higgs bosons*, *Phys. Rev. D* **88** (2013) 072012 [[1303.0571](#)].
- [23] Belle Collaboration, *Measurement of the branching ratio of $\bar{B} \rightarrow D^{(*)}\tau^-\bar{\nu}_\tau$ relative to $\bar{B} \rightarrow D^{(*)}\ell^-\bar{\nu}_\ell$ decays with hadronic tagging at Belle*, *Phys. Rev. D* **92** (2015) 072014 [[1507.03233](#)].
- [24] LHCb Collaboration, *Measurement of the ratio of branching fractions $\mathcal{B}(\bar{B}^0 \rightarrow D^{*+}\tau^-\bar{\nu}_\tau)/\mathcal{B}(\bar{B}^0 \rightarrow D^{*+}\mu^-\bar{\nu}_\mu)$* , *Phys. Rev. Lett.* **115** (2015) 111803 [[1506.08614](#)].
- [25] Belle Collaboration, *Measurement of the τ lepton polarization and $R(D^*)$ in the decay $\bar{B} \rightarrow D^*\tau^-\bar{\nu}_\tau$* , *Phys. Rev. Lett.* **118** (2017) 211801 [[1612.00529](#)].
- [26] LHCb Collaboration, *Test of Lepton Flavor Universality by the measurement of the $B^0 \rightarrow D^{*-}\tau^+\nu_\tau$ branching fraction using three-prong τ decays*, *Phys. Rev. D* **97** (2018) 072013 [[1711.02505](#)].
- [27] LHCb Collaboration, *Measurement of the ratio of the $B^0 \rightarrow D^{*-}\tau^+\nu_\tau$ and $B^0 \rightarrow D^{*-}\mu^+\nu_\mu$ branching fractions using three-prong τ -lepton decays*, *Phys. Rev. Lett.* **120** (2018) 171802 [[1708.08856](#)].
- [28] Muon g-2 Collaboration, *Measurement of the Positive Muon Anomalous Magnetic Moment to 0.46 ppm*, *Phys. Rev. Lett.* **126** (2021) 141801 [[2104.03281](#)].
- [29] *Measurement of lepton universality parameters in $B^+ \rightarrow K^+\ell^+\ell^-$ and $B^0 \rightarrow K^{*0}\ell^+\ell^-$ decays*, [**2212.09153**](#).
- [30] D. J. Gross and F. Wilczek, *Ultraviolet Behavior of Nonabelian Gauge Theories*, *Phys. Rev. Lett.* **30** (1973) 1343.
- [31] H. D. Politzer, *Reliable Perturbative Results for Strong Interactions?*, *Phys. Rev. Lett.* **30** (1973) 1346.
- [32] S. L. Glashow, *Partial Symmetries of Weak Interactions*, *Nucl. Phys.* **22** (1961) 579.

- [33] S. Weinberg, *A Model of Leptons*, *Phys. Rev. Lett.* **19** (1967) 1264.
- [34] A. Salam, *Weak and Electromagnetic Interactions*, *Conf. Proc. C* **680519** (1968) 367.
- [35] G. Arnison et al., *Experimental Observation of Isolated Large Transverse Energy Electrons with Associated Missing Energy at $\sqrt{s} = 540$ GeV*, *Phys. Lett. B* **122** (1983) 103.
- [36] M. Banner et al., *Observation of Single Isolated Electrons of High Transverse Momentum in Events with Missing Transverse Energy at the CERN anti- p Collider*, *Phys. Lett. B* **122** (1983) 476.
- [37] F. Englert and R. Brout, *Broken symmetry and the mass of gauge vector mesons*, *Phys. Rev. Lett.* (1964) 321.
- [38] P. W. Higgs, *Broken symmetries, massless particles and gauge fields*, *Phys. Lett.* (1964) 132.
- [39] P. W. Higgs, *Broken symmetries and the masses of gauge bosons*, *Phys. Rev. Lett.* (1964) 508.
- [40] P. W. Higgs, *Spontaneous symmetry breakdown without massless bosons*, *Phys. Rev.* (1966) 1156.
- [41] G. S. Guralnik, C. R. Hagen and T. W. B. Kibble, *Global conservation laws and massless particles*, *Phys. Rev. Lett.* (1964) 585.
- [42] T. W. B. Kibble, *Symmetry breaking in non-Abelian gauge theories*, *Phys. Rev.* (1967) 1554.
- [43] J. Goldstone, *Field Theories with Superconductor Solutions*, *Nuovo Cim.* **19** (1961) 154.
- [44] CMS Collaboration, *A measurement of the Higgs boson mass in the diphoton decay channel*, *Phys. Lett. B* **805** (2020) 135425 [[2002.06398](#)].
- [45] S. P. Martin, *A Supersymmetry Primer*, *Advanced Series on Directions in High Energy Physics* (1998) 1–98.
- [46] B. Diaz, M. Schmaltz and Y.-M. Zhong, *The leptoquark hunter's guide: Pair production*, *JHEP* **10** (2017) 097 [[1706.05033](#)].

- [47] M. Schmaltz and Y.-M. Zhong, *The leptoquark hunter's guide: Large coupling*, *JHEP* **01** (2019) 132 [[1810.10017](#)].
- [48] Muon g-2 Collaboration, *Final Report of the Muon E821 Anomalous Magnetic Moment Measurement at BNL*, *Phys. Rev. D* **73** (2006) 072003 [[hep-ex/0602035](#)].
- [49] V. Ilisie, *New Barr-Zee contributions to $(g - 2)_\mu$ in two-Higgs-doublet models*, *JHEP* **04** (2015) 077 [[1502.04199](#)].
- [50] S. M. Barr and A. Zee, *Electric Dipole Moment of the Electron and of the Neutron*, *Phys. Rev. Lett.* **65** (1990) 21.
- [51] L. Evans and P. Bryant (Eds.), *LHC Machine*, *JINST* **3** (2008) S08001.
- [52] LEP Injector Study Group, *LEP Design Report: Vol.2. The LEP Main Ring, Technical Design Report* **CERN-LEP-84-01** (1984) .
- [53] ALICE Collaboration, *The ALICE experiment at the CERN LHC*, *JINST* **3** (2008) S08002.
- [54] ATLAS Collaboration, *The ATLAS Experiment at the CERN Large Hadron Collider*, *JINST* **3** (2008) S08003.
- [55] LHCb Collaboration, *The LHCb Detector at the LHC*, *JINST* **3** (2008) S08005.
- [56] CMS Collaboration, *Operation and performance of the CMS tracker*, *PoS TIPP2014* (2014) 347.
- [57] CMS Collaboration, *CMS Technical Design Report for the Pixel Detector Upgrade*, *Technical Design Report* **CERN-LHCC-2012-016, CMS-TDR-011** (2012) .
- [58] CMS Collaboration, *Energy Calibration and Resolution of the CMS Electromagnetic Calorimeter in pp Collisions at $\sqrt{s} = 7$ TeV*, *JINST* **8** (2013) P09009 [[1306.2016](#)].
- [59] CMS Collaboration, *Description and performance of track and primary-vertex reconstruction with the CMS tracker*, *JINST* **9** (2014) P10009 [[1405.6569](#)].

- [60] R. Fruhwirth, *Application of Kalman filtering to track and vertex fitting*, *Nucl. Instrum. Meth. A* **262** (1987) 444.
- [61] CMS Collaboration, *Measurement of Tracking Efficiency*, *CMS Physics Analysis Summary CMS-PAS-TRK-10-002* (2010) .
- [62] K. Rose, *Deterministic annealing for clustering, compression, classification, regression, and related optimization problems*, *IEEE Proc. 86* (1998) 2210.
- [63] R. Fruhwirth, W. Waltenberger and P. Vanlaer, *Adaptive vertex fitting*, *J. Phys. G* **34** (2007) N343.
- [64] CMS Collaboration, *Particle-flow reconstruction and global event description with the CMS detector*, *JINST* **12** (2017) P10003 [[1706.04965](#)].
- [65] CMS Collaboration, *Commissioning of the Particle-flow Event Reconstruction with the first LHC collisions recorded in the CMS detector*, *CMS Physics Analysis Summary CMS-PAS-PFT-10-001*, *CMS-PAS-PFT-10-001* (2010) .
- [66] CMS Collaboration, *Commissioning of the Particle-Flow reconstruction in Minimum-Bias and Jet Events from pp Collisions at 7 TeV*, *CMS Physics Analysis Summary CMS-PAS-PFT-10-002*, *CMS-PAS-PFT-10-002* (2010) .
- [67] T. Speer, W. Adam, R. Fruhwirth, A. Strandlie, T. Todorov and M. Winkler, *Track reconstruction in the CMS tracker*, *Nucl. Instrum. Meth. A* **559** (2006) 143.
- [68] CMS Collaboration, *Performance of CMS Muon Reconstruction in pp Collision Events at $\sqrt{s} = 7$ TeV*, *JINST* **7** (2012) P10002 [[1206.4071](#)].
- [69] CMS Collaboration, *Performance of the CMS muon detector and muon reconstruction with proton-proton collisions at $\sqrt{s} = 13$ TeV*, *JINST* **13** (2018) P06015 [[1804.04528](#)].
- [70] CMS Collaboration, *Performance of CMS Muon Reconstruction in Cosmic-Ray Events*, *JINST* **5** (2010) T03022 [[0911.4994](#)].
- [71] G. P. Salam, *Towards Jetography*, *Eur. Phys. J. C* **67** (2010) 637 [[0906.1833](#)].

- [72] M. Cacciari, G. P. Salam and G. Soyez, *The anti- k_t jet clustering algorithm*, *JHEP* **04** (2008) 063 [[0802.1189](#)].
- [73] M. Cacciari, G. P. Salam and G. Soyez, *FastJet user manual*, *Eur. Phys. J. C* **72** (2012) 1896 [[1111.6097](#)].
- [74] CMS Collaboration, *Jet algorithms performance in 13 TeV data*, *CMS Physics Analysis Summary CMS-PAS-JME-16-003* (2017) .
- [75] CMS Collaboration, *Pileup Jet Identification*, *CMS Physics Analysis Summary CMS-PAS-JME-13-005* (2013) .
- [76] CMS Collaboration, *Identification of heavy-flavour jets with the CMS detector in pp collisions at 13 TeV*, *JINST* **13** (2018) P05011 [[1712.07158](#)].
- [77] E. Bols, J. Kieseler, M. Verzetti, M. Stoye and A. Stakia, *Jet flavour classification using DeepJet*, *JINST* **15** (2020) P12012 [[2008.10519](#)].
- [78] CMS Collaboration, *Performance of missing energy reconstruction in 13 TeV pp collision data using the CMS detector*, *CMS Physics Analysis Summary CMS-PAS-JME-16-004* (2016) .
- [79] CMS Collaboration, *Pileup mitigation at CMS in 13 TeV data*, *JINST* **15** (2020) P09018 [[2003.00503](#)].
- [80] E. Bagnaschi, H. Bahl, E. Fuchs, T. Hahn, S. Heinemeyer, S. Liebler et al., *MSSM Higgs boson searches at the LHC: Benchmark scenarios for Run 2 and beyond*, *Eur. Phys. J. C* **79** (2019) 617 [[1808.07542](#)].
- [81] H. Bahl, P. Bechtle, S. Heinemeyer, S. Liebler, T. Stefaniak and G. Weiglein, *HL-LHC and ILC sensitivities in the hunt for heavy Higgs bosons*, *Eur. Phys. J. C* **80** (2020) 916 [[2005.14536](#)].
- [82] H. Bahl, S. Liebler and T. Stefaniak, *MSSM Higgs benchmark scenarios for Run 2 and beyond: the low $\tan \beta$ region*, *Eur. Phys. J. C* **79** (2019) 279 [[1901.05933](#)].
- [83] E. A. Bagnaschi, S. Heinemeyer, S. Liebler, P. Slavich and M. Spira, *Benchmark scenarios for MSSM Higgs boson searches at the LHC*, *LHC Higgs Cross Section Working Group Public Note LHCHWG-2021-001* (2021) .

- [84] CMS Collaboration, *Search for beyond the standard model Higgs bosons decaying into a $b\bar{b}$ pair in pp collisions at $\sqrt{s} = 13$ TeV*, *JHEP* **08** (2018) 113 [[1805.12191](#)].
- [85] P. Nason, *A new method for combining NLO QCD with shower Monte Carlo algorithms*, *JHEP* **11** (2004) 040 [[hep-ph/0409146](#)].
- [86] S. Frixione, P. Nason and C. Oleari, *Matching NLO QCD computations with parton shower simulations: the POWHEG method*, *JHEP* **11** (2007) 070 [[0709.2092](#)].
- [87] S. Alioli, P. Nason, C. Oleari and E. Re, *A general framework for implementing NLO calculations in shower Monte Carlo programs: the POWHEG BOX*, *JHEP* **06** (2010) 043 [[1002.2581](#)].
- [88] T. Ježo and P. Nason, *On the treatment of resonances in next-to-leading order calculations matched to a parton shower*, *JHEP* **12** (2015) 065 [[1509.09071](#)].
- [89] R. D. Ball et al., *Parton distributions for the LHC Run II*, *JHEP* **04** (2015) 040 [[1410.8849](#)].
- [90] R. D. Ball et al., *Parton distributions from high-precision collider data*, *Eur. Phys. J. C* **77** (2017) 663 [[1706.00428](#)].
- [91] CMS Collaboration, *Extraction and validation of a new set of CMS PYTHIA8 tunes from underlying-event measurements*, *Eur. Phys. J. C* **80** (2020) 4 [[1903.12179](#)].
- [92] T. Sjöstrand, S. Ask, J. R. Christiansen, R. Corke, N. Desai, P. Ilten et al., *An introduction to PYTHIA 8.2*, *Comput. Phys. Commun.* **191** (2015) 159 [[1410.3012](#)].
- [93] S. Agostinelli et al., GEANT4—a simulation toolkit, *Nucl. Instrum. Meth. A* **506** (2003) 250.
- [94] CMS Collaboration, *The search for a third-generation leptoquark coupling to a τ lepton and a b quark through single, pair and nonresonant production at $\sqrt{s} = 13$ TeV*, *CMS Physics Analysis Summary CMS-PAS-EXO-19-016* (2022) .

- [95] J. Alwall, M. Herquet, F. Maltoni, O. Mattelaer and T. Stelzer, *MadGraph 5: Going beyond*, *JHEP* **06** (2011) 128 [[1106.0522](#)].
- [96] R. Frederix and S. Frixione, *Merging meets matching in MC@NLO*, *JHEP* **12** (2012) 061 [[1209.6215](#)].
- [97] L. Bianchini, J. Conway, E. K. Friis and C. Veelken, *Reconstruction of the Higgs mass in $H \rightarrow \tau\tau$ events by dynamical likelihood techniques*, *J. Phys. Conf. Ser.* **513** (2014) 022035.
- [98] CMS Collaboration, *Measurements of Higgs boson production in the decay channel with a pair of τ leptons in proton-proton collisions at $\sqrt{s} = 13$ TeV*, *CMS Physics Analysis Summary CMS-HIG-19-010, CERN-EP-2022-027* (2022) [[2204.12957](#)].
- [99] CMS Collaboration, *Search for additional neutral MSSM higgs bosons in the $\tau\tau$ final state in proton-proton collisions at $\sqrt{s} = 13$ TeV*, *JHEP* **09** (2018) 007 [[1803.06553](#)].
- [100] CMS Collaboration, *Measurement of the $Z/\gamma^* \rightarrow \tau\tau$ cross section in pp collisions at $\sqrt{s} = 13$ TeV and validation of τ lepton analysis techniques*, *Eur. Phys. J. C* **78** (2018) 708 [[1801.03535](#)].
- [101] J. Alwall, S. Höche, F. Krauss, N. Lavesson, L. Lönnblad, F. Maltoni et al., *Comparative study of various algorithms for the merging of parton showers and matrix elements in hadronic collisions*, *Eur. Phys. J. C* **53** (2008) 473 [[0706.2569](#)].
- [102] S. Alioli, S.-O. Moch and P. Uwer, *Hadronic top-quark pair-production with one jet and parton showering*, *JHEP* **01** (2012) 137 [[1110.5251](#)].
- [103] R. Frederix, E. Re and P. Torrielli, *Single-top t-channel hadroproduction in the four-flavour scheme with POWHEG and aMC@NLO*, *JHEP* **09** (2012) 130 [[1207.5391](#)].
- [104] E. Re, *Single-top Wt-channel production matched with parton showers using the POWHEG method*, *Eur. Phys. J. C* **71** (2011) 1547 [[1009.2450](#)].
- [105] K. Melnikov and F. Petriello, *Electroweak gauge boson production at hadron colliders through $\mathcal{O}(\alpha_s^2)$* , *Phys. Rev. D* **74** (2006) 114017 [[hep-ph/0609070](#)].

- [106] M. Czakon and A. Mitov, *Top++: A program for the calculation of the top-pair cross-section at hadron colliders*, *Comput. Phys. Commun.* **185** (2014) 2930 [[1112.5675](#)].
- [107] N. Kidonakis, *Top quark production*, in *Helmholtz International Summer School on Physics of Heavy Quarks and Hadrons*, p. 139, 2014, [1311.0283](#), DOI.
- [108] J. M. Campbell, R. K. Ellis and C. Williams, *Vector boson pair production at the LHC*, *JHEP* **07** (2011) 018 [[1105.0020](#)].
- [109] T. Gehrmann, M. Grazzini, S. Kallweit, P. Maierhöfer, A. von Manteuffel, S. Pozzorini et al., *W^+W^- production at hadron colliders in next to next to leading order QCD*, *Phys. Rev. Lett.* **113** (2014) 212001 [[1408.5243](#)].
- [110] CMS Collaboration, *Measurements of Inclusive W and Z Cross Sections in pp Collisions at $\sqrt{s} = 7$ TeV*, *JHEP* **01** (2011) 080 [[1012.2466](#)].
- [111] CMS Collaboration, *Performance of the CMS Level-1 trigger in proton-proton collisions at $\sqrt{s} = 13$ TeV*, *JINST* **15** (2020) P10017 [[2006.10165](#)].
- [112] CMS Collaboration, *Measurement of the differential cross section for top quark pair production in pp collisions at $\sqrt{s} = 8$ TeV*, *Eur. Phys. J. C* **75** (2015) 542 [[1505.04480](#)].
- [113] R. J. Barlow and C. Beeston, *Fitting using finite Monte Carlo samples*, *Comput. Phys. Commun.* **77** (1993) 219.
- [114] J. S. Conway, *Incorporating Nuisance Parameters in Likelihoods for Multisource Spectra*, in *PHYSTAT 2011*, pp. 115–120, 2011, [1103.0354](#), DOI.
- [115] CMS Collaboration, *Performance of missing transverse momentum reconstruction in proton-proton collisions at $\sqrt{s} = 13$ TeV using the CMS detector*, *JINST* **14** (2019) P07004 [[1903.06078](#)].
- [116] T. Junk, *Confidence level computation for combining searches with small statistics*, *Nucl. Instrum. Meth. A* **434** (1999) 435 [[hep-ex/9902006](#)].
- [117] A. L. Read, *Presentation of search results: The CL_s technique*, *J. Phys. G* **28** (2002) 2693.

- [118] G. Cowan, K. Cranmer, E. Gross and O. Vitells, *Asymptotic formulae for likelihood-based tests of new physics*, *Eur. Phys. J. C* **71** (2011) 1554 [[1007.1727](#)].
- [119] LHC Higgs Cross Section Working Group Collaboration, *Handbook of LHC Higgs Cross Sections: 3. Higgs Properties*, [1307.1347](#).
- [120] LHC Higgs Cross Section Working Group Collaboration, *Handbook of LHC Higgs Cross Sections: 4. Deciphering the Nature of the Higgs Sector*, [1610.07922](#).
- [121] A. Denner, S. Heinemeyer, I. Puljak, D. Rebuzzi and M. Spira, *Standard model Higgs-boson branching ratios with uncertainties*, *Eur. Phys. J. C* **71** (2011) 1753 [[1107.5909](#)].
- [122] M. Krause, M. Mühlleitner and M. Spira, *2HDECAY —A program for the calculation of electroweak one-loop corrections to Higgs decays in the Two-Higgs-Doublet Model including state-of-the-art QCD corrections*, *Comput. Phys. Commun.* **246** (2020) 106852 [[1810.00768](#)].
- [123] S. Catani and M. Grazzini, *Next-to-next-to-leading-order subtraction formalism in hadron collisions and its application to higgs-boson production at the large hadron collider*, *Phys. Rev. Lett.* **98** (2007) 222002.
- [124] A. Rogozhnikov, *Reweighting with Boosted Decision Trees*, *J. Phys. Conf. Ser.* **762** (2016) 012036 [[1608.05806](#)].
- [125] CMS Collaboration, *Analysis of the CP structure of the Yukawa coupling between the Higgs boson and τ leptons in proton-proton collisions at $\sqrt{s} = 13$ TeV*, *JHEP* **06** (2022) 012 [[2110.04836](#)].
- [126] F. J. Massey, *The kolmogorov-smirnov test for goodness of fit*, *Journal of the American Statistical Association* **46** (1951) 68.
- [127] H. Bahl, T. Biekötter, S. Heinemeyer, C. Li, S. Paasch, G. Weiglein et al., *HiggsTools: BSM scalar phenomenology with new versions of HiggsBounds and HiggsSignals*, [2210.09332](#).

List of acronyms

2HDM two Higgs doublet model

BDT boosted decision tree

BEH Brout-Englet-Higgs

BSM Beyond Standard Model

CERN the European Organization for Nuclear Research

CHS charged hadron subtraction

CL confidence level

CMS Compact Muon Solenoid

CP charge-parity

CSC cathode strip chambers

CTF Combinatorial Track Finder

DM decay mode

DNN deep neural network

DT drift tube

EB ECAL barrel

ECAL electromagnetic calorimeter

EE ECAL endcap

FS flavour scheme

GSF Gaussian sum filter

HB hadron barrel

HCAL hadronic calorimeter

HE hadron endcap

HF hadron forward

HLT high-level trigger

HO hadron outer

HPS hadron-plus-strips

KF Kalman filter

L1 Level-1

LEP Large Electron-Positron

LHC Large Hadron Collider

LO leading-order

LS1 Long Shutdown 1

LS2 Long Shutdown 2

MC monte carlo

MET missing transverse energy

ML machine learning

MSSM Minimal Supersymmetric Standard Model

MVA multivariate analysis

NLO next-to-leading-order

NN neural network

NNLO next-to-next-to-leading-order

PDF probability distribution function

PF particle flow

PS Proton Synchrotron

PSB Proton Synchrotron Booster

PU pileup

PV primary vertex

QCD Quantum Chromodynamics

RF radio frequency

RPC resistive plate chambers

SC supercluster

SM Standard Model

SPS Super Proton Synchrotron

SUSY supersymmetry

SV secondary vertex

TEC tracker endcaps

TIB tracker inner barrel

TID tracker inner disks

TOB tracker outer barrel

VEV vacuum expectation value

WLCG Worldwide LHC Computing Grid

WP working points