KUN Counseling And Training
Well Of Awareness For Thirsty People

**Patient' Name:**_____ **Reg.No:**_____

## Brief History Of Patient

# Pregnancy Guide Book

**Dr. Sameera Bilal**

**Senior Gynacologist**

**Share my experience:**

If you are trying to have a baby or are just thinking about it, it is not too early to prepare for a safe pregnancy and a healthy baby. You should see your health care provider for preconception care

Preconception care is a care you receive before you get pregnant. By taking action on health issues before pregnancy, you can prevent many future problems for yourself and your baby. Your health care provider can help you figure out the steps you need to take before getting pregnant, such as

1) Finding and taking care of any problems that might affect you and your baby later, like diabetes , high blood pressure , thyroid problem or sexually transmitted diseases.
2) 500 micro gram folic acid daily for all women and 5 mg folic acid for those who are at high risk such as previous h/o Neural tube defects , diabetics , BMI>30 , taking antiepileptics drugs .it is recommended to start folic acid at least 01 moth before conception and should be continued up to first 12 weeks of pregnancy .
3) Getting up to date on vaccines, such as Hepatitis B, influenza, tetanus. For Rubella vaccination one must have to do contraception for one month.
4) Avoiding smoking, drinking alcohol, or taking drugs.
5) Trying to reach a healthy weight is necessary before planning pregnancy. People who are overweight or obese have a higher risk for many serious conditions, including complications during pregnancy, heart disease, type 2 diabetes, and certain cancers (endometrial, breast, and colon). The key to achieving and maintaining a healthy weight isn't about short-term dietary changes. It's about a lifestyle that includes healthy eating and regular physical activity [1]
6) Avoid harmful chemicals, environmental contaminants, and other toxic substances such as synthetic chemicals, metals, fertilizer, bug spray, and handling of cat or rodent faeces.[2]
7) If you are taking any drugs such as antihypertensives, anti-epileptics or anti-thyroids and many more, it is essential to discuss with your health care to check if they are teratogenic and harmful for your upcoming pregnancy, and how to replace them with safer options.
8) Avoid vit a supplements or retinol topical applications as it's teratogenic.
9) You should take enough calcium, vit D and iron supplements. Also, it is important to correct anemia…

**Comments on Manual:**

I must say that this manual is comprehensive yet up to date and provides all the necessary information for those who are planning to face this beautiful as well as challenging phase of life. In my view, it's a complete guide for new mothers. This manual would help many couples to get through this phase of parenthood successfully.

**Dr. Shazia Qadir**

**DPT(D) Doctor of Physiotherapy**

**Assalaam o alaikum ,**

"PREGNANCY ", in my opinion is one of the most difficult yet beautiful phase of a woman 's life. Unfortunately in our society we still hesitate to discuss pregnancy related issues (pre and post pregnancy both),which end up in many problems & complications faced by women. It's a great effort of Kun consultancy ----- Sir Sagheer to design such a book,which is going to help all the women whether pregnant or non pregnant. It is a concise yet detailed book about pregnancy. Although it's a fact that every pregnancy is different from other and symptoms as well as problems & complications vary from person to person but there are many common things which are same in all pregnancies which are covered in this book. It almost covers all the sign & symptoms, stages of pregnancy as well as labour, diet, exercises, myths, problems & solutions related to pregnancy.

In sha Allah this book will be a source of great help for not only women but also for men.

I feel very privileged for been asked to comment on this book.

May Allah bless you with all the success always,ameen.

Jazakallah.

**Dr(H). Sara Zahid**

**Vice President Kun Consultancy**

Due to massive hormonal changes in pregnancy women usually don't consider how to manage their health with this blessing.

So they suffer from some major complications like miscarriage, high BP issue, preeclampsia, Gestational Diabetes, Anemia, infection, Pre term labour ,Breech position.

Here it is important to Educate yourself and stay connected with your consultant properly over the period of nine month.

Otherwise 80 percent of females suffering from first trimester miscarriage or reoccurring miscarriage s as per Clinical facts .

Now I feel this Manual under the guidance of consultant plays an Excellent role to educate lovely women.

You just need to hang in front of your bed and understand properly this Manual month by month under the shadow of Consultant you will not only enjoy your pregnancy infact you will teach other females how to spend pregnancy precious time .

Remember Manual and Mentor will change your pains in pleasure .

May Allah bless you all with this precious pregnancy gift Ameen.

**Dr. Zona Irfan**

**MBBS (DOW)**

**FCPS 2 trainee**

Undoubtedly an exceptional book by Our Mentor ..Father of Time Topology ..It not only covers the whole topic but also each detail is highlighted and elaborated properly.. Eloquent explanation we haven't read until now also breaking the myths related to this delicate period which every female has to go at some point in her life..

This book is indeed a much needed discussion that we usually don't see in our asian setups where talking about pregnancy and sex is considered a taboo..

The most beautiful part of this book is an entirely different concept of pregnancy management that not only includes the application of reflexology but also gives a pertinent view on diet and exercise.

On the top of it , all the myths related to pregnancy are also answered with logical explanations ..of course this is  the uniqueness of our Mentor.

I will recommend it to all the married females..a must read for them!


FEW DAYS
AFTER CONCEPTION
SYMPTOMS OF EARLY
PREGNANCY
TWO WEEKS
AFTER CONCEPTION
SENSITIVE BREASTS
DROWSINESS AND FATIGUE
IRRITABILITY
HEADACHE
INSOMNIA
NAUSEA, VOMITING AND DIZZINESS
CHANGE IN TASTE PREFERENCES
INCREASED BODY TEMPERATURE
DELAY IN MENSTRUATION
MINOR BLEEDING
BLOATED CONSTIPATION HEARTBURN
FREQUENCY OF URINATION
BREAST SWELLING
DARKENING OF THE NIPPLE OF THE NIPPLES
EXACERBATION OF CHRONIC DISEASES
INCREASED SENSE OF SMELL

1-_____ 5-_____

2-_____ 6-_____

3-_____ 7-_____

4-_____ 8-_____

9-_____

## ADVICE:

**1-Diet:**

**2-Exercise:**

**3-Massage:**

**4: NLP:**

**5: Pressure Points:**

## ADVICE:

**1-Diet:**

**2-Exercise:**

**3-Massage:**

**4: NLP:**

**5: Pressure Points:**




## ADVICE:

**1-Diet:**

**2-Exercise:**

**3-Massage:**

**4: NLP:**

**5: Pressure Points:**




## ADVICE:

**1-Diet:**

**2-Exercise:**

**3-Massage:**

**4: NLP:**

**5: Pressure Points:**

## ADVICE:

**1-Diet:**

**2-Exercise:**

**3-Massage:**

**4: NLP:**

**5: Pressure Points:**

## ADVICE:

**1-Diet:**

**2-Exercise:**

**3-Massage:**

**4: NLP:**

**5: Pressure Points:**




## ADVICE:

**1-Diet:**

**2-Exercise:**

**3-Massage:**

**4: NLP:**

**5: Pressure Points:**




## ADVICE:

**1-Diet:**

**2-Exercise:**

**3-Massage:**

**4: NLP:**

**5: Pressure Points:**




## ADVICE:

**1-Diet:**

**2-Exercise:**

**3-Massage:**

**4: NLP:**

**5: Pressure Points:**

# PREPARATION FOR NORMAL DELIVERY:

Prof DR MOHD AZHAR – "NOMAL LABOUR & DELIVERY"

## NORMAL LABOUR AND DELIVERY

### STAGES OF LABOUR

Labour can be divided into three stages, which are unequal in length.

| FIRST STAGE | SECOND STAGE | THIRD STAGE |
|---|---|---|
| It begins with the onset of true labour contractions and ends when the cervix is fully dilated (10 cm). | The second stage of labour begins with complete dilatation of the cervix and ends with the birth of the baby. | The third stage is that of separation and expulsion of placenta and membranes and also involves the control of bleeding. |
| Cervical effacement and dilatation occur in the first stage | The duration is about 1 to 1½ hours in nulliparas and about 30 to 45 minutes in parous women. | It begins after the birth of the baby and ends with the expulsion of the placenta and membranes. |
| First stage of labour consists of two phases:- **latent** and **active**. | | This is the shortest stage, lasting up to 30 minutes, with an average length of 5 to 10 minutes. There is no difference in duration for nulliparous and parous. |
| The first stage of labour is the longest for both nulliparous and parous women. | | |



**QUICK GUIDE TO SUCCESSFUL APM**

**→ Start the practice at 34 weeks of gestation.**

**→ Use fingers or thumbs inserted about an inch on each side into the vaginal canal to massage from the side of the vaginal opening down to the back of the opening. Use a repeated slow motion to glide from the sides of the vagina to the back portion (near the rectum), and then up again to the sides. The goal is to pull your fingers or thumbs apart from each other as you glide around the sides and back area of the tissues, stretching and widening this U-shaped portion at the back of the birth canal.**

**→ Use non-irritating products to increase slipperiness and decrease trauma during the APM (see below).**

**→ Massage daily for 4 to 5 minutes each day.**

**→ Feel a slight (but not overwhelming) burning sensation the first week or two especially as the perineal tissue is stretched backwards.**

**→ Have confidence that if you feel this burn or stretching, you ARE doing it right**

## How to do perineal massage

- Sit with your legs apart and your back supported. ...
- Put some massage oil on your fingers. ...
- Place a thumb or finger about 5 cm (2 in.) ...
- Still pressing and stretching outward, sweep down to 6 o'clock and over to 9 o'clock.
- Repeat for a total of 4 or 5 minutes



**Stages of labor**

**There are 3 stages of labor.**

The 1st stage—from onset of labor to full dilation of the cervix (about 10 cm)—has 2 phases, latent and active.

During the latent phase, irregular contractions become progressively better coordinated, discomfort is minimal, and the cervix effaces and dilates to 4 cm. The latent phase is difficult to time precisely, and duration varies, averaging 8 hours in nulliparas and 5 hours in multiparas; duration is considered abnormal if it lasts > 20 hours in nulliparas or > 12 hours in multiparas.

During the active phase, the cervix becomes fully dilated, and the presenting part descends well into the mid-pelvis. On average, the active phase lasts 5 to 7 hours in nulliparas and 2 to 4 hours in multiparas. Traditionally, the cervix was expected to dilate about 1.2 cm/hour in nulliparas and 1.5 cm/hour in multiparas. However, recent data suggest that slower progression of cervical dilation from 4 to 6 cm may be normal (1). Pelvic examinations are done every 2 to 3 hours to evaluate labor progress. Lack of progress in dilation and descent of the presenting part may indicate dystocia (fetopelvic disproportion).

Standing and walking shorten the first stage of labor by > 1 hour and reduce the rate of cesarean delivery (1).

If the membranes have not spontaneously ruptured, some clinicians use amniotomy (artificial rupture of membranes) routinely during the active phase. As a result, labor may progress more rapidly, and meconium-stained amniotic fluid may be detected earlier. Amniotomy during this stage may be necessary for specific indications, such as facilitating internal fetal monitoring to confirm fetal well-being. Amniotomy should be avoided in women with HIV infection or hepatitis B or C, so that the fetus is not exposed to these organisms.

During the 1st stage of labor, maternal heart rate and blood pressure and fetal heart rate should be checked continuously by electronic monitoring or intermittently by auscultation, usually with a portable Doppler ultrasound device (see fetal monitoring). Women may begin to feel the urge to bear down as the presenting part descends into the pelvis. However, they should be discouraged from bearing down until the cervix is fully dilated so that they do not tear the cervix or waste energy

The 2nd stage is the time from full cervical dilation to delivery of the fetus. On average, it lasts 2 hours in nulliparas (median 50 minutes) and 1 hour in multiparas (median 20 minutes). It may last another hour or more if conduction (epidural) analgesia or intense opioid sedation is used. For spontaneous delivery, women must supplement uterine contractions by expulsively bearing down. In the 2nd stage, women should be attended constantly, and fetal heart sounds should be checked continuously or after every contraction. Contractions may be monitored by palpation or electronically.

During the 2nd stage of labor, perineal massage with lubricants and warm compresses may soften and stretch the perineum and thus reduce the rate of 3rd- and 4th-degree perineal tears (2). These techniques are widely used by midwives and birth attendants. Precautions should be taken to reduce risk of infection with perineal massage.

During the 2nd stage (in contrast to the 1st stage), the mother's position does not affect duration or mode of delivery or maternal or neonatal outcome in deliveries without epidural anesthesia (3). Also, the pushing technique (spontaneous versus directed and delayed versus immediate) does not affect the mode of delivery or maternal or neonatal outcome. Use of epidural anesthesia delays pushing and may lengthen the 2nd stage by an hour (4).

The 3rd stage of labor begins after delivery of the infant and ends with delivery of the placenta. This stage usually lasts only a few minutes but may last up to 30 minutes.

**Birth-Massage:**

In order to stimulate the soft tissues and lower the buildup of fluids in swollen joints, deep tissue massage is recommended for pregnant women. At the same time, massage is able to improve the lymph system's ability to remove tissue waste in the body. Alleviates muscle pain and cramp...

We recommend getting a labor encouraging massage starting at 39 weeks and then every week after that until you go into labor. To make the protocol more potent you can come 2 or 3 times a week.

And, for most patients, the force of a 20- to 40-pound child bumping your belly is not enough to harm the baby. That said, toddlers can be unpredictable, and a hug could quickly turn into flailing arms and legs, which might cause abdominal injury or a fall. Consider explaining a safer way to hug you...

**Pressure Points for Normal Delivery:**

Reflexology is very relaxing and can help with stress related symptoms. Most women sleep well following a treatment and many reports a sense of wellbeing. Although relaxing many women report feeling energized. Other benefits include help with headaches, constipation, backache and symphysis pubis pain.







Certain parts of your feet and lower legs become "switches" during pregnancy, being able to trigger reactions regarding your pregnancy. The 3 spots on the legs to be avoided are: the reflexology areas around the ankles, SP6 acupressure point, urinary bladder 60, and urinary bladder 67...

**Exercises for birth:**

During the first trimester (weeks 1 to 13) you can keep doing whatever you were doing before you became pregnant, unless there's a risk that you could be hit, get too hot or have a fall. Read about exercises to avoid in pregnancy. If you're not used to exercising, start gently and build up slowly.




Most exercises are safe to perform during pregnancy, as long as you exercise with caution and do not overdo it. The safest and most productive activities are swimming, brisk walking, indoor stationary bicycling and low-impact aerobics (taught by certified aerobics instructor).

**Tips for a healthy and happy pregnancy**

- Avoid smoking. Quitting smoking is the single most preventable cause of illness and death among mothers and infants.
- Avoid alcohol consumption. When you drink alcohol, so does your developing baby. There is no known safe amount of alcohol to drink while pregnant.
- Avoid marijuana. The chemicals in marijuana pass through your system to your baby and can harm their development.
- Learn the ins and outs of a healthy pregnancy eating plan. Protein, fiber, vitamins, minerals, and more are essential.
- Take 400 micrograms (mcg) of folic acid every day. Folic acid can help prevent some major birth abnormalities.




- Seek help for depression. Depression is common and treatable. If you think you have depression, seek treatment from your health care provider as soon as possible.
- Talk to your doctor about traveling. Traveling might cause problems during pregnancy, particularly air travel later in pregnancy, so discuss your options with your doctor.
- See your doctor before starting or stopping any medication. If you are planning to become pregnant, discuss your current medicines with your doctor, midwife, or pharmacist.
- Get up to date on all of your vaccines. This will help protect you and your developing baby against serious diseases.
- Seek help for depression. Depression is common and treatable. If you think you have depression, seek treatment from your health care provider as soon as possible.

- Talk to your doctor about traveling. Traveling might cause problems during pregnancy, particularly air travel later in pregnancy, so discuss your options with your doctor.
- See your doctor before starting or stopping any medication. If you are planning to become pregnant, discuss your current medicines with your doctor, midwife, or pharmacist.
- Get up to date on all of your vaccines. This will help protect you and your developing baby against serious diseases.

## -Which month of pregnancy is the hardest?

The first trimester of pregnancy can often be the hardest. Pregnancy hormones, extreme fatigue, nausea and vomiting, tender breasts, and perpetually needing to wee make life growing a human no easy feat.

## -How do I know my baby is healthy in the womb?

- Top 7 signs of a healthy pregnancy
- The right blood pressure and blood sugar levels. ...
- The placenta's position. ...
- The right growth of the foetus. ...
- Gaining the right weight. ...
- The levels of progesterone and estrogen.

## -What are the symptoms if baby dies in womb?

The most common symptom of stillbirth is when you stop feeling your baby moving and kicking. Others include cramps, pain or bleeding from the vagina. Call your health care provider right away or go to the emergency room if you have any of these conditions.

## -Which pregnancy is the hardest Boy or girl?

Compared to girls, boys had 27 percent higher odds of preterm birth between 20- and 24-weeks' gestation; 24 percent greater risk for birth between 30 and 33 weeks; and 17 percent higher odds for delivery at 34 to 36 weeks, the study found...

## -What week is OK to give birth?




Pregnancy lasts for about 280 days or 40 weeks. A preterm or premature baby is delivered before 37 weeks of your pregnancy. Extremely preterm infants are born 23 through 28 weeks. Moderately preterm infants are born between 29 and 33 weeks.

## -Which trimester is most weight gain?

Third trimester weight gain is an important part of later pregnancy and is not usually a cause for concern. Many women will experience rapid weight gain during their third trimester. This is because the fetus typically gains the most weight in this time, according to the Office on Women's Health (OWH).

## -Can I hurt my baby by pressing on my stomach?

And, for most patients, the force of a 20- to 40-pound child bumping your belly is not enough to harm the baby. That said, toddlers can be unpredictable, and a hug could quickly turn into flailing arms and legs, which might cause abdominal injury or a fall. Consider explaining a safer way to hug you...

-What can I do to make my baby intelligent during pregnancy?

Are you pregnant? 8 simple things you can do to have an intelligent baby

- Start a story- time habit.
- Eat healthy.
- Stay fit and active.
- Recite Holy Quran and get talking.
- Keep thyroid levels in check.
- Don't ignore the supplements.
- Get a little sunshine.
- Gently massage your tummy.




-What are the safe exercises during pregnancy?

## These activities usually are safe during pregnancy:

- Walking. Taking a brisk walk is a great workout that doesn't strain your joints and muscles. ...
- Swimming and water workouts. ...
- Riding a stationary bike. ...
- Yoga and Pilates classes. ...
- Low-impact aerobics classes. ...
- Strength training.

-How can I check my baby's heartbeat at home?

How to use a stethoscope to hear your baby's heartbeat

- Find a quiet location. The quieter your surroundings, the easier it'll be to hear your baby's heartbeat. ...
- Lie down on a soft surface. ...

- Feel around your stomach and find your baby's back. ...
- Place the chest piece on this area of your stomach.

-What is an abnormal pregnancy?

Abnormal Pregnancy Conditions that develop during pregnancy are often related to the pregnancy itself (early pregnancy loss or ectopic and molar pregnancies) but not always. Sometimes issues can be caused by events unrelated to the pregnancy, like if a mother develops a disease such as measles.

-Which fruits should avoid during pregnancy?

Fruits to Avoid During Pregnancy Diet

- **Papaya**– It tops the list for obvious reasons. ...
- **Pineapple**– These are also not recommended to the pregnant women as they contain certain enzymes that alters the texture of cervix which could induce premature contractions. ...
- **Grapes**– Grapes aren't recommended for consumption during the final trimester.

-What if you are not gaining weight during pregnancy?

If a woman does not gain weight throughout pregnancy, complications such as low birth weight or premature delivery could occur. Babies who are born to mothers who do not gain more than 20 pounds are often considered small for gestational age (SGA), meaning they may have been malnourished during pregnancy.

-Why shouldn't you cross your legs when pregnant?

That said, muscles strains, backaches, and cramps are all common during pregnancy. While sitting with your legs crossed won't hurt your baby, it may contribute to ankle swelling or leg cramps. If you find your ankles swelling or your legs cramping, try sitting with both feet on the floor or elevated on a stool.

-Is it bad to sit all day while pregnant?

Researchers say they've uncovered a link between depression during pregnancy and sitting down for long periods during the second trimester. The findings also show an

increased risk for weight gain and gestational diabetes in pregnant women who sit too much.

-What should I eat to have a beautiful baby?

So, just for you, we have compiled a list of 10 foods that you can eat when you are pregnant!

- **Dairy products. ...**
- **Eggs** are considered as superfoods by many as they are a great source of vitamins, proteins and minerals. ...
- **Sweet potatoes**. ...
- **Legumes**. ...
- **Nuts**. ...
- **Orange juice**. ...
- **Leafy vegetables**. ...
- **Oatmeal**...

-How can I improve my baby's brain during pregnancy?

But here are six simple, research-supported ways to help boost your baby's brain development in utero.

- **Take a Hike**. Well, it doesn't have to be a hike, a 30-minute walk will do the trick! ...
- **Food as Medicine**. ...
- **Supplement A Healthy Diet**. ...
- **Read to Your Bump**. ...
- **Get More Sleep**. ...
- **Get Geared Up**...

-Which trimester is most critical?

The first trimester is the most critical time in your pregnancy. Although the fetus at the end of three months is only about 4 inches long and weighs less than 1 ounce, all of its functions have begun to form — major organs and nervous system, heartbeat, arms, fingers, legs, toes, hair, and buds for future teeth.

-What should I eat everyday while pregnant?

## Goals for Healthy Eating When Pregnant

Recommended daily servings include 6-11 servings of breads and grains, two to four servings of fruit, four or more servings of vegetables, four servings of dairy products, and three servings of protein sources (meat, poultry, fish, eggs or nuts). Consume fats and sweets sparingly.

-What is the reason of weak pregnancy?

In these instances, a faint positive can be caused by low levels of the pregnancy hormone human chorionic gonadotropin (hCG). As soon as you become pregnant, your body begins producing hCG. The hormone level increases as your pregnancy progresses. Home pregnancy tests are designed to detect this hormone.

-What causes abnormalities in pregnancy?

Some birth defects are caused by genes that can be passed from parents to children. Others result from a problem with chromosomes. A small number of birth defects are caused by exposure during pregnancy to certain medications, infections, and chemicals. For many birth defects, the cause is not known.

-Which fruit is best for pregnancy?

**7 nutritious** fruits you should eat during pregnancy

- **Oranges**. Oranges help you stay hydrated. ...
- **Mangoes**. Mangoes are another great source of vitamin C. ...
- **Avocados**. Avocados have more folate than other fruits. ...
- **Lemons**. ...
- **Bananas**. ...

- **Berries**. ...
- **Apples**...

-What drinks to avoid while pregnant?

## Beverages to avoid when pregnant

- Alcoholic beverages.
- Unpasteurized (raw) milk.
- Unpasteurized or "fresh squeezed" juice from a juice bar or grocery store.
- More than 200 mg of caffeine per day (12 ounces of coffee)

-What are some bad signs during pregnancy?

## DANGER SIGNS DURING PREGNANCY

- vaginal bleeding.
- convulsions/fits.
- severe headaches with blurred vision.
- fever and too weak to get out of bed.
- severe abdominal pain.
- fast or difficult breathing.

-Is dark yellow urine a sign of pregnancy?

Pregnancy. Anecdotal evidence suggests that bright-yellow urine may be an early symptom of pregnancy. However, there are no studies to support these claims.

-What is a good age to get pregnant?

Experts say the best time to get pregnant is between your late 20s and early 30s. This age range is associated with the best outcomes for both you and your baby. One study pinpointed the ideal age to give birth to a first child as 30.5. Your age is just one factor that should go into your decision to get pregnant.

## -Summary of Study Session 10

In Study Session 10, you have learned that:

- Measuring the fundal height tells you the duration of the pregnancy, how fast the baby is growing, and the probable due date.
- Remember to position the woman correctly before measuring the fundal height. The fundus of the uterus grows on average two finger-widths for each month of pregnancy.
- If the fundal height is not equal to the gestational age, you need to check the duration of pregnancy from the last normal menstrual period (LNMP). Having the wrong date is one of the main reasons for discrepancy between fundal height and gestational age.
- If the fundal height is bigger than expected for gestational age, the mother may have given you the wrong LNMP, or she may have twins, diabetes, too much water in the uterus, or a molar pregnancy.
- If the fundal height is smaller than expected for gestational age, the mother may have given you the wrong LNMP, she may have too little amniotic fluid surrounding the fetus, raised blood pressure, poor nutrition, she may be drinking alcohol or taking other harmful drugs, or the baby may be dead.

## -How can I get pregnant easily?

"In general, every other night around the time of ovulation helps increase your chance of getting pregnant," Goldfarb says. Sperm can live up to 5 days inside your body. The best suggestion is to have sex regularly -- when you're ovulating, and when you're not.

## -How much sperm is required for getting pregnant?

How many sperm do you need to get pregnant? It takes just one sperm to fertilize a woman's egg. Keep in mind, though, for each sperm that reaches the egg, there are millions that don't. On average, each

## - time men Can you feel pregnant after 2 days?

Some women may notice symptoms as early as 5 DPO, although they won't know for certain that they are pregnant until much later. Early signs and symptoms include

implantation bleeding or cramps, which can occur 5–6 days after the sperm fertilizes the egg. Other early symptoms include breast tenderness and mood changes. Ejaculate they release nearly 100 million sperm...

KUN WORLD

KUN Counseling And Training

Well Of Awareness For Thirsty People

Health Management Plan (Without Medicine)

Only a healthy body has quality thinking. Today, 90% of diseases are due to our thinking, so if you don't understand the following points, you too will become a person with a sick body and brain:

1- Phsychological and Physical knowledge
2-Diagonosis pattern and steps...
3-Fittness exercise for individuals...
4-Training for pressure points of whole body...
5- Short course of NLP...
6- Body Alignment with chiropractor....
7- Diet plan according to your body
8- Training of massage therapy...
9- Solution with Ailm-e-Noorani...
10- One hour session weekly
11- Awareness and training of "First Aid"
12- Live therapic session

and many many more... ...

Contact for registration:

0333-2278049
033-52257085

www.kunworld.com