

# Mismatch between IUCN range maps and species interactions data illustrated using the Serengeti food web

Gracielle T. Higinio<sup>1</sup> Francis Banville<sup>2,3,4</sup> Gabriel Dansereau<sup>2,4</sup> Norma R. Forero-Muñoz<sup>2,4</sup>  
Fredric M. Windsor<sup>5</sup> Timothée Poisot<sup>2,4</sup>

<sup>1</sup> Biodiversity Research Centre, University of British Columbia <sup>2</sup> Université de Montréal <sup>3</sup> Université de Sherbrooke <sup>4</sup> Quebec Centre for Biodiversity Science <sup>5</sup> School of Natural and Environmental Sciences, Newcastle University

## Correspondance to:

Gracielle T. Higinio — [graciellehiginio@gmail.com](mailto:graciellehiginio@gmail.com)

Timothée Poisot — [timothee.poisot@umontreal.ca](mailto:timothee.poisot@umontreal.ca)

**Abstract: Background.** Range maps are a useful tool to describe the spatial distribution of species. However, they need to be used with caution, as they essentially represent a rough approximation of a species' suitable habitats. When stacked together, the resulting communities in each grid cell may not always be realistic, especially when species interactions are taken into account. Here we show the extent of the mismatch between range maps, provided by the International Union for Conservation of Nature (IUCN), and species interactions data. More precisely, we show that local networks built from those stacked range maps often yield unrealistic communities, where species of higher trophic levels are completely disconnected from primary producers. **Methodology.** We used the well-described Serengeti food web of mammals and plants as our case study, and provide updated range maps for all predators by taking into account food-web structure. We then used occurrence data from the Global Biodiversity Information Facility (GBIF) to investigate where data is most lacking. **Results.** We found that most predator ranges comprised large areas without any overlapping distribution of their preys. However, many of these areas contained GBIF occurrences of the predator. **Conclusions.** Our results suggest that the mismatch between both data sources could be due either to the lack of information about ecological interactions or the geographical occurrence of preys. We finally discuss general guidelines to help identify defective data among distributions and interactions data, and we recommend this method as a valuable way to assess whether the occurrence data that are being used, even if incomplete, are ecologically accurate.

# 1 Introduction

2 Finding a species in a certain location is like finding an encrypted message that traveled through  
3 time. It carries the species' evolutionary history, migration patterns, as well as any direct and  
4 indirect effects generated by other species (some of which we may not even know exist). Ecol-  
5 ogists have been trying to decode this message with progressively more powerful tools, from  
6 their field notes to highly complex computational algorithms. However, to succeed in this chal-  
7 lenge it is important to have the right clues in hand. There are many ways we can be misled  
8 by data - or the lack of it: taxonomic errors (e.g., due to updates in the taxonomy of a species),  
9 geographic inaccuracy (e.g., approximate coordinates or lack of documentation about their ac-  
10 curacy), or sampling biases (e.g. data clustered near roads or research centers) (Ladle and Hortal  
11 2013; Hortal et al. 2015; Poisot et al. 2021). One way to identify - and potentially fix - these  
12 errors is to combine many different pieces of information about the occurrence of a species, so  
13 agreements and mismatches can emerge. Although previous studies have combined different  
14 types of occurrence data to measure the accuracy of datasets (Hurlbert and Jetz 2007; Hurlbert  
15 and White 2005; Ficetola et al. 2014), none have used different types of information so far  
16 (i.e., ecological characteristics other than geographical distribution). Here we suggest jointly  
17 analysing species occurrence (range maps and point occurrences) and ecological interactions to  
18 identify mismatches between datasets.

19 Interactions form complex networks that shape ecological structures and maintain the essential  
20 functions of ecosystems, such as seed dispersal, pollination, and biological control (Albrecht  
21 2018; Fricke et al. 2022) that ultimately affect the composition, richness, and successional pat-  
22 terns of communities across biomes. Yet, the connection between occurrence and interaction  
23 data is a frequent debate in ecology (Blanchet, Cazelles, and Gravel 2020; Wisz et al. 2013). For  
24 instance, macroecological models are often used with point or range occurrence data in order  
25 to investigate the dynamics of a species with its environment. However, these models do not  
26 account for ecological interactions, although it has been demonstrated that they might largely  
27 affect species distribution (Abrego et al. 2021; Afkhami, McIntyre, and Strauss 2014; Araújo,  
28 Marcondes-Machado, and Costa 2014; Godsoe et al. 2017; Godsoe and Harmon 2012; Gotelli,  
29 Graves, and Rahbek 2010; Wisz et al. 2013). Some researchers argue that occurrence data can

also capture real-time interactions (see Roy et al. 2016; Ryan et al. 2018), and, because of that, it would not be necessary to include ecological interaction dynamics in macroecological models. On the other hand, many mechanistic simulation models in ecology have considered the effect of competition and facilitation in range shifts. For example, Gotelli *et al.* (2010) demonstrate how conspecific attraction might be the main factor driving the distribution of migratory birds; Afkhami *et al.* (2014) explores how mutualistic fungal endophytes are responsible for expanding the range of native grass; many other examples are discussed in Wisz *et al.* (2013). Although interactions across trophic levels are demonstrated to determine species range (Wisz et al. 2013), the use of these interactions in mechanistic simulation models in macroecology remains insufficient (as discussed in Cabral, Valente, and Hartig 2017).

A significant challenge in this debate is the quality and quantity of species distribution and ecological data (Boakes et al. 2010; Ronquillo et al. 2020; Meyer, Weigelt, and Kreft 2016) - a gap that can lead to erroneous conclusions in macroecological research (Hortal et al. 2008). Amongst the geographical data available are the range maps provided by the International Union for the Conservation of Nature (IUCN). Such maps consist of simplified polygons, often created as alpha or convex hulls around known species locations, refined by expert knowledge about the species (IUCN Red List Technical Working Group 2019). These maps can be used in macroecological inferences in the lack of more precise information (Fourcade 2016; Alhajeri and Fourcade 2019), but it has been recommended that they are used with caution since they tend to underestimate the distribution of species that are not well-known (Herkt, Skidmore, and Fahr 2017) (especially at fine scale resolutions; Hurlbert and Jetz (2007); Hurlbert and White (2005)), do not represent spatial variation in species occurrence and abundance (Dallas, Pironon, and Santini 2020), and can include inadequate areas within the estimated range. Another source of species distribution information is the Global Biodiversity Information Facility (GBIF), which is an online repository of georeferenced observational records that come from various sources, including community science programs, museum collections, and long-term monitoring schemes. A great source of bias in these datasets is the irregular sampling effort, with more occurrences originating from attractive and accessible areas and observation of charismatic species (Alhajeri and Fourcade 2019). As for ecological data, a complete assessment is difficult and is aggravated by

59 biased sampling methods and data aggregation (Poisot et al. 2020; Hortal et al. 2015). Never-  
60 theless, we have witnessed an increase in the availability of biodiversity data in the last decades,  
61 including those collected through community science projects (Callaghan et al. 2019; Pocock et  
62 al. 2015) and dedicated databases, such as Mangal (Poisot et al. 2016). This provides an oppor-  
63 tunity to merge species distribution and ecological interaction data to improve our predictions  
64 of where a species may be found across large spatial scales (e.g., continental and global).

65 It has been demonstrated that the agreement between range maps and point data varies geograph-  
66 ically (Hurlbert and Jetz 2007; Hurlbert and White 2005; Ficetola et al. 2014). Adding ecolog-  
67 ical interaction data to this comparison might help to elucidate where these (dis)agreements are  
68 more likely to be true and which dataset better represent the actual distribution of species. In  
69 this context, we elaborate a method that allows us to detect areas of potential misestimation of  
70 species' distribution data (more precisely range maps) based on interaction data. This method is  
71 based on the assumption that organisms cannot persist in an area unless they are directly or in-  
72 directly connected to a primary producer within their associated food web (Power 1992). Thus,  
73 given that herbivores are the main connection between plant resources (directly limited by en-  
74 vironmental conditions) and predators (Dobson 2009; Scott et al. 2018), the range of a predator  
75 (omnivore or carnivore) depends on the overlapping ranges of its herbivore preys. If sections  
76 of a predator's range does not overlap with at least one of its prey it will become disconnected  
77 from primary producers, and therefore we would not expect the predator to occur in this area.

78 This mismatch can be the result of different mechanisms, like the misestimation of both the  
79 predator's and the preys' ranges (Ladle and Hortal 2013; Rondinini et al. 2006), taxonomic  
80 errors (Isaac, Mallet, and Mace 2004; Ladle and Hortal 2013), or the lack of information about  
81 trophic links (i.e., the lack of connection between the ranges of a predator and a primary producer  
82 may be due a third species we don't know is connected to both). Here we discuss the mechanisms  
83 that can lead to the lack of agreement between data, and build from that a vision for the next  
84 steps, reinforcing the importance of geographically explicit interaction data.

## Methods

We identified areas of uncertainty on the ranges of predators based on a simple rule: we removed any part of a predator's range that did not intersect with the range of at least one prey herbivore species, which in turn is directly connected to a primary producer (plants). To do that, we used a Serengeti food web dataset (Baskerville et al. 2011) (which comprises carnivores, herbivores, and plants from Tanzania) and its species ranges from IUCN. Then, we calculated the difference in range sizes between the original IUCN ranges of predators and those without the areas where they would be alone, based on species interaction data. Finally, we added the GBIF occurrence points for the Serengeti species to investigate whether the results would be different if we used another source of distribution data. ## Data

We investigated the mismatch between savannah species ranges and interactions in Africa (fig. 1). These ecosystems host a range of different species, including the well-characterized predator-prey dynamics between iconic predators (e.g., lions, hyenas, and leopards) and large herbivores (e.g., antelopes, wildebeests, and zebras), as well as a range of herbivorous and carnivorous small mammals. The Serengeti ecosystem has been extensively studied and its food web is one of the most complete we have to date, including primary producers identified to the species level. Here we focus on six groups of herbivores and carnivores from the Serengeti Food Web Data Set (Baskerville et al. 2011). These species exhibit direct antagonistic (predator-prey) interactions with one another and are commonly found across savannah ecosystems on the African continent (McNaughton 1992). Plants in the network were included indirectly in our analyses as we do not expect the primary producers to significantly influence the range of herbivores for several reasons. Firstly, many savannah plants are functionally similar (i.e., grasses, trees and shrubs) and cooccur across the same habitats (Baskerville et al. 2011). Secondly, herbivores in the network are broadly generalists feeding on a wide range of different plants across habitats. Indeed, out of 129 plants in our dataset, herbivores ( $n = 23$ ) had a mean out degree (mean number of preys) of around 22 (std = 17.5). There is also an absence of global range maps for many plant species (Daru 2020), which prevents their direct inclusion in our analysis. Therefore, we assume that plants consumed by herbivores are present across their ranges, and as such the ranges of herbivores are not expected to be significantly constrained by the availability of food plants.

From the wider ecological network presented in Baskerville (2011), we sampled interaction data for herbivores and carnivores. This subnetwork contained 32 taxa (23 herbivores and 9 carnivores) and 84 interactions and had a connectance of 0.08. Although self-loops are informative, we removed these interactions to allow for the original IUCN ranges of predators with cannibalistic interactions to be adjusted. We treated this overall network as a metaweb since it *should* contain all potential species interactions between mammalian taxa occurring across savannah ecosystems such as the Serengeti.

We compiled IUCN range maps for the 32 species included in the metaweb from the Spatial Data Download portal ([www.iucnredlist.org/resources/spatial-data-download](http://www.iucnredlist.org/resources/spatial-data-download)), which we rasterized at a 0.5 degrees resolution (~50 km at the equator). We restricted the rasters to a spatial extent comprised between latitudes 35°S and 40°N and longitudes 20°W and 55°E. We then combined interaction data from the metaweb and cooccurrence data generated from species ranges to create networks for each raster pixel. This generated a total of 11,308 pixel-level networks. These networks describe potential predation, not actual interactions: the former is derived information from the metaweb, and the latter is contingent on the presence of herbivores.

### Range overlap measurement

We calculated the geographical overlap, i.e. the extent to which interacting predator and prey species co-occurred across their ranges, as  $a/(a + c)$ , where  $a$  is the number of pixels where predator and prey cooccur and  $c$  is the number of pixels where only the focal species occur. This index of geographical overlap can be calculated with prey or predators as the focal species. Values vary between 0 and 1, with values closer to 1 indicating that there is a large overlap in the ranges of the two species and values closer to 0 indicating low cooccurrence across their ranges. For each predator species, we calculated its generality to understand whether the level of trophic specialization (i.e., number of prey items per predator) affects the extent to which the ranges of the species were altered. One would assume that predators with a greater number of prey taxa (i.e., a higher generality) are less likely to have significant changes in their range as it is more likely that at least one prey species is present across most of their range.

## Validation

For each species in the dataset we collated point observation data from GBIF ([www.gbif.org](http://www.gbif.org)). We used the GBIF download API to retrieve all species occurrences on November 22nd 2022 (GBIF.org 2022). We restricted our query to the data with spatial coordinates and which were inside the spatial extent of our rasters. We did not use continental or date filters to retrieve as much data as possible. However, a few observations were localized in the ocean near latitude 0° and longitude 0°. We assumed these were errors and removed all observations falling in the extent between latitudes 2°S and 2°N and longitudes 2°W and 2°E to keep only mainland sites. We then converted the occurrence data into raster format by determining which pixels had a least one GBIF occurrence. This allowed us to remove the effect of repeated sampling in some locations. These data were used to validate the range adjustments made based on species interactions (see beginning of Methods section). To do so, we calculated the proportion of GBIF presence pixels occurring within both the original IUCN species range and the adjusted one. We then compared these proportions for the predators to verify if the range adjustments removed locations with GBIF observations, hence likely true habitats.

## Software

We performed all analyses using *Julia* v1.7.2 (Bezanson et al. 2017). We used the packages `SimpleSDMLayers.jl` (Dansereau and Poisot 2021) to manipulate the raster layers, `EcologicalNetworks.jl` (Poisot et al. 2019) to construct and manipulate the interaction networks, and `GBIF.jl` (Dansereau and Poisot 2021) to reconcile species names with the GBIF backbone taxonomy (GBIF Secretariat 2021). We also used *GDAL* (GDAL/OGR contributors 2021) to rasterize the IUCN range maps (initially available as shapefiles from the Spatial Data Download portal). All the scripts required to reproduce the analyses are available at <https://doi.org/10.5281/zenodo.6842861>.



## Results

Mammal species found in the Serengeti food web are widespread in Africa, especially in grasslands and savannahs (first panel of fig. 1). However, most local networks (83.2%) built using the original IUCN range maps had at least one mammal species without a path to a primary producer (second panel of fig. 1). On average, local food webs had almost the third of their mammal species (mean = 30.5%, median = 14.3%) disconnected from basal species. In addition, many networks (16.6%) only had disconnected mammals; these networks however all had a very low number of mammal species, specifically between 1 and 4 (from a total of 32). As expected, the proportion of carnivores with a path to a primary producer was conditional on the total number of mammal species in each local network (third panel of fig. 1).

[Figure 1 about here.]

### Specialized predators lose more range

[Figure 2 about here.]

Predators with fewer prey lose more range with our method (fig. 2). For instance, both *Leptailurus serval* and *Canis mesomelas* have only one prey in the Serengeti food web (tbl. 1), each of them with a very small range compared to those of their predator. This discrepancy between range sizes promotes significant range loss. On the other hand, predators of the genus *Panthera* are some of the most connected species, and they also lose the least proportion of their ranges. This mismatch between predators and preys can also be a result of taxonomic disagreement between the geographical and ecological data. Although *Canis aureus* has the same number of prey as *Caracal caracal*, none of the prey taxa of the former occurs inside its original range (tbl. 1), which results in complete range loss.

[Figure 3 about here.]

There was a high variation in the overlap of predator and prey ranges (fig. 3). The high density of points on the left-hand side of fig. 3 indicates that most preys have small ranges in comparison to

189 those of the set of carnivores in the networks, resulting in either low overlap between both ranges  
190 (bottom) or high overlap of ranges because much of that of the prey is within predators' range  
191 (top). The top-right side of the plot encompasses situations where the ranges of both predator  
192 and prey are similar and overlapping, while the bottom-right part of the plot represents a situation  
193 where the range of the predator is smaller than that of its prey and much of it occurs within the  
194 preys' range. For example, *Panthera pardus* had many preys occurring inside its range, with  
195 highly variable levels of overlap (tbl. 1). In general, species exhibited more consistent values of  
196 prey-predator overlap, than predator-prey overlap – indicated by the spread of points along the  
197 x-axis, yet more restricted variation on the y-axis (fig. 3). There was also no overall relationship  
198 between the two metrics, or for any predator species.

Table 1: List of species analysed, their out and in degrees, total original range size (in pixels), and proportion of their ranges occupied by their preys and predators (values between 0 and 1). Species are sorted according to the groups identified by Baskerville et al. (2011). Notice how some species are isolated in the network (*Loxodonta africana*) and how *Canis aureus*'s range does not overlap with any of its preys.

Species	Number		Total range size	Proportion	Proportion
	Number of preys	of predators		of range	of range
				occupied by preys	occupied by predators
Large carnivores					
Acinonyx jubatus	8	1	9250	0.437	0.618
Crocuta crocuta	12	1	4822	0.844	0.253
Lycaon pictus	14	0	427	0.918	-
Panthera leo	18	0	1274	0.935	-
Panthera pardus	22	0	7563	0.766	-
Small carnivores					
Canis aureus	4	1	816	0.000	0.782
Canis mesomelas	1	1	2201	0.190	0.994
Caracal caracal	4	0	5239	0.833	-

Species	Number of preys	Number of predators	Total range size	Proportion of range occupied by preys	Proportion of range occupied by predators
Leptailurus serval	1	1	4319	0.011	0.978
<b>Small herbivores</b>					
Damaliscus lunatus	0	4	626	-	1
Hippopotamus amphibius	0	0	419	-	-
Kobus ellipsiprymnus	0	4	2961	-	1
Ourebia ourebi	0	5	2484	-	1
Pedetes capensis	0	2	1318	-	1
Phacochoerus africanus	0	5	3331	-	1
Redunca redunca	0	5	1935	-	1
Rhabdomys pumilio	0	5	53	-	1
Tragelaphus oryx	0	2	2316	-	0.990
Tragelaphus scriptus	0	3	3999	-	0.985
<b>Large grazers</b>					
Aepyceros melampus	0	5	1167	-	1
Alcelaphus buselaphus	0	4	2307	-	1
Connochaetes taurinus	0	6	1074	-	1
Equus quagga	0	5	786	-	1
Eudorcas thomsonii	0	6	51	-	1
Nanger granti	0	6	261	-	1
<b>Hyraxes</b>					
Heterohyrax brucei	0	1	1961	-	0.973
Procavia capensis	0	1	5312	-	0.647

Species	Number of preys	Number of predators	Total range size	Proportion of range occupied by preys	Proportion of range occupied by predators
<b>Others</b>					
<i>Giraffa camelopardalis</i>	0	1	607	-	0.473
<i>Loxodonta africana</i>	0	0	1078	-	-
<i>Madoqua kirkii</i>	0	7	443	-	1
<i>Papio anubis</i>	0	1	2571	-	0.937
<i>Syncerus caffer</i>	0	1	2808	-	0.251

## Validation with GBIF occurrences

The proportion of GBIF pixels (pixels with at least one GBIF occurrence) falling within the IUCN ranges varied from low to high depending on the species (fig. 4, left). The lowest proportions occurred for species with small ranges. Amongst herbivores, *Rhabdomys pumilio* has a proportion of 22.6% of its presence pixels within its IUCN range, while predators have this proportion above 55% (such as *Lycaon pictus*, with 55.1%, and *Canis aureus*, with 56.2%). Nevertheless, some species with smaller ranges showed high data overlap (such as *Canis mesomelas*, with 94.9%, and many herbivores). Overall, predators and preys displayed similar overlap variations, and species with median and large ranges had higher proportions of occurrences falling into their IUCN range.

The proportion of GBIF pixels in updated ranges can only be equal to or lower than that of the original ranges, as our analysis removes pixels from the original range and does not add new ones. Rather, the absence of a difference between the two types of ranges indicates that no pixels with GBIF observations, hence likely true habitats, were removed by our analysis. Here this proportion was mostly similar to that of the original IUCN ranges for most predator species (fig. 4). Two species showed no difference in proportion while four species showed only

small differences (*Crocota crocuta* lost 1.3% of the original data overlap; *Acinonyx jubatus* lost 1.9%; *Panthera pardus* lost 8.8%; and *Caracal caracal* lost 12.3%). On the other hand, three species, *Canis aureus*, *Canis mesomelas*, and *Leptailurus serval* showed very high differences, with overlaps lowered by 100%, 57.4%, and 100% respectively. These last two species are also the only predators with a single prey in our metaweb. *Canis aureus* has four preys, but it has one of the smallest ranges in IUCN, which is not covered by any of its preys. This result reinforces the concern raised in the literature on the use of IUCN range maps for species that are not well known (Herkt, Skidmore, and Fahr 2017), demonstrating how small range species are likely to have their distribution underestimated in the IUCN database. Additionally, the fact that *Canis aureus* had such a conspicuous mismatch between both the original and updated IUCN range maps, and between GBIF and IUCN data, may indicate a taxonomic incongruency between the three databases used here, which we explore in the Discussion section. Our results delineate how a mismatch between GBIF and IUCN databases differ greatly with small changes in herbivore species ranges, and it is somewhat positively related to range size for predator species. Moreover, we show that accounting for interactions does not necessarily aggravates this dissimilarity, but it is relevant for species with little ecological information or specialists.

[Figure 4 about here.]

## Discussion

Here we identify areas of data mismatch between species range maps by using ecological interaction data (predator-prey interactions within food webs). Our results did show a significant mismatch in the IUCN range areas of specialized and generalist predatory organisms and their prey, which highlights the importance of accounting for species interactions when estimating the range of a species. Although this type of data mismatch can result of actual ecological processes, outdated occurrence data, taxonomic errors and more, we argue that, here, they rather indicate a lack of interaction sampling data.

The case of the golden jackal (*Canis aureus*) is a good illustration of how the taxonomic, geographical and ecological data can be used to validate one another. The jackal is a widespread

taxon in northern Africa, Europe, and Australasia, generally well adapted to local conditions due to its largely varied diet (Tsunoda and Saito 2020; Krofel et al. 2021). Because of that, we expected that the *Canis* species in our dataset would be the ones losing the least amount of range, with a higher value of the proportion of GBIF pixels within their IUCN range maps. However, the taxonomy of this group is a matter of intense discussion, as molecular and morphological data seem to disagree in the clustering of species and subspecies (Krofel et al. 2021; Stoyanov 2020). This debate probably influenced our results: with originally only 56.2% of the GBIF pixels of the golden jackal overlapping with its IUCN data, we suspect that many of the GBIF occurrences refer to other *Canis* species, and that its taxonomic identification in the network database is probably outdated. This led to a complete exclusion of *Canis aureus* from its original range in our analysis, despite the fact that this species has four documented preys in our metaweb.

#### **Geographical mismatch and data availability**

The lack of superposition between IUCN range maps and GBIF occurrences in our results suggests that we certainly miss geographical information about the distribution of either the prey or the predator. On the other hand, if both GBIF and IUCN occurrences tended to superpose and the species was still locally removed, this indicates that we don't have information about all its interactions (e.g., predators may be feeding on different species than the ones in our dataset outside the Serengeti ecosystem). This rationale can be illustrated with three types of mismatches identified in our results.

First, *Panthera leo* was one of the species with no difference between ranges before and after our analysis, but 40.7% of the GBIF occurrences for this species fell outside its IUCN range (fig. 4). In this particular case, the IUCN maps seem to agree with species interaction data. However, the disagreement between the IUCN and the GBIF databases is concerning and suggests that the IUCN maps might underestimate the lion's distribution.

On the other hand, *Leptailurus serval* and *Canis mesomelas* are two of the three species that have the higher proportion of mismatched range due to the lack of paths to a herbivore, but are also some of the species with the higher proportion of GBIF occurrences inside IUCN range

maps (fig. 4). This indicates that the information we are missing for these two species is related to either an additional interaction or to the presence of interacting species. To illustrate that, we mapped the GBIF data for the prey of *Leptailurus serval*, with a mobility buffer around each point (fig. 5). When considering GBIF data, approximately 42% of the prey's occurrences are within the portion of the predator's range that was divergent from its original IUCN data. With the buffer area, this corresponds to 15% of the mismatched area. By adding GBIF information for the prey, we could therefore reduce the discrepancy of the range (or information) for the predator by 15% since its distribution is conditional on the occurrence of its preys. In other words, the range mismatch was exaggerated because we were missing information on the presence of an interacting species (i.e., this also indicates that there is a mismatch - or complementarity - between the IUCN and GBIF data for their prey).

[Figure 5 about here.]

Finally, the extreme case of *Canis aureus* illustrates a lack of both geographical and ecological information: only half of its GBIF presence pixels and none of its preys occur inside its IUCN range. We believe, therefore, that the validation of species distribution based on ecological interaction is a relevant method that can further fill in information gaps. Nevertheless, it is imperative that more geographically explicit data about ecological networks and interactions become available. This would help clarify when cooccurrences can be translated into interactions (Windsor et al. 2022) and help the development of more advanced validation methods for occurrence data.

## Next steps

Here we demonstrated how we can detect uncertainty in species distribution data using ecological interactions. Knowing where questionable occurrence data are can be crucial in ecological modelling (Hortal 2008; Ladle and Hortal 2013), and accounting for these errors can improve model outputs by diminishing the error propagation (Draper 1995). For instance, we believe this is a way to account for ecological interactions in habitat suitability models without making the models more complex, but by making sure (not assuming) that the input data - the species occurrence - actually accounts for ecological interactions. It is important to notice, however,

that the quality and usefulness of this method are highly correlated with the amount and quality of data available about species' occurrences **and** interactions. With this paper, we hope to add to the collective effort to decode the encrypted message that is the occurrence of a species in space and time. A promising avenue that adds to our method is the prediction of networks and interactions in large scales (Strydom et al. 2021; Windsor et al. 2022), for they can add valuable information about ecological interactions where they are missing. Additionally, in order to achieve a robust modelling framework towards actual species distribution models we should invest in efforts to collect and combine open data on species occurrence and interactions (Windsor et al. 2022), especially because we may be losing ecological interactions at least as fast as we are losing species (Valiente-Banuet et al. 2015).

## Acknowledgements

We acknowledge that this study was conducted on land within the traditional unceded territory of the Saint Lawrence Iroquoian, Anishinabewaki, Mohawk, Huron-Wendat, and Omàmiwininiwak nations. We thank the editor and reviewer for their thoughtful comments, which considerably improved this manuscript.

## References

- Abrego, Nerea, Tomas Roslin, Tea Huotari, Yinqiu Ji, Niels Martin Schmidt, Jiaxin Wang, Douglas W. Yu, and Otso Ovaskainen. 2021. "Accounting for Species Interactions Is Necessary for Predicting How Arctic Arthropod Communities Respond to Climate Change." *Ecography* 44 (6): 885–96. <https://doi.org/10.1111/ecog.05547>.
- Afkhami, Michelle E., Patrick J. McIntyre, and Sharon Y. Strauss. 2014. "Mutualist-Mediated Effects on Species' Range Limits Across Large Geographic Scales." *Ecology Letters* 17 (10): 1265–73. <https://doi.org/10.1111/ele.12332>.
- Albrecht, Jörg. 2018. "Plant and Animal Functional Diversity Drive Mutualistic Network Assembly Across an Elevational Gradient." *NATURE COMMUNICATIONS*, 10.



- Alhajeri, Bader H, and Yoan Fourcade. 2019. “High Correlation Between Species-Level Environmental Data Estimates Extracted from IUCN Expert Range Maps and from GBIF Occurrence Data.” *Journal of Biogeography*, 13. <https://doi.org/10.1111/jbi.13619>.
- Araújo, Carlos B. de, Luiz Octavio Marcondes-Machado, and Gabriel C. Costa. 2014. “The Importance of Biotic Interactions in Species Distribution Models: A Test of the Eltonian Noise Hypothesis Using Parrots.” *Journal of Biogeography* 41 (3): 513–23. <https://doi.org/10.1111/jbi.12234>.
- Baskerville, Edward B., Andy P. Dobson, Trevor Bedford, Stefano Allesina, T. Michael Anderson, and Mercedes Pascual. 2011. “Spatial Guilds in the Serengeti Food Web Revealed by a Bayesian Group Model.” *PLOS Computational Biology* 7 (12): e1002321. <https://doi.org/10.1371/journal.pcbi.1002321>.
- Bezanson, Jeff, Alan Edelman, Stefan Karpinski, and Viral B. Shah. 2017. “Julia: A Fresh Approach to Numerical Computing.” *SIAM Review* 59 (1): 65–98. <https://doi.org/10.1137/141000671>.
- Blanchet, F. Guillaume, Kevin Cazelles, and Dominique Gravel. 2020. “Co-Occurrence Is Not Evidence of Ecological Interactions.” *Ecology Letters* 23 (7): 1050–63. <https://doi.org/10.1111/ele.13525>.
- Boakes, Elizabeth H., Philip J. K. McGowan, Richard A. Fuller, Ding Chang-qing, Natalie E. Clark, Kim O’Connor, and Georgina M. Mace. 2010. “Distorted Views of Biodiversity: Spatial and Temporal Bias in Species Occurrence Data.” *PLOS Biology* 8 (6): e1000385. <https://doi.org/10.1371/journal.pbio.1000385>.
- Cabral, Juliano Sarmiento, Luis Valente, and Florian Hartig. 2017. “Mechanistic Simulation Models in Macroecology and Biogeography: State-of-Art and Prospects.” *Ecography* 40 (2): 267–80. <https://doi.org/10.1111/ecog.02480>.
- Callaghan, Corey T., Jodi J. L. Rowley, William K. Cornwell, Alistair G. B. Poore, and Richard E. Major. 2019. “Improving Big Citizen Science Data: Moving Beyond Haphazard Sampling.” *PLOS Biology* 17 (6): e3000357. <https://doi.org/10.1371/journal.pbio.3000357>.

350 Dallas, Tad, Samuel Pironon, and Luca Santini. 2020. “The Abundant-Centre Is Not All That  
 351 Abundant: A Comment to Osorio-Olvera Et Al. 2020,” 2020.02.27.968586. [https://doi.  
 352 org/10.1101/2020.02.27.968586](https://doi.org/10.1101/2020.02.27.968586).

353 Dansereau, Gabriel, and Timothée Poisot. 2021. “SimpleSDMLayers.jl and GBIF.jl: A Frame-  
 354 work for Species Distribution Modeling in Julia.” *Journal of Open Source Software* 6 (57):  
 355 2872. <https://doi.org/10.21105/joss.02872>.

356 Daru, Barnabas H. 2020. “GreenMaps: A Tool for Addressing the Wallacean Shortfall in the  
 357 Global Distribution of Plants.” *bioRxiv*, 2020.02.21.960161. [https://doi.org/10.1101/  
 358 2020.02.21.960161](https://doi.org/10.1101/2020.02.21.960161).

359 Dobson, Andy. 2009. “Food-Web Structure and Ecosystem Services: Insights from the Serengeti.”  
 360 *Philosophical Transactions of the Royal Society B: Biological Sciences* 364 (1524): 1665–  
 361 82. <https://doi.org/10.1098/rstb.2008.0287>.

362 Draper, D. 1995. “Assessment and Propagation of Model Uncertainty.” *Journal of the Royal  
 363 Statistical Society Series B-Statistical Methodology* 57 (1): 45–97. [https://doi.org/10.  
 364 1111/j.2517-6161.1995.tb02015.x](https://doi.org/10.1111/j.2517-6161.1995.tb02015.x).

365 Ficetola, Gentile Francesco, Carlo Rondinini, Anna Bonardi, Vineet Katariya, Emilio Padoa-  
 366 Schioppa, and Ariadne Angulo. 2014. “An Evaluation of the Robustness of Global Am-  
 367 phibian Range Maps.” *Journal of Biogeography* 41 (2): 211–21. [https://doi.org/10.  
 368 1111/jbi.12206](https://doi.org/10.1111/jbi.12206).

369 Fourcade, Yoan. 2016. “Comparing Species Distributions Modelled from Occurrence Data  
 370 and from Expert-Based Range Maps. Implication for Predicting Range Shifts with Climate  
 371 Change.” *Ecological Informatics* 36: 8–14. [https://doi.org/10.1016/j.ecoinf.2016.  
 372 09.002](https://doi.org/10.1016/j.ecoinf.2016.09.002).

373 Fricke, Evan C., Alejandro Ordonez, Haldre S. Rogers, and Jens-Christian Svenning. 2022.  
 374 “The Effects of Defaunation on Plants’ Capacity to Track Climate Change.” *Science*. [https:  
 375 //doi.org/10.1126/science.abk3510](https://doi.org/10.1126/science.abk3510).

376 GBIF.org. 2022. “GBIF Occurrence Download.” The Global Biodiversity Information Facility.  
 377 <https://doi.org/10.15468/DL.PF4586>.

378 GBIF Secretariat. 2021. “GBIF Backbone Taxonomy.” <https://doi.org/10.15468/39omei>.

379 GDAL/OGR contributors. 2021. *GDAL/OGR Geospatial Data Abstraction Software Library*.

380 Manual. Open Source Geospatial Foundation.

381 Godsoe, William, and Luke J. Harmon. 2012. “How Do Species Interactions Affect Species Dis-

382 tribution Models?” *Ecography* 35 (9): 811–20. [https://doi.org/10.1111/j.1600-0587.](https://doi.org/10.1111/j.1600-0587.2011.07103.x)

383 [2011.07103.x](https://doi.org/10.1111/j.1600-0587.2011.07103.x).

384 Godsoe, William, Jill Jankowski, Robert D. Holt, and Dominique Gravel. 2017. “Integrating

385 Biogeography with Contemporary Niche Theory.” *Trends in Ecology and Evolution* 32 (7):

386 488–99. <https://doi.org/10.1016/j.tree.2017.03.008>.

387 Gotelli, Nicholas J., Gary R. Graves, and Carsten Rahbek. 2010. “Macroecological Signals

388 of Species Interactions in the Danish Avifauna.” *Proceedings of the National Academy of*

389 *Sciences* 107 (11): 5030–35. <https://doi.org/10.1073/pnas.0914089107>.

390 Herkt, K. Matthias B., Andrew K. Skidmore, and Jakob Fahr. 2017. “Macroecological Conclu-

391 sions Based on IUCN Expert Maps: A Call for Caution.” *Global Ecology and Biogeography*

392 26 (8): 930–41. <https://doi.org/10.1111/geb.12601>.

393 Hortal, Joaquín. 2008. “Uncertainty and the Measurement of Terrestrial Biodiversity Gradi-

394 ents.” *Journal of Biogeography* 35 (8): 1335–36. [https://doi.org/10.1111/j.1365-2699.](https://doi.org/10.1111/j.1365-2699.2008.01955.x)

395 [2008.01955.x](https://doi.org/10.1111/j.1365-2699.2008.01955.x).

396 Hortal, Joaquín, Francesco de Bello, José Alexandre F. Diniz-Filho, Thomas M. Lewinsohn,

397 Jorge M. Lobo, and Richard J. Ladle. 2015. “Seven Shortfalls That Beset Large-Scale

398 Knowledge of Biodiversity.” *Annual Review of Ecology, Evolution, and Systematics* 46 (1):

399 523–49. <https://doi.org/10.1146/annurev-ecolsys-112414-054400>.

400 Hortal, Joaquín, Alberto Jiménez-Valverde, José F. Gómez, Jorge M. Lobo, and Andrés Baselga.

401 2008. “Historical Bias in Biodiversity Inventories Affects the Observed Environmental

402 Niche of the Species.” *Oikos* 117 (6): 847–58. [https://doi.org/10.1111/j.0030-1299.](https://doi.org/10.1111/j.0030-1299.2008.16434.x)

403 [2008.16434.x](https://doi.org/10.1111/j.0030-1299.2008.16434.x).

404 Hurlbert, Allen H., and Walter Jetz. 2007. “Species Richness, Hotspots, and the Scale Depen-

405 dence of Range Maps in Ecology and Conservation.” *Proceedings of the National Academy*

of Sciences 104 (33): 13384–89. <https://doi.org/10.1073/pnas.0704469104>.

Hurlbert, Allen H., and Ethan P. White. 2005. “Disparity Between Range Map- and Survey-Based Analyses of Species Richness: Patterns, Processes and Implications.” *Ecology Letters* 8 (3): 319–27. <https://doi.org/10.1111/j.1461-0248.2005.00726.x>.

Isaac, Nick J. B., James Mallet, and Georgina M. Mace. 2004. “Taxonomic Inflation: Its Influence on Macroecology and Conservation.” *Trends in Ecology & Evolution* 19 (9): 464–69. <https://doi.org/10.1016/j.tree.2004.06.004>.

IUCN Red List Technical Working Group. 2019. “Mapping Standards and Data Quality for IUCN Red List Spatial Data.” Prepared by the Standards and Petitions Working Group of the IUCN SSC Red . . . .

Krofel, M., J. Hatlauf, W. Bogdanowicz, L. a. D. Campbell, R. Godinho, Y. V. Jhala, A. C. Kitchener, et al. 2021. “Towards Resolving Taxonomic Uncertainties in Wolf, Dog and Jackal Lineages of Africa, Eurasia and Australasia.” *Journal of Zoology* n/a (n/a): 1–14. <https://doi.org/10.1111/jzo.12946>.

Ladle, Richard, and Joaquín Hortal. 2013. “Mapping Species Distributions: Living with Uncertainty.” *Frontiers of Biogeography* 5 (1): 4–6.

McNaughton, S. J. 1992. “The Propagation of Disturbance in Savannas Through Food Webs.” *Journal of Vegetation Science* 3 (3): 301–14. <https://doi.org/10.2307/3235755>.

Meyer, Carsten, Patrick Weigelt, and Holger Kreft. 2016. “Multidimensional Biases, Gaps and Uncertainties in Global Plant Occurrence Information.” *Ecology Letters* 19 (8): 992–1006. <https://doi.org/10.1111/ele.12624>.

Pocock, Michael J. O., Helen E. Roy, Chris D. Preston, and David B. Roy. 2015. “The Biological Records Centre: A Pioneer of Citizen Science.” *Biological Journal of the Linnean Society* 115 (3): 475–93. <https://doi.org/10.1111/bij.12548>.

Poisot, Timothée, Benjamin Baiser, Jennifer A Dunne, Sonia Kéfi, François Massol, Nicolas Mouquet, Tamara N Romanuk, Daniel B Stouffer, Spencer A Wood, and Dominique Gravel. 2016. “Mangal - Making Ecological Network Analysis Simple.” *Ecography* 39 (4): 384–90.

- Poisot, Timothée, Gabriel Bergeron, Kevin Cazelles, Tad Dallas, Dominique Gravel, Andrew MacDonald, Benjamin Mercier, Clément Violet, and Steve Vissault. 2021. “Global Knowledge Gaps in Species Interaction Networks Data.” *Journal of Biogeography* 48 (7): 1552–63. <https://doi.org/10.1111/jbi.14127>.
- Poisot, Timothée, Gabriel Bergeron, Kevin Cazelles, Tad Dallas, Dominique Gravel, Andrew Macdonald, Benjamin Mercier, Clément Violet, and Steve Vissault. 2020. “Environmental Biases in the Study of Ecological Networks at the Planetary Scale.” *bioRxiv*, 2020.01.27.921429. <https://doi.org/10.1101/2020.01.27.921429>.
- Poisot, Timothée, Zachary Bélisle, Laura Hoebeke, Michiel Stock, and Piotr Szefer. 2019. “EcologicalNetworks.jl: Analysing Ecological Networks of Species Interactions.” *Ecography* 42 (11): 1850–61. <https://doi.org/10.1111/ecog.04310>.
- Power, Mary E. 1992. “Top-Down and Bottom-Up Forces in Food Webs: Do Plants Have Primacy.” *Ecology* 73 (3): 733–46. <https://doi.org/10.2307/1940153>.
- Rondinini, Carlo, Kerrie A. Wilson, Luigi Boitani, Hedley Grantham, and Hugh P. Possingham. 2006. “Tradeoffs of Different Types of Species Occurrence Data for Use in Systematic Conservation Planning.” *Ecology Letters* 9 (10): 1136–45. <https://doi.org/10.1111/j.1461-0248.2006.00970.x>.
- Ronquillo, Cristina, Fernanda Alves-Martins, Vicente Mazimpaka, Thadeu Sobral-Souza, Bruno Vilela-Silva, Nagore G. Medina, and Joaquín Hortal. 2020. “Assessing Spatial and Temporal Biases and Gaps in the Publicly Available Distributional Information of Iberian Mosses.” *Biodiversity Data Journal* 8: e53474. <https://doi.org/10.3897/BDJ.8.e53474>.
- Roy, Helen E., Elizabeth Baxter, Aoiné Saunders, and Michael J. O. Pocock. 2016. “Focal Plant Observations as a Standardised Method for Pollinator Monitoring: Opportunities and Limitations for Mass Participation Citizen Science.” *PLOS ONE* 11 (3): e0150794. <https://doi.org/10.1371/journal.pone.0150794>.
- Ryan, S. F., N. L. Adamson, A. Aktipis, L. K. Andersen, R. Austin, L. Barnes, M. R. Beasley, et al. 2018. “The Role of Citizen Science in Addressing Grand Challenges in Food and Agriculture Research.” *Proceedings of the Royal Society B: Biological Sciences* 285 (1891).

<https://doi.org/10.1098/rspb.2018.1977>.

Scott, Abigail L., Paul H. York, Clare Duncan, Peter I. Macreadie, Rod M. Connolly, Megan T.

Ellis, Jessie C. Jarvis, Kristin I. Jinks, Helene Marsh, and Michael A. Rasheed. 2018. “The Role of Herbivory in Structuring Tropical Seagrass Ecosystem Service Delivery.” *Frontiers in Plant Science* 9: 127. <https://doi.org/10.3389/fpls.2018.00127>.

Stoyanov, S. 2020. “Cranial Variability and Differentiation Among Golden Jackals (*Canis Aureus*) in Europe, Asia Minor and Africa.” *ZooKeys*. <https://doi.org/10.3897/zookeys.917.39449>.

Strydom, Tanya, Michael D. Catchen, Francis Banville, Dominique Caron, Gabriel Dansereau, Philippe Desjardins-Proulx, Norma R. Forero-Muñoz, et al. 2021. “A Roadmap Towards Predicting Species Interaction Networks (across Space and Time).” *Philosophical Transactions of the Royal Society B: Biological Sciences* 376 (1837): 20210063. <https://doi.org/10.1098/rstb.2021.0063>.

Tsunoda, Hiroshi, and Masayuki U. Saito. 2020. “Variations in the Trophic Niches of the Golden Jackal *Canis Aureus* Across the Eurasian Continent Associated with Biogeographic and Anthropogenic Factors.” *Journal of Vertebrate Biology* 69 (4): 20056.1. <https://doi.org/10.25225/jvb.20056>.

Valiente-Banuet, Alfonso, Marcelo A. Aizen, Julio M. Alcántara, Juan Arroyo, Andrea Cocucci, Mauro Galetti, María B. García, et al. 2015. “Beyond Species Loss: The Extinction of Ecological Interactions in a Changing World.” Edited by Marc Johnson. *Functional Ecology* 29 (3): 299–307. <https://doi.org/10.1111/1365-2435.12356>.

Windsor, Fredric M., Johan van den Hoogen, Thomas W. Crowther, and Darren M. Evans. 2022. “Using Ecological Networks to Answer Questions in Global Biogeography and Ecology.” *Journal of Biogeography* n/a (n/a). <https://doi.org/10.1111/jbi.14447>.

Wisz, Mary Susanne, Julien Pottier, W Daniel Kissling, Loïc Pellissier, Jonathan Lenoir, Christian F Damgaard, Carsten F Dormann, et al. 2013. “The Role of Biotic Interactions in Shaping Distributions and Realised Assemblages of Species: Implications for Species Distribution Modelling.” *Biological Reviews of the Cambridge Philosophical Society* 88 (1):

15–30. <https://doi.org/10.1111/j.1469-185X.2012.00235.x>.




Figure 1: (a) Spatial distribution of species richness according to the original IUCN range maps of all 32 mammal species of the Serengeti food web. (b) Proportion of mammal species remaining in each local network (i.e., each pixel) after removing all species without a path to a primary producer. (c) Proportion of mammal species remaining in each local network as a function of the number of species given by the original IUCN range maps.






Figure 2: Negative relationship between the out degree of predator species and their relative range loss. More specialized predators lose a higher proportion of their ranges due to mismatches with the ranges of their preys.




Figure 3: Geographical similarity between the original IUCN range maps of predators and preys. Dots represent predator-prey pairs, with different symbols corresponding to different predators. For a given pair of species, the number  $c$  of pixels where the focal species is present but not the other and the number  $a$  of pixels where the predator and prey cooccur, were calculated. Geographic similarities were given by  $a/(a + c)$ , with the predator being the focal species in the predator to prey similarity (x-axis), while the prey is the focal one in the prey to predator similarity (y-axis). One of the predators, *Canis aureus*, is not represented in the image because it is an extreme case (where all its range is suppressed by the absence of preys) and it would make the interpretation of the data more difficult.




Figure 4: Left panel: Distribution of the proportion of GBIF pixels (pixels with at least one occurrence in GBIF) falling into the IUCN range for different range sizes. Right panel: Differences between the proportion of GBIF pixels falling into the IUCN and the updated ranges for every predator species. Arrows go from the proportion inside the original range to the proportion inside the updated range, which can only be equal or lower. Overlapping markers indicate no difference between the types of layers. Species markers are the same on both figures, with predators presented in distinct colored markers and all herbivores grouped in a single grey marker. Pixels represent a resolution of 10 arc-minutes.




Figure 5: Mismatch between serval's range loss and GBIF occurrence of its prey. The left panel shows the reduction of serval's range when we consider the IUCN data on its prey. On the right panel, we added GBIF data on both serval and its prey, with a buffer for the prey to account for species mobility.