


Recap exercises

Excuse: An inherently nonlinear model


This is a model for death rates in certain situations:

$$y = \beta_0 + \lambda \exp(\beta_1^2 x) + \epsilon$$

It cannot be made linear in terms of parameters:



This can be verified using the TA plots:



Data wrangling

Read in the data set `wrangle_1.csv`. Transform the data set such that it can be considered tidy.

Then create a new data set that contains the means for all variables for each country. Missing values should be ignored when computing the means.

```
T4_df <- fread(here("data/wrangle_1.csv"), header = TRUE) %>%
  pivot_longer(
    cols = -all_of(c("country", "name")),
    names_to = "year",
    values_to = "value") %>%
  pivot_wider(names_from = "name", values_from = "value")
head(T4_df)
```

```
# A tibble: 6 x 5
  country year Growth EducationSpending HealthSpending
  <chr>   <chr>   <dbl>         <dbl>         <dbl>
1 Germany 2005   0.732          NA           10.3
2 Germany 2006   3.82           4.29          10.2
3 Germany 2007   2.98           4.37          10.1
```

4	Germany	2008	0.960	4.44	10.3
5	Germany	2009	-5.69	4.91	11.2
6	Germany	2010	4.18	4.94	11.1

```
T4_summary <- T4_df %>%
  group_by(country) %>%
  summarise(across(where(is.double), ~ mean(.x, na.rm=TRUE)))
T4_summary
```

```
# A tibble: 4 x 4
  country      Growth EducationSpending HealthSpending
  <chr>      <dbl>          <dbl>          <dbl>
1 Germany      1.17            4.79            11.0
2 Italy       -0.528            4.19             8.73
3 Netherlands  1.15             5.29            10.0
4 Spain        0.479            4.41             8.90
```


Linear regression

Consider the data set `reg_data_1.csv`. It contains the following variables:


- `y`: ice cream consumption in litres per year
- `x1`: Temperature in 10 degrees Celsius
- `x2`: Income in 1000 EUR
- `x3`: Height in cm

Study how ice cream consumption is associated with the explanatory variables and derive a sensitive linear regression model. Briefly justify your model specification.

```
dist_y <- ggplot(data = reg_data, mapping = aes(x=y)) +
  geom_histogram() + theme_bw()
dist_y
```



```
linlin_plot_x1 <- ggplot(data = reg_data, mapping = aes(x=x1, y=y)) +  
  geom_point() +  
  labs(title = "Original data") +  
  theme_bw()  
  
linlin_plot_x2 <- ggplot(data = reg_data, mapping = aes(x=x2, y=y)) +  
  geom_point() +  
  labs(title = "Original data") +  
  theme_bw()  
  
linlin_plot_x3 <- ggplot(data = reg_data, mapping = aes(x=x3, y=y)) +  
  geom_point() +  
  labs(title = "Original data") +  
  theme_bw()  
  
ggarrange(  
  dist_y, linlin_plot_x1, linlin_plot_x2,  
  linlin_plot_x3, ncol = 2, nrow = 2)
```



```
lm_correct <- lm(y~x1+I(x1**2)+log(x2), data = reg_data)
summary(lm_correct)
```

Call:

```
lm(formula = y ~ x1 + I(x1^2) + log(x2), data = reg_data)
```

Residuals:

Min	1Q	Median	3Q	Max
-0.032602	-0.007347	-0.000002	0.007745	0.026055

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	1.400e+01	7.435e-04	18832	<2e-16 ***
x1	1.499e+00	8.354e-04	1795	<2e-16 ***
I(x1^2)	1.500e+00	6.967e-04	2153	<2e-16 ***
log(x2)	2.199e+00	3.650e-04	6026	<2e-16 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 0.01008 on 496 degrees of freedom

Multiple R-squared: 1, Adjusted R-squared: 1

F-statistic: 2.121e+07 on 3 and 496 DF, p-value: < 2.2e-16

```
TA_correct <- ggplot(  
  tibble("Fitted"=lm_correct$fitted.values,  
         "Residuals"=lm_correct$residuals),  
  mapping = aes(x=Fitted, y=Residuals)  
  ) +  
  labs(title = "TA plot for correct specification") +  
  geom_point() +  
  theme_bw()  
TA_correct
```

