

Name: ADOO CHARLENE KUUKUA SAA

Id: 190951

Instructions:

- Make sure you are working on your problem set as each problem set is different.
- The answers to the questions of this problem set are to be given exclusively in the answer sheet
- The answers sheet MUST be printed and not photocopied. Photocopies will not be accepted.
- Questions marked with the symbol & admit more than one correct answer
- Please fill the boxes in the answer sheet completely using a black pen as follows

Question 1: B C D E

- The answer sheet must not be creased or folded otherwise your problem set won't be graded.
- You can hand back your problem set at the END of class on Friday, April 29th.



With a sample of 706 observations, we estimate the following model:

$$ln(hwage_i) = \beta_0 + \beta_1 age_i + \beta_2 age_i^2 + \beta_3 educ_i + \beta_4 yngkid_i + u_i$$

and obtain these results:

where lhwage is the logarithm of the hourly wage in euro, age is measured in years, educ is years of education and yngkid is a variable equal to 1 in case the person has a child younger than three years. **Question 1** What is the interpretation of β_4 ?

- A If a person has one small kid more, he/she earns about 9.5% more per hour with respect to someone who does not have small kids, ceteris paribus.
- B If a person has small kids (< 3 years old), he/she earns about 0.095 euros more per hour with respect to someone who does not have small kids, ceteris paribus.
- The first of the control of the cont
- D If a person has one small kid more, he/she earns about 0.095 more per hour with respect to someone who does not have small kids, ceteris paribus.

Question 2 What is the interpretation of β_2 ?


- A Increasing age by one year, the hourly wage decreases by 0.077% on average, ceteris paribus.
- B Increasing the square of age by one year, the hourly wage decreases by 0.00077 euros on average, ceteris paribus.
- C By itself does not have a proper interpretation.
- D Increasing the square of age by one year, the hourly wage decreases by 0.077% on average, ceteris paribus.

Question 3 \clubsuit Is β_3 statistically higher than 0.05 at 5%?

- A No, it is not since the t-value is larger than 1.96.
- B Yes, it is, since the t-value is larger than 1.64.
- C No, it is not, since the t-value is smaller than 1.96.
- D Yes, it is, since the t-value is smaller than 1.64.
- E None of these answers are correct.

Question 4 Is β_1 statistically significant?

- A It is at 10% level.
- B We cannot check for this, it makes no sense.
- C It is not at 1% level
- D It is not at 5% level.



$$\boxed{\mathbf{B}} \ ln(hwage_i) = \beta_0 + \beta_1 ln(age_i) + \beta_2 educ_i + u_i$$

$$\boxed{C}$$
 $ln(hwage_i) = \beta_0 + \beta_1 age_i + \beta_2 ln(educ_i) + u_i$

$$\boxed{D} ln(hwage_i) = \beta_0 + \beta_1 age + ln(\beta_2 educ_i) + u_i$$

$$\boxed{\text{E}} \ hwage_i = \beta_0 + \beta_1 age_i + \beta_2 ln(educ_i) + u_i$$

Question 6 Keeping other variables fixed, at what age the logarithm of hourly wage is maximized?

- At about 0, but this makes no sense.
- B At about 93.3 years.
- C At about 46.7 years.
- D At about 56.3 years.

Let us define with Y the amount of cholesterol in mlg in the blood and with Med a dummy variable which takes the value of 1 for medication B and 0 for medication A, where A and B are two different medications that lower cholesterol. Female is a dummy variable which takes the value of 1 for females and 0 otherwise.

Consider the following regression:

$$Y = \beta_0 + \beta_1 \times med + \beta_2 \times female + \beta_3 \times med \times female + u$$

Question 7 What is the effect of using medication B with respect to no medication for men?

- A None of the others.
- $\boxed{\mathbf{B}} \beta_0 + \beta_1$
- C β_1 - β_0
- $D \beta_0$
- $[E] \beta_1$

Question 8 Suppose you use this model: $Y = \beta_0 + \beta_1 \times med + \beta_2 \times female + u$ In this case, what is the effect of medication B for man and women, respectively?

- $\boxed{\mathbf{A}} \beta_1 \text{ and } \beta_1 + \beta_2.$
- B None of the others.
- C $\beta_0 + \beta_1$ and $\beta_0 + \beta_1 + \beta_2$.
- D β_1 for both genders.
- $[E] \beta_1 \beta_0$ for both genders.

These data are taken from the Medical Expenditure Panel Survey survey conducted in 1996. These data were provided by Professor Harvey Rosen of Princeton University and were used in his paper with Craig Perry "The Self-Employed Are Less Likely Than Wage-Earners to Have Health Insurance. So What?" in Douglas Holtz-Eakin and Harvey S. Rosen, eds., Entrepeneurship and Public Po licy, MIT Press 2004.

Among the variables in the dataset, ins is a dummy equal to one if the interviewee has the insurance; selfemp is equal to one if the interviewee is a self-employed workers; gender is equal to one if the in dividual is a male; married is one if the individual is married; health is one if the individual reports to be in good health; educ is 0 if the person has no education, 1 if he/she achieved middle school diploma, 2 for the high school diploma, 3 for the bachelor degree, 4 for the master degree and 5 for the PhD; age is in years and age2 is the square of age.

We estimate two models:

$$Pr(ins = 1|X) = \beta_0 + \beta_1 \times selfemp + \beta_2 \times married + \beta_3 \times gender + \beta_4 \times health + \beta_5 \times gender * health + \beta_6 \times educ + \beta_7 \times age + \beta_8 \times age^2$$

Coefficients:

	Estimate St	d. Error t	value H	Pr(> t)
(Intercept)	0.2974634	0.0580248	5.13	0.000003
selfemp	-0.1742361	0.0141740	-12.29	< 2e-16
married	0.1181062	0.0094187	12.54	< 2e-16
gender	-0.0232270	0.0343575	-0.68	0.49903
health	0.0744310	0.0247243	3.01	0.00262
genderxhealth	-0.0206248	0.0353131	-0.58	0.55920
educ	0.0529807	0.0029210	18.14	< 2e-16
age	0.0105315	0.0027482	3.83	0.00013
age2	-0.0000788	0.0000333	-2.37	0.01796

Heteroskadasticity robust standard errors used

$$Pr(ins = 1|X) = \Phi(\beta_0 + \beta_1 \times selfemp + \beta_2 \times married + \beta_3 \times gender + \beta_4 \times health + \beta_5 \times gender * health + \beta_6 \times educ + \beta_7 \times age + \beta_8 \times age^2)$$
 (II)

Coefficients:

	Estimate Std.	Error z v	alue Pro	(> z)
(Intercept)	-0.844932	0.195991	-4.31	0.000016
selfemp	-0.651923	0.046842	-13.92	< 2e-16
married	0.455241	0.034845	13.06	< 2e-16
gender	-0.040238	0.111653	-0.36	0.71856
health	0.300503	0.082988	3.62	0.00029
genderxhealth	n -0.124880	0.116613	-1.07	0.28422
education	0.226139	0.012852	17.60	< 2e-16
age	0.029150	0.009899	2.94	0.00323
age2	-0.000162	0.000126	-1.29	0.19821

Question 9 What is the interpretation of β_2 in model (II)?

- A On average, married individuals are 45.5% less likely than others to have an insurance, controlling for all other factors.
- B On average, a married worker has a probability of 45.5% to have an insurance, ceteris paribus.
- C It does not have a proper interpretation in terms of magnitude.
- D On average, married individuals are 45.5% more likely than others to have an insurance, controlling for all other factors.



Question 10 Is being married significantly linked to having an insurance under model (II)?

- A Yes, since the model includes the variable "married".
- \fbox{B} Yes, since the coefficient β_2 is significant.
- $\boxed{\mathbb{C}}$ No, since the coefficient β_2 is not significant.
- D It depends on the values of all other covariates.





Name: ADOO CHARLENE KUUKUA SAA Id: 190951

- Answers must be given exclusively on this sheet: answers given on the other sheets will be ignored.
- This sheet MUST be printed out and not photocopied. Photocopies will not be accepted.
- Please fill the boxes below completely using a black pen.
- Do not crease or fold.
- You can hand back your problem set by putting it into my mailbox on the fifth floor of the viale Romania campus by noon of Friday, March 25 at noon.

Question 1: A B C D

Question 2: A B C D

Question 3: A B C D E

Question 4: A B C D

Question 5: A B C D E

Question 6: A B C D

Question 7: A B C D E

Question 8: A B C D E

Question 9: A B C D

Question 10: A B C D