www.Padasalai.Net

Virudhunagar District
Common Quarterly Examination - 2023

www.CBSEtips.in

V9M

Standard 9 MATHEMATICS PART - 1

Marks: 100 14x1=14

I. Choose the correct answer:

1) The set  $P = \{ x \mid x \in \mathbb{Z}, -1 < x < 1 \}$  is a

a) singleton set b) Power set c) N

c) Null set

d) Subset

2) If  $A \cup B = A \cap B$  then

a) A≠B

b) A = B

c) ⋅ A ⊂ B

c) B ⊂ A

3) For any three sets A, B, C (A-B)  $\bigcap$  (B-C) is equal to ...........

a)  $\phi$ 

Time: 3.00 Hours

b) Conly

c) B only

d) A only

4) Which of the following is true?

a)  $A - B = A \cap B$ 

b) A - B = B - A

c)  $(A \cup B)' = A' \cup B'$ 

d)  $(A \cap B)' = A' \cup B'$ 

5) Which one of the following is an irrational number .........

a)  $\sqrt{25}$ 

b)  $\sqrt{\frac{9}{4}}$ 

c)  $\frac{7}{11}$ 

d) π

6)  $0.\overline{34} + 0.34 = \dots$ 

a) 0.687

b)  $0.\overline{68}$ 

c) 0.68

**d) 0.6**87

7)  $\sqrt{27} + \sqrt{12} = \dots$ 

a)  $5\sqrt{3}$ 

b)  $5\sqrt{6}$ 

c)  $\sqrt{39}$ 

d)  $3\sqrt{5}$ 

8)  $(0.000729)^{-\frac{3}{4}} \times (0.09)^{-\frac{3}{4}} = \dots$ 

a)  $\frac{10^{4}}{3^{3}}$ 

b)  $\frac{10^{6}}{3^{6}}$ 

c)  $\frac{10^{6}}{3^{5}}$ 

d)  $\frac{10}{3}$ 

9) 2x + 3 = 0 is the root of the polynomial equation

a)  $\frac{-3}{2}$ 

b)  $\frac{-2}{3}$ 

c)  $\frac{1}{3}$ 

d)  $\frac{-}{3}$ 

10) If P(a) = 0 then (x - a) is a .......... of P(x)

a) Remainder

b) factor

c) quotient

d) Divisor

Find the value of m from the equation 2x + 3y = m. If the solution is x = 2 and y = -2.

a) 10

b) 0

c) -2

d) 2

12) The exterior angle of a triangle is equal to the sum of two

a) Interrior opposite angles

b) Alternative anglesd) Interior angles

c) Exterior angles d) Interior and 13) If  $x^{40} + 40$  is divisible by x + 1, then the remainder is

a) 39

b) 40

c) -40

d) 41

14) For any set A

 $A \cup A = A \& A \cap A = A$ 

b) Identity Law

a) Idempotent Lawsc) De Morgan's Law

d) Symmetric Law

PART - II

II. Answer any 10 questions: (Q.No. 28 is compulsory)

10x2=20

15) Find the number of subsets and the number of proper subsets of a set X = {a, b, c, x, y, z}

16)  $X = \{5, 6, 7\} Y = \{5, 7, 9, 10\}$ . Find the symmetric difference between the sets.

17) A={b, e, f, g} and B ={c, e, g, h}, then verify the commutative property of union of sets

18) If n(A) = 25, n(B) = 40  $n(A \cup B) = 50$  and n(B') = 25 find  $n(A \cap B)$  and n(U)


19) Find any two rational number between  $\frac{1}{2}$  and  $\frac{2}{3}$ 

20) Express the following decimal expression into rational numbers  $0.\overline{24}$ 

21) Give any two rational numbers lying between 0-51-511115 and 0-535335335 ...

VSIVI 22) MWW Radasalai Net 3/16 www.CBSEtips.in

- 23) Find the roots of the polynomial equation 5x 3 = 0
- 24) Factorise :  $8x^3 + 27y^3 + 64z^3 72 xyz$
- 25) The base of the parallelogram is (5x + 4)Find its height, if the area is  $25x^2 - 16$ .
- 26) Evaluate  $10^3 15^3 + 5^3$ .
- 27) In the figure AB is parallel to CD, find x.
- 28) Find G.C.D  $(y^3 + 1)$  and  $(y^{3-1})$ .




10x5=50

## PART - III

## III. Answer any 10 questions: (Q.No. 42 is compulsory)

29) Verify  $(A \cup B)' = A' \cap B'$  Using Venn diagrams.

- 30) In a group of 100 students, 85 students speak Tamil, 40 students speak English, 20 students speak French, 32 speak Tamil and English, 13 speaks English and French and 10 speak Tamil and French. In each student knows atleast any one of these languages then find the number of students who speak all these three languages.
- 31) If  $A = \{0, 2, 4, 6, 8\}$   $B = \{x : x \text{ is a prime number and } x < 11\}$  and  $C = \{x : x \in N\}$ and  $5 \le x < 9$ } then verify  $A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$
- 32) If  $A = \{x: x = 6n \ n \in W \ and \ n < 6\} B = \{x: x = 2n \ n \in N \ and \ 2 < n \le 9\} \ and \ C = \{x: x = 2n \ n \in N \ and \ 2 < n \le 9\} \ and \ C = \{x: x = 2n \ n \in N \ and \ 2 < n \le 9\} \ and \ C = \{x: x = 2n \ n \in N \ and \ 2 < n \le 9\} \ and \ C = \{x: x = 2n \ n \in N \ and \ 2 < n \le 9\} \ and \ C = \{x: x = 2n \ n \in N \ and \ 2 < n \le 9\} \ and \ C = \{x: x = 2n \ n \in N \ and \ 2 < n \le 9\} \ and \ C = \{x: x = 2n \ n \in N \ and \ 2 < n \le 9\} \ and \ C = \{x: x = 2n \ n \in N \ and \ 2 < n \le 9\} \ and \ C = \{x: x = 2n \ n \in N \ and \ 2 < n \le 9\} \ and \ C = \{x: x = 2n \ n \in N \ and \ 2 < n \le 9\} \ and \ C = \{x: x = 2n \ n \in N \ and \ 2 < n \le 9\} \ and \ C = \{x: x = 2n \ n \in N \ and \ 2 < n \le 9\} \ and \ C = \{x: x = 2n \ n \in N \ and \ 2 < n \le 9\} \ and \ C = \{x: x = 2n \ n \in N \ and \ 2 < n \le 9\} \ and \ C = \{x: x = 2n \ n \in N \ and \ 2 < n \le 9\} \ and \ C = \{x: x = 2n \ n \in N \ and \ 2 < n \le 9\} \ and \ C = \{x: x = 2n \ n \in N \ and \ 2 < n \le 9\} \ and \ C = \{x: x = 2n \ n \in N \ and \ 2 < n \le 9\} \ and \ C = \{x: x = 2n \ n \in N \ and \ 2 < n \le 9\} \ and \ C = \{x: x = 2n \ n \in N \ and \ 2 < n \le 9\} \ and \ C = \{x: x = 2n \ n \in N \ and \ 2 < n \le 9\} \ and \ C = \{x: x = 2n \ n \in N \ and \ 2 < n \le 9\} \ and \ C = \{x: x = 2n \ n \in N \ and \ 2 < n \le 9\} \ and \ C = \{x: x = 2n \ n \in N \ and \ 2 < n \le 9\} \ and \ C = \{x: x = 2n \ n \in N \ and \ 2 < n \le 9\} \ and \ C = \{x: x = 2n \ n \in N \ and \ 2 < n \le 9\} \ and \ C = \{x: x = 2n \ n \in N \ and \ 2 < n \le 9\} \ and \ C = \{x: x = 2n \ n \in N \ and \ 2 < n \le 9\} \ and \ C = \{x: x = 2n \ n \in N \ and \ 2 < n \le 9\} \ and \ C = \{x: x = 2n \ n \in N \ and \ 2 < n \le 9\} \ and \ C = \{x: x = 2n \ n \in N \ and \ 2 < n \le 9\} \ and \ C = \{x: x = 2n \ n \in N \ and \ 2 < n \le 9\} \ and \ C = \{x: x = 2n \ n \in N \ and \ 2 < n \le 9\} \ and \ C = \{x: x = 2n \ n \in N \ and \ 2 < n \le 9\} \ and \ C = \{x: x = 2n \ n \in N \ and \ 2 < n \le 9\} \ and \ C = \{x: x = 2n \ n \in N \ and \ 2 < n \le 9\} \ and \ C = \{x: x = 2n \ n \in N \ and \ 2 < n \le 9\} \ and \ C = \{x: x = 2n \ n \in N \ and \ 2 < n \le 9\} \ and \ C = \{x: x = 2n \ n \in N \ and \ 2 < n \le 9\} \ and \$  $\{x: x = 3n, n \in \mathbb{N} \text{ and } 4 \le n < 10\}$  then show that  $A - (B \cap C) = (A - B) \cup (A - C)$ 33) Convert the decimal numbers in the form of  $\frac{p}{q}$  (p,q \in z and q \neq 0)  $0.5\overline{68}$
- 34) Represent  $3.\overline{45}$  on the number line upto 4 decimal places.
- 35) Arrange in ascending order:  $\sqrt[3]{2}$ ,  $\sqrt[3]{4}$ ,  $\sqrt[4]{3}$
- 36) Simplify:  $(7\sqrt{a} 5\sqrt{b})(7\sqrt{a} + 5\sqrt{b})$
- 37) If  $\left(y \frac{1}{y}\right)^3 = 27$  then find the value of  $y^3 \frac{1}{y^3}$
- 38) Find quotient and the remainder when f(x) is divided by g(x)i)  $f(x) = (8x^3 - 6x^2 + 15x - 7) g(x) = 2x + 1$ .
- 39) Is (3x 2) a factor of  $3x^3 + x^2 20x + 12$ ?
- 40) If the quotient on dividing  $x^4+10x^3+35x^2+50x+29$  by (x+4) is  $x^3-ax^2+bx+6$ , then find the value of a,b and also remainder.
- 41) Find all the three angles of the  $\triangle$ ABC



42) Expand: (x-12)(x+7)(x-18)

## PART - IV

## IV. Answer both questions:

2x8 = 16

- 43) a) Draw the  $\triangle$  ABC where AB = 6cm  $B = 110^{\circ}$  and AC = 9cm and construct the centroid. (OR)
  - b) Draw an equilateral triangle of sides 6.5cm and locate its Orthocentre.
- 44) a) Let  $U = \{0, 1, 2, 3, 4, 5, 6, 7\}$  A= $\{1, 3, 5, 7\}$  and B= $\{0, 2, 3, 5, 7\}$  find the following sets i) A' ii) B' iii)  $A' \cup B'$  iv)  $A' \cap B'$  v)  $(A \cup B)'$  vi)  $(A \cap B)'$ vii) (A)' viii) (B')'

(OR)

- b) Factorise the following:
  - i)  $x^2 + 10x + 24$

iii) a<sup>2</sup> + 10a -600 ii) a<sup>2</sup> + 10a -600 kindiy sendeme your study materials to padagalai ne god medi 2 yom