

YIELD LOSS DUE TO CROP DISEASES AND PEST INFESTATION

A Comprehensive Statistical and Technical Documentation
Global and Regional Data, Year-wise Outbreaks, Economic Losses,
Causal Organisms, and Recommended Remedial Strategies

Compiled:	September 2026
Scope:	Global overview with focused regional case studies (South Asia, Sub-Saharan Africa, Latin America)
Coverage period:	Historical outbreaks (1840s) through 2026
Primary sources:	FAO, ICAR, CABI, ICO, CIMMYT, USAID, peer-reviewed journals, government records

Note: Figures in this document are drawn from publicly available reports by the Food and Agriculture Organization (FAO), the Indian Council of Agricultural Research (ICAR), CABI, the International Coffee Organization (ICO), CIMMYT, USAID and government/parliamentary records, supplemented by peer-reviewed studies. Estimates of yield loss vary by methodology, season and source; ranges are given wherever the underlying reports diverge, and every figure is attributed to its origin.

TABLE OF CONTENTS

1. Executive Summary
2. Introduction
3. Global Statistical Overview of Yield Loss
4. Major Crop Diseases — Year-wise Case Studies and Remedies
 - 4.1 Wheat Diseases (Rusts, Blast, Karnal Bunt)
 - 4.2 Rice Diseases (Blast, Bacterial Blight, Sheath Blight, Tungro)
 - 4.3 Potato Late Blight and the Historical Irish Famine
 - 4.4 Coffee Leaf Rust (La Roya)
 - 4.5 Banana Fusarium Wilt (Panama Disease TR4)
 - 4.6 Citrus Greening (Huanglongbing)
 - 4.7 Cotton Diseases and Other Field-Crop Diseases
5. Major Pest Infestations — Year-wise Case Studies and Remedies
 - 5.1 Fall Armyworm (*Spodoptera frugiperda*)
 - 5.2 Desert Locust (*Schistocerca gregaria*)
 - 5.3 Pink Bollworm and Cotton Pest Complex (India)
 - 5.4 Brown Planthopper and Rice Pest Complex
 - 5.5 Plant-Parasitic Nematodes
 - 5.6 Fruit Flies, Aphids, Whiteflies and Other Sucking Pests
6. Regional Case Studies
 - 6.1 India — National Loss Estimates
 - 6.2 Sub-Saharan Africa
 - 6.3 Central America and the Coffee Belt
 - 6.4 South Asia — Transboundary Threats
7. Economic Impact Analysis
8. Climate Change and the Changing Geography of Pests and Diseases
9. Integrated Pest and Disease Management (IPDM) — Remedies in Detail
 - 9.1 Cultural and Agronomic Control
 - 9.2 Host-Plant Resistance and Breeding
 - 9.3 Biological Control
 - 9.4 Chemical Control and Judicious Pesticide Use
 - 9.5 Digital Agriculture, Early Warning and Forecasting Systems
10. Policy Frameworks and Institutional Response
11. Recommendations
12. Conclusion
13. References and Data Sources

1. Executive Summary

Plant pests and diseases remain one of the single largest, most persistent threats to global food security. The Food and Agriculture Organization of the United Nations (FAO) estimates that **between 20 and 40 percent of global crop production is lost every year** to plant pests and diseases, with associated trade losses exceeding **USD 220 billion annually** and invasive pests alone causing at least **USD 70 billion** in global economic damage each year. In India, government and ICAR estimates converge on a comparable figure: crop losses from pests, diseases, weeds and rodents range between **10 and 30 percent** of potential production annually, valued at roughly **Rs 50,000–90,000 crore (USD 6–14 billion)** depending on the year and the crops assessed.

This documentation compiles year-wise, disease-wise and pest-wise loss statistics drawn from FAO, ICAR, CABI, the International Coffee Organization (ICO), CIMMYT, USAID and peer-reviewed literature. It covers major outbreaks across cereals (wheat, rice, maize), cash crops (cotton, coffee, banana, citrus) and vegetables (potato), spanning historical events such as the Irish Potato Famine (1845–1852) to contemporary crises such as the Fall Armyworm invasion of Africa (2016–present), the Desert Locust upsurge of 2019–2021, wheat blast in Bangladesh (2016), and pink bollworm resurgence in Indian cotton (2017–2022). For each major disease and pest, the document records the causal organism, affected crop, geography, year of outbreak, quantified yield or economic loss, and the recommended remedy — spanning cultural, biological, chemical, genetic and digital control measures assembled under Integrated Pest and Disease Management (IPDM).

Key findings include: (a) invasive transboundary pests such as fall armyworm and desert locust cause disproportionate damage because affected regions lack prior exposure, resistant germplasm or established natural enemies; (b) climate change is expanding the geographic range and seasonal window of many pests and pathogens, including wheat rusts, fall armyworm and locusts; (c) resistance breakdown — as seen with Ug99 wheat stem rust and Bt-resistant pink bollworm — repeatedly erodes genetic and biotechnological control gains, making continuous breeding and resistance-management essential; and (d) Integrated Pest Management (IPM), early warning systems, and coordinated international response (such as FAO's Global Action for Fall Armyworm Control and the Desert Locust response of 2020–2021) can avert millions of tonnes of loss when funded and mobilised promptly — the 2020–2021 desert locust response alone is credited with averting an estimated 4.5 million tonnes of cereal loss worth approximately USD 1.77 billion.

2. Introduction

Agricultural production sits at the intersection of biology, climate and economics. Every crop grown by humans is simultaneously a food source for thousands of co-evolved and invasive fungi, bacteria, viruses, nematodes, insects, mites and rodents. Yield loss — the gap between the theoretically attainable yield of a crop under a given set of environmental conditions and the yield actually harvested — is driven in large part by this biotic pressure. Humans depend on plants for roughly 80 percent of the food they eat and 98 percent of the oxygen they breathe, which is why the FAO designated plant health a foundational pillar of food security when it established the International Plant Protection Convention (IPPC) framework and marked 2020 as the International Year of Plant Health.

Losses from pests and diseases are typically classified into three categories: (i) **pre-harvest losses**, occurring in the field during crop growth due to pathogen infection, insect feeding, or nematode damage; (ii) **post-harvest losses**, occurring during storage, transport and processing due to storage pests, moulds and rodents; and (iii) **quality losses**, where the crop survives but its market value is degraded (for example, mycotoxin contamination in maize, or pest-damaged cotton lint). This document focuses primarily on pre-harvest yield losses caused by diseases and pest infestation, while referencing post-harvest and quality impacts where the source data reports them.

Understanding these losses matters for four practical reasons. First, food security: with the world population projected to approach 9.7 billion by 2050 and global food demand expected to rise by 59–98 percent over 2050 levels, closing the yield gap caused by pests and diseases is one of the most cost-effective ways to expand food supply without expanding cultivated area. Second, farmer livelihoods: smallholder farmers, who produce a large share of the food consumed in developing countries, are disproportionately exposed to catastrophic loss from a single outbreak. Third, trade and macroeconomic stability: outbreaks such as coffee leaf rust in Central America or pink bollworm in Indian cotton move global commodity prices and export volumes. Fourth, environmental sustainability: over-reliance on chemical pesticides in response to pest pressure carries its own costs in soil health, pollinator decline, and human health, making evidence-based Integrated Pest Management essential rather than optional.

The sections that follow present statistics organised first globally, then by individual disease and pest (with year, location, quantified loss and remedy), then by region, and finally as a consolidated discussion of management strategies, policy response and recommendations.

3. Global Statistical Overview of Yield Loss

The table below summarises headline global and national statistics on crop loss attributable to pests, diseases, weeds and rodents, as published by FAO, ICAR, CABI and allied bodies over the past two decades. Figures differ by methodology (attainable-yield modelling vs. field-survey vs. expert elicitation) and by whether weeds and post-harvest losses are included, which is why ranges rather than single numbers are common in the literature.

Year / Period	Scope	Reported Loss	Source
Recurring (FAO estimate)	Global, all crops, pests+diseases only	20% – 40% of annual crop yield	FAO Plant Production & Protection Division
Recurring (FAO/OECD)	Global, pests+diseases+weeds	26% – 40% of potential production (could double without crop protection)	FAO–OECD / CropLife International
Recurring	Global trade losses from plant pests	USD 220 billion/year	FAO, International Day of Plant Health briefings
Recurring	Global economic loss from invasive pests specifically	at least USD 70 billion/year	FAO One Health / Plant Health programme
2007	India, all crops (pest-associated)	10% – 30% of production, valued at Rs 1,400 billion (~USD 17 bn at 2007 rates); pesticide cost Rs 68 billion	ICAR – National Agricultural Innovation Project (NAIP)
2009	India, all crops (CARE report cited in Parliament)	~18% of crop value, Rs 90,000 crore	CARE Ratings, cited in Lok Sabha reply (14 Jul 2009)
2006 & recurring	India, all crops	10% – 30% of annual production depending on season/severity	Ministry of Agriculture, Lok Sabha replies (2006)
Recent (2018–2023)	India, all crops (ICAR-AICRP)	30% – 35% of annual yield to pests; nematodes alone account for ~60 million tonnes (10–12% of production), valued at Rs 21,068 million across 24 major crops	ICAR – AICRP on Nematodes / P.K. Chakrabarty, ADG Plant Protection
Recent estimate	India, all crops (industry study)	Rs 500 billion (~USD 8 billion) lost annually	Industry study cited in Business Standard / pesticide-sector press
2025	Global hunger context	~645 million people affected by hunger	FAO, One Health highlight (2026)

Table 3.1 — Headline global and national (India) crop loss statistics attributable to pests and diseases.

3.1 Why Estimates Vary

Three structural reasons explain the spread in reported figures. First, attainable-yield baselines differ: some studies model the yield a crop would achieve with zero biotic stress under ideal management, while others compare against realistic on-farm best-practice yields, producing systematically different percentages for the same underlying damage. Second, scope differs: FAO's headline 20–40% figure includes weeds in some citations and excludes them in others, and post-harvest losses are sometimes bundled with pre-harvest field losses. Third, crop and regional composition differs: national aggregates blend high-loss crops (pulses, oilseeds, vegetables) with lower-loss cereals under intensive management, so the same country can report anywhere from 10% to 35% loss depending on which crop basket and which year's climatic severity is being described.

3.2 Loss by Crop Category (Illustrative Global Ranges)

Crop Category	Typical Pre-harvest Loss Range	Dominant Biotic Stress
Cereals (wheat, rice, maize)	10% – 25%	Rusts, blast, blights, stem borers, armyworms, aphids
Pulses and oilseeds	15% – 30%	Pod borers, wilts, aphids, whiteflies
Cotton and fibre crops	20% – 50% (outbreak years higher)	Bollworms (pink, American), whitefly, jassids
Vegetables	20% – 40%	Late blight, fruit borers, nematodes, viral complexes
Fruits (citrus, banana)	Up to 100% at orchard level in epidemic zones	Huanglongbing, Fusarium wilt (TR4), fruit flies
Plantation crops (coffee)	20% – 50% in epidemic years	Coffee leaf rust, coffee berry borer

Table 3.2 — Illustrative pre-harvest loss ranges by crop category, compiled from FAO, ICAR and crop-specific studies cited throughout this document.

4. Major Crop Diseases — Year-wise Case Studies and Remedies

This section documents the most economically significant crop diseases worldwide, organised by crop. For each disease, the causal organism, representative outbreak years, quantified loss, and current recommended remedy are presented in summary tables followed by narrative context.

4.1 Wheat Diseases — Rusts, Blast and Karnal Bunt

Wheat (*Triticum aestivum*), grown on over 217 million hectares globally, is threatened by three rust diseases (stem/black rust — *Puccinia graminis* f. sp. *tritici*; stripe/yellow rust — *P. striiformis*; and leaf/brown rust — *P. triticea*), by wheat blast (*Magnaporthe oryzae* pathotype *Triticum*), and by Karnal bunt (*Tilletia indica*), among others.

Year	Disease	Location	Loss Reported	Remedy
1998 onward	Ug99 stem rust (new virulent race)	First detected Uganda; spread to East Africa, Yemen, Iran	Historically, stem rust epidemics have caused up to 100% loss in susceptible fields; Ug99 defeated most resistance genes deployed at the time	Deployment of Sr-gene resistant varieties (e.g., via the Borlaug Global Rust Initiative), fungicide seed/foliar treatment, international surveillance
Recurring	Stripe/yellow rust	South Asia, East Africa, temperate wheat belts	Can cause 10%-70% yield loss in susceptible varieties under favourable cool, moist conditions	Resistant cultivars, triazole fungicides, early sowing to escape peak disease pressure
2016	Wheat blast (first Asian outbreak)	Eight south-western districts of Bangladesh (Meherpur, Kushtia, Chuadanga, Pabna, Jhenaidah, Jessore, Barisal, Bhola)	Affected >15,000 ha; localized yield loss up to 90-100%; average district loss ~51%; total production loss ~8,205 tonnes worth USD 2.1 million	Fungicide (triazole+strobilurin) application at flowering, avoidance of late sowing, quarantine on seed movement; no fully resistant variety yet available
Projected risk	Wheat blast — South Asia spread risk	6.9 million ha of vulnerable wheat area across Bangladesh, India, Pakistan	A modelled 5% severity scenario projects potential loss of 886,000 tonnes worth USD 132 million in a single year	Early-warning weather-based forecasting, restricted wheat cultivation near rice fallows in high-risk windows
2020-21 & 2021-22 (ICAR-IA RI study)	Aphid complex + Karnal bunt + black point	New Delhi, India (variety HD 3086)	Avoidable yield loss of 6.1%-7.8% under higher aphid pressure; 4.8%-6.9% under lower pressure	Imidacloprid-based aphid management, timely sowing, resistant/tolerant varieties

Table 4.1 — Wheat disease outbreaks, losses and remedies.

Stem rust historically devastated wheat crops before the Green Revolution; the emergence of race Ug99 in 1998 in Uganda was globally significant because it overcame Sr31, a resistance gene deployed in the majority of the world's wheat varieties at the time, prompting the international Durable Rust Resistance in Wheat (DRRW) and Borlaug Global Rust Initiative programmes. Wheat blast is a newer and arguably more alarming threat: it jumped continents from South America to Bangladesh in 2016, and no wheat variety currently carries full resistance, making fungicide timing and cropping-calendar adjustments the primary lines of defence.

4.2 Rice Diseases — Blast, Bacterial Blight, Sheath Blight and Tungro

Disease	Causal Organism	Typical Loss	Remedy
Rice blast	Magnaporthe oryzae (Pyricularia oryzae)	Considered the most destructive rice disease worldwide; capable of 10%-30% average loss and up to 100% in severe neck-blast epidemics; estimated to destroy enough rice annually to feed 60 million people	Resistant varieties (Pi-gene stacking), tricyclazole/azoxystrobin fungicide, balanced nitrogen fertilisation, seed treatment
Bacterial leaf blight	Xanthomonas oryzae pv. oryzae	10%-20% average loss; up to 50%-70% in epidemic years on susceptible varieties, especially after storm damage	Xa-gene resistant varieties, avoiding excess nitrogen, field sanitation, copper-based bactericides, balanced irrigation
Sheath blight	Rhizoctonia solani	5%-50% depending on canopy density and nitrogen use; a major constraint in high-input irrigated rice	Wider spacing, silicon fertilisation, validamycin/hexaconazole fungicide, resistant lines
Rice tungro disease	Rice tungro bacilliform + spherical viruses, transmitted by green leafhopper	Can cause complete crop failure in hopper-heavy seasons; historically severe in Philippines/South-East Asia	Vector (leafhopper) control, tungro-tolerant varieties, synchronized planting

Table 4.2 — Major rice diseases, typical losses and remedies.

Rice, as the staple food of more than half of humanity, carries unusually high stakes when disease strikes. Blast disease alone is estimated by international rice research bodies to destroy a volume of rice sufficient to feed tens of millions of people annually, and its ability to evolve new races that defeat single-gene resistance means breeding programmes must continually pyramid multiple resistance genes to stay ahead of the pathogen.

4.3 Potato Late Blight and the Historical Irish Famine

Late blight, caused by the oomycete *Phytophthora infestans*, remains the most important disease of potato worldwide and is historically significant as the trigger of the Great Irish Famine.

Year	Event	Location	Impact
1845-1852	Irish Potato Famine (An Gorta Mor)	Ireland	Potato blight destroyed a large share of Ireland's staple crop across successive seasons; roughly one million people died of starvation and disease, and roughly one million more emigrated, reducing Ireland's population by about 20-25%
Recurring (modern era)	Late blight epidemics	Global potato-growing regions, especially cool humid zones	Without fungicide protection, late blight can destroy an entire potato crop within 1-2 weeks under favourable cool, wet weather; global crop-protection spend on blight control is among the highest of any single plant disease

Table 4.3 — Potato late blight: historical and contemporary impact.

Remedy: Certified disease-free seed potatoes, resistant/tolerant cultivars, protectant and systemic fungicides (mancozeb, metalaxyl, cymoxanil-based products) applied on a forecasting-model schedule (e.g., Blitecast, Hutton Criteria), destruction of volunteer plants and cull piles that harbour inoculum between seasons, and wider crop rotation.

4.4 Coffee Leaf Rust (La Roya)

Caused by the fungus *Hemileia vastatrix*, coffee leaf rust triggered the most severe regional coffee crisis of the 21st century when it swept Central America and Mexico starting in the 2011-2012 season.

Year	Region	Extent	Economic/Social Loss	Remedy
2012-13	Central America (Guatemala, Honduras, Costa Rica, El Salvador, Nicaragua) + Mexico	53% average incidence across the region; up to 70% of planted area affected in worst-hit countries	~2.5-3 million 60-kg bags of coffee lost in 2012/13 alone; USD ~499-500 million in foreign-currency losses; ~374,000-437,000 rural jobs at risk	Emergency ICO/national phytosanitary response, fungicide (copper + triazole) spraying, shade management
2011-2016 (cumulative epidemic)	Central America region-wide	Affected 70% of farms across the region	Over 1.7 million coffee workers lost jobs; USD 3.2 billion in cumulative damage and lost income over the epidemic	Replanting with rust-resistant varieties (e.g., Lempira, Catimor derivatives), farmer training manuals (WCR/CATIE), improved agronomy and fertilisation

Table 4.4 — Coffee leaf rust outbreak (2011-2016), Central America.

Because coffee is a perennial crop, rust damage does not merely reduce a single season's harvest — defoliated bushes require one to three years to recover full bearing capacity, and severely affected smallholders often cannot afford replanting, leading to permanent exit from coffee farming. Climate change (rising minimum temperatures at higher elevations) is cited as a contributing factor in the rust's spread into previously rust-free high-altitude zones.

4.5 Banana Fusarium Wilt (Panama Disease, Tropical Race 4)

Fusarium wilt of banana, caused by the soil-borne fungus *Fusarium oxysporum* f. sp. *cubense*, nearly wiped out the export banana industry once before (the Gros Michel cultivar collapse of the mid-20th century, caused by Race 1) and is now threatened again by Tropical Race 4 (TR4), which attacks the Cavendish cultivar that replaced Gros Michel and today accounts for roughly half of global banana production.

Period	Strain	Spread	Impact	Remedy
1950s-1960s	Race 1	Central America, global Gros Michel plantations	Effectively ended commercial viability of the Gros Michel cultivar; industry-wide replacement by Cavendish	Cultivar replacement (no cure exists for Fusarium wilt once soil is infested)
1990s-present (TR4 first found in Taiwan, 1960s; global spread accelerating since 1990s)	Tropical Race 4 (TR4)	South-East Asia, Australia, Middle East, Mozambique (2013), Colombia (2019, first Latin American detection), Peru (2021)	Can cause total loss of infected plantations; no chemical or biological cure; infested soil remains unusable for susceptible cultivars for decades	Strict phytosanitary quarantine, disinfection protocols for farm equipment/footwear, use of TR4-tolerant cultivars (e.g., GCTCV-derived lines), diversification away from Cavendish monoculture

Table 4.5 — Banana Fusarium wilt (Panama disease), historical and current strain impact.

4.6 Citrus Greening (Huanglongbing, HLB)

Huanglongbing (HLB), associated with the bacterium *Candidatus Liberibacter* spp. and transmitted by the Asian citrus psyllid (*Diaphorina citri*), is considered the most destructive citrus disease in the world. Since its detection in Florida in 2005, HLB has been linked to a decline of Florida orange production from roughly 240 million boxes in the early 2000s to under 20 million boxes by the early 2020s — a drop of more than 90 percent over two decades — with the state citrus industry's annual economic contribution falling by billions of dollars. **Remedy:** aggressive psyllid vector control (systemic and foliar insecticides), removal of infected trees, use of disease-free certified nursery stock, thermal and nutritional therapy to prolong productive life of infected trees, and long-term

breeding for HLB-tolerant rootstocks and scions.

4.7 Cotton and Other Field-Crop Diseases

Crop/Disease	Organism	Typical Loss	Remedy
Cotton bacterial blight (Angular leaf spot)	Xanthomonas citri pv. malvacearum	10%-30% in susceptible varieties in humid seasons	Resistant varieties, acid-delinted/treated seed, crop rotation, avoiding overhead irrigation
Maize/rice grain mould & mycotoxin complex (Aflatoxin, Fumonisin)	Aspergillus flavus, Fusarium spp.	Quality/marketability loss can approach total crop rejection in contaminated lots; major post-harvest concern in humid storage	Timely harvest, proper drying (<14% moisture), aflatoxin-resistant hybrids, biocontrol (Aflasafe-type atoxigenic strains)
Groundnut/pulse wilt complex	Fusarium spp., Sclerotium rolfsii	10%-40% in continuous mono-cropped fields	Crop rotation, Trichoderma-based seed treatment, resistant varieties
Downy mildew (pearl millet, maize)	Sclerospora graminicola / Peronosclerospora spp.	Localized epidemics up to 30%-40% loss	Metalaxyl seed treatment, resistant hybrids, roguing infected plants early

Table 4.6 — Other economically important field-crop diseases.

5. Major Pest Infestations — Year-wise Case Studies and Remedies

5.1 Fall Armyworm (*Spodoptera frugiperda*)

Native to the Americas, fall armyworm (FAW) was first confirmed in Africa in January 2016 in south-western Nigeria. Within roughly two years it had spread to more than 40 African countries and subsequently to India (2018), South-East Asia, China and Australia, becoming one of the fastest-spreading invasive insect pests on record. It feeds on more than 80 plant species but strongly prefers maize, Africa's most important staple.

Year	Geography	Reported/Projected Loss	Source
2017 (CABI evidence note)	12 major African maize-producing countries	Potential maize yield loss of 8.3-20.6 million tonnes/year (21%-53% of annual production) if unmanaged; value USD 2,481-6,187 million/year	CABI 'Fall Armyworm: Impacts and Implications for Africa' (DFID-commissioned)
2018	12 African countries (post-outbreak analysis)	Cost farmers 40% of harvests in 2018; potential maize losses up to 17.7 million tonnes/year; economic value USD 2.5-6.3 billion/year for maize alone (averaging ~USD 4.6 bn); plus USD 6.7 bn (rice), USD 2.8 bn (sugarcane), USD 0.8 bn (sorghum)	CABI / Impakter analysis
2018	Ghana	98% of surveyed farmers reported maize affected; ~USD 117 million in crop loss	CABI country assessment
2018	Zambia	98% of surveyed farmers reported maize affected; ~USD 159 million in crop loss	CABI country assessment
2017-18 season	Uganda	450,000 tonnes of maize destroyed, worth USD 192.8 million (~12% of national yield)	Uganda Ministry of Agriculture
2017-18 season	Kenya	>250,000 hectares attacked, ~11% of national maize crop affected	Kenya Ministry of Agriculture
2017-18 season	Tanzania / Ethiopia	Estimated yield losses of 3,238,980 tonnes (Tanzania) and 3,054,727 tonnes (Ethiopia); Uganda 1,391,109 tonnes	CABI/regional press analysis
2021 (comprehensive CABI/University of Minnesota study)	Africa-wide, all invasive alien species	Total invasive-species yield loss of USD 65.58 billion/year; fall armyworm alone responsible for the single highest share at USD 9.4 billion/year	CABI & University of Minnesota (Senay, Pardey et al.)
2018 onward	India	First reported July 2018 in Karnataka; spread to most maize-growing states within a year, prompting national advisory-based IPM response	ICAR-NBAIR national pest surveillance

Table 5.1 — Fall armyworm outbreak chronology and quantified losses, 2016-2021.

Remedy: FAO launched a USD 500 million Global Action for Fall Armyworm Control in December 2019, a three-year (2020-2022) coordination initiative. Field-level control combines early scouting (egg mass and early-instar detection), biopesticides (*Bacillus thuringiensis*-based products, entomopathogenic fungi/viruses), selective insecticides (emamectin benzoate, chlorantraniliprole) applied against young larvae before they bore into the whorl, push-pull companion cropping (e.g., *Desmodium* intercropping with Napier grass border, developed originally for stemborer control and now validated against FAW), conservation of natural enemies (parasitoid wasps, predatory ants), and community-based Farmer Field School training for early detection.

5.2 Desert Locust (*Schistocerca gregaria*)

The 2019-2021 desert locust upsurge was the worst to hit East Africa and Yemen in 25 years, and the worst in Kenya specifically in 70 years, driven by an unusual sequence of cyclones and heavy rains in the Arabian Peninsula that allowed several uncontrolled breeding cycles before detection.

Year	Milestone	Figures
Jan 2020	GIEWS Special Alert issued	Worst outbreak in 25 years across Ethiopia, Somalia, Kenya; swarms up to 60 km long and 40 km wide reported in Kenya
By Apr 2020 (Ethiopia)	National impact assessment	1.35 million ha of pasture and 197,163 ha of cropland affected; ~3.5 million quintals of cereal lost by March 2020; 2020/21 agricultural output forecast to fall 8% (~26.8 million quintals, enough to feed 12 million people for a year)
2020 (World Bank estimate)	Regional economic projection	Potential damages and losses of up to USD 8.5 billion across East Africa and Yemen by end of 2020
Jan 2020-Jun 2020 (early response)	FAO control operations	759,718 ha treated; 1.52 million tonnes of crop loss averted by June harvest; USD 456 million worth of crop secured; 9.87 million people met annual cereal needs as a result; 684,000 pastoral households' livestock fed
Jan 2020-Dec 2021 (full response)	FAO cumulative response	USD 230.5 million mobilised; 2.3 million ha treated via ground and aerial operations; 4.5 million tonnes of crop losses averted (commercial value ~USD 1.77 billion); 900 million litres of milk production saved; food secured for 41.5 million people
Jun 2022	Upsurge declared over	East Africa and Yemen region declared free of the Desert Locust upsurge

Table 5.2 — Desert locust upsurge (2019-2021): impact and FAO-coordinated response.

Remedy: Desert locust control relies on continuous ground and aerial surveillance (satellite-linked eLocust3 reporting devices), rapid deployment of ultra-low-volume pesticide spraying (organophosphates and, increasingly, biopesticides such as *Metarhizium acridum*) targeted at hopper bands before they fledge into mobile swarms, and — critically — sustained international funding, since locust plagues originate in remote breeding grounds that cross national boundaries and require coordinated regional response bodies such as the Desert Locust Control Organization for Eastern Africa (DLCO-EA).

5.3 Pink Bollworm and the Cotton Pest Complex (India)

India is the world's largest cotton producer and grower of Bt cotton, engineered for resistance to the earlier dominant pest, the American bollworm (*Helicoverpa armigera*). However, field-evolved resistance to Bt toxins in the pink bollworm (*Pectinophora gossypiella*) triggered a major resurgence beginning in Gujarat (2014-15) and escalating sharply in Maharashtra and Telangana in the 2017-18 season.

Year	Location	Impact	Remedy
2014-15	Gujarat	First major Bt-resistant pink bollworm outbreak recorded in India	Refugia planting mandates reinforced, pheromone trap monitoring introduced
2017-18	Maharashtra (state-wide)	State output forecast cut 37% from September outlook, down to 6 million bales (170 kg each) from 10.7 million bales the previous year	State-advised uprooting of infested plants in January to break the pest cycle

Year	Location	Impact	Remedy
2017-18	National (India)	National cotton production estimate revised down from a record 40 million bale forecast to 36.2 million bales (Cotton Association of India); exportable surplus fell from an expected 7.5 million bales to ~6 million bales	Cotton Corporation of India procurement support; Gujarat announced Rs 500/quintal bonus above MSP
2018 (Yavatmal district, Maharashtra)	District-level	Pesticide misuse to combat resurgent bollworm and sap-sucking secondary pests linked to a spate of farmer pesticide-poisoning deaths in Yavatmal	Push for licensed spray-service operators, safer molecule recommendations, KVK (Krishi Vigyan Kendra) advisory
2019-2022 (recurring)	Multiple cotton states	Pink bollworm can inflict up to 55% locule (boll) damage and reduce seed-cotton yield by up to 90% in severely infested fields	Dense pheromone trapping (8-10 traps/acre for mass trapping), early and synchronized sowing, timely crop termination after last picking, mandatory refugia (non-Bt border rows), resistant/tolerant hybrid development

Table 5.3 — Pink bollworm resurgence in Indian cotton, 2014-2022.

The pink bollworm case illustrates a recurring theme in pest management: resistance breakdown. Bt cotton transformed India into the world's second-largest cotton exporter after its 2006 approval, but continuous cultivation without adequate refugia, coupled with the availability of spurious/unapproved seed and long-duration hybrids that extend the pest's feeding window, allowed resistance to build up within roughly a decade — underscoring why resistance-management protocols must be enforced alongside any single-gene biotechnology solution.

5.4 Brown Planthopper and the Rice Pest Complex

Pest	Scientific Name	Typical Loss	Remedy
Brown planthopper (BPH)	<i>Nilaparvata lugens</i>	Can cause complete 'hopperburn' crop failure in localized patches within days under outbreak (biotype-shift) conditions; historically responsible for major Asian rice production crises in the 1970s-1980s	Resistant varieties (Bph-gene lines), avoiding excess nitrogen and insecticide-induced resurgence (avoid synthetic pyrethroids early season), conserving spiders/mirid bug predators, synchronized planting
Yellow stem borer	<i>Scirpophaga incertulas</i>	5%-60% depending on stage of attack (deadheart in vegetative stage vs. whitehead at reproductive stage, which is a total loss for the affected tiller)	Pheromone mass trapping, resistant varieties, need-based cartap hydrochloride/chlorantraniliprole application, early planting
Rice gundhi bug (Gundhi/stink bug)	<i>Leptocorisa</i> spp.	Grain discoloration and shriveling; 5%-25% yield/quality loss in affected panicles	Field sanitation, light traps, need-based insecticide at grain-filling stage

Table 5.4 — Rice pest complex: brown planthopper, stem borer and gundhi bug.

5.5 Plant-Parasitic Nematodes

Nematodes (roundworms, threadworms, eelworms) are frequently underestimated because damage occurs below ground and symptoms (stunting, chlorosis, wilting) are easily mistaken for nutrient deficiency. ICAR's All India Co-ordinated Research Project (AICRP) on Nematodes estimates that nematodes cause a loss of approximately **60 million tonnes of crop production annually in India (roughly 10-12% of total production)**, with the annual monetary loss across 24 major crops estimated at **Rs 21,068 million**. Globally, root-knot nematodes (*Meloidogyne* spp.) and cyst nematodes (*Heterodera* and *Globodera* spp.) are considered among the most damaging, affecting virtually every cultivated crop. **Remedy:** crop rotation with non-host/antagonistic crops (marigold, mustard), resistant rootstocks/varieties, soil solarisation, organic amendments and biocontrol agents (*Paecilomyces lilacinus*, *Pochonia chlamydosporia*), and targeted nematicide application where economically justified.

5.6 Fruit Flies, Aphids, Whiteflies and Other Sucking Pests

Pest	Crops Affected	Typical Loss	Remedy
Fruit flies (<i>Bactrocera</i> spp.)	Mango, guava, citrus, cucurbits	Up to 30%-100% fruit loss in unmanaged orchards; major quarantine/export-restriction pest	Methyl eugenol/cue-lure traps, bagging of fruit, field sanitation (destroying fallen fruit), sterile insect technique in area-wide programmes
Whitefly (<i>Bemisia tabaci</i>)	Cotton, tomato, cassava, cucurbits	Direct sap-sucking loss 10%-30%; far larger indirect loss as vector of viral diseases (e.g., cotton leaf curl virus, tomato yellow leaf curl virus) which can reach 40%-100%	Yellow sticky traps, reflective/silver mulch, systemic neonicotinoid or diamide seed treatment, resistant varieties, avoiding indiscriminate insecticide use that kills natural enemies
Aphids (multiple species)	Wheat, mustard, vegetables	5%-30% direct feeding loss; additional losses from virus transmission (e.g., Barley Yellow Dwarf Virus)	Imidacloprid/thiamethoxam seed treatment, timely sowing, conservation of ladybird beetle and lacewing predators
Coffee berry borer (<i>Hypothenemus hampei</i>)	Coffee (Arabica, Robusta)	Can destroy 5%-90% of berries in an infested field if untreated; considered the most damaging insect pest of coffee worldwide	Strip-picking and field sanitation to remove residual berries, <i>Beauveria bassiana</i> biopesticide, Broca traps, resistant Robusta-derived rootstocks

Table 5.5 — Sucking pests and fruit flies across major crops.

6. Regional Case Studies

6.1 India — National Loss Estimates

India presents a particularly well-documented national case because the Ministry of Agriculture has repeatedly reported loss figures to Parliament in response to formal questions. Successive replies (2006, 2009, 2013, 2018) place total loss from pests, diseases, rodents and weeds at a stable **10-30% of annual production**, while more granular ICAR institute-level studies (AICRP on Nematodes, AICRP on wheat pests at IARI) show that specific pest complexes alone can account for 6-13% avoidable loss even under managed research-station conditions — implying substantially higher losses under average farmer field conditions with less optimal pest monitoring and input access. India's pesticide consumption is comparatively low, applied to roughly one-fourth of its ~180 million hectares of arable land, which several industry analyses cite as a structural factor behind the country's estimated USD 8 billion (Rs 500 billion) in annual pest-and-disease-related crop loss.

The 2017-18 Maharashtra cotton crisis additionally triggered a state disaster-relief claim: the Government of Maharashtra sought Rs 2,425.53 crore from the National Disaster Response Fund (NDRF) for combined pink bollworm and Cyclone Ockhi damage, later managed through the State Disaster Response Fund (SDRF) after central clarifications were sought — illustrating how pest outbreaks can escalate into formal disaster-management events with fiscal consequences at the state level.

6.2 Sub-Saharan Africa

Africa's experience is dominated by transboundary and invasive-species pressure. Beyond fall armyworm (Section 5.1), a 2021 CABI/University of Minnesota study calculated that five invasive alien species alone cause USD 0.9-1.1 billion in annual losses to smallholder farmers across six eastern African countries — equivalent to 1.8-2.2% of regional agricultural GDP — a figure projected to grow to USD 1.0-1.2 billion per year. Malawi has been identified as the country facing the highest relative risk from fall armyworm, with forecasts of 12.5-30% of its agricultural economy at risk absent effective management. The continent's dependence on maize as the primary staple magnifies the food-security consequences of any single-pest outbreak far beyond what the same percentage loss would mean in a more diversified cropping system.

6.3 Central America and the Coffee Belt

The 2012-2016 coffee leaf rust epidemic (Section 4.4) demonstrated how a single fungal disease can destabilise a regional economy built around one export commodity. Coffee contributed measurably to GDP in every affected country (an estimated 1.40% of GDP in Honduras, 0.53% in Nicaragua, 0.31% in El Salvador, 0.19% in Guatemala and 0.03% in Costa Rica in 2012 alone), and more than 80% of affected farmers were smallholders in already vulnerable, poverty-prone areas — meaning the nutritional and health consequences of the epidemic extended well beyond the immediate export-revenue loss.

6.4 South Asia — Transboundary Threats

South Asia's wheat belt (India, Pakistan, Bangladesh, Nepal) is now recognised as a frontline zone for wheat blast following the 2016 Bangladesh outbreak, given the region's dense wheat cultivation, warm humid pockets conducive to *Magnaporthe oryzae* Triticum, and proximity to the site of first Asian detection. International wheat research bodies (CIMMYT, ICAR, Bangladesh Wheat and Maize Research Institute) have since coordinated cross-border surveillance and rapid-response fungicide protocols to prevent a repeat epidemic from spreading into India's much larger wheat area, which underpins national food-grain reserves.

7. Economic Impact Analysis

Consolidating the case studies above, three tiers of economic consequence are evident. **Tier 1 — Global structural cost:** USD 220 billion in annual trade losses and at least USD 70 billion in invasive-pest damage, borne diffusely across all producing and importing nations. **Tier 2 — National/regional shock events:** single-country or single-region outbreaks in the USD 1-10 billion range (fall armyworm in Africa, desert locust in East Africa/Yemen, coffee rust in Central America, cotton pink bollworm in India), which disrupt specific commodity markets, government budgets (disaster relief claims) and rural employment over a 1-3 year window. **Tier 3 — Localized/farm-level catastrophe:** losses of 50-100% at the level of an individual field, orchard or plantation (banana TR4 infestation, citrus HLB decline, wheat blast hot-spots), which may be statistically small in national aggregate terms but represent complete livelihood loss for the affected farming household.

Loss Tier	Illustrative Case	Approx. Value	Time Horizon
Global structural	FAO global trade loss estimate	USD 220 billion/year	Ongoing/recurring
Global structural	Invasive pests specifically	USD 70 billion/year (minimum)	Ongoing/recurring
Regional shock	Fall armyworm, Africa maize (2018 estimate)	USD 2.5-6.3 billion/year (maize only)	2016-present
Regional shock	All invasive alien species, Africa (2021 study)	USD 65.58 billion/year	Ongoing
Regional shock	Desert locust, East Africa/Yemen (2020 projection)	Up to USD 8.5 billion	2019-2021
Regional shock	Coffee leaf rust, Central America (2011-2016)	USD 3.2 billion cumulative	2011-2016
National shock	India, all-crop annual pest/disease loss (industry estimate)	USD ~8 billion/year (Rs 500 billion)	Recurring
National shock	Maharashtra cotton, pink bollworm (2017-18)	37% state output cut	Single season
Localized/farm	Wheat blast hot-spot fields, Bangladesh (2016)	Up to 100% field-level loss	Single season
Localized/farm	Florida citrus, HLB cumulative decline	>90% production decline over ~20 years	2005-present

Table 7.1 — Three-tier framework for classifying pest/disease economic impact, with representative figures from this document.

8. Climate Change and the Changing Geography of Pests and Diseases

FAO explicitly links climate change to worsening pest pressure, noting that variable weather patterns are likely to make plant pests more damaging in intensity, distribution and spread. Several mechanisms are documented across the case studies in this report:

- **Range expansion:** Warmer minimum temperatures have allowed coffee leaf rust to establish at higher elevations in Central America that were previously too cool for the fungus to persist year-round.
- **Extreme-weather-triggered outbreaks:** The 2019-2021 desert locust upsurge was directly linked to an unusual sequence of cyclones in the Arabian Sea (eight cyclones in the 2018-19 season, compared with a long-term average of fewer than one per year), which created ideal wet-soil breeding conditions in the Empty Quarter that went undetected due to the region's remoteness.
- **Faster pest generation cycles:** Rising temperatures shorten the developmental time of insect pests such as fall armyworm and brown planthopper, allowing more generations per season and faster resistance evolution against both host-plant resistance and insecticides.
- **Altered disease windows:** Changes in monsoon timing and humidity affect the onset and duration of favourable conditions for fungal diseases such as wheat rust, wheat blast and late blight, sometimes extending the window of vulnerability or shifting it earlier/later relative to the crop's most susceptible growth stage.
- **New crop-pest combinations:** As cultivation zones shift poleward and to higher elevations in response to warming, crops are increasingly grown in regions with pest and pathogen communities to which local varieties have no co-evolved resistance — a dynamic already visible in wheat blast's jump from South America to South Asia and in the poleward creep of fall armyworm into temperate maize-growing zones of China and southern Europe.

These dynamics mean that historical loss statistics should be read as a floor rather than a ceiling: several of the case studies documented here (fall armyworm, desert locust, wheat blast, citrus HLB) are recent arrivals in their respective regions, and modelling by CIMMYT, CABI and FAO consistently projects further geographic expansion of high-risk zones under most emissions scenarios.

9. Integrated Pest and Disease Management (IPDM) — Remedies in Detail

No single remedy addressed in the case studies above is sufficient in isolation. FAO defines Integrated Pest Management (IPM) as the careful consideration of all available pest-control techniques, integrating measures that discourage pest population development while keeping pesticide use and other interventions at levels that are economically justified and minimise risk to human health and the environment. The following subsections organise remedies by category, cross-referencing the specific diseases/pests discussed earlier.

9.1 Cultural and Agronomic Control

The first and often cheapest line of defence. Includes: crop rotation (breaks the life cycle of soil-borne pathogens and nematodes — critical for *Fusarium* wilts, nematode complexes, and groundnut/pulse wilt disease); timely and synchronized sowing across a locality (reduces the window during which pests such as stem borer, brown planthopper and bollworm can build up continuous generations); field sanitation (removal of crop residue, volunteer plants and fallen fruit — essential for late blight, fruit flies and coffee berry borer); balanced fertilisation (excess nitrogen favours sheath blight, brown planthopper and aphid outbreaks); optimal plant spacing (improves airflow, reducing humidity-dependent fungal disease pressure); and trap/border cropping such as the push-pull system (*Desmodium* intercrop + Napier grass border), which was originally developed against stemborers in East Africa and has since shown effectiveness against fall armyworm.

9.2 Host-Plant Resistance and Breeding

Genetic resistance remains the most cost-effective, environmentally benign control method once developed, though it requires continuous investment because pathogens and pests evolve to overcome single-gene resistance. Examples referenced in this document include: gene-pyramided rice varieties combining multiple blast-resistance (Pi) and bacterial-blight-resistance (Xa) genes; wheat lines carrying stacked Sr (stem rust) genes developed through the Borlaug Global Rust Initiative in response to Ug99; Bt cotton (Cry gene insertion for bollworm resistance), now requiring parallel resistance-management (refugia) strategies as pink bollworm resistance has emerged; TR4-tolerant banana cultivars derived from somaclonal variation and conventional breeding programmes (e.g., GCTCV lines); and rust-resistant coffee cultivars (Catimor/Lempira-type hybrids combining Robusta-derived rust resistance with Arabica cup quality) used in post-epidemic replanting across Central America.

9.3 Biological Control

Biological control uses natural enemies, antagonistic organisms or microbial agents to suppress pest/pathogen populations, and has grown in importance as chemical resistance and environmental concerns limit purely chemical approaches. Documented examples include: *Bacillus thuringiensis* (Bt)-based biopesticides and entomopathogenic fungi (*Metarhizium*, *Beauveria*) against fall armyworm and coffee berry borer; *Metarhizium acridum* biopesticide formulations increasingly used alongside chemical sprays in desert locust control; *Trichoderma*-based seed and soil treatments against *Fusarium* and *Sclerotium* wilts; *Paecilomyces lilacinus* and *Pochonia chlamydosporia* as nematode-antagonistic biocontrol agents; and conservation biological control — simply avoiding broad-spectrum insecticide use early in the season to protect naturally occurring predators (spiders, mirid bugs, ladybird beetles, lacewings, parasitoid wasps) that otherwise suppress brown planthopper, aphid and whitefly populations without any input cost to the farmer.

9.4 Chemical Control and Judicious Pesticide Use

Chemical control remains indispensable for rapid, large-scale response, particularly for migratory pests (desert locust aerial spraying) and fast-moving fungal epidemics (wheat rust and blast fungicide response). However, FAO and the case studies in this document underline that indiscriminate use carries three documented risks: (a)

resurgence of secondary pests when natural enemies are killed alongside the target pest (observed in Indian cotton after early-season pyrethroid overuse against bollworm); (b) resistance evolution, as seen in Bt-resistant pink bollworm and in fungicide-tolerant *Phytophthora infestans* strains; and (c) direct human health risk, tragically illustrated by farmer pesticide-poisoning deaths in Yavatmal, Maharashtra (2017), linked to indiscriminate tank-mixing of multiple pesticide products against resurgent bollworm and sucking pests. Judicious use principles include: spraying only after economic threshold levels (ETL) are crossed (as determined by field scouting, not calendar-based spraying); rotating chemical classes/modes of action to delay resistance; targeting the most vulnerable pest life stage (e.g., young larvae, not mature caterpillars); and using licensed applicators with proper protective equipment.

9.5 Digital Agriculture, Early Warning and Forecasting Systems

Several of the case studies in this document demonstrate the value of early detection. FAO's eLocust3 satellite-linked field reporting system allows real-time mapping of desert locust hopper bands, enabling targeted spraying before swarms form and disperse. Weather-driven disease-forecasting models (such as Blitecast and the Hutton Criteria for potato late blight, and similar humidity/temperature-triggered models for wheat rust and blast) allow farmers to time fungicide sprays precisely rather than on a fixed calendar, reducing both cost and resistance-selection pressure. Mobile-based pest identification and advisory tools (including ICAR's Farmer Field School networks and CABI's Plantwise/PlantwisePlus plant-clinic network) extend expert diagnosis to smallholder farmers who previously had no rapid access to it. Pheromone trap networks (used against pink bollworm, fruit flies and stem borer) provide low-cost, real-time population monitoring that triggers timely intervention before economic damage thresholds are crossed.

10. Policy Frameworks and Institutional Response

International and national institutions play a decisive role in every case study documented above. Key frameworks include:

- **International Plant Protection Convention (IPPC):** the FAO-hosted international treaty framework (with its Secretariat under FAO's Plant Production and Protection Division, established 1947) that coordinates phytosanitary standards and transboundary pest reporting among member countries, and which designated 2020 the International Year of Plant Health and 12 May as annual International Day of Plant Health.
- **FAO Global Action for Fall Armyworm Control (2020-2022):** a USD 500 million coordination mechanism launched in December 2019 covering Africa, the Near East and Asia-Pacific, linking technical partners to deliver science-based control solutions to farmers.
- **FAO Desert Locust Response Plan (2020-2021):** a phased regional response (USD 230.5 million mobilised against a stated requirement of USD 311.4 million) built on three pillars — curbing the spread, safeguarding livelihoods and early recovery, and coordination/preparedness — delivered in partnership with national governments and the Desert Locust Control Organization for Eastern Africa (DLCO-EA).
- **India's Strengthening and Modernization of Pest Management Approach (SMPMA) Scheme:** implemented through 31-35 Central Integrated Pest Management Centres (CIPMCs) across 28 states and one union territory, responsible for pest/disease surveillance, biocontrol-agent production and release, and Farmer Field School training; IPM packages have been developed for 77 major crops in collaboration with State Agricultural Universities and ICAR institutions.
- **ICAR All India Co-ordinated Research Projects (AICRP):** dedicated national networks such as AICRP on Nematodes in Cropping Systems and crop-specific AICRPs (wheat, rice, cotton) that generate the yield-loss and management data referenced throughout this document.
- **Borlaug Global Rust Initiative / Durable Rust Resistance in Wheat (DRRW):** the international wheat-breeding response to Ug99 stem rust, coordinating CIMMYT, ICARDA and national programmes to develop and disseminate stacked-resistance varieties globally.
- **CABI's Plantwise/PlantwisePlus programme:** a global network of plant health clinics run by trained "plant doctors" who provide diagnosis and management advice directly to smallholder farmers, alongside CABI's ongoing Global Burden of Crop Loss (GBCL) initiative aimed at building the first comprehensive, standardised dataset on global pre-harvest crop loss.

11. Recommendations

1. Invest in early-warning infrastructure.

The desert locust and fall armyworm case studies show that early detection converts a potential multi-billion-dollar loss into a controllable, cost-effective response. Expanding satellite-based surveillance, pheromone trap networks and mobile diagnostic tools to more regions and more crops should be a funding priority for governments and international donors alike.

2. Enforce resistance-management protocols alongside any single-gene technology.

Bt cotton and single-gene disease resistance both eventually face resistance breakdown (pink bollworm, Ug99 stem rust). Mandatory refugia, gene-stacking/pyramiding, and rotation of resistance sources should be treated as non-negotiable components of deployment, not optional add-ons.

3. Strengthen regional and international coordination for transboundary pests.

Fall armyworm, desert locust and wheat blast all crossed continents within a few growing seasons. No single country's response is sufficient; sustained funding for coordination bodies (IPPC, DLCO-EA, FAO Global Action initiatives) needs to be maintained even in years without an active outbreak, so that surveillance capacity does not atrophy between crises.

4. Expand farmer-level training and diagnostic access.

A recurring theme across the Indian, African and Central American case studies is that smallholder farmers often detect outbreaks late or respond with inappropriate/excessive chemical use. Farmer Field Schools, plant clinics (CABI Plantwise model) and mobile advisory services should be scaled to reach the last-mile smallholder, not just commercial or export-oriented growers.

5. Diversify cropping systems in high-risk monoculture zones.

Banana Cavendish monoculture (TR4 vulnerability), extensive single-hybrid maize cultivation (fall armyworm) and large contiguous Bt cotton belts (pink bollworm) all illustrate how genetic and species uniformity amplifies outbreak severity. Diversification, intercropping (e.g., push-pull systems) and varietal mosaics reduce the speed at which a single pest or pathogen can spread across a landscape.

6. Continue funding long-term breeding pipelines even between crises.

The multi-decade lag between the emergence of Ug99 (1998) and widespread deployment of durable-resistance wheat varieties, and the ongoing search for a true HLB-tolerant citrus variety two decades after Florida's 2005 detection, both show that breeding solutions take 10-20 years. Funding must be sustained through the quiet years, not concentrated only during acute outbreaks.

7. Build robust, standardised global crop-loss data infrastructure.

The wide range in cited figures throughout this document (20-40% FAO global estimate; 10-30% India estimate; differing methodologies across studies) reflects a genuine data gap. Initiatives such as CABI's Global Burden of Crop Loss project deserve wider institutional and donor support to produce comparable, standardised loss estimates across crops, regions and years.

12. Conclusion

Across every case examined in this document — from the 19th-century Irish Potato Famine to the 2019-2021 desert locust upsurge and the ongoing spread of banana Tropical Race 4 — a consistent pattern emerges. Yield loss from pests and diseases is never purely a biological phenomenon; it is shaped by cropping-system diversity, breeding investment, surveillance infrastructure, farmer knowledge, international coordination and the pace of climate change. The FAO's headline figure that up to 40 percent of global crop production is lost to pests and diseases every year is not a fixed natural constant but a number that has been measurably pushed down in specific times and places through resistant varieties, early warning systems and coordinated response — and pushed back up whenever any of those defences lapse, resistance breaks down, or a new invasive species crosses a border undetected. Sustained, well-funded Integrated Pest and Disease Management, backed by robust data and rapid international cooperation, remains the most reliable path to narrowing this persistent gap between potential and actual harvest.

13. References and Data Sources

- [1] Food and Agriculture Organization of the United Nations (FAO). 'Understanding the Context.' Pest and Pesticide Management, FAO. <https://www.fao.org/pest-and-pesticide-management/>
- [2] FAO Plant Production and Protection Division. Overview and historical brief, FAO Open Knowledge Repository.
- [3] FAO. 'The Hidden Health Crisis: How Plant Diseases Threaten Global Food Security.' One Health Highlights, 2026.
- [4] FAO. 'FAO Report Reveals Shocking Crop Losses: Up to 40% Due to Pests and Diseases Annually.' International Day of Plant Health briefings, 2023.
- [5] CropLife International. 'Crop Protection Industry Supports FAO on IPM,' citing FAO-OECD joint reporting.
- [6] CABI. 'Global Burden of Crop Loss' project overview, CABI.org.
- [7] CABI. 'Fall Armyworm: Impacts and Implications for Africa' (evidence note commissioned by UK DFID).
- [8] CABI / University of Minnesota (Senay, Pardey et al.). Study on invasive alien species economic impact on African agriculture, 2021.
- [9] The East African. 'Fall Armyworm to Cost More Than \$2bn in Losses,' 2017.
- [10] Impakter. 'Fall Armyworm Escalates Africa's Vulnerability,' analysis of 12-country FAW impact, 2018.
- [11] Down to Earth. 'Combating Fall Armyworm Attack Amid Confluence of Challenges in Africa.'
- [12] FAO. 'Fall Armyworm: Scaling Up Response Across the Globe' — Director-General remarks on Global Action for Fall Armyworm Control.
- [13] FAO Emergencies and Resilience. 'Desert Locust Crisis (2020-2021)' progress reports.
- [14] FAO GIEWS Special Alert No. 347, 'East Africa: The Worst Desert Locust Outbreak in Decades,' 29 January 2020.
- [15] USAID. East Africa Desert Locust Crisis Fact Sheets #4 and #7, 2020.
- [16] Al Jazeera. 'In Pictures: Ethiopia Struggles to Suppress Desert Locust Infestation,' 2020 (citing World Bank USD 8.5bn estimate).
- [17] Ethiopian Business Review. 'Desert Locust' impact assessment, Ethiopia, 2020.
- [18] IPPC / FAO Regional Workshop Africa. Desert Locust situation reports, September 2020 and April 2021.
- [19] Indian Council of Agricultural Research (ICAR). 'Pests Cause 30% Loss of Crop Yields in India,' AICRP on Nematodes annual group meeting remarks (P.K. Chakrabarty, ADG Plant Protection & Biosafety).
- [20] ICAR-Indian Agricultural Research Institute. Jadhav, M.M., Chander, S., Saharan, M.S., Suroshe, S.S., Rajna, S. 'Assessment of Avoidable Yield Losses Due to Insect Pests and Diseases in Wheat (*Triticum aestivum*).' Indian Journal of Agricultural Sciences 93(11): 1186-1190, 2023.
- [21] ICAR National Agricultural Innovation Project (NAIP), Component 4 project abstract, krishi.icar.gov.in.
- [22] Government of India, Lok Sabha Unstarred Question No. 986 (27 Feb 2006), No. 1012 (27 Feb 2006), No. 1305 (14 Jul 2009), No. 5914 (3 Apr 2018) — Ministry of Agriculture and Farmers Welfare replies on crop loss due to pests and diseases.
- [23] Business Standard / industry press. 'India: Crops Worth \$8 Billion Lost Every Year Because of Pest, Disease Attacks.'
- [24] Mottaleb, K.A., Hodson, D.P., Krupnik, T.J., Sonder, K. 'Quantifying Wheat Blast Disease Induced Yield and Production Losses of Wheat: A Quasi-Natural Experiment.' Journal of Agricultural and Applied Economics 55(1): 34-56, 2023 (CIMMYT).
- [25] Chowdhury, A.K. et al. 'Occurrence of Wheat Blast in Bangladesh and Its Implications for South Asian Wheat Production.' Indian Journal of Genetics and Plant Breeding.
- [26] Encyclopedia.pub. 'Wheat Blast and Food Security' entry, summarising outbreak history since 1985 (Brazil) and 2016 (Bangladesh).
- [27] International Coffee Organization (ICO). Resolution 451, 110th Session, London, March 2013 — 'Critical Situation in Central America Caused by Coffee Leaf Rust Outbreak.'
- [28] Bloomberg, cited in Revista Cafeicultura. 'Coffee Leaf Rust Threatens Production in Central America,' January 2013.
- [29] World Coffee Research. 'World Coffee Research Publishes Manual for Coffee Leaf Rust' (citing USD 3.2 billion cumulative 2011-2016 damage figure).
- [30] Graphic News. 'Rust Fungus Sweeps Central America' infographic, July 2013.

[31] Cotton Association of India (CAI); Reuters; TextileExcellence.com; FashionNetwork.com; Cottongrower.com — India pink bollworm/cotton production reporting, 2017-2018.

[32] Mongabay / India Environment Portal. 'The Pink Bollworm Menace Adds to Maharashtra Cotton Farmers' Distress,' 2018.

[33] Business Standard. 'Pink Bollworm' topic archive — cotton yield and lint-quality impact figures, 2017-2022.

[34] USAID/Center for Disease Control historical reference and general agronomic literature on the Irish Potato Famine (1845-1852).

End of document. This compilation is intended as an educational and planning reference. For operational decisions (fungicide/pesticide selection, quarantine action, or investment decisions), readers should consult current advisories from their national plant protection organisation, ICAR/State Agricultural University extension services, or FAO's Global Information and Early Warning System (GIEWS).