

The Grand Meson

Shuhei Ihara

June 23, 2026

Abstract

This paper introduces the Grand Meson, not as a theory of everything, but as a safe $n = 3$ general-purpose mesonic object. The starting point is the General Meson: a two-port Yukawa-style mediator whose readable structure is the arrow $T \rightarrow S$. The Grand Meson opens the two remaining readable axes and thereby becomes a three-axis, six-port mediator. The extension is conservative: the original two-port meson is recovered by projection, while the new ports are registered as defined, null, or active instead of being left outside the type of the theory. The central claim is modest and structural: the Grand Meson is the first complete and last safe mesonic case for a three-dimensional observer.

1 Opening

A meson is usually approached through its physical role: it mediates an interaction. In the present paper the same word is first read through its directional role. A meson stands between a presented side and a readable side. It is therefore written as

$$T \longrightarrow S.$$

This arrow is not yet a full dynamics. It is the minimum shape of a mediator: something is presented at T , something is read at S , and the middle term makes the relation nameable.

The former working nickname “Ethon” is retired here. The technical object of this paper is the Grand Meson. The name is structural, not honorific: “grand” means that a two-port meson has been closed into the safe three-axis case.

2 The General Meson

Let the first readable axis be

$$A_1 = \{T, S\}.$$

Let Pol be a class of admissible mesonic polymers. The smallest implementation may be generated from

$$\mathbb{P}_3 = \{-\mathbf{1}, \mathbf{0}, +\mathbf{1}\},$$

but no physical charge, amplitude, or particle identity is assigned to these pseudo-values at this level.

Definition 1 (General Meson). A General Meson is a two-port mesonic record

$$m : A_1 \rightarrow \text{Pol}.$$

Equivalently,

$$\text{Gen} := \text{Pol}^{A_1}.$$

The port T is the presented boundary. The port S is the readable boundary.

This definition deliberately keeps the Yukawa-style mediator idea narrow. It does not say that a meson is universal. It says only that the basic mesonic act is readable mediation from T to S . In that sense a General Meson is the optimal eye for the nuclear-force problem: the source and sink boundaries are already specified by the problem itself.

3 The Hidden Cost of Undefined Ports

The decisive distinction is between null and undefined. A null port is known to be empty. An undefined port has not entered the type of the theory.

$$\mathbf{0} = \text{registered null}, \quad \perp = \text{unregistered undefined}.$$

Definition 2 (Undefined lift). Let

$$A_3 = \{T, S, C, E, U, D\}, \quad \text{Pol}_\perp := \text{Pol} \cup \{\perp\}.$$

The undefined lift of a General Meson $m \in \text{Gen}$ is

$$L_\perp(m) : A_3 \rightarrow \text{Pol}_\perp$$

defined by

$$L_\perp(m)(T) = m(T), \quad L_\perp(m)(S) = m(S),$$

and

$$L_\perp(m)(d) = \perp \quad (d \in A_3 \setminus A_1).$$

Thus the remaining axes are not false, not zero, and not disproved. They are absent from the original type. The two-port theory remains locally correct while it cannot register consequences that travel through the untyped directions.

Remark 3 (Meta-theoretic reading). The undefined ports are not a defect in the original General Meson. They are the price of a successful restriction. A narrow mediator may solve its local problem precisely because it does not carry the full burden of every other axis.

4 The Grand Meson

The Grand Meson is obtained by replacing undefined ports with readable ports. A port may be active, or it may be null, but it may not be outside the type.

Definition 4 (Grand Meson). A Grand Meson is a six-port mesonic record

$$g : A_3 \rightarrow \text{Pol}.$$

Equivalently,

$$\text{Grand} := \text{Pol}^{A_3}.$$

The six ports are grouped into three readable axes:

$$T \leftrightarrow S, \quad C \leftrightarrow E, \quad U \leftrightarrow D.$$

In this notation the Grand Meson is the safe $n = 3$ object:

$$\text{Grand} = \text{Pol}^{2 \cdot 3}.$$

The adjective “safe” means that all three axes are typed. It does not mean that every transition is allowed, nor that every physical theory has been solved.

Definition 5 (Registered support). For $g \in \text{Grand}$, define

$$\text{supp}(g) = \{d \in A_3 \mid g(d) \neq \mathbf{0}\}.$$

A Grand Meson may have inactive ports, but inactivity is registered by $\mathbf{0}$, not by $\perp$.

5 Conservative Recovery

The Grand Meson must not destroy the General Meson. It must contain it as the two-port face that was already working.

Definition 6 (Projection to the General Meson). Define the T, S -projection

$$p_{TS} : \text{Grand} \rightarrow \text{Gen}$$

by

$$p_{TS}(g) = g|_{\{T, S\}}.$$

Proposition 7 (Conservative recovery). *Every General Meson is recovered from any Grand Meson whose T, S -face agrees with it. In particular, if $g(T) = m(T)$ and $g(S) = m(S)$, then*

$$p_{TS}(g) = m.$$

Proof. The projection p_{TS} discards the four ports outside $\{T, S\}$ and keeps exactly the T - and S -values. Therefore agreement on those two ports is sufficient and necessary for recovery. $\square$

This is the main discipline of the construction. The Grand Meson does not rewrite the older meson. It keeps the working $T \rightarrow S$ face and opens two additional faces under explicit registration.

6 The Einstein Gate

The Grand Meson must also avoid the classical aether failure. It is not a hidden medium with an absolute state of rest. It is a local six-port mediator whose outputs must be readable without introducing an absolute frame.

Convention 8 (No absolute mesonic frame). No equation in the Grand Meson theory may depend on a privileged global rest frame of the mediator. Only typed ports, local registration, and observer-readable projections may be used.

This convention is the Einstein gate. It does not ban mediation. It bans absolute mediation. The Grand Meson is therefore not a return to the classical aether. It is a typed local mediator that must survive projection into admissible observations.

7 Why $n = 3$ Is the Safe Case

Formally one may define an n -meson by

$$\text{Mes}_n := \text{Pol}^{2n}.$$

The case $n = 1$ is the General Meson. The case $n = 3$ is the Grand Meson.

The reason to stop at $n = 3$ is not that higher n is impossible. It is that a three-dimensional observer already has the first complete rotational reading at $n = 3$. Three axes close the ordinary posture problem:

$$\text{Read}_{O_3}(\text{Mes}_3) = \text{Read}_{O_3}(\text{Pol}^6).$$

For $n > 3$, additional axes require additional readout structure. Such structure may be meaningful in a later theory, but it is not required for the present observer class:

$$\text{Read}_{O_3}(\text{Mes}_n) \not\approx \text{Read}_{O_3}(\text{Mes}_3) \quad (n > 3).$$

Thus $n = 3$ is the first complete and last safe case: complete because the three axes are readable as a closed posture, safe because the theory does not ask the reader to chase an infinite ascent of unread axes.

8 Safe Transition

Safety is not the absence of change. Safety is the registration of change. Let a transition be written

$$u : g \rightarrow g'.$$

The transition may change ports, support, or observable posture. It is unsafe only when the change crosses a typed boundary without registration.

Definition 9 (Grand-safe transition). Let $\text{Reg}(u)$ be the registration record of a transition $u : g \rightarrow g'$. The transition is Grand-safe when every changed port is registered:

$$\{d \in A_3 \mid g(d) \neq g'(d)\} \subseteq \text{Reg}(u).$$

It is Grand-unsafe when a changed port is absent from $\text{Reg}(u)$.

Equivalently,

$$\text{unsafe} = \text{unregistered discontinuity}.$$

The discontinuity itself is not the violation. The violation is that the boundary crossing occurred while the theory failed to enter it into the ledger.

9 Conclusion

The Grand Meson is not proposed as an omnipotent meson. It is proposed as a safe $n = 3$ general-purpose meson:

$$\text{Grand} = \text{Pol}^6.$$

It conservatively recovers the General Meson by projection, replaces undefined ports with typed ports, avoids an absolute aether frame, and treats safety as registration rather than immobility.

The path is therefore narrow. Begin with the $T \rightarrow S$ eye that already works. Open $C \leftrightarrow E$ and $U \leftrightarrow D$ without breaking it. Stop at $n = 3$ until a future theory can make higher axes readable without turning them into a burden of infinite explanation.

References

- [1] H. Yukawa. On the interaction of elementary particles. *Proceedings of the Physico-Mathematical Society of Japan*, 17:48–57, 1935.
- [2] A. Einstein. On the electrodynamics of moving bodies. *Annalen der Physik*, 17:891–921, 1905.