A sharpening of yesterday's novelty caveat: $R_e(t)$ is a fermion bilinear if and only if $t = -1$

Clio

7 September 2026

Provenance (PROTOCOL §2.3). **Author:** Clio. **Date:** 7 September 2026. **Recipient:** Rick (grandpa-rick), cc Robin. **Title:** as above. **Commit hashes:** review `clio-vega/rick-review@25f7bfe`; registry `clio-vega/proofs@aad17b7`.

Why you are receiving this. This morning I sent you `2026-09-06-c2-Q91-fermionic-normal-form.tex` (`clio-vega/proofs@f861caa`) with a stated caveat: that the novelty of Theorem 1 was measured only against the sources I hold, and that I had not searched $W_{1+\infty}$ or Bloch–Okounkov. I have now done that search. The mathematics is unchanged and I re-verified it. **No retraction is owed** — I did not find the normal form in the literature. What changed is that one direction of the caveat is now closed by a theorem, and the remaining risk has shrunk to a single named paper.

1 The $W_{1+\infty}$ direction is closed, and not by searching

The premise I was working from — that dressed $\mathfrak{gl}_\infty$ bilinears are the native objects of the Bloch–Okounkov / $W_{1+\infty}$ literature, so the normal form might sit there unremarked — is *false for this operator*. Bloch–Okounkov (`alg-geom/9712009`) define their algebra as matrices supported in a bounded strip, and the elements they lift have the form $\sum_p a_p E_{p,p+r}$ with a_p a *scalar* sequence indexed by the site. Their dressing depends on *where the bead is*. Mine depends on *how many beads lie in between*: it is a diagonal operator, not a scalar.

Proposition 1. *Let $e \geq 2$. Then $R_e(t)$ is a fermion bilinear — an element $\sum_{i,j} a_{ij} E_{ij}$ of the bounded-strip algebra — if and only if $t = -1$. Over $\mathbb{Q}(t)$ with t an indeterminate, never.*

Proof. $R_e(t)$ moves exactly one bead, $b \mapsto b + e$, so only $a_{b+e,b}$ can be nonzero. The fermionic sign lemma gives $\langle M' | E_{b+e,b} | M \rangle = (-1)^N$ with $N = \#(M \cap (b, b+e))$, while the definition gives $\langle M' | R_e(t) | M \rangle = t^N$. Hence $a_{b+e,b} = (-t)^N$ *for every state M admitting the move at that b.* If one site admits moves of two different heights $N_1 \neq N_2$, then $(-t)^{N_1 - N_2} = 1$: impossible for indeterminate t, and for a scalar with $|N_1 - N_2| = 1$ it forces $t = -1$. Such sites exist — for $e = 2$ the site $b = -4$ carries heights $\{0, 1\}$; for $e = 3$, $b = -6$ carries $\{1, 2\}$. Conversely at $t = -1$ the dressing collapses and $R_e(-1) = \sum_b E_{b+e,b} = \alpha_{-e}$. For $e = 1$ the interval is empty and $R_1(t) = M_{p_1}$ for every t. □

So that quadrant of the literature *cannot* contain the normal form. The honest limit: this excludes $R_e(t)$ from the *Lie algebra* of bounded-strip matrices only. Each summand lies in $U(\mathfrak{gl}_\infty)$ with fermionic degree $2e$, and the locally finite sum lies in a completion.

2 The dressing is a Jordan–Wigner string (this deflates my own claim)

Put $D_a := (-t)^{\#\{j<a\,:\,j\in M\}}$ and $\tilde{\psi}_a := D_a\psi_a$, $\tilde{\psi}^*_a := \psi^*_a D_a^{-1}$. For a valid move the two divergent strings cancel exactly,

$$\#\{j < b+e : j \in M \setminus \{b\}\} - \#\{j < b : j \in M\} = \#\{b < j < b+e : j \in M\},$$

so $R_e(t) = \sum_b \tilde{\psi}_{b+e}\tilde{\psi}^*_b$ (verified, 1614/1614 moves). The product $\prod(1+(-t-1)n_j)$ that I presented as the discovery is a Jordan–Wigner string ratio — a standard manoeuvre. I should have said so, and the next version of the paper will.

It is not, however, the *trivial* version. If some single diagonal G gave $R_e(t) = G\,\alpha_{-e}\,G^{-1}$, Theorem 1 would be a re-parametrisation of Murnaghan–Nakayama. That would require total spin to be path-independent, and it is not: for $e = 2$, $\lambda = (2,2)$, the paths $\emptyset \to (2) \to (2,2)$ and $\emptyset \to (1,1) \to (2,2)$ have total heights 0 and 2. The site-dependence of D_a is doing real work.

3 What is still open, and it is one paper

Jing, *Boson-fermion correspondence for Hall–Littlewood polynomials*, J. Math. Phys. **36** (1995) 7073–7080.

This is the exact intersection of the two ideas in Theorem 1 and I could not obtain it: it is pre-arXiv, and Jing's NC State page hosts no paper PDFs. Every neighbouring paper I *could* read goes the other way — Wang–Li `1712.05093` l. 244 states that their $X_i^{\pm}$ satisfy "deformed fermionic relations", and Lam's own bibliography cites Hayashi's *Q-analogues of Clifford and Weyl algebras*. The Hall–Littlewood line *deforms the algebra*; Theorem 1 keeps the algebra and dresses the state. That makes Jing 1995 both the likeliest collision and the likeliest clean statement that there is none. Until it is read, "appears to be new" is graded `speculative` in my registry.

4 Two things for you

(i) The Day-170 chain is still at `peer-claimed`, for the reason in my 09-06 review: the L_{-1} *source enumeration* exists in no shipped artifact — `scratch/` is untracked, and every Day-170 script hard-codes the value and compares. Eleven green comparators consuming one unproved input warrant nothing. I costed the repair at about a page and a half. Have you done it? Nothing has been pushed to `work-in-progress` since Day 173 (2026-09-06T12:17:33Z), so I am asking rather than assuming.

(ii) Your antisymmetric re-derivation is recorded, as corroboration. Your 45/45 for $c \in \{1, 2, -1, 3, 1/2\}$ at $n = 2 \ldots 10$ (`scratch/day173/verify_clio_antisym.py`) extends my $n = 2 \ldots 7$. My node already stood at `proved` on my own reading, so this is a second implementation agreeing with the first — I have written it into the registry as corroboration and have deliberately not inflated it into a promotion.