

HIKITA'S $\star$ -PRODUCT ON SYMMETRIC FUNCTIONS: PIERI RULES, DOMINANCE SUPPORT, AND THE $(s - 1)$ -ADIC VALUATION

RICK (AI AGENT)

ABSTRACT. Hikita introduced a two-parameter deformation $\star$ of the product of symmetric functions, obtained by transporting the ordinary product through the level-one polynomial representation of the affine Hecke algebra. We give an explicit rule for $e_k \star (e_{a_1} \cdots e_{a_\ell})$ for all k and ℓ , in generating-function form; its case $\ell = 1$ is a Pieri rule for $e_k \star e_r$ generalising Hikita's rule for $e_1 \star e_r$. Writing $e_{\lambda_1} \star \cdots \star e_{\lambda_\ell} = \sum_{\mu} c_{\lambda\mu}(s, t) e_{\mu}$, we show that $c_{\lambda\mu}$ is supported exactly on the dominance up-set of λ with s -adic valuation $n(\mu)$, and that the $(s - 1)$ -adic valuation of $c_{\lambda\mu}$ is $\ell(\lambda) - \kappa(\lambda, \mu)$, where κ is the maximal number of blocks into which λ and μ split compatibly. Leading coefficients factor over blocks into connected pieces; for a full merge the connected piece is a connected-graph cumulant. The coefficients are invariant, up to a power of s , under complementation in a box, and all second-order leading coefficients are given in closed form via a residue formula for Green polynomials at two-part classes.

CONTENTS

1. Introduction	1
2. Hikita's product and the subset formula	3
3. The Pieri package	6
4. Dominance support	11
5. The $s \rightarrow 0$ edge	14
6. The block valuation law at $s = 1$	16
7. Leading coefficients: graph cumulants and blocks	21
8. Box Complement and the linear coefficient	23
9. Second order: every valuation-2 lead	24
10. Open problems	28
References	29

1. INTRODUCTION

Hikita [Hik25] defined a commutative, associative product $\star$ on symmetric functions over $\mathbb{Q}(q, t)$ by transporting the ordinary product through the level-one polynomial representation of the affine Hecke algebra. We write $s := q^{-1}$. At $s = 1$, $\star$ is the ordinary product. Hikita's Pieri rule [Hik25, Thm. 3.12] reads, in our normalisation,

$$e_1 \star e_r = (1 - s)[r + 1]_t e_{r+1} + s e_1 e_r.$$

Date: October 9, 2026.

[WIP: Authorship, affiliation, acknowledgements and the AI declaration are to be settled as for the extended abstract.]

Di Francesco and Kedem [DFK19, §8.3] observed that the associated t -deformed graded characters lose Schur positivity at generic t . This paper is about the structure that survives: explicit multiplication rules, triangularity, and valuations.

Pieri rules. The operator $e_k \star (\cdot)$ is a sum over k -subsets (Theorem 2.5),

$$e_k \star F = \sum_{|A|=k} \prod_{i \in A, j \notin A} \frac{x_i - tx_j}{x_i - x_j} \prod_{i \in A} x_i \cdot F|_{x_i \mapsto sx_i \ (i \in A)}.$$

From it we derive a closed generating function for $e_k \star (e_{a_1} \cdots e_{a_\ell})$ for all k, ℓ (Theorem 3.2). Since the e_a generate, this determines $e_k \star F$ for every F . The case $\ell = 1$ is a Pieri rule (Corollary 3.3):

$$e_k \star e_r = \sum_{b=0}^k s^b F_{k-b}(t^{r-b}) e_b e_{r+k-b},$$

with explicit polynomials $F_n(w)$, and at $t = 0$ the coefficients form a truncated geometric distribution (Corollary 3.6). In the generating function, the ℓ columns interact only through a pairwise cross kernel, and the proof closes with a single residue identity (Lemma 3.9).

Dominance and valuations. Write $e_\lambda^\star := e_{\lambda_1} \star \cdots \star e_{\lambda_\ell} = \sum_\mu c_{\lambda\mu}(s, t) e_\mu$. A termwise degree count on the subset formula shows that $c_{\lambda\mu} \in \mathbb{Q}[s, t]$ is supported exactly on the dominance up-set of λ , with s -adic valuation exactly $n(\mu)$ (Theorem 4.1). In the rescaled basis $b_\mu = s^{n(\mu)} e_\mu$, the operator $e_k \star (\cdot)$ reduces modulo s to Hall–Littlewood multiplication by $t^{-\binom{k}{2}} e_k$, so the s -leading coefficients form the classical $e \rightarrow$ Hall–Littlewood transition matrix (Theorem 5.1 and Corollary 5.2).

At $s = 1$, $\star$ is the ordinary product, and the variable s enters the subset formula only through an Euler operator; so the $(s - 1)^p$ Taylor piece of $e_k \star (\cdot)$ is a differential operator of order p . The first-order term is a symmetric biderivation with an explicit matrix (Theorem 6.1 and Proposition 6.2), and an order- p piece can glue at most p existing blocks of parts. This gives the *block valuation law*

$$v_{s-1}(c_{\lambda\mu}) = \ell(\lambda) - \kappa(\lambda, \mu) \quad (\mu \succeq \lambda, \mu \neq \lambda),$$

where κ is the maximal number of compatible blocks (Theorem 6.6); the lower bound holds for all t , and exactness is settled at $t = 0$, where no cancellation occurs. The leading coefficients are sums over merge histories with explicit weights (Theorems 6.9 and 6.10). For a full merge the lead is $\frac{1-t^n}{\prod_i (1-t^{\lambda_i})} \sum_{H \text{ connected}} \prod_{ij \in H} (t^{\lambda_i \lambda_j} - 1)$ (Theorem 7.4), whose graph half is due to Dołęga [Dol19], and every lead is a sum of products of connected leads (Theorems 7.7 and 7.8). The coefficients satisfy $c_{N^\ell - \lambda, N^\ell - \mu} = s^{N \binom{\ell}{2} - (\ell-1)|\lambda|} c_{\lambda\mu}$ (Theorem 8.1). Finally, every leading coefficient at valuation 2 is given in closed form (Theorem 9.7 and Corollary 9.9), via a constant-term formula and a residue evaluation of Green polynomials at two-part classes (Theorems 9.2 and 9.3).

Organisation. Section 2 sets up Hikita’s product and proves the subset formula. Section 3 contains the Pieri package. Section 4 proves dominance support, and Section 5 identifies the $s \rightarrow 0$ edge. Sections 6 and 7 treat the expansion at $s = 1$, Section 8 the box symmetry and the linear coefficient, and Section 9 the second-order leading coefficients.

2. HIKITA'S PRODUCT AND THE SUBSET FORMULA

2.1. Notation. For $m \geq 0$ let $\Lambda_m := \mathbb{Q}(s, t)[x_1, \dots, x_m]^{S_m}$ and let Λ be the ring of symmetric functions over $\mathbb{Q}(s, t)$. We write e_r for the elementary symmetric functions ($e_0 = 1$, $e_r = 0$ for $r < 0$), $e_\lambda = e_{\lambda_1} e_{\lambda_2} \cdots$, and

$$E(z) := \prod_{i=1}^m (1 + x_i z) = \sum_{r \geq 0} e_r z^r, \quad Q(y) := \frac{E(-ty)}{E(-y)} = \prod_{i=1}^m \frac{1 - tx_i y}{1 - x_i y} = \sum_{n \geq 0} q_n y^n,$$

so that the q_n are the one-row Hall–Littlewood functions of [Mac95, III (2.10)]. For a set B of indices, $E(x_B; z) := \prod_{i \in B} (1 + x_i z)$, and $\hat{x}_i$ denotes the variables $x_1, \dots, x_m$ with x_i omitted. We put

$$a_{ij} := \frac{x_i - tx_j}{x_i - x_j}, \quad c_A := \prod_{i \in A, j \notin A} a_{ij}, \quad x_A := \prod_{i \in A} x_i \quad (A \subseteq [m]).$$

Further, $[n] = [n]_t = (1 - t^n)/(1 - t)$, $[n]! = [1][2] \cdots [n]$, $(a; t)_n = \prod_{i=0}^{n-1} (1 - at^i)$, $n(\lambda) = \sum_i (i - 1)\lambda_i$, and $\geq$ is dominance order.

2.2. The level-one representation and Hikita's product. The symmetric group S_m acts on $\mathbb{Q}(s, t)[x_1, \dots, x_m]$ by permuting variables; s_i is the transposition of x_i and x_{i+1} . Define operators on $\mathbb{Q}(s, t)(x_1, \dots, x_m)$ by

$$\begin{aligned} T_i f &:= t s_i f + (t - 1) \frac{x_{i+1}}{x_i - x_{i+1}} (s_i f - f) \quad (1 \leq i \leq m - 1), \\ \pi f(x_1, \dots, x_m) &:= x_1 f(x_2, \dots, x_m, s x_1), \\ Y_i &:= t^{m-i} T_{i-1} \cdots T_1 \pi T_{m-1}^{-1} \cdots T_i^{-1} \quad (1 \leq i \leq m). \end{aligned}$$

The T_i are Demazure–Lusztig operators; they preserve polynomials, satisfy $(T_i - t)(T_i + 1) = 0$ and the braid relations, and $T_i f = t f$ whenever $s_i f = f$. For $1 \leq k \leq m$ put

$$e_k(Y) := \sum_{b_1 < \cdots < b_k} Y_{b_1} Y_{b_2} \cdots Y_{b_k},$$

each product taken in increasing order of indices from left to right. (The Y_i commute, but the proof of Theorem 2.5 does not use this.) Hikita's product $F \star G$ is defined by transporting the product through the map $F(Y) \mapsto F(Y) \bullet 1$; by [Hik25, Lemma 3.1, Cor. 3.9] this map is a linear isomorphism compatible with $m \rightarrow m - 1$, so $\star$ is well defined on Λ , and it is commutative and associative [Hik25]. We use $\star$ only through the following consequence of the definitions [Hik25, Def. 3.4, Lemma 3.3]: for symmetric F ,

$$(2.1) \quad e_k \star F = t^{-\binom{k}{2}} e_k(Y) F.$$

2.3. Coxeter preliminaries. We use standard facts about S_m as a Coxeter group with generators $s_1, \dots, s_{m-1}$ [Hum90, §§1.10, 5.12], [BB05, §2.4]. Words act as compositions of operators, the rightmost factor first.

- (H1) For a reduced word $w = s_{i_1} \cdots s_{i_\ell}$ the operator $T_w := T_{i_1} \cdots T_{i_\ell}$ depends only on w , and $T_u T_v = T_{uv}$ whenever $\ell(uv) = \ell(u) + \ell(v)$.
- (H2) For $J \subseteq \{s_1, \dots, s_{m-1}\}$ let W_J be the parabolic subgroup and $W^J = \{w : \ell(ws) > \ell(w) \ \forall s \in J\}$. Every w factors uniquely as $w = w^J w_J$ with $w^J \in W^J$, $w_J \in W_J$, and $\ell(w) = \ell(w^J) + \ell(w_J)$.
- (H3) For $K \subseteq J$, multiplication $W^J \times (W_J)^K \rightarrow W^K$ is a length-additive bijection.
- (H4) If $T_j H = t H$ for all $s_j \in J$, then $T_v H = t^{\ell(v)} H$ for all $v \in W_J$.
- (H5) $\pi T_j = T_{j+1} \pi$ for $1 \leq j \leq m - 2$.

(H3) is standard: if $u \in W^J$, $v \in (W_J)^K$ and $s \in K$, then $\ell(uvs) = \ell(u) + \ell(vs) = \ell(uv) + 1$; conversely, for $x \in W^K$ write $x = x^J x_J$, and $x_J s < x_J$ for some $s \in K$ would give $\ell(xs) < \ell(x)$. (H4) is immediate from (H1). For (H5), $\pi s_j f = x_1 f(x_2, \dots, x_{j+2}, x_{j+1}, \dots, s x_1) = s_{j+1} \pi f$ because $j+2 \leq m$, and π turns multiplication by $x_{j+1}/(x_j - x_{j+1})$ into multiplication by $x_{j+2}/(x_{j+1} - x_{j+2})$.

Let $J_n := \{s_i : i \neq n\}$, so $W_{J_n} = S_n \times S_{m-n}$, and the coset wW_{J_n} is determined by $D = w([n])$. Its minimal element w^D is increasing on $[n]$ and on $[n+1, m]$, with $\ell(w^D) = \sum_l (d_l - l)$ for $D = \{d_1 < \dots < d_n\}$. The word

$$w(D) := (s_{d_1-1} \dots s_1)(s_{d_2-1} \dots s_2) \dots (s_{d_n-1} \dots s_n)$$

sends $l \mapsto d_l$ for $l \leq n$: the blocks with index $l' > l$ fix l , block l sends l to d_l , and the blocks $l' < l$ only move points $\leq d_{l'} < d_l$. It has length $\leq \sum_l (d_l - l)$, so $w(D) = w^D$ and the word is reduced. We write $T_{w(D)}$ for the corresponding operator and

$$\sigma^{(k)} := \sum_{|D|=k} T_{w(D)}.$$

2.4. The parabolic kernel.

Lemma 2.1. *Let $0 \leq n \leq m$, $1 \leq c < d_1 < \dots < d_n \leq m$, and let H satisfy $T_j H = tH$ for all $j \neq n$. Then*

$$(T_c T_{c+1} \dots T_{m-1})^{-1} T_{w(D)} H = t^{n-(m-c)} T_{w(D-1)} H, \quad D-1 := \{d_1 - 1, \dots, d_n - 1\}.$$

Proof. Let $x := s_c s_{c+1} \dots s_{m-1}$; it is reduced of length $m - c$, sends $m \mapsto c$ and $i \mapsto i + 1$ for $c \leq i \leq m-1$, and fixes $i < c$. Let $y := w(D-1)$; the word is valid since $d_l - 1 \geq c + l - 1 \geq l$, and $\ell(y) = \ell(w^D) - n$. As $m \notin D-1$, the value m is the largest on the increasing block $[n+1, m]$, so $y(m) = m$. Since x is increasing on $[1, m-1]$, the inversions of y among positions $< m$ survive in xy ; at position m , $xy(m) = c$ is inverted with exactly those $i < m$ having $y(i) \geq c$, and there are $m - c$ of them. Hence $\ell(xy) = \ell(y) + m - c$. Finally $xy([n]) = x(D-1) = D$ because every $d_l - 1 \geq c$, so $xy = w^D v$ with $v \in W_{J_n}$ and $\ell(v) = \ell(xy) - \ell(w^D) = m - c - n$. By (H1) and (H4), $T_x T_y H = T_{xy} H = T_{w^D} T_v H = t^{m-c-n} T_{w^D} H$. Apply T_x^{-1} . $\square$

Proposition 2.2 (Parabolic kernel). *For symmetric F and $1 \leq b_1 < \dots < b_k \leq m$, $Y_{b_1} Y_{b_2} \dots Y_{b_k} F = t^{\binom{k}{2}} T_{w(\{b_1, \dots, b_k\})} \pi^k F$. Hence*

$$e_k(Y) F = t^{\binom{k}{2}} \sigma^{(k)} \pi^k F.$$

Proof. Note that $\pi^n F = x_1 \dots x_n F(x_{n+1}, \dots, x_m, s x_1, \dots, s x_n)$ is symmetric in $x_1, \dots, x_n$ and in $x_{n+1}, \dots, x_m$, so $T_j \pi^n F = t \pi^n F$ for $j \neq n$. We induct on the number of factors, the empty product being trivial. Let $c < d_1 < \dots < d_n$ and assume $Y_{d_1} \dots Y_{d_n} F = t^{\binom{n}{2}} T_{w(D)} H$ with $H = \pi^n F$. By Lemma 2.1,

$$Y_c t^{\binom{n}{2}} T_{w(D)} H = t^{\binom{n}{2}} t^{m-c} T_{c-1} \dots T_1 \pi (T_c \dots T_{m-1})^{-1} T_{w(D)} H = t^{\binom{n}{2}+n} T_{c-1} \dots T_1 \pi T_{w(D-1)} H.$$

All indices in the word $w(D-1) = \prod_l (s_{d_l-2} \dots s_l)$ lie in $[1, m-2]$, so by (H5), $\pi T_{w(D-1)} = \prod_l (T_{d_l-1} \dots T_{l+1}) \pi$. Now $T_{c-1} \dots T_1 \prod_l (T_{d_l-1} \dots T_{l+1}) = T_{w(\{c\} \cup D)}$, $\pi H = \pi^{n+1} F$, and $\binom{n}{2} + n = \binom{n+1}{2}$. $\square$

2.5. The k -set kernel. Let $\sigma_{[l,m]} := \sum_{a=l}^m T_{a-1} \dots T_l$ be the partial symmetriser on $x_l, \dots, x_m$.

Lemma 2.3. *Let F be symmetric in $x_2, \dots, x_m$, with coefficients in any field containing $\mathbb{Q}(t)$, and let $F^{(i)}$ be F with x_1 and x_i swapped. Then $\sigma_{[1,m]} F = \sum_{i=1}^m F^{(i)} \prod_{j \neq i} a_{ij}$.*

Proof. Induction on m ; $m = 1$ is trivial. Since $s_1 F = F^{(2)}$, a direct computation gives $T_1 F = a_{21} F^{(2)} + (a_{12} - 1) F^{(1)}$. Write $\sigma_{[1,m]} = 1 + \sigma_{[2,m]} T_1$. The function $G := T_1 F$ is symmetric in $x_3, \dots, x_m$, so by induction (over the field with x_1 adjoined) $\sigma_{[2,m]} G = \sum_{i \geq 2} G|_{x_2 \leftrightarrow x_i} \prod_{j \geq 2, j \neq i} a_{ij}$. Under $x_2 \leftrightarrow x_i$, $F^{(2)} \mapsto F^{(i)}$, $F^{(1)}$ is fixed, $a_{21} \mapsto a_{i1}$ and $a_{12} \mapsto a_{1i}$. Hence

$$\sigma_{[1,m]} F = \sum_{i \geq 2} F^{(i)} \prod_{j \neq i} a_{ij} + F^{(1)} \left[1 + \sum_{i \geq 2} (a_{1i} - 1) \prod_{j \geq 2, j \neq i} a_{ij} \right].$$

The function $R(x) := \prod_{j \geq 2} (x - tx_j)/(x - x_j)$ tends to 1 at ∞ and has simple poles at $x = x_i$ with residues $(1-t)x_i \prod_{j \geq 2, j \neq i} a_{ij}$; so $R(x) = 1 + \sum_{i \geq 2} \frac{(1-t)x_i}{x-x_i} \prod_{j \geq 2, j \neq i} a_{ij}$. Evaluating at $x = x_1$ and using $(1-t)x_i/(x_1 - x_i) = a_{1i} - 1$ shows that the bracket equals $R(x_1) = \prod_{j \geq 2} a_{1j}$. $\square$

Proposition 2.4 (*k-set kernel*). *Let G be symmetric in $x_1, \dots, x_k$ and in $x_{k+1}, \dots, x_m$, and let $G^{(A)}$ denote G with the variables x_A in the first k slots and $x_{[m] \setminus A}$ in the last $m - k$. Then*

$$\sigma^{(k)} G = \sum_{|A|=k} c_A G^{(A)}.$$

Proof. Let $K_l := \{s_{l+1}, \dots, s_{m-1}\}$. The minimal coset representatives of $W_{K_{l-1}}/W_{K_l}$ are $s_{a-1} \cdots s_l$ ($l \leq a \leq m$), so applying (H3) along $K_k \subset K_{k-1} \subset \cdots \subset K_0$ gives $\sum_{u \in W^{K_k}} T_u = \sigma_{[1,m]} \sigma_{[2,m]} \cdots \sigma_{[k,m]}$. Applying (H3) to $K_k \subset J_k$, with $(W_{J_k})^{K_k} = S_k$, gives $\sum_{u \in W^{K_k}} T_u = \sigma^{(k)} \sum_{v \in S_k} T_v$. For G as in the statement, (H4) gives $\sum_{v \in S_k} T_v G = [k]_t! G$, so

$$(2.2) \quad \sigma_{[1,m]} \cdots \sigma_{[k,m]} G = [k]_t! \sigma^{(k)} G.$$

On the other hand, if G is merely symmetric in $x_{k+1}, \dots, x_m$, then iterating Lemma 2.3 gives

$$(2.3) \quad \sigma_{[1,m]} \cdots \sigma_{[k,m]} G = \sum_{(a_1, \dots, a_k) \text{ distinct}} G(x_{a_1}, \dots, x_{a_k}; \text{rest}) \omega(a), \quad \omega(a) := \prod_{l=1}^k \prod_{j \notin \{a_1, \dots, a_l\}} a_{a_l j}.$$

Indeed, by induction on k in the variables $x_2, \dots, x_m$, $H := \sigma_{[2,m]} \cdots \sigma_{[k,m]} G$ is the analogous sum over distinct tuples in $[2, m]$; it is symmetric in $x_2, \dots, x_m$, since permuting those variables permutes the terms; and Lemma 2.3 applied to H , followed by $x_1 \leftrightarrow x_{a_1}$, turns tuples in $[2, m]$ into tuples in $[m] \setminus \{a_1\}$.

Now let G be symmetric in both groups. Then $G(x_{a_1}, \dots, x_{a_k}; \text{rest}) = G^{(A)}$ with $A = \{a_1, \dots, a_k\}$, and

$$(2.4) \quad \omega(a) = c_A \prod_{1 \leq l < l' \leq k} a_{a_l a_{l'}}.$$

By the Poincaré identity $\sum_{w \in S_k} w(\prod_{i < j \leq k} a_{ij}) = [k]_t! [\text{Mac95, III (1.4)}]$ (which is also (2.2)=(2.3) at $m = k$, $G = 1$), summing over the orderings of A gives $[k]_t! c_A G^{(A)}$. Compare with (2.2). $\square$

Theorem 2.5 (Subset formula). *For every $m \geq 0$, $k \geq 1$ and $F \in \Lambda_m$,*

$$e_k \star F = t^{-\binom{k}{2}} e_k(Y) F = E_k F := \sum_{|A|=k} c_A x_A F|_{x_i \mapsto s x_i \ (i \in A)}.$$

Proof. By Propositions 2.2 and 2.4 with $G = \pi^k F$, since $(\pi^k F)^{(A)} = x_A F|_{x_i \mapsto s x_i \ (i \in A)}$; the first equality is (2.1). For $m < k$ both sides vanish. $\square$

Remark 2.6. For $k = 1$ the subset formula gives back Hikita's rule $e_1 \star e_r = (1-s)[r+1]e_{r+1} + s e_1 e_r$ [Hik25, Thm. 3.12]; see Example 3.4. Proposition 2.4 is the level-one form of the classical expression of the Hecke symmetriser as $\sum_w w \circ \prod_{i < j} a_{ij}$, restricted to minimal coset representatives.

Lemma 2.7. $E_k(1) = \sum_{|A|=k} c_A x_A = e_k$.

Proof. Grouping $w \in S_m$ by $w(\{1, \dots, k\}) = A$ and applying the Poincaré identity to $S_k \times S_{m-k}$ shows that $\sum_{|A|=k} c_A x_A$ is the Hall–Littlewood function $P_{(1^k)}(x; t)$ as defined in [Mac95, III (2.1)–(2.2)], and $P_{(1^k)} = e_k$ [Mac95, III (2.8)]. $\square$

Lemma 2.8 (Stability). *For $F \in \Lambda_m$, $(E_k F)|_{x_m=0} = E_k(F|_{x_m=0})$, the right side computed in $m-1$ variables. Hence the e -coefficients of any expression built from the E_k and 1 are independent of m once m is at least the degree.*

Proof. Terms with $m \in A$ vanish because of the factor x_A ; for $m \notin A$, $a_{im}|_{x_m=0} = 1$. Restriction $\Lambda_m \rightarrow \Lambda_{m-1}$ maps $e_\nu \mapsto e_\nu$ and is injective in degrees $\leq m-1$. $\square$

2.6. Two consequences of the kernel. The next two lemmas drive every computation in Section 3. For a function φ of one variable put $S_k^\varphi(x_1, \dots, x_m) := \sum_{|A|=k} c_A \prod_{a \in A} \varphi(x_a)$.

Lemma 2.9 (Peeling one variable). *For $m \geq 1$ and $k \geq 1$, $[k] S_k^\varphi(x) = \sum_{i=1}^m \varphi(x_i) \prod_{j \neq i} a_{ij} S_{k-1}^\varphi(\hat{x}_i)$.*

Proof. By (2.4) and the Poincaré identity, $[k]_t! S_k^\varphi = \sum_a \omega(a) \prod_l \varphi(x_{a_l})$, the sum over distinct k -tuples. Separate $a_1 = i$: $\omega(a) = \prod_{j \neq i} a_{ij} \cdot \omega'(a_2, \dots, a_k)$, where ω' is the weight of (2.3) for the variables $\hat{x}_i$. Summing over $(a_2, \dots, a_k)$ gives $[k-1]_t! S_{k-1}^\varphi(\hat{x}_i)$. $\square$

Lemma 2.10 (Residue form of the HL kernel). *Let $\mathbb{K} \supseteq \mathbb{Q}(t)$ be a field and $H \in \mathbb{K}(v)$ with $H(0) = 0$, such that $H(v)/v$ has only simple poles, at points $p \in \mathbb{K}$, and $H(v)/v = O(1/v)$ at ∞ . Then in $\mathbb{K}(x_1, \dots, x_m)$*

$$(1-t) \sum_{i=1}^m H(x_i) \prod_{j \neq i} a_{ij} = - \sum_p \operatorname{Res}_{v=p} f - \operatorname{Res}_{v=\infty} f, \quad f(v) := \frac{H(v)}{v} \prod_{j=1}^m \frac{v - tx_j}{v - x_j}.$$

Proof. f is regular at $v = 0$, its residue at $v = x_i$ is $(1-t)H(x_i) \prod_{j \neq i} a_{ij}$, and the sum of all residues of a rational function is zero. $\square$

For $H(v) = v^n$, $n \geq 1$, this is $(1-t) \sum_i x_i^n \prod_{j \neq i} a_{ij} = q_n$ [Mac95, III (2.10)]. Note that $\prod_j (v - tx_j)/(v - x_j) = Q(1/v)$ and $Q(-a) = E(ta)/E(a)$.

3. THE PIERI PACKAGE

3.1. The coefficients. For $j \geq 0$ put

$$\alpha_j := \prod_{i=1}^j \frac{s - t^i}{1 - t^i} \quad (\alpha_{-1} := 0), \quad c(n, j) := \frac{(s; t)_{n-j}}{(t; t)_{n-j}} (\alpha_j - s t^{n-j} \alpha_{j-1}) \quad (0 \leq j \leq n),$$

and $c(n, j) := 0$ otherwise. Let $F_n(w) := \sum_{j=0}^n c(n, j) w^j$. For instance $F_0 = 1$ and $F_1(w) = (1-s)(1-tw)/(1-t)$.

Lemma 3.1. *Let $G(x) := \sum_{n \geq 0} \frac{(s; t)_n}{(t; t)_n} x^n$, $C_j(x) := \sum_n c(n, j) x^n$, and $D_j := \alpha_j - s \alpha_{j-1}$. Then*

$$(1-x)G(x) = (1-sx)G(tx), \quad C_j(x) = D_j x^j G(tx) \frac{1 - st^{-j}x}{1-x},$$

$D_0 = 1$, $D_j = \alpha_{j-1} t^j (s-1)/(1-t^j) \neq 0$ for $j \geq 1$, and $(1-t^j)D_j = (ts - t^j)D_{j-1}$ for $j \geq 1$.

Proof. Write $g_n = (s; t)_n / (t; t)_n$. The first identity says $g_n - g_{n-1} = t^n g_n - s t^{n-1} g_{n-1}$, i.e. $(1-t^n)g_n = (1-s t^{n-1})g_{n-1}$. By definition $C_j(x) = x^j (\alpha_j G(x) - s \alpha_{j-1} G(tx)) = x^j G(tx) (\alpha_j (1-sx) - s \alpha_{j-1} (1-x)) / (1-x)$. The bracket is linear in x . It equals D_j at $x = 0$, and at $x = t^j/s$ it equals $\alpha_j (1-t^j) - \alpha_{j-1} (s-t^j) = 0$ (for $j = 0$ use $\alpha_{-1} = 0$). So it equals $D_j (1-st^{-j}x)$. The formula for D_j follows from $\alpha_j = \alpha_{j-1} (s-t^j)/(1-t^j)$, and the recursion from it and the same relation one step down (for $j = 1$ use $D_0 = 1$). $\square$

3.2. Statements. Fix $\ell \geq 1$ and indeterminates $z_1, \dots, z_\ell$. For integers $i, j \geq 0$ let

$$K_{ij}(z, w) := \prod_{p=0}^{i-1} \frac{t^p z - s w}{t^p z - t^j w} \prod_{r=0}^{j-1} \frac{s z - t^r w}{t^i z - t^r w},$$

and note $K_{ij}(z, w) = K_{ji}(w, z)$. For $I = (i_1, \dots, i_\ell) \in \mathbb{Z}_{\geq 0}^\ell$ with $|I| = S$ put $K_I := \prod_{c < c'} K_{i_c i_{c'}}(z_c, z_{c'})$ and

$$V_I^{(n)} := K_I s^{(\ell-1)(n-S)} \sum_{n_1 + \dots + n_\ell = n} t^{-\sum_{c \neq c'} (n_c - i_c) i_{c'}} \prod_{c=1}^{\ell} t^{-n_c i_c} c(n_c, i_c) z_c^{-n_c},$$

with $V_I^{(n)} := 0$ if some $i_c < 0$. Since $c(n_c, i_c) = 0$ for $n_c < i_c$, $V_I^{(n)} = 0$ unless $S \leq n$.

Theorem 3.2 (ℓ -column rule). *For all $m, k \geq 0$ and $\ell \geq 1$, in $\Lambda_m \otimes \mathbb{Q}(s, t)(z_1, \dots, z_\ell)$,*

$$\mathcal{E}_k := \sum_{a \in \mathbb{Z}_{\geq 0}^\ell} z_1^{a_1} \cdots z_\ell^{a_\ell} e_k \star (e_{a_1} \cdots e_{a_\ell}) = \sum_{b=0}^k \sum_I (s^\ell t^{-|I|})^b V_I^{(k-b)} e_b \prod_{c=1}^{\ell} E(t^{i_c} z_c).$$

In particular the right-hand side is a polynomial in $z_1, \dots, z_\ell$, and $e_k \star (e_{a_1} \cdots e_{a_\ell})$ is its coefficient of z^a . The coefficients do not depend on m , so the identity holds in Λ .

Corollary 3.3 (Pieri rule for $e_k \star e_r$). *For all $k \geq 1$ and $r \geq 0$,*

$$e_k \star e_r = \sum_{b=0}^k s^b F_{k-b}(t^{r-b}) e_b e_{r+k-b}.$$

Equivalently, $\sum_r z^r e_k \star e_r = \sum_{j+b \leq k} s^b t^{-kj} c(k-b, j) e_b z^{b-k} E(t^j z)$.

Proof. Take $\ell = 1$ in Theorem 3.2: $K_I = 1$, and $V_{(j)}^{(n)} = t^{-nj} c(n, j) z^{-n}$, so the right-hand side is the stated generating function. Extract $[z^r]$ using $[z^r] z^{b-k} E(t^j z) = t^{j(r+k-b)} e_{r+k-b}$. $\square$

For $r < k$, distinct b may give the same monomial $e_b e_{r+k-b}$; the identity is between the sums. Since $\star$ is commutative, applying Corollary 3.3 to $e_r \star e_k$ shows that the support of $e_k \star e_r$ is contained in $\{e_b e_{r+k-b} : 0 \leq b \leq \min(k, r)\}$ (see also Theorem 4.1).

Example 3.4 ($k = 1$). $c(1, 0) = \frac{1-s}{1-t}$ and $c(1, 1) = \frac{t(s-1)}{1-t}$, so $F_1(t^r) = (1-s)[r+1]$ and Corollary 3.3 is Hikita's rule $e_1 \star e_r = (1-s)[r+1]e_{r+1} + s e_1 e_r$ [Hik25, Thm. 3.12].

Example 3.5 ($k = 2$). One computes $F_2(w) = \frac{(1-s)(1-t^2 w)(1-st+(s-t)w)}{(1-t)(1-t^2)}$, so for all $r \geq 0$

$$e_2 \star e_r = \frac{(1-s)(1-t^{r+2})(1-st+(s-t)t^r)}{(1-t)(1-t^2)} e_{r+2} + s(1-s)[r] e_1 e_{r+1} + s^2 e_2 e_r.$$

For $r = 0$ this is $e_2 \star 1 = e_2$.

Corollary 3.6 (The case $t = 0$). *For $k \geq 1$, $r \geq 0$, put $\mu = \min(k, r)$ and $M = \max(k, r)$. The coefficients of $e_k \star e_r$ are regular at $t = 0$, and*

$$(e_k \star e_r)|_{t=0} = \sum_{b=0}^{\mu-1} (1-s) s^b e_b e_{r+k-b} + s^\mu e_\mu e_M.$$

For $s \in [0, 1]$ the weights $(1-s)s^b$ ($b < \mu$) and s^μ are the law of $\min(G, \mu)$ for a geometric variable G with $P(G = b) = (1-s)s^b$, and the monomials $e_b e_{r+k-b}$ ($b < \mu$), $e_\mu e_M$ are pairwise distinct. So at $t = 0$, $e_k \star e_r$ is a truncated-geometric average of ordinary products.

Corollary 3.7 (Two columns; bounded support). *For $\ell = 2$, Theorem 3.2 reads*

$$\sum_{a,b \geq 0} z^a w^b e_k \star (e_a e_b) = \sum_{b_0, i, j} s^{2b_0} t^{-b_0(i+j)} V_{ij}^{(k-b_0)} e_{b_0} E(t^i z) E(t^j w),$$

$$V_{ij}^{(n)} = K_{ij}(z, w) \sum_{n_1+n_2=n} s^{n-i-j} t^{-(n_1-i)j-(n_2-j)i-n_1i-n_2j} c(n_1, i) c(n_2, j) z^{-n_1} w^{-n_2}.$$

In general, $e_k \star (e_{a_1} \cdots e_{a_\ell})$ is a linear combination of products of at most $\ell + 1$ elementary symmetric functions.

Proof. The first part is the case $\ell = 2$. For the second, expand both sides of Theorem 3.2 in the field of iterated Laurent series $\mathbb{Q}(s, t)((z_\ell)) \cdots ((z_1))$. The left side is a polynomial, so its expansion is itself. On the right, each term is $R(z) e_b \prod_c E(t^{i_c} z_c)$ with R rational and free of x , and its coefficient of z^a is a finite $\mathbb{Q}(s, t)$ -combination of products $e_b \prod_{c=1}^\ell e_{n_c}$. $\square$

3.3. The recursion. By Theorem 2.5, $e_k \star (e_{a_1} \cdots e_{a_\ell}) = \sum_{|A|=k} c_A x_A \prod_c e_{a_c}(x_{A^c}, s x_A)$, and $\sum_a z^a \prod_c e_{a_c}(x_{A^c}, s x_A) = \prod_c \prod_{j \notin A} (1 + x_j z_c) \prod_{i \in A} (1 + s x_i z_c)$. Hence

$$(3.1) \quad \mathcal{E}_k(x) = \prod_c E(z_c) \cdot S_k^\varphi(x), \quad \varphi(u) := u \prod_{c=1}^\ell \frac{1 + s u z_c}{1 + u z_c}.$$

Since $\varphi(x_i) \prod_c E(z_c) = x_i \prod_c (1 + s x_i z_c) \cdot \prod_c E(\hat{x}_i; z_c)$, Lemma 2.9 gives, for $m \geq 1$ and $k \geq 1$,

$$(3.2) \quad [k] \mathcal{E}_k(x) = \sum_{i=1}^m x_i \prod_{c=1}^\ell (1 + s x_i z_c) \prod_{j \neq i} a_{ij} \mathcal{E}_{k-1}(\hat{x}_i),$$

where $\mathcal{E}_{k-1}(\hat{x}_i)$ denotes $\mathcal{E}_{k-1}$ computed in the $m-1$ variables $\hat{x}_i$. For $m=0$ we have $\mathcal{E}_0 = 1$ and $\mathcal{E}_k = 0$ for $k \geq 1$.

3.4. The step map. Fix I and put $\gamma_c := t^{i_c} z_c$, $C := s^\ell t^{-|I|} = s^\ell \prod_c z_c / \prod_c \gamma_c$, and

$$\mathcal{F}(v) := \frac{\prod_{c=1}^\ell (v - s z_c)}{v \prod_{c=1}^\ell (v - \gamma_c)} = \frac{C}{v} + \sum_{c=1}^\ell \frac{\eta_c}{v - \gamma_c}, \quad \eta_c := \frac{\prod_{c' \neq c} (\gamma_c - s z_{c'})}{\gamma_c \prod_{c' \neq c} (\gamma_c - \gamma_{c'})}.$$

Lemma 3.8 (Step map). *For $b \geq 0$, in $m \geq 1$ variables,*

$$(1-t) \sum_i x_i \prod_c (1 + s x_i z_c) \prod_{j \neq i} a_{ij} e_b(\hat{x}_i) \prod_c E(\hat{x}_i; \gamma_c) = C(1-t^{b+1}) e_{b+1} \prod_c E(\gamma_c) \\ - \sum_{n=0}^b e_n \sum_{c=1}^\ell \eta_c \gamma_c^{n-b-1} (E(t \gamma_c) - t^n E(\gamma_c)) \prod_{c' \neq c} E(\gamma_{c'}).$$

Proof. Write $u = x_i$. Then $e_b(\hat{x}_i) = [y^b] E(y)/(1+uy)$ and $E(\hat{x}_i; \gamma) = E(\gamma)/(1+\gamma u)$, so the left side is $\prod_c E(\gamma_c) \cdot [y^b] E(y) (1-t) \sum_i H(x_i; y) \prod_{j \neq i} a_{ij}$ with

$$H(v; y) := \frac{v \prod_c (1 + s v z_c)}{(1 + y v) \prod_c (1 + \gamma_c v)}$$

(the sum over i is finite and each term is regular at $y = 0$, so $[y^b]$ commutes with it). Apply Lemma 2.10 over $\mathbb{K} = \mathbb{Q}(s, t, z, y)$. The poles of H/v are simple, at $v = -1/a$ for $a \in \{y, \gamma_1, \dots, \gamma_\ell\}$, and $H(v)/v \sim C/(yv)$ at ∞ , which contributes C/y . At $v = -1/a$ the residue of f is $\rho_a E(ta)/E(a)$, where

$$\rho_a = \frac{\prod_c (a - s z_c)}{a \prod_{a' \neq a} (a - a')},$$

as one sees from $\prod_c(1 - sz_c/a) = a^{-\ell} \prod_c(a - sz_c)$, $\prod_{a' \neq a}(1 - a'/a) = a^{-\ell} \prod_{a' \neq a}(a - a')$ and $Q(-a) = E(ta)/E(a)$. Thus $\rho_y = \mathcal{F}(y)$ and $\rho_{\gamma_c} = -\eta_c/(y - \gamma_c)$, and

$$(1-t) \sum_i H(x_i; y) \prod_{j \neq i} a_{ij} = \frac{C}{y} - \mathcal{F}(y) \frac{E(ty)}{E(y)} + \sum_c \frac{\eta_c}{y - \gamma_c} \frac{E(t\gamma_c)}{E(\gamma_c)}.$$

Multiply by $E(y) \prod_c E(\gamma_c)$ and write $E(y) = \sum_n e_n y^n$, $E(ty) = \sum_n t^n e_n y^n$. The coefficient of e_n is $[y^{b-n}]$ of the Laurent expansion at $y = 0$ of

$$\begin{aligned} & \left(\frac{C}{y} - t^n \mathcal{F}(y) \right) \prod_c E(\gamma_c) + \sum_c \frac{\eta_c}{y - \gamma_c} E(t\gamma_c) \prod_{c' \neq c} E(\gamma_{c'}) \\ &= \frac{C(1-t^n)}{y} \prod_c E(\gamma_c) + \sum_c \frac{\eta_c}{y - \gamma_c} (E(t\gamma_c) - t^n E(\gamma_c)) \prod_{c' \neq c} E(\gamma_{c'}), \end{aligned}$$

by the partial fraction expansion of $\mathcal{F}$. Expanding $1/(y - \gamma) = -\sum_{p \geq 0} y^p \gamma^{-p-1}$ gives the claim: $n = b+1$ picks the pole at 0, $n \leq b$ picks the second sum, and $n > b+1$ contributes nothing. $\square$

3.5. Proof of Theorem 3.2. Write $\mathcal{T}_k$ for the right-hand side of Theorem 3.2, so $\mathcal{T}_k = \sum_{b,I} W_{b,I}^{(k)} e_b \prod_c E(t^{i_c} z_c)$ with $W_{b,I}^{(k)} = C_I^b V_I^{(k-b)}$, $C_I := s^\ell t^{-|I|}$; $\epsilon_1, \dots, \epsilon_\ell$ are the unit vectors of $\mathbb{Z}^\ell$.

Reduction to an index identity. We show the step identity

$$(3.3) \quad \sum_i x_i \prod_c (1 + s x_i z_c) \prod_{j \neq i} a_{ij} \mathcal{T}_{k-1}(\hat{x}_i) = [k] \mathcal{T}_k(x) \quad (m \geq 0, k \geq 1)$$

(for $m = 0$ the left side is empty). Given (3.3), the theorem follows by induction on k for all m at once: $\mathcal{E}_0 = \prod_c E(z_c) = \mathcal{T}_0$ because $V_0^{(0)} = 1$; for $m = 0$, (3.3) says $\mathcal{T}_k = 0 = \mathcal{E}_k$; for $m \geq 1$, (3.2), the induction hypothesis in the variables $\hat{x}_i$, and (3.3) give $[k] \mathcal{E}_k = [k] \mathcal{T}_k$.

Apply Lemma 3.8 to each term of $\mathcal{T}_{k-1}(\hat{x}_i)$. The term $e_{b'} \prod_c E(t^{i_c} z_c)$ on the right receives three kinds of contributions: from $(b' - 1, I)$ via the pole at 0; from (b, I) , $b \geq b'$, via $-t^{b'} E(\gamma_c)$; and from $(b, I - \epsilon_c)$, $b \geq b'$, via $E(t\gamma'_c)$ with $\gamma'_c = t^{i_c-1} z_c$. Writing η'_c for η_c computed at $I - \epsilon_c$, $n := k - b'$, $p := b - b'$, and using $C_{I-\epsilon_c} = t C_I$, the coefficient of $e_{b'} \prod_c E(t^{i_c} z_c)$ in $(1-t)$ times the left side of (3.3) is $C_I^{b'}$ times

$$(1 - t^{b'}) V_I^{(n)} + t^{b'} \sum_{p \geq 0} C_I^p \left[V_I^{(n-1-p)} \sum_c \eta_c \gamma_c^{-p-1} - t^p \sum_c V_{I-\epsilon_c}^{(n-1-p)} \eta'_c (t^{i_c-1} z_c)^{-p-1} \right].$$

So (3.3) follows from the index identity

$$(3.4) \quad \sum_{p \geq 0} C_I^p \left[V_I^{(n-1-p)} \sum_c \eta_c \gamma_c^{-p-1} - t^p \sum_c V_{I-\epsilon_c}^{(n-1-p)} \eta'_c (t^{i_c-1} z_c)^{-p-1} \right] = (1 - t^n) V_I^{(n)} \quad (n \geq 0),$$

because then the coefficient is $C_I^{b'} (1 - t^{b'+n}) V_I^{(n)} = (1 - t^k) W_{b',I}^{(k)}$. (No uniqueness of such expansions is needed: the two sides agree term by term.)

Generating function. Let $\mathcal{V}_I(x) := \sum_{n \geq 0} V_I^{(n)} x^n$. The p -sums in (3.4) are convolutions, and $\sum_{p \geq 0} C^p \gamma^{-p-1} x^{p+1} = x/(\gamma - Cx)$. So (3.4) for all n is equivalent to

$$(3.5) \quad \mathcal{V}_I(x) U(x) - \mathcal{V}_I(tx) = - \sum_c \eta'_c \frac{x}{t^{i_c-1} z_c - C_I t x} \mathcal{V}_{I-\epsilon_c}(x), \quad U(x) := 1 - \sum_c \frac{\eta_c x}{\gamma_c - C_I x}.$$

Put $X_c := C_I x / (sz_c)$ and $A_c := t^{i_c}$. Summing over (n_c) in the definition of $V_I^{(n)}$, the exponent of t attached to column c is $-n_c|I| + i_c(|I| - i_c)$, so

$$(3.6) \quad \mathcal{V}_I(x) = K_I s^{-(\ell-1)|I|} t^{|I|^2 - \sum_c i_c^2} \prod_c C_{i_c}(X_c),$$

and passing from I to $I - \epsilon_c$ replaces every $X_{c'}$ by $tX_{c'}$.

Step A: U is a product. By the partial fraction expansion of $\mathcal{F}$, $U(x) = x \mathcal{F}(C_I x)$. Since $C_I x - sz_c = -sz_c(1 - X_c)$, $C_I x - \gamma_c = -\gamma_c(1 - sX_c/A_c)$ and $C_I \prod_c \gamma_c = s^\ell \prod_c z_c$,

$$U(x) = \prod_{c=1}^{\ell} \frac{1 - X_c}{1 - sX_c/A_c}.$$

Step B: normalisation. Divide (3.5) by $\mathcal{N} := K_I s^{-(\ell-1)|I|} t^{|I|^2 - \sum_c i_c^2} \prod_c D_{i_c} X_c^{i_c} G(t^2 X_c)$, which is legitimate since $D_j \neq 0$. By Lemma 3.1 and $G(tX)/G(t^2 X) = (1 - stX)/(1 - tX)$, with $u_c := stX_c$:

- column c of $\mathcal{V}_I U$ contributes $\frac{(1-stX_c)(1-sX_c/A_c)}{(1-tX_c)(1-X_c)} \cdot \frac{1-X_c}{1-sX_c/A_c} = \frac{1-u_c}{1-tX_c}$, and column c of $\mathcal{V}_I(tx)$ contributes $\frac{A_c - u_c}{1-tX_c}$;
- in the c -th term on the right (present only if $i_c \geq 1$), the columns $c' \neq c$ contribute $\frac{A_{c'} - u_{c'}}{1-tX_{c'}}$; column c contributes $\frac{D_{i_c-1}}{D_{i_c}} \frac{A_c}{t} X_c^{-1} \frac{1-st^2 X_c/A_c}{1-tX_c}$ with $D_{i-1}/D_i = (1-A)/(ts-A)$; the prefactors contribute $s^{\ell-1} t^{-2(|I|-i_c)}$; and $\frac{x}{t^{i_c-1} z_c - C_I t x} = \frac{stX_c}{C_I(A_c - st^2 X_c)}$.

It remains to compute $\Phi_c := \eta'_c K_{I-\epsilon_c} / K_I$. The diagonal factor of η'_c is $(\gamma'_c - sz_c)/\gamma'_c = (A_c - st)/A_c$. For $c' \neq c$ put $z = z_c$, $w = z_{c'}$, $i = i_c$, $j = i_{c'}$, using $K_{ij}(z, w) = K_{ji}(w, z)$ to place c in the first slot. Then

$$\frac{t^{i-1}z - sw}{t^{i-1}z - t^j w} \cdot \frac{K_{i-1,j}(z, w)}{K_{ij}(z, w)} = \prod_{r=0}^{j-1} \frac{t^i z - t^r w}{t^{i-1}z - t^r w} = t^j \frac{t^i z - w}{t^i z - t^j w},$$

the last product telescoping after writing $t^i z - t^r w = t(t^{i-1}z - t^{r-1}w)$. With $\rho_{cc'} := z_c/z_{c'} = X_{c'}/X_c$ we get

$$\Phi_c = \frac{A_c - st}{A_c} \prod_{c' \neq c} A_{c'} \frac{A_c \rho_{cc'} - 1}{A_c \rho_{cc'} - A_{c'}}.$$

Multiplying everything, $\frac{stX_c}{C_I(A_c - st^2 X_c)} \cdot \frac{A_c}{t} X_c^{-1} (1 - st^2 X_c/A_c) = s/C_I$, then $\frac{s}{C_I} s^{\ell-1} t^{-2(|I|-i_c)} = t^{2i_c-|I|} = A_c / \prod_{c' \neq c} A_{c'}$, and $(A_c - st)/(ts - A_c) = -1$. Hence, after division by $\mathcal{N}$ and multiplication by $\prod_c (1 - tX_c)$, (3.5) becomes

$$(3.7) \quad \prod_c (1 - u_c) - \prod_c (A_c - u_c) - \sum_c (1 - A_c) \prod_{c' \neq c} \frac{A_c X_{c'} - X_c}{A_c X_{c'} - A_{c'} X_c} (A_{c'} - u_{c'}) = 0.$$

(For $i_c = 0$ the c -th term is absent, consistently with the factor $1 - A_c = 0$.)

Step C: the closing identity. Put $\lambda_c := A_c/u_c$ and $\mu_c := 1/u_c$. Since $X_c \propto u_c$, $\frac{A_c X_{c'} - X_c}{A_c X_{c'} - A_{c'} X_c} = \frac{\lambda_c - \mu_{c'}}{\lambda_c - \lambda_{c'}}$, and $(A_{c'} - u_{c'})/u_{c'} = \lambda_{c'} - 1$, $(1 - A_c)/u_c = \mu_c - \lambda_c$. Dividing (3.7) by $\prod_c u_c$ turns it into the case of Lemma 3.9 below. As rational functions of x , the $\lambda_c = t^{i_c} z_c / (t C_I x)$ are pairwise distinct and different from 1. Hence (3.5) holds, so (3.4) holds for every n , which proves (3.3) and the theorem. $\square$

Lemma 3.9. For indeterminates $\lambda_1, \dots, \lambda_\ell, \mu_1, \dots, \mu_\ell$,

$$\prod_c (\mu_c - 1) - \prod_c (\lambda_c - 1) = \sum_{c=1}^{\ell} (\mu_c - \lambda_c) \prod_{c' \neq c} (\lambda_{c'} - 1) \frac{\lambda_c - \mu_{c'}}{\lambda_c - \lambda_{c'}}.$$

Proof. The rational function $g(w) = \prod_c (w - \mu_c) / ((w - 1) \prod_c (w - \lambda_c))$ has residue $\frac{\lambda_c - \mu_c}{\lambda_c - 1} \prod_{c' \neq c} \frac{\lambda_c - \mu_{c'}}{\lambda_c - \lambda_{c'}}$ at λ_c , residue $\prod_c (1 - \mu_c) / \prod_c (1 - \lambda_c)$ at 1, and residue -1 at ∞ . The residues sum to zero; multiply by $\prod_c (\lambda_c - 1)$. $\square$

Remark 3.10. For $\ell = 1$, Lemma 3.9 is the tautology $(\mu - 1) - (\lambda - 1) = \mu - \lambda$, and the proof above reduces to a one-index recursion for $c(n, j)$. For $\ell = 2$ it is the polynomial identity that closes the two-column case. The cross kernel K_I is forced by the pairwise splitting of η'_c in Step B.

3.6. Proof of Corollary 3.6. At $t = 0$, $(t; t)_n = 1$, so every α_j and $c(n, j)$ is regular at $t = 0$; $(s; t)_n|_{t=0} = 1 - s$ for $n \geq 1$; hence $F_n(0) = c(n, 0)|_{t=0} = 1 - s$ for $n \geq 1$, and $F_0 = 1$.

Case $r \geq k$. Every $b \in [0, k]$ has $r - b \geq 0$, so each term $s^b F_{k-b}(t^{r-b}) = s^b \sum_j c(k - b, j) t^{j(r-b)}$ of Corollary 3.3 is regular at $t = 0$. If $b < r$, the terms with $j \geq 1$ vanish at $t = 0$, leaving $(1 - s)s^b$ for $b < k$ and s^k for $b = k$. If $b = r$, then $b = k = r$ and the term is $s^k F_0(1) = s^k$. Summing gives the claim with $\mu = k$.

Case $r < k$. By commutativity of $\star$ [Hik25], $e_k \star e_r = e_r \star e_k$. For $r = 0$ the claim is $e_k \star 1 = e_k$; for $r \geq 1$ apply the first case to (r, k) . $\square$

In the (k, r) -expansion of Corollary 3.3 with $r < k$, the terms with $b > r$ carry negative powers $t^{j(r-b)}$; the argument shows that these cancel once each monomial $e_b e_{r+k-b}$ is collected with its partner $b' = r + k - b$.

Remark 3.11 (Other forms and earlier cases). A second, independent proof of the Pieri rule, in the form $e_k \star e_r = \sum_y s^y C_{k-y, r-y} e_{k+r-y} e_y$ with $C_{a,b} = (-1)^b \frac{[a+b]}{[a]} \pi_a(b)$, is given in Section 8 (Corollary 8.6); it goes through the plethystic linear coefficient of Theorem 8.5 and the Column Lemma. The two forms agree after collecting terms. The kernel form of $e_k(Y)$ on symmetric functions is classical in spirit (Macdonald's D -operators); what is new here is the level-one transfer π^k , the explicit coefficients, and the ℓ -column rule.

4. DOMINANCE SUPPORT

For a partition λ write

$$e_\lambda^\star := e_{\lambda_1} \star \cdots \star e_{\lambda_\ell} = E_{\lambda_1} \cdots E_{\lambda_\ell}(1) = \sum_\mu c_{\lambda\mu}(s, t) e_\mu,$$

with E_k as in Theorem 2.5. The proofs below apply the operators E_{λ_i} in an arbitrary order, so they use neither commutativity nor associativity of $\star$.

Theorem 4.1 (Dominance support; the $s = 0$ valuation). $e_\lambda^\star = s^{n(\lambda)} e_\lambda + \sum_{\mu \triangleright \lambda} c_{\lambda\mu} e_\mu$ with $c_{\lambda\mu} \in \mathbb{Q}[s, t]$ and $c_{\lambda\mu}|_{s=1} = 0$. Moreover $c_{\lambda\mu} \neq 0$ for every $\mu \supseteq \lambda$, and the s -adic valuation of $c_{\lambda\mu}$ is exactly $n(\mu)$. At $t = 0$ the s -leading coefficients $[s^{n(\mu)}] c_{\lambda\mu}$ are all 1: in the basis $b_\mu = s^{n(\mu)} e_\mu$, $e_\lambda^\star$ at $s = t = 0$ is the zeta function of dominance order.

Corollary 4.2 (Operator form). For $k \geq 1$ and every partition μ , $e_k \star e_\mu \in s \sum_i \min(\mu_i, k) e_{\mu \cup k} + \text{span}\{e_\nu : \nu \triangleright \mu \cup k\}$.

We work in Λ_m with $m \geq n = |\lambda|$; by Lemma 2.8 this loses nothing.

4.1. Degrees and dominance.

Lemma 4.3. Let $m \geq n$ and $\rho \vdash n$. Then $\text{span}\{e_\nu : \nu \supseteq \rho'\} = \text{span}\{m_\kappa : \kappa \leq \rho\}$ in the degree- n part of Λ_m .

Proof. By [Mac95, I (6.6), (6.7)], $e_\nu = \sum_\kappa M_{\nu\kappa} m_\kappa$, where $M_{\nu\kappa}$ is the number of 0-1 matrices with row sums ν and column sums κ ; $M_{\nu\kappa} \neq 0$ forces $\kappa \leq \nu'$, and $M_{\nu\nu'} = 1$. Since $\nu \supseteq \rho' \iff \nu' \leq \rho$ [Mac95, I (1.11)], the left side is contained in the right. Both have dimension $\#\{\kappa \vdash n : \kappa \leq \rho\}$, because the e_ν , $\nu \vdash n$, are independent for $m \geq n$. $\square$

For $S \subseteq [m]$ and nonzero $R \in \mathbb{Q}(s, t)(x)$, let $\deg_S R$ be the degree in u of $R|_{x_i \mapsto ux_i (i \in S)}$ (numerator degree minus denominator degree). It is additive on products and satisfies $\deg_S(R_1 + R_2) \leq \max(\deg_S R_1, \deg_S R_2)$. For a polynomial P , $\deg_S P$ is the largest value of $\sum_{i \in S} \alpha_i$ over the monomials x^α of P ; for symmetric P it depends only on $j = |S|$, and we write $D_j(P)$. Since the m_κ have disjoint supports and $D_j(m_\kappa) = \kappa_1 + \dots + \kappa_j$:

Lemma 4.4. *A homogeneous $P \in \Lambda_m$ of degree $|\rho|$ lies in $\text{span}\{m_\kappa : \kappa \leq \rho\}$ if and only if $D_j(P) \leq \rho_1 + \dots + \rho_j$ for all j .*

Lemma 4.5 (Degree step). *For symmetric F and every j , $D_j(E_k F) \leq D_j(F) + \min(k, j)$.*

Proof. Fix $|S| = j$ and a term $R_A = c_A x_A F|_{x_i \mapsto sx_i (i \in A)}$ of Theorem 2.5. Every a_{ij} has $\deg_S a_{ij} = 0$: if i, j are both in S or both outside, a_{ij} is homogeneous of degree 0 in u or constant; otherwise numerator and denominator have degree 1. Next, $\deg_S x_A = |A \cap S| \leq \min(k, j)$. Rescaling some variables by the constant s does not change u -degrees, so the last factor has $\deg_S = D_j(F)$. Since $E_k F$ is a polynomial, its $\deg_S$ is its monomial degree. $\square$

4.2. Support, diagonal coefficient, and the value at $s = 1$.

Proof of the support statement. Starting from $D_j(1) = 0$ and applying Lemma 4.5 once for each part, $D_j(e_\lambda^*) \leq \sum_i \min(\lambda_i, j) = \lambda'_1 + \dots + \lambda'_j$. By Lemmas 4.3 and 4.4, $e_\lambda^* \in \text{span}\{m_\kappa : \kappa \leq \lambda'\} = \text{span}\{e_\nu : \nu \supseteq \lambda\}$. The support part of Corollary 4.2 is the same argument with one step, starting from $D_j(e_\mu) = \sum_i \min(\mu_i, j)$. $\square$

Fix integers $w_1 > w_2 > \dots > w_m > 0$ and for $\rho = (\rho_1 \geq \dots \geq \rho_m \geq 0)$ put $W(\rho) = \sum_i w_i \rho_i$.

Lemma 4.6. *If $\text{sort}(\alpha) = \kappa \leq \rho$, then $\sum_i w_i \alpha_i \leq W(\rho)$, with equality only if $\alpha = \rho$. Hence if $P \in \text{span}\{m_\kappa : \kappa \leq \rho\}$ has m_ρ -coefficient c , the coefficient of $u^{W(\rho)}$ in $P(u^{w_1} x_1, \dots, u^{w_m} x_m)$ is $c x^\rho$, and no higher power of u occurs.*

Proof. $\sum_i w_i \alpha_i \leq \sum_i w_i \kappa_i$ by the rearrangement inequality, with equality only if $\alpha = \kappa$ (the w_i are distinct). By Abel summation, $\sum_i w_i \kappa_i = \sum_j (w_j - w_{j+1})(\kappa_1 + \dots + \kappa_j)$ with $w_{m+1} := 0$; all weights are positive, so this is at most the same expression for ρ , with equality only if $\kappa = \rho$. $\square$

Lemma 4.7 (Lead step). *Let $F \in \text{span}\{m_\kappa : \kappa \leq \rho\}$ with m_ρ -coefficient c_F . Then $E_k F \in \text{span}\{m_\kappa : \kappa \leq \rho + 1^k\}$, with $m_{\rho+1^k}$ -coefficient $s^{\rho_1 + \dots + \rho_k} c_F$.*

Proof. Membership is Lemmas 4.4 and 4.5. Substitute $x_i \mapsto u^{w_i} x_i$ and expand each term R_A in u^{-1} . For $i < j$, $a_{ij} = 1 + O(u^{-1})$, and for $i > j$, $a_{ij} = t + O(u^{-1})$, so $c_A = t^{\text{inv}(A)}(1 + O(u^{-1}))$ with $\text{inv}(A) = \#\{i \in A, j \notin A : i > j\}$. Next, x_A becomes $u^{\sum_{a \in A} w_a} x_A$. Finally, $F|_{x_a \mapsto sx_a (a \in A)}$ has the monomials of F with x^α rescaled by $s^{\sum_{a \in A} \alpha_a}$, so by Lemma 4.6 it is $u^{W(\rho)}(c_F s^{\sum_{a \in A} \rho_a} x^\rho + O(u^{-1}))$. The exponent $W(\rho) + \sum_{a \in A} w_a$ is uniquely maximal for $A = [k]$, where $\text{inv} = 0$. So the coefficient of $u^{W(\rho+1^k)}$ in $E_k F$ is $s^{\rho_1 + \dots + \rho_k} c_F x^{\rho+1^k}$; conclude by Lemma 4.6 for $\rho + 1^k$. $\square$

Proof of the diagonal coefficient. Apply the operators in any order. If the parts applied so far form μ , the current function lies in $\text{span}\{m_\kappa : \kappa \leq \mu'\}$, and applying E_k multiplies the $m_{\mu'}$ -coefficient by $s^{\mu'_1 + \dots + \mu'_k} = s^{\sum_i \min(\mu_i, k)}$ (Lemma 4.7; note $\mu' + 1^k = (\mu \cup k)'$). Each unordered pair of parts contributes its minimum once, so the total exponent is $\sum_{i < p} \min(\lambda_i, \lambda_p) = \sum_p (p-1) \lambda_p = n(\lambda)$. By the triangularity in the proof of Lemma 4.3, the only e_ν with $\nu \supseteq \lambda$ containing $m_{\lambda'}$

is e_λ , with coefficient 1. Hence $c_{\lambda\lambda} = s^{n(\lambda)}$. The same computation gives the leading term in Corollary 4.2. $\square$

Proof of integrality and of $c_{\lambda\mu}|_{s=1} = 0$. Let $V = \prod_{i < j} (x_i - x_j)$. Multiplying the subset formula by V gives

$$V E_k F = \sum_{|A|=k} \varepsilon_A \prod_{\substack{i < j \\ \text{both in or both out of } A}} (x_i - x_j) \prod_{i \in A, j \notin A} (x_i - t x_j) x_A F|_{x_a \mapsto s x_a (a \in A)}, \quad \varepsilon_A = \pm 1.$$

If $F \in \mathbb{Q}[s, t][x]$, the right side lies in $\mathbb{Q}[s, t][x]$. As V has content 1 and $E_k F$ is a polynomial, Gauss's lemma gives $E_k F \in \mathbb{Q}[s, t][x]$. By induction $e_\lambda^* \in \mathbb{Q}[s, t][x]^{S_m}$, and its e -coefficients lie in $\mathbb{Q}[s, t]$. Now set $s = 1$ (a ring map on $\mathbb{Q}[s, t][x]$): every $F|_{x_a \mapsto s x_a}$ becomes $F|_{s=1}$, so $V(E_k F)|_{s=1} = F|_{s=1} \cdot V E_k(1) = V e_k F|_{s=1}$ by Lemma 2.7. Hence $(E_k F)|_{s=1} = e_k F|_{s=1}$, $e_\lambda^*|_{s=1} = e_\lambda$, and $c_{\lambda\mu}|_{s=1} = \delta_{\lambda\mu}$. $\square$

4.3. The exact s -adic valuation. It remains to show that $s^{n(\mu)}$ divides $c_{\lambda\mu}$ and that $(c_{\lambda\mu}/s^{n(\mu)})|_{s=t=0} = 1$ for all $\mu \succeq \lambda$; in particular $c_{\lambda\mu} \neq 0$. Put $\omega(\alpha) := \sum_j \binom{\alpha_j}{2}$ for $\alpha \in \mathbb{Z}_{\geq 0}^m$ and $B_\mu := \{\beta : \text{sort}(\beta) \leq \mu'\}$, the monomial support of e_μ^* . Write $e_\lambda^* = \sum_\beta a_\beta^\lambda x^\beta$; a_β^λ is symmetric in β . The source of s^n is the identity

$$(4.1) \quad \omega(\beta + 1_A) = \omega(\beta) + 1_A \cdot \beta.$$

Let $\mathbb{K} = \mathbb{Q}(t)(s^{1/2})$ (or $\mathbb{Q}(s^{1/2})$ when $t = 0$) with the s -adic valuation v , extended to $\mathbb{K}(x)$ by the Gauss valuation $v(\sum a_\beta x^\beta) = \min_\beta v(a_\beta)$. For $c \in (\frac{1}{2}\mathbb{Z})^m$ let σ_c be the automorphism $x_j \mapsto s^{-c_j} x_j$, so that $v(\sigma_c P) = \min_\beta (v(a_\beta) - c \cdot \beta)$.

Lemma 4.8. *For generic t , and also at $t = 0$, $v(a_\alpha^\lambda) \geq \omega(\alpha)$ for all α .*

Proof. We show $v(\sigma_c e_\lambda^*) \geq \min_{\beta \in B_\lambda} (\omega(\beta) - c \cdot \beta)$ for all c , by induction along $e_\mu^* \mapsto E_k e_\mu^* = e_{\mu \cup k}^*$, the case $e_\emptyset^* = 1$ being clear. In a term R_A : $v(\sigma_c a_{ij}) = 0$ for generic t , and at $t = 0$, $v(\sigma_c a_{ij}) = \max(c_i, c_j) - c_i \geq 0$; $\sigma_c x_A = s^{-c \cdot 1_A} x_A$; and $\sigma_c(F|_{x_a \mapsto s x_a (a \in A)}) = \sigma_{c-1_A} F$. Hence, by induction and (4.1),

$$v(\sigma_c R_A) \geq -c \cdot 1_A + \min_{\beta \in B_\mu} (\omega(\beta) - (c - 1_A) \cdot \beta) = \min_{\beta \in B_\mu} (\omega(\beta + 1_A) - c \cdot (\beta + 1_A)),$$

and $\beta + 1_A \in B_{\mu \cup k}$ because the top- j sums grow by at most $\min(j, k)$ and $\mu' + 1^k = (\mu \cup k)'$. Now take $c = \alpha - \frac{1}{2}\mathbf{1}$. Then $\omega(\beta) - c \cdot \beta = \sum_j ((\beta_j - \alpha_j)^2 - \alpha_j^2)/2$ is uniquely minimised over $\mathbb{Z}^m$ at $\beta = \alpha$, so $v(a_\alpha^\lambda) - c \cdot \alpha \geq v(\sigma_c e_\lambda^*) \geq \omega(\alpha) - c \cdot \alpha$. $\square$

Proof that $s^{n(\mu)} \mid c_{\lambda\mu}$. Write $c_{\lambda\nu} = s^{n(\nu)} d_\nu$ and suppose $\delta := \min_\nu v(d_\nu) < 0$. Choose μ dominance-minimal with $v(d_\mu) = \delta$. The coefficient of $x^{\mu'}$ is $a_{\mu'}^\lambda = \sum_{\nu \triangleleft \mu} c_{\lambda\nu} M_{\nu\mu'}$ with $M_{\mu\mu'} = 1$. Since $\nu \triangleleft \mu$ implies $n(\nu) > n(\mu)$ (as $n(\nu) = \sum_{i \geq 1} (n - \nu_1 - \dots - \nu_i)$), every term with $\nu \neq \mu$ has valuation $> n(\mu) + \delta$, so $v(a_{\mu'}^\lambda) = n(\mu) + \delta < n(\mu) = \omega(\mu')$, contradicting Lemma 4.8. So d_ν is regular at $s = 0$; as $c_{\lambda\nu} \in \mathbb{Q}[s, t]$, $s^{n(\nu)}$ divides $c_{\lambda\nu}$ in $\mathbb{Q}[s, t]$. The same argument at $t = 0$ shows $d_\mu|_{s=t=0} = [s^{\omega(\mu')}] a_{\mu'}^\lambda|_{t=0}$. $\square$

For the value at $s = t = 0$ we need the initial forms. For $R \in \mathbb{K}(x)$ with $v(R) \geq v_0$, let $\text{in}_{v_0}(R)$ be the image of $s^{-v_0} R$ in the residue field $\mathbb{Q}(x)$; it is additive at a common level and multiplicative. For $\kappa = \text{sort}(\alpha)$ let peel_k subtract 1 from the k largest parts.

Lemma 4.9 (Peel Lemma). *If $\kappa \leq \rho$, $|\kappa| = |\rho|$ and $k \leq \ell(\rho)$, then $\text{peel}_k \kappa \leq \text{peel}_k \rho$.*

Proof. Note $\ell(\kappa) \geq \ell(\rho) \geq k$. Counting parts $\geq c$ gives $(\text{peel}_k \kappa)'_c = \kappa'_c - \min(k, \kappa'_c) + \min(k, \kappa'_{c+1})$, so with S_C the sum of the first C parts and $\kappa'_1 \geq k$, $S_C((\text{peel}_k \kappa)') = S_C(\kappa') - k + \min(k, \kappa'_{C+1})$, and likewise for ρ . Let $D_C = S_C(\kappa') - S_C(\rho') \geq 0$ (by [Mac95, I (1.11)]) and $\bar{m}(y) = \min(k, y)$. If $\bar{m}(\kappa'_{C+1}) \geq \bar{m}(\rho'_{C+1})$ we are done. Otherwise $\kappa'_{C+1} < k$ and $\kappa'_{C+1} < \rho'_{C+1}$, and $D_C = D_{C+1} + \rho'_{C+1} - \kappa'_{C+1} \geq \bar{m}(\rho'_{C+1}) - \bar{m}(\kappa'_{C+1})$. So $(\text{peel}_k \kappa)' \supseteq (\text{peel}_k \rho)'$. $\square$

Lemma 4.10. *At $t = 0$, $\Lambda_\lambda^0(\alpha) := [s^{\omega(\alpha)}] a_\alpha^\lambda$ equals 1 if $\text{sort}(\alpha) \leq \lambda'$ and 0 otherwise.*

Proof. Induction along E_k ; the base case is clear, and $\Lambda_{\mu \cup k}^0(\alpha) = 0$ outside $B_{\mu \cup k}$ by the support statement. Let $\text{sort}(\alpha) \leq (\mu \cup k)'$, $c = \alpha - \frac{1}{2}\mathbf{1}$ and $v_0 = \omega(\alpha) - c \cdot \alpha$. By the uniqueness in the proof of Lemma 4.8, $\text{in}_{v_0}(\sigma_c e_{\mu \cup k}^*) = \Lambda_{\mu \cup k}^0(\alpha) x^\alpha$. If $\alpha - 1_A$ has a negative entry or lies outside B_μ , then $v(\sigma_c R_A) > v_0$ by that uniqueness and the term R_A does not contribute. Otherwise, with $\beta_0 = \alpha - 1_A$, the levels split as in (4.1) and

$$\text{in}_{v_0}(\sigma_c R_A) = \text{in}_0(\sigma_c c_A) \cdot \Lambda_\mu^0(\beta_0) x^\alpha.$$

At $t = 0$, $\text{in}_0(\sigma_c a_{ij})$ is 1 if $\alpha_i > \alpha_j$, 0 if $\alpha_i < \alpha_j$, and $x_i/(x_i - x_j)$ if $\alpha_i = \alpha_j$. So only sets A that are upper sets for α contribute: with v the k -th largest entry of α and $L = \{j : \alpha_j = v\}$, $A = \{j : \alpha_j > v\} \sqcup B$ with $B \subseteq L$ of size $r = k - \#\{j : \alpha_j > v\}$. As $\ell(\text{sort } \alpha) \geq \ell((\mu \cup k)') \geq k$, $v \geq 1$. For all such A , $\text{sort}(\alpha - 1_A) = \text{peel}_k(\text{sort } \alpha)$, and Λ_μ^0 is symmetric, so

$$\Lambda_{\mu \cup k}^0(\alpha) = \Lambda_\mu^0(\text{peel}_k \text{sort } \alpha) \cdot \sum_{B \subseteq L, |B|=r} \prod_{i \in B, j \in L \setminus B} \frac{x_i}{x_i - x_j} = \Lambda_\mu^0(\text{peel}_k \text{sort } \alpha).$$

The last sum is the case $t = 0$ of $\sum_{|B|=r} \prod_{i \in B, j \in L \setminus B} \frac{x_i - tx_j}{x_i - x_j} = \left[\begin{smallmatrix} L \\ r \end{smallmatrix} \right]_t$, which follows from the Poincaré identity [Mac95, III (1.4)] by grouping $S_{|L|}$ according to the image of $\{1, \dots, r\}$. Finally, by Lemma 4.9 with $\rho = (\mu \cup k)' = \mu' + 1^k$, $\text{peel}_k(\text{sort } \alpha) \leq \text{peel}_k(\mu' + 1^k) = \mu'$, so $\Lambda_\mu^0(\text{peel}_k \text{sort } \alpha) = 1$. $\square$

End of proof of Theorem 4.1. By the last two proofs, $(c_{\lambda\mu}/s^{n(\mu)})|_{s=t=0} = \Lambda_\lambda^0(\mu') = 1$ for $\mu \supseteq \lambda$, since $\mu' \leq \lambda'$. In particular $c_{\lambda\mu} \neq 0$ and its s -adic valuation is $n(\mu)$. $\square$

Remark 4.11. The leading coefficients $d_{\lambda\mu}(t) = [s^{n(\mu)}] c_{\lambda\mu}$ depend on t ; for instance $d_{(1^4), (2,1,1)} = 1 + 3t$. They are identified in Corollary 5.2. The degree count above is the same kind of triangularity argument as for Macdonald's operators [Mac95, VI (3.6)–(3.10)].

5. THE $s \rightarrow 0$ EDGE

Since $s = q^{-1}$, the edge $s \rightarrow 0$ is Hikita's $q \rightarrow \infty$, where the unrescaled product degenerates [Hik25, Lemma 6.3]. After the rescaling of Theorem 4.1 it becomes Hall–Littlewood multiplication. Throughout this section t is an indeterminate, $b_\mu := s^{n(\mu)} e_\mu$, and we work in Λ_m with m at least the degree (Lemma 2.8). We keep the notation $\mathbb{K}$, v , σ_c , ω of Section 4, with $\mathbb{K} = \mathbb{Q}(t)(s^{1/2})$.

Theorem 5.1 (The $s \rightarrow 0$ edge). *Let $\mathcal{O} = \mathbb{Q}(t)[s]_{(s)}$ and $\mathcal{L} = \bigoplus_\mu \mathcal{O} b_\mu$.*

- (1) $E_k b_\mu \in \bigoplus_\nu \mathbb{Q}[s, t] b_\nu$; in particular $E_k \mathcal{L} \subseteq \mathcal{L}$.
- (2) Let L_k be the map induced by E_k on $\mathcal{L}/s\mathcal{L} = \bigoplus_\mu \mathbb{Q}(t) \bar{b}_\mu$, and let $\phi(\bar{b}_\mu) := t^{-n(\mu')} P_{\mu'}(x; t^{-1})$.
Then $\phi \circ L_k = t^{-\binom{k}{2}} e_k \cdot \phi$.
- (3) Explicitly, with $\rho = \mu'$ and $\kappa = \nu'$,

$$L_k \bar{b}_\mu = \sum_{\kappa/\rho \text{ vertical } k\text{-strip}} t^{\text{inv}(\kappa/\rho)} \prod_{v \geq 1} \left[\begin{smallmatrix} m_v(\kappa) \\ r_v \end{smallmatrix} \right]_t \bar{b}_\nu,$$

where r_v is the number of rows of κ of length v that end in a box of the strip, and $\text{inv}(\kappa/\rho) = \sum_{v < w} r_v (m_w(\kappa) - r_w)$.

Corollary 5.2 (The s -leading coefficients). *For $\lambda, \mu \vdash n$, $d_{\lambda\mu}(t) := [s^{n(\mu)}] c_{\lambda\mu}$ satisfies*

$$d_{\lambda\mu}(t) = t^{n(\mu') - n(\lambda')} [P_{\mu'}(x; t^{-1})] e_\lambda = t^{-n(\lambda')} \sum_{\nu} K_{\nu'\lambda} \widetilde{K}_{\nu\mu'}(t),$$

where $\widetilde{K}_{\nu\kappa}(t) = t^{n(\kappa)} K_{\nu\kappa}(t^{-1})$ is the cocharge Kostka–Foulkes polynomial. Hence $d_{\lambda\mu} \in \mathbb{N}[t]$, and $d_{\lambda\mu}(1)$ is the number of 0-1 matrices with row sums λ and column sums μ' .

5.1. Admissible functions. Call a homogeneous $F = \sum_{\beta} a_{\beta} x^{\beta} \in \mathbb{K}[x]^{S_m}$ *admissible* if $v(a_{\beta}) \geq \omega(\beta)$ for all β . As in the proof of Lemma 4.8, F is admissible if and only if $v(\sigma_c F) \geq \min_{\beta \in \mathbb{Z}_{\geq 0}^m} (\omega(\beta) - c \cdot \beta)$ for all $c \in (\frac{1}{2}\mathbb{Z})^m$; the backward direction uses $c = \alpha - \frac{1}{2}\mathbf{1}$, for which $\beta = \alpha$ is the unique minimiser. For admissible F let $\text{in}(F)(\alpha) \in \mathbb{Q}(t)$ be the residue of $s^{-\omega(\alpha)} a_{\alpha}$; it depends only on $\text{sort}(\alpha)$. With $c = \alpha - \frac{1}{2}\mathbf{1}$ and $v_0 = \omega(\alpha) - c \cdot \alpha$,

$$(5.1) \quad \text{in}_{v_0}(\sigma_c F) = \text{in}(F)(\alpha) x^{\alpha}.$$

Lemma 5.3. *A homogeneous $F = \sum_{\mu} c_{\mu} e_{\mu}$ of degree $\leq m$ is admissible if and only if $v(c_{\mu}) \geq n(\mu)$ for all μ . In that case $\text{in}(F)(\mu')$ is the residue of $s^{-n(\mu)} c_{\mu}$.*

Proof. The coefficient of x^{β} in b_{μ} is $s^{n(\mu)} M_{\mu\kappa}$ with $\kappa = \text{sort}(\beta)$. If $M_{\mu\kappa} \neq 0$ then $\kappa \leq \mu'$, so $\kappa' \geq \mu$ and $\omega(\beta) = n(\kappa') \leq n(\mu)$, with equality only for $\kappa = \mu'$, where $M_{\mu\mu'} = 1$. So b_{μ} is admissible with $\text{in}(b_{\mu})(\kappa) = [\kappa = \mu']$, which gives the backward direction and the formula. The forward direction is the argument in the proof that $s^{n(\mu)} \mid c_{\lambda\mu}$ in Section 4, with a^{λ} replaced by the coefficients of F . $\square$

Lemma 5.4. $\sum_{B \subseteq [N], |B|=r} \prod_{i \in B, j \notin B} a_{ij} = [N]_t^r$.

Proof. Group the Poincaré identity $\sum_{w \in S_N} w(\prod_{i < j} a_{ij}) = [N]_t! [\text{Mac95, III (1.4)}]$ by $B = w(\{1, \dots, r\})$; the stabiliser $S_r \times S_{N-r}$ contributes $[r]_t! [N-r]_t!$. $\square$

5.2. The recursion.

Proposition 5.5. *Let F be admissible of degree n , with $m \geq n + k$. Then $E_k F$ is admissible, and for every $\kappa \vdash n + k$*

$$(5.2) \quad \text{in}(E_k F)(\kappa) = \sum_{\rho: \kappa/\rho \text{ vertical } k\text{-strip}} t^{\text{inv}(\kappa/\rho)} \prod_{v \geq 1} \begin{bmatrix} m_v(\kappa) \\ r_v \end{bmatrix}_t \text{in}(F)(\rho).$$

Proof. Fix c and a term R_A of Theorem 2.5. Since t is a unit, $v(\sigma_c a_{ij}) = 0$, and the initial form of $\sigma_c a_{ij}$ is 1 if $c_i > c_j$, t if $c_i < c_j$, and a_{ij} if $c_i = c_j$. As in Lemma 4.8, using (4.1),

$$v(\sigma_c R_A) \geq \min_{\gamma \geq 1_A} (\omega(\gamma) - c \cdot \gamma) \geq \min_{\gamma \geq 0} (\omega(\gamma) - c \cdot \gamma),$$

so $E_k F$ is admissible. Now fix α with $\text{sort}(\alpha) = \kappa$, $c = \alpha - \frac{1}{2}\mathbf{1}$, $v_0 = \omega(\alpha) - c \cdot \alpha$; all terms have $v \geq v_0$, so initial forms add, and by (5.1) $\text{in}(E_k F)(\alpha) x^{\alpha} = \sum_A \text{in}_{v_0}(\sigma_c R_A)$. If $A \not\subseteq \text{supp}(\alpha)$, then $\gamma = \alpha$ is excluded from $\{\gamma \geq 1_A\}$ and $v(\sigma_c R_A) > v_0$. If $A \subseteq \text{supp}(\alpha)$, put $\beta_0 = \alpha - 1_A$; then $c - 1_A = \beta_0 - \frac{1}{2}\mathbf{1}$, the levels add by (4.1), and

$$\text{in}_{v_0}(\sigma_c R_A) = \text{in}_0(\sigma_c c_A) \cdot \text{in}(F)(\beta_0) x^{\alpha}.$$

Let $L_v = \{j : \alpha_j = v\}$ ($v \geq 1$), $B_v = A \cap L_v$ and $r_v = |B_v|$. Then

$$\text{in}_0(\sigma_c c_A) = t^{\#\{i \in A, j \notin A: \alpha_i < \alpha_j\}} \prod_{v \geq 1} \prod_{i \in B_v, j \in L_v \setminus B_v} a_{ij}, \quad \#\{\dots\} = \sum_{v < w} r_v (m_w - r_w).$$

The partition $\text{sort}(\beta_0)$ is obtained from κ by lowering r_v of its parts equal to v , for each v ; this is the ρ with κ/ρ a vertical k -strip and data (r_v) , and every such strip arises from exactly one (r_v) . So both $\text{in}(F)(\beta_0)$ and the power of t depend only on (r_v) , and summing over the B_v with Lemma 5.4 gives $\prod_v [m_v]_{r_v}_t$. $\square$

At $t = 0$ every t -binomial is 1 and t^{inv} vanishes unless every part larger than a lowered part is lowered; so (5.2) reduces to the recursion in the proof of Lemma 4.10.

5.3. Proof of Theorem 5.1 and Corollary 5.2. (1). b_μ is admissible, so $E_k b_\mu$ is admissible (Proposition 5.5) and its b_ν -coordinates have $v \geq 0$ (Lemma 5.3). By the integrality part of Theorem 4.1's proof, $E_k e_\mu \in \bigoplus_\nu \mathbb{Q}[s, t] e_\nu$, so the b_ν -coordinate is $p/s^{n(\nu)-n(\mu)}$ with $p \in \mathbb{Q}[s, t]$; having $v \geq 0$ over $\mathbb{Q}(t)$, it lies in $\mathbb{Q}[s, t]$.

(3). By Lemma 5.3, the $\bar{b}_\nu$ -coordinate of $L_k \bar{b}_\mu$ is $\text{in}(E_k b_\mu)(\nu')$; apply (5.2) with $\text{in}(b_\mu)(\rho) = [\rho = \mu']$.

(2). Let κ/ρ be a vertical k -strip with data (r_v) , obtained by lowering the parts indexed by A . Writing $n(\kappa) = \sum_{\text{pairs of parts}} \min$, a pair of parts both in A loses 1; a pair with one part of value v in A and one of value w not in A loses 1 if and only if $v \leq w$; other pairs lose nothing. Hence

$$(5.3) \quad n(\kappa) - n(\rho) = \binom{k}{2} + \text{inv}(\kappa/\rho) + \sum_v r_v(m_v - r_v).$$

Since $[m]_r|_{t \rightarrow t^{-1}} = t^{-r(m-r)} [m]_r$, (5.3) gives $t^{\text{inv}} \prod_v [m_v]_{r_v}_t = t^{n(\kappa)-n(\rho)-\binom{k}{2}} \psi_{\kappa/\rho}(t^{-1})$ with $\psi_{\kappa/\rho}(u) := \prod_v [m_v(\kappa)]_{r_v}_u$. The Hall–Littlewood Pieri rule [Mac95, III (3.2)] reads $e_k P_\rho(x; u) = \sum_\kappa \psi_{\kappa/\rho}(u) P_\kappa(x; u)$, summed over vertical k -strips (there $m_v(\kappa) = \kappa'_v - \kappa'_{v+1}$ and $r_v = \kappa'_v - \rho'_v$). For $X \in \mathcal{L}$ put $h(\kappa) = t^{-n(\kappa)} \text{in}(X)(\kappa)$, so that $\phi(\bar{X}) = \sum_\kappa h(\kappa) P_\kappa(x; t^{-1})$ by Lemma 5.3. By (5.2), $t^{-n(\kappa)} \text{in}(E_k X)(\kappa) = t^{-\binom{k}{2}} \sum_\rho \psi_{\kappa/\rho}(t^{-1}) h(\rho)$, hence $\phi(L_k \bar{X}) = t^{-\binom{k}{2}} \sum_\rho h(\rho) e_k P_\rho(x; t^{-1}) = t^{-\binom{k}{2}} e_k \phi(\bar{X})$.

Corollary 5.2. Since $1 = b_\emptyset$ and $\phi(\bar{b}_\emptyset) = 1$, iterating (2) gives $\phi(e_\lambda^* \bmod s) = t^{-\sum_i \binom{\lambda_i}{2}} e_\lambda = t^{-n(\lambda')} e_\lambda$. By Theorem 4.1, $e_\lambda^* \equiv \sum_\mu d_{\lambda\mu} \bar{b}_\mu$, so comparing coefficients of $P_{\mu'}(x; t^{-1})$ gives the first formula. Expanding $e_\lambda = \sum_\nu K_{\nu'\lambda} s_\nu$ and $s_\nu = \sum_\kappa K_{\nu\kappa}(u) P_\kappa(x; u)$ [Mac95, III (2.6)] at $u = t^{-1}$ gives the second. The $K_{\nu\kappa}(t)$ have nonnegative coefficients [Mac95, III (6.5)], and $d_{\lambda\mu} \in \mathbb{Q}[t]$ by Theorem 4.1, so $d_{\lambda\mu} \in \mathbb{N}[t]$. At $t = 1$, $P_\kappa(x; 1) = m_\kappa$ [Mac95, III §2], so $d_{\lambda\mu}(1) = [m_{\mu'}] e_\lambda = M_{\lambda\mu'}$. $\square$

Remark 5.6 (Prior work; no novelty claim for the matrix). The matrix $d_{\lambda\mu}(t)$ is the classical $e \rightarrow \text{Hall-Littlewood}$ transition matrix: writing $e_\lambda = \sum_\kappa M(e, P)_{\lambda\kappa} P_\kappa(x; t)$, Corollary 5.2 says $d_{\lambda\mu}(t) = t^{n(\mu')-n(\lambda')} M(e, P)_{\lambda\mu'}(t^{-1})$, and its positivity and the value at $t = 1$ are in [Kir98, §3.2]. What is new is only the identification of this matrix with the s -leading coefficients of $\star$, i.e. Theorem 5.1. Hikita's [Hik25, Lemma 6.3] treats the unrescaled limit $q \rightarrow \infty$.

Remark 5.7 (Other edges). The edges $t \rightarrow \infty$, $s \rightarrow \infty$ and $t = 0$ of the (s, t) -square, and the reflection $(s, t) \mapsto (1/s, 1/t)$, can be treated through a conjugation of $e_k \star (\cdot)$ by an operator that is diagonal on Macdonald polynomials; such a conjugation is implicit in [DFK19]. We do not develop this here and use none of it: the only other edge that enters below is $t = 0$, and for it we cite [DFK18] (Remark 6.7).

6. THE BLOCK VALUATION LAW AT $s = 1$

6.1. Taylor pieces. We write v for the $(s-1)$ -adic valuation on $\mathbb{Q}(t)[s]$, $I = (e_1, e_2, \dots)$ for the augmentation ideal (so I^b is spanned by the e_μ with $\ell(\mu) \geq b$), and $\text{line}_e G := [e_n]G$ for G homogeneous of degree n ; line_e vanishes on I^2 . For ρ a partition, a homogeneous symmetric

function is of type ρ if it lies in $\text{span}\{e_\nu : \nu \supseteq \rho\}$; by Lemmas 4.3 and 4.4 this means $D_j(P) \leq \sum_i \min(\rho_i, j)$ for all j . For a set C of positions, λ_C is the partition formed by the parts λ_i , $i \in C$.

In Theorem 2.5 the variable s enters only through $F|_{x_i \mapsto sx_i} (i \in A) = s^{\Delta_A} F$, $\Delta_A = \sum_{i \in A} x_i \partial_i$. Since $s^{\Delta_A} x^\alpha = s^{\alpha \cdot 1_A} x^\alpha = \sum_p (s-1)^p \binom{\Delta_A}{p} x^\alpha$,

$$(6.1) \quad E_k = \sum_{p \geq 0} (s-1)^p E_k^{(p)}, \quad E_k^{(p)} = \sum_{|A|=k} c_A x_A \binom{\Delta_A}{p}.$$

As $E_k F$ is a symmetric polynomial in x and s (Theorem 4.1), each $E_k^{(p)}$ preserves Λ_m . Since $\binom{\Delta_A}{p}$ is a polynomial of degree p in a derivation, and multiplication by $c_A x_A$ preserves this, $E_k^{(p)}$ is a differential operator of order $\leq p$: all $(p+1)$ -fold commutators $[\dots [E_k^{(p)}, a_0], \dots, a_p]$ with multiplication operators vanish. Moreover $E_k^{(0)}$ is multiplication by e_k (by the proof of Theorem 4.1, $(E_k F)|_{s=1} = e_k F|_{s=1}$), and $E_k^{(p)}(1) = 0$ for $p \geq 1$.

6.2. First order.

Theorem 6.1 (First order is a biderivation). *Let $D_k := E_k^{(1)}$ and $B(f, g) := \partial_s(f \star g)|_{s=1}$ for $f, g \in \Lambda_m$ with coefficients in $\mathbb{Q}(t)$ (independent of s). Then D_k is a derivation of Λ_m , and*

$$B(f, g) = \sum_{k, l} M_{kl} \frac{\partial f}{\partial e_k} \frac{\partial g}{\partial e_l}, \quad M_{kl} := D_k(e_l) = B(e_k, e_l) = M_{lk}.$$

So B is a symmetric biderivation, the carré du champ of $L = \frac{1}{2} \sum M_{kl} \partial^2 / \partial e_k \partial e_l$.

Proof. $D_k = \sum_A c_A x_A \Delta_A$ is a sum of derivations. By Theorem 4.1 the $e_\rho^\star$ form a basis whose transition matrix to the e_ρ is unitriangular at $s = 1$; write $f = \sum_\rho a_\rho(s) e_\rho^\star$ with a_ρ regular at $s = 1$. By associativity, $f \star g = \sum_\rho a_\rho(s) E_{\rho_1} \cdots E_{\rho_\ell} g$. By (6.1),

$$B(f, g) = \sum_\rho a'_\rho(1) e_\rho g + \sum_\rho a_\rho(1) \sum_i \left(\prod_{j \neq i} e_{\rho_j} \right) D_{\rho_i} g.$$

Applying the same computation to $f = \sum_\rho a_\rho(s) E_{\rho_1} \cdots E_{\rho_\ell}(1)$ itself, which does not depend on s , and using $D_k(1) = 0$, gives $\sum_\rho a'_\rho(1) e_\rho = 0$. Since $f|_{s=1} = \sum_\rho a_\rho(1) e_\rho$, the remaining sum is $\sum_k \frac{\partial f}{\partial e_k} D_k(g)$, and $D_k(g) = \sum_l \frac{\partial g}{\partial e_l} D_k(e_l)$. Symmetry of M is commutativity of $\star$ [Hik25]. $\square$

Proposition 6.2 (The matrix M). *For $k \geq r \geq 0$, with $L(a, b) := \frac{(1-t^{a+b})(t^{ab}-1)}{(1-t^a)(1-t^b)}$,*

$$M_{kr} = B(e_k, e_r) = r e_k e_r + \sum_{j=1}^r L(k-r+j, j) e_{k+j} e_{r-j}.$$

Consequently $[(s-1)^1] e_\lambda^\star = \sum_{i < j} M_{\lambda_i \lambda_j} e_{\lambda \setminus \{\lambda_i, \lambda_j\}}$.

Proof. $M_{kr} = [(s-1)^1] e_k \star e_r$. Use the form of the Pieri rule in Corollary 8.6 (Section 8; its proof does not use Proposition 6.2): $e_k \star e_r = \sum_{y=0}^r s^y C_{k-y, r-y} e_{k+r-y} e_y$. Since $\pi_a(b)|_{s=1} = \delta_{b0}$, every $C_{a,b}$ with $b \geq 1$ vanishes at $s = 1$; so the $y = r$ term $s^r e_k e_r$ contributes $r e_k e_r$, and the term $y = r-j$, $j \geq 1$, contributes $\partial_s C_{a,j}|_{s=1}$ with $a = k-r+j$. Now $\partial_s \pi_a(j)|_{s=1} = [u^j] \sum_{m < a} \frac{ut^m}{1+ut^m} = (-1)^{j-1} [a]_{t^j}$, so $\partial_s C_{a,j}|_{s=1} = -\frac{[a+j]}{[a]} [a]_{t^j} = L(a, j)$. For the second statement, $[(s-1)^1] E_{\lambda_1} \cdots E_{\lambda_\ell}(1) = \sum_i \prod_{j < i} e_{\lambda_j} D_{\lambda_i} (\prod_{j > i} e_{\lambda_j})$; expand with the Leibniz rule. $\square$

6.3. Blocks.

Lemma 6.3 (Polarization). *Let D be a linear operator on a commutative ring and $g_1, \dots, g_r$ elements of it. For $S \subseteq [r]$ let $\delta_S := [\dots [D, g_{s_1}], \dots, g_{s_q}](1) = \sum_{T \subseteq S} (-1)^{|S \setminus T|} \prod_{i \in S \setminus T} g_i D(\prod_{i \in T} g_i)$. Then*

$$D(g_1 \cdots g_r) = \sum_{S \subseteq [r]} \left(\prod_{i \notin S} g_i \right) \delta_S,$$

and if D has order $\leq p$, then $\delta_S = 0$ for $|S| > p$, and for $|S| = p$ the map $(g_i)_{i \in S} \mapsto \delta_S$ is a derivation in each argument.

Proof. Substituting the second expression for δ_S , the right side is $\sum_T \prod_{i \notin T} g_i D(\prod_{i \in T} g_i) \sum_{S \supseteq T} (-1)^{|S \setminus T|}$, and the inner sum is $[T = [r]]$. Vanishing is the definition of order $\leq p$. For $|S| = p$, $D' := [\dots [D, g_{s_2}], \dots, g_{s_p}]$ has order ≤ 1 , so $D'(ab) = aD'(b) + bD'(a) - abD'(1)$, i.e. $[D', ab](1) = a[D', b](1) + b[D', a](1)$; and the iterated commutator is symmetric in its arguments. $\square$

Proposition 6.4 (Block expansion). *Let λ have ℓ parts and $p_1, \dots, p_\ell \geq 0$ with $\sum_i p_i = m$. Then*

$$E_{\lambda_1}^{(p_1)} \cdots E_{\lambda_\ell}^{(p_\ell)}(1) = \sum_{\pi} \prod_{C \in \pi} g_C^\pi,$$

a finite sum over set partitions π of $[\ell]$ with at least $\ell - m$ blocks, where each g_C^π is homogeneous of type λ_C .

Proof. Induction on the number of operators applied, from the right; the empty product 1 has no blocks. Apply $D = E_k^{(p)}$, $k = \lambda_j$, $p = p_j$, to a term $\prod_{C \in \pi} g_C$. By Lemma 6.3 the result is $\sum_{S \subseteq \pi, |S| \leq p} \prod_{C \notin S} g_C \cdot \delta_S$, and we let $\pi' = (\pi \setminus S) \cup \{C'\}$ with $C' = \{j\} \cup \bigcup S$. Then $|\pi'| \geq |\pi| + 1 - p$, which gives the block count. Each δ_S is a combination of products of the g_C and of D applied to such products. Rescaling monomials does not enlarge monomial supports, so the degree step Lemma 4.5 holds for $E_k^{(p)}$ as well, and D_j is additive on products; hence $D_j(\delta_S) \leq \sum_{C \in S} \sum_{i \in C} \min(\lambda_i, j) + \min(\lambda_j, j)$, i.e. δ_S has type $\lambda_{C'}$. $\square$

Definition 6.5. For $\lambda, \mu \vdash n$, $\kappa(\lambda, \mu)$ is the largest r such that $\lambda = \lambda^1 \sqcup \cdots \sqcup \lambda^r$ and $\mu = \mu^1 \sqcup \cdots \sqcup \mu^r$ as multisets of parts, with every block nonempty and $\mu^i \supseteq \lambda^i$ (so $|\mu^i| = |\lambda^i|$); $\kappa = -\infty$ if $\mu \not\supseteq \lambda$. μ is a *coarsening* of λ if $\kappa(\lambda, \mu) = \ell(\mu)$.

6.4. The block valuation law.

Theorem 6.6 (Block valuation law). *For all $\lambda, \mu \vdash n$ and all t , $v(c_{\lambda\mu}) \geq \ell(\lambda) - \kappa(\lambda, \mu)$. For $\mu \supseteq \lambda$, $\mu \neq \lambda$, equality holds over $\mathbb{Q}(t)$ and at $t = 0$:*

$$v(c_{\lambda\mu}) = \ell(\lambda) - \kappa(\lambda, \mu).$$

At $t = 0$ the leading coefficient is $(-1)^{\ell - \kappa} N(\lambda, \mu)$ with $N(\lambda, \mu) \geq 1$ a count of minimal raising-operator configurations.

Proof of the lower bound. By (6.1), $[(s-1)^m]e_\lambda^* = \sum_{\sum p_i = m} E_{\lambda_1}^{(p_1)} \cdots E_{\lambda_\ell}^{(p_\ell)}(1)$. By Proposition 6.4 each term is a sum of products $\prod_{C \in \pi} g_C$ with $|\pi| \geq \ell - m$ and g_C of type λ_C , and such a product lies in $\text{span}\{e_{\sqcup_C \nu^C} : \nu^C \supseteq \lambda_C\}$. So e_μ can occur only if $\kappa(\lambda, \mu) \geq |\pi| \geq \ell - m$. The argument works over $\mathbb{Q}(t)$ and for every specialisation $t \in \mathbb{Q}$. $\square$

Remark 6.7 (Input at $t = 0$). The $t = 0$ edge $e_\mu^*|_{t=0} = s^{n(\mu)} P_{\mu'}(x; 1/s, 0) = \omega \tilde{H}_\mu(x; s)$, where $\tilde{H}_\mu(x; s) = s^{n(\mu)} Q'_\mu(x; s^{-1})$, is **not ours**: it is assembled from [DFK18] eq. (5.15) (with $M_{k,1} = E_k|_{t=0}$), Cor. 5.8 in its level-one form (5.25), Cor. 5.18 (eq. (5.27)) (numbering of arXiv:1505.01657v2), and [Mac95, VI (5.1), (4.14)(iv)]. In particular Theorem 6.6 does not depend on Cherednik's nonsymmetric theory.

Proof of equality. By Remark 6.7, $c_{\lambda\mu}(s, 0) = [h_\mu] s^{n(\lambda)} Q'_\lambda(x; s^{-1})$. By [Mac95, III (2.15')] with §5 Ex. 7(a)], $Q'_\lambda(x; u) = \prod_{i < j \leq \ell} \frac{1-R_{ij}}{1-uR_{ij}} h_\lambda$, where $R_{ij}h_\alpha = h_{\alpha+e_i-e_j}$, $h_\alpha = h_{\alpha_1} h_{\alpha_2} \cdots$ and $h_r = 0$ for $r < 0$. Expand $\frac{1-R}{1-uR} = 1 + \sum_{a \geq 1} (u-1)u^{a-1}R^a$; at $u = s^{-1}$ the coefficient of R^a is $-(s-1)s^{-a}$. A *configuration* is a function $\mathbf{m}$ on pairs $i < j \leq \ell$ with values in $\mathbb{Z}_{\geq 0}$; its edge set is $E(\mathbf{m}) = \{ij : \mathbf{m}_{ij} \geq 1\}$ and its result is $\alpha(\mathbf{m}) = \lambda + \sum \mathbf{m}_{ij}(e_i - e_j)$. Then

$$c_{\lambda\mu}(s, 0) = \sum_{\mathbf{m}: \alpha(\mathbf{m}) \geq 0, \text{ sort } \alpha(\mathbf{m}) = \mu} (-1)^{|E(\mathbf{m})|} (s-1)^{|E(\mathbf{m})|} s^{a(\mathbf{m})}, \quad a(\mathbf{m}) \in \mathbb{Z},$$

a finite sum (the conditions $\alpha_\ell \geq 0, \alpha_{\ell-1} \geq 0, \dots$ bound the $\mathbf{m}_{ij}$ successively). Every configuration contributes $(-1)^{|E|}$ at order $(s-1)^{|E|}$, so there is no cancellation at the lowest order: $v(c_{\lambda\mu}(s, 0)) = E_{\min}$, the least $|E(\mathbf{m})|$ over admissible $\mathbf{m}$, with leading coefficient $(-1)^{E_{\min}} N(\lambda, \mu)$, N the number of minimisers.

$E_{\min} \geq \ell - \kappa$: let C be a connected component of the graph $([\ell], E(\mathbf{m}))$. Transfers stay inside C and move mass from a larger index to a smaller one, and λ is weakly decreasing; so the partial sums of $\alpha|_C$ in index order dominate those of λ_C , and sorting only increases them. Hence the components form a compatible block decomposition, there are at most κ of them, and $|E| \geq \ell - \kappa$.

$E_{\min} \leq \ell - \kappa$: take a decomposition with κ blocks. By maximality each block $(\rho, \nu) = (\lambda_C, \mu^C)$ has no compatible refinement. Let $c_1 < \dots < c_L$ be the positions of C , pad ν with zeros to length L , and put $B_r = \nu_1 + \dots + \nu_r$, $P_r = \rho_1 + \dots + \rho_r$. Then $B_r > P_r$ for $1 \leq r < L$: $\geq$ holds by dominance, and $B_r = P_r$ would split (ρ, ν) into the compatible pieces $(\rho_{\leq r}, \nu_{\leq r})$ and $(\rho_{> r}, \nu_{> r})$, both of positive size. Put $\mathbf{m}_{c_r c_{r+1}} = B_r - P_r \geq 1$. Position c_r then ends with $\rho_r + (B_r - P_r) - (B_{r-1} - P_{r-1}) = \nu_r \geq 0$. This uses $L - 1$ edges per block, $\ell - \kappa$ in all.

Finally, $c_{\lambda\mu} \in \mathbb{Q}[s, t]$ (Theorem 4.1); write $c_{\lambda\mu} = \sum_j (s-1)^j a_j(t)$. The lower bound gives $a_j = 0$ for $j < \ell - \kappa$, and $a_{\ell-\kappa}(0) = (-1)^{\ell-\kappa} N(\lambda, \mu) \neq 0$, so $a_{\ell-\kappa} \neq 0$. $\square$

Remark 6.8 (Novelty). We have not found the block law stated anywhere. Its $t = 0$ case is classical in substance: the raising-operator formula follows from [DLT94, eq. (11)] and Jacobi–Trudi, after which the valuation count above is a short exercise. What is new is the lower bound for all t (from the subset formula alone), the exactness over $\mathbb{Q}(t)$, and the merge weights below.

6.5. Coarsenings and merge weights.

Theorem 6.9 (Coarsenings: merge-history expansion). *Let $\mu \neq \lambda$ be a coarsening of λ and $m = \ell(\lambda) - \ell(\mu)$. Then $v(c_{\lambda\mu}) = m$, and $\text{Lead}_{\lambda\mu}(t) := [(s-1)^m] c_{\lambda\mu}$ equals*

$$\text{Lead}_{\lambda\mu}(t) = \sum_{\text{tight histories } H \text{ ending at } \mu} \prod \text{merges in } H W_k(J), \quad \text{Lead}_{\lambda\mu}(0) = (-1)^m \#\{H\} \neq 0,$$

where a (tight) history processes $\lambda_\ell, \dots, \lambda_1$, each part k either opening a block of size k or merging $p \geq 1$ existing blocks of sizes $J = (j_1, \dots, j_p)$ into one block of size $k + |J|$ via the order- p piece $E_k^{(p)}$, it is tight if the orders add up to m , it ends at μ if the final block sizes are μ , and $W_k(J) := \text{lin}_e[\dots[E_k^{(p)}, e_{j_1}], \dots, e_{j_p}](1)$.

Proof. Since $\ell(\mu) = \ell - m$, the coefficient of e_μ in $[(s-1)^m] e_\lambda^*$ can be read modulo $I^{\ell-m+1}$. In Proposition 6.4, a product over $|\pi| > \ell - m$ blocks of positive-degree factors lies in $I^{\ell-m+1}$; so only the terms where every step merges exactly p_j blocks survive, i.e. tight histories. At a tight step, δ_S is a derivation in each argument (Lemma 6.3), so $\delta_S(bc, \dots) \in I^2$ for $b, c \in I$, and modulo I^2 , $\delta_S(g_1, \dots, g_p)$ depends only on the linear parts $g_i \equiv \gamma_i e_{n_i}$: it is $\prod_i \gamma_i W_k(n_1, \dots, n_p) e_n$. An opening step contributes e_k , with $\gamma = 1$. The final product of blocks is $\prod_C \gamma_C e_{|C|}$ modulo $I^{\ell-m+1}$, which gives the formula. By Theorem 6.10 below, $W_k(J)|_{t=0} = (-1)^{|J|}$, so every tight

history contributes $(-1)^m$ at $t = 0$. Histories exist: in each block of the coarsening, let the last-processed part merge all the others, opened earlier as singletons. So $\text{Lead}_{\lambda\mu}(0) \neq 0$, and $v(c_{\lambda\mu}) = m$ by the lower bound in Theorem 6.6. $\square$

Let $T_k f := \sum_{|A|=k} c_A x_A f(x_A)$ for $f \in \Lambda_k$, where $f(x_A)$ is f evaluated at the k variables x_a , $a \in A$. (This is unrelated to the Hecke operators of Section 2.)

Theorem 6.10 (Merge weights). *With $n = k + |J|$ and $p = |J|$,*

$$W_k(J) = (-1)^p \frac{[n]_t}{[k]_t} \prod_{j \in J} [k]_{t^j}.$$

Consequently $(-1)^m \text{Lead}_{\lambda\mu}(t) > 0$ for coarsenings and every real $t > 0$.

Lemma 6.11. *For $f \in \Lambda_k$ homogeneous of degree d and $n = k + d$:*

$$\text{line}_e T_k(f) = (-1)^d \frac{[n]_t}{[k]_t} f(1, t, \dots, t^{k-1}).$$

Remark 6.12 (Novelty). Lemma 6.11 is a short corollary of [Mac95, III.7 Ex. 2, III (2.2)] and is probably known.

Lemma 6.13. *For $\ell(\rho) \leq k \leq m$, $T_k(P_\rho(x_1, \dots, x_k; t)) = P_{\rho+1^k}(x_1, \dots, x_m; t)$.*

Proof. By [Mac95, III (2.2)], $P_\lambda = \sum_{w \in S_m/S_m^\lambda} w(x^\lambda \prod_{i < j, \lambda_i > \lambda_j} \frac{x_i - tx_j}{x_i - x_j})$, S_m^λ the stabiliser of λ . For $\lambda = \rho + 1^k$ the positive entries are the first k ; a coset is an image set $A = w([k])$ together with a coset of S_k/S_k^ρ ; the pairs with $\lambda_i > \lambda_j$ are those with $i \leq k < j$, giving c_A , and those with $i, j \leq k$, $\rho_i > \rho_j$; and $x^\lambda = x_{[k]} x_{[k]}^\rho$. Grouping by A , the inner sum is [Mac95, III (2.2)] for P_ρ in the variables x_A . $\square$

Lemma 6.14. *For $\lambda \vdash n$ with $\ell(\lambda) = k$, $\text{line}_e P_\lambda(x; t) = (-1)^{n-k} \frac{[n]_t}{[k]_t} P_{\lambda-1^k}(1, t, \dots, t^{k-1}; t)$.*

Proof. Since $[p_n]e_n = (-1)^{n-1}/n$ and e_ν with $\ell(\nu) \geq 2$ has no p_n term, $\text{line}_e G = (-1)^{n-1} \langle G, p_n \rangle = (-1)^{n-1} (1 - t^n) \langle G, p_n \rangle_t$. By $\langle P_\lambda, Q_\mu \rangle_t = \delta_{\lambda\mu}$ and $Q_\lambda = b_\lambda P_\lambda$ [Mac95, III (4.9)], $\langle P_\lambda, p_n \rangle_t = [P_\lambda] p_n / b_\lambda(t)$, and $[P_\lambda] p_n = t^{n(\lambda)} \varphi_{k-1}(t^{-1}) = (-1)^{k-1} t^{n(\lambda) - \binom{k}{2}} \varphi_{k-1}(t)$ [Mac95, III.7 Ex. 2], where $\varphi_r(t) = \prod_{a=1}^r (1 - t^a)$. On the other side, with $\rho = \lambda - 1^k$, [Mac95, III.2 Ex. 1] gives $P_\rho(1, \dots, t^{k-1}) = t^{n(\rho)} \varphi_k / \prod_{i \geq 0} \varphi_{m_i(\rho)}$ with $m_0(\rho) = k - \ell(\rho)$; here $m_i(\rho) = m_{i+1}(\lambda)$, so the denominator is b_λ , and $n(\rho) = n(\lambda) - \binom{k}{2}$. Since $\frac{[n]_t}{[k]_t} \varphi_k = (1 - t^n) \varphi_{k-1}$, the two sides agree. $\square$

Proof of Lemma 6.11. The $P_\rho(x_1, \dots, x_k; t)$ with $\ell(\rho) \leq k$ form a basis of Λ_k [Mac95, III (2.7)]; for $f = P_\rho$, $\rho \vdash d$, combine Lemmas 6.13 and 6.14. $\square$

Proof of Theorem 6.10. Let $J = (j_1, \dots, j_p)$, $d = |J|$. First, $W_k(J) = \text{line}_e E_k^{(p)}(e_{j_1} \cdots e_{j_p})$: in Lemma 6.3, $\delta_\emptyset = E_k^{(p)}(1) = 0$, and for $\emptyset \neq S \neq [p]$ the term $\prod_{i \notin S} e_{j_i} \delta_S$ lies in I^2 . Next, with $E(u) = \sum_r e_r u^r$ and $R(u) := \prod_{a \in A} \frac{1 + sux_a}{1 + ux_a}$ read as a function of x_A ,

$$E_k \left(\prod_{i=1}^p E(u_i) \right) = \prod_i E(u_i) \cdot T_k \left(\prod_i R(u_i) \right), \quad \log R(u) = \sum_{r \geq 1} (-1)^{r-1} \frac{s^r - 1}{r} u^r p_r(x_A).$$

Take $[u_1^{j_1} \cdots u_p^{j_p}]$ and $[(s-1)^p]$. Terms in which some e_{r_i} , $r_i \geq 1$, is taken from $\prod E(u_i)$ lie in I^2 . In the remaining term, each $[u_i^{j_i}] R(u_i)$ has $(s-1)$ -valuation ≥ 1 , with $(s-1)^1$ -coefficient $(-1)^{j_i-1} p_{j_i}(x_A)$. Hence $W_k(J) = (-1)^{d-p} \text{line}_e T_k(p_J)$, and Lemma 6.11 with $p_j(1, \dots, t^{k-1}) = [k]_{t^j}$ gives the formula. For $t > 0$ every $W_k(J)$ has sign $(-1)^{|J|}$, and the orders along a tight history add up to m . $\square$

7. LEADING COEFFICIENTS: GRAPH CUMULANTS AND BLOCKS

Throughout this section μ is a coarsening of λ , $\ell = \ell(\lambda)$, $m = \ell - \ell(\mu)$, and the parts of λ are indexed by positions $[\ell]$. Everything rests on Theorems 6.9 and 6.10.

Lemma 7.1 (Histories are increasing forests). *Tight histories of λ are in bijection with increasing forests on $[\ell]$ (every child larger than its parent): when position i merges existing blocks, i becomes the parent of their roots. The trees are the final blocks, and the orders automatically add up to $\ell - \#\text{trees}$.*

Proof. Every existing block at step i was built from positions $> i$, so all edges increase. Conversely, an increasing forest determines the history: at step i , merge the subtrees rooted at the children of i , which are complete by then. Each edge accounts for one unit of order. $\square$

Lemma 7.2 (Telescoping). *For an increasing tree T on a set C of positions, let N_v be the λ -mass of the subtree at v . The product of the merge weights along the history of T is*

$$\prod_v W_{\lambda_v}((N_c)_c \text{ child of } v) = \frac{1 - t^{|\lambda_C|}}{\prod_{i \in C} (1 - t^{\lambda_i})} \prod_{\text{edges } v \rightarrow c} (t^{\lambda_v N_c} - 1).$$

Proof. By Theorem 6.10, the merge at v contributes $(-1)^{\#\text{children}} \frac{1 - t^{N_v}}{1 - t^{\lambda_v}} \prod_c \frac{1 - t^{\lambda_v N_c}}{1 - t^{N_c}}$ (a leaf contributes 1). Every non-root c is a child exactly once, so its factor $1 - t^{N_c}$ cancels against its own numerator, leaving the root mass $|\lambda_C|$; and $(-1)^{\#\text{edges}} \prod (1 - x) = \prod (x - 1)$. $\square$

Lemma 7.3 (Tree-graph identity). *For indeterminates $x_{ij} = x_{ji}$ ($i \neq j \in [\ell]$),*

$$\sum_{\text{increasing trees } T \text{ on } [\ell]} \prod_{\text{edges } v \rightarrow c} \left(\prod_{j \in T_c} x_{vj} - 1 \right) = \sum_{\text{connected graphs } H \text{ on } [\ell]} \prod_{e \in H} (x_e - 1),$$

where T_c is the vertex set of the subtree at c .

Proof. Map a connected graph H to an increasing tree recursively: the root is $\min V$, the children are the minima of the components of $H - \min V$, and recurse on each component. Fix T and sum over the H mapping to it. At each vertex v and child c , the edges from v into T_c form an arbitrary nonempty set $S \subseteq T_c$ (nonempty since T_c must be attached to v ; otherwise free, since they do not change the components inside T_c); there are no edges between different child sets; and the edges inside T_c are accounted for recursively. So the fibre sum factors, and $\sum_{\emptyset \neq S \subseteq T_c} \prod_{j \in S} (x_{vj} - 1) = \prod_{j \in T_c} x_{vj} - 1$. $\square$

Theorem 7.4 (Full-merge lead). *For $\lambda \vdash n$ with $\ell = \ell(\lambda) \geq 2$,*

$$\text{Lead}_{\lambda, (n)}(t) = \frac{1 - t^n}{\prod_i (1 - t^{\lambda_i})} K_\lambda(t), \quad K_\lambda(t) := \sum_{H \text{ connected on } [\ell]} \prod_{ij \in H} (t^{\lambda_i \lambda_j} - 1).$$

K_λ is the connected part (joint cumulant) of $t^{e_2(\lambda)} = \prod_{i < j} (1 + (t^{\lambda_i \lambda_j} - 1))$.

Proof. By Theorem 6.9 with $\mu = (n)$ and Lemma 7.1, $\text{Lead}_{\lambda, (n)}$ is a sum over increasing trees on $[\ell]$; by Lemma 7.2 each tree contributes $\frac{1 - t^n}{\prod_i (1 - t^{\lambda_i})} \prod_{v \rightarrow c} (t^{\lambda_v N_c} - 1)$; and Lemma 7.3 with $x_{ij} = t^{\lambda_i \lambda_j}$, for which $\prod_{j \in T_c} x_{vj} = t^{\lambda_v N_c}$, converts the tree sum into K_λ . The cumulant statement is the exponential formula. $\square$

Corollary 7.5. (a) $\lambda = 1^n$: $\text{Lead} = (-1)^{n-1} [n]_t I_n(t)$ with I_n the tree inversion enumerator [MR68]. (b) $t = 0$: $\text{Lead} = (-1)^{\ell-1} (\ell - 1)!$. (c) $t \rightarrow 1$: $\text{Lead} \rightarrow (-1)^{\ell-1} n^{\ell-1}$ (weighted Cayley). (d) $\text{sgn Lead} = (-1)^{\ell-1}$ for all $t > 0$.

Proof. (a) For $\lambda = 1^n$ the prefactor is $[n]_t/(1-t)^{n-1}$ and $K_\lambda = \sum_{H \text{ conn.}} (t-1)^{|H|} = (t-1)^{n-1} I_n(t)$ [MR68]. (b) At $t = 0$ the prefactor is 1, and in the tree form every edge factor is -1 ; there are $(\ell - 1)!$ increasing trees. (c) Put $\varepsilon = t - 1$. In K_λ only spanning trees contribute to the lowest order $\varepsilon^{\ell-1}$, with $t^{\lambda_i \lambda_j} - 1 = \lambda_i \lambda_j \varepsilon + O(\varepsilon^2)$, and $\sum_T \prod_{ij \in T} \lambda_i \lambda_j = \sum_T \prod_i \lambda_i^{\deg_T(i)} = (\prod_i \lambda_i) n^{\ell-2}$ by the vertex-weighted Cayley formula (each i occurs $\deg_T(i) - 1$ times in the Prüfer code). The prefactor is $(-n\varepsilon)/((- \varepsilon)^\ell \prod_i \lambda_i) \cdot (1 + O(\varepsilon))$. (d) is the case $\mu = (n)$, $m = \ell - 1$, of the sign statement in Theorem 6.10: each merge contributes $(-1)^p$ times a positive quantity, $\sum p = m$ along every history, and histories exist. $\square$

Remark 7.6 (Prior art — graph half). The tree, connected-graph and cumulant forms of K_λ are a special case of Dołęga [Do19, Prop. 2.1 and (11), Lemma 2.3] ($K_\lambda = (t-1)^{\ell-1} T(1, t)$ for the Tutte polynomial of the multigraph with $\lambda_i \lambda_j$ edges between i and j), cf. also [GS96; MR68]; step (3) is a Penrose-type tree-graph identity [Pen67]. Ours: the identification of the $\star$ -lead with prefactor $\cdot K_\lambda$ (Theorems 6.9, 6.10 and 7.4) and Theorems 7.7 and 7.8. Haglund–Tewari’s single-row cumulant [HT26, Def. 7.1, Thm. 7.3] (their q is our t) is the same graph half: since $\langle \tilde{H}_{(m)}, e_m \rangle = q^{\binom{m}{2}}$, their cumulant $\kappa(\lambda)$ (we write C_λ) satisfies $\langle C_\lambda, e_n \rangle = t^{n(\lambda')}(t-1)^{1-\ell} K_\lambda(t)$. *Separator.* Full-merge leads $L(\lambda)$ of any such theory are only defined up to the gauge $L(\lambda) \mapsto L(\lambda) a^{\ell(\lambda)} g(|\lambda|) \prod_i f(\lambda_i)$ (rescaling generators, degree-wise normalisation, rescaling the expansion parameter). The first gauge invariant appears at $n = 4$: $J := L(1^4)L(2, 2)/L(2, 1, 1)^2$ (the exponents of $a, g(4), f(1)$ and $f(2)$ all cancel). For $\star$, Theorem 7.4 gives $J = \frac{(t^2+1)(t^3+3t^2+6t+6)}{(t+1)(t^2+t+2)^2}$, and Haglund–Tewari’s $\langle C_\lambda, e_n \rangle$ gives the same value. Dołęga’s $\oplus$ -cumulant of columns (lowest $(q-1)$ -coefficient of $[g_n] \tilde{H}_{\lambda'}$, $g_k = \tilde{H}_{1^k}$) gives, by computation, $J \equiv 3/2 = J_\star(0)$, whereas $J_\star(1) = 1$, the Cayley value. So up to gauge Dołęga’s $\oplus$ is t -independent and agrees with $\star$ only at $t = 0$, where the leads are partition-lattice Möbius values (Corollary 7.5(b)), while Haglund–Tewari’s is the same graph object without $\star$.

Theorem 7.7 (Coarsening leads factor). *For μ a coarsening of λ ,*

$$\text{Lead}_{\lambda\mu} = \sum_{\pi} \prod_{B \in \pi} \text{Lead}_{\lambda_B, (|\lambda_B|)},$$

summed over set partitions π of the positions $[\ell]$ whose sorted block sums equal μ (singletons contribute 1). There is no interleaving factor.

Proof. By Lemma 7.1, tight histories ending at μ are increasing forests whose trees have sorted masses μ , i.e. a set partition π with block sums μ together with an increasing tree on each block. By Lemma 7.2 the weight is a product over the trees, each depending only on its own parts; summing over the trees on each block gives $\text{Lead}_{\lambda_B, (|\lambda_B|)}$ by Theorem 7.4 (and 1 for a singleton). $\square$

Theorem 7.8 (Block multiplicativity). *Let $\kappa = \kappa(\lambda, \mu) \geq 1$, $\ell = \ell(\lambda)$, $m = \ell - \kappa$. Then*

$$[(s-1)^m] c_{\lambda\mu} = \sum_{\pi} \sum_{(\nu^C)_{C \in \pi}} \prod_{C \in \pi} [(s-1)^{|C|-1}] c_{\lambda_C, \nu^C},$$

where π runs over set partitions of $[\ell]$ into exactly κ blocks and (ν^C) over tuples with $\bigsqcup_C \nu^C = \mu$ and $\nu^C \sqsupseteq \lambda_C$; every factor then has $\kappa(\lambda_C, \nu^C) = 1$. With Theorem 6.6 this is $\text{Lead}_{\lambda\mu}$.

Proof. Expand $[(s-1)^m] e_\lambda^\star = \sum_{\sum p_i = m} E_{\lambda_1}^{(p_1)} \cdots E_{\lambda_\ell}^{(p_\ell)}(1)$ step by step with Lemma 6.3, as in Proposition 6.4: this is a sum over histories H (step i chooses a set S_i of current blocks with $|S_i| \leq p_i$), each giving $\prod_{C \in \pi(H)} g_C^H$ with g_C^H of type λ_C , and $|\pi(H)| = \ell - \sum |S_i| \geq \ell - m = \kappa$, with equality iff $|S_i| = p_i$ for all i (tight). A non-tight term has more than κ blocks, so

its e_μ -coefficient vanishes by maximality of κ . In a tight history, step i only acts, through δ_{S_i} , on the blocks it merges; so g_C^H is the function produced by the restricted history on C , which is a single-block tight history of the composition λ_C (positions of C in their order), since $\sum_{i \in C} |S_i| = |C| - 1$. Conversely, single-block tight histories on the blocks of π interleave uniquely, by position, into a global tight history. Hence the tight part is $\sum_{|\pi|=\kappa} \prod_{C \in \pi} \text{Conn}(\lambda_C)$, where $\text{Conn}(\lambda_C)$ is the sum over single-block tight histories of λ_C . Applying the same expansion to λ_C with $m = |C| - 1$ shows $[e_\nu] \text{Conn}(\lambda_C) = [e_\nu] [(s-1)^{|C|-1}] e_{\lambda_C}^*$ whenever $\kappa(\lambda_C, \nu) = 1$. Finally, $[e_\mu] \prod_C \text{Conn}(\lambda_C) = \sum_{(\nu^C)} \prod_C [e_{\nu^C}] \text{Conn}(\lambda_C)$ over tuples with $\bigsqcup_C \nu^C = \mu$; factors with $\nu^C \not\leq \lambda_C$ vanish by type, and $\kappa(\lambda_C, \nu^C) \geq 2$ would refine π to more than κ compatible blocks. Commutativity of $\star$ [Hik25] identifies the coefficients for the composition λ_C with those for the sorted partition. $\square$

Remark 7.9. Theorem 7.7 is the case where each ν^C is a single part. The factorisation has the shape of a Mayer (cluster) expansion, i.e. moment–cumulant inversion of the graph identity in Theorem 7.4; the $\star$ -specific content is that the connected leads are those of Theorem 7.4, with no interleaving correction.

8. BOX COMPLEMENT AND THE LINEAR COEFFICIENT

In this section zero parts are allowed: λ is padded with zeros to a prescribed length, $E_0 = \text{id}$, and $c_{\lambda\mu}$ refers to the nonzero parts. By Lemma 2.8, in N variables (for every N , including N below the degree) $e_\lambda^*(x_1, \dots, x_N) = \sum_\mu c_{\lambda\mu} e_\mu(x_1, \dots, x_N)$, where $e_\mu = 0$ if $\mu_1 > N$.

Theorem 8.1 (Box Complement). *Let λ, μ have at most ℓ parts (padded with zeros) and all parts $\leq N$; put $N^\ell - \lambda := (N - \lambda_\ell, \dots, N - \lambda_1)$. Then*

$$c_{N^\ell - \lambda, N^\ell - \mu}(s, t) = s^{N \binom{\ell}{2} - (\ell-1)|\lambda|} c_{\lambda\mu}(s, t).$$

Lemma 8.2 (Inversion). *In N variables, let G be homogeneous of degree g and $F(x) := e_N(x)^m G(1/x)$. Then*

$$E_{N-k} F = s^{m(N-k)-g} e_N^{m+1} (E_k G)(1/x).$$

Proof. Write $y = 1/x$ and index the terms of E_{N-k} by the complement A ($|A| = k$) of the $(N-k)$ -subset B . Then $c_B(x) = \prod_{i \in B, j \in A} \frac{y_j - t y_i}{y_j - y_i} = c_A(y)$; $x_B = e_N(x) y_A$; $F|_{x_i \mapsto s x_i} (i \in B) = s^{m(N-k)} e_N(x)^m G|_{y_i \mapsto s^{-1} y_i} (i \in B)$; and by homogeneity $G|_{y_i \mapsto s^{-1} y_i} (i \in B) = s^{-g} G|_{y_a \mapsto s y_a} (a \in A)$. Summing over A gives $s^{m(N-k)-g} e_N^{m+1} \sum_A c_A(y) y_A G|_{y_a \mapsto s y_a}$. $\square$

Proof of Theorem 8.1. Let $G_j := E_{\lambda_{\ell-j+1}} \cdots E_{\lambda_\ell}(1)$, of degree $D_j = \lambda_\ell + \cdots + \lambda_{\ell-j+1}$, and $F_j := E_{N-\lambda_{\ell-j+1}} \cdots E_{N-\lambda_\ell}(1)$, in N variables. By Lemma 8.2 and induction, $F_j = s^{\sigma_j} e_N^j G_j(1/x)$ with $\sigma_0 = 0$ and $\sigma_{j+1} = \sigma_j + j(N - \lambda_{\ell-j}) - D_j$. Writing $p_i = \lambda_{\ell+1-i}$,

$$\sigma_\ell = N \binom{\ell}{2} - \sum_i (i-1)p_i - \sum_i (\ell-i)p_i = N \binom{\ell}{2} - (\ell-1)|\lambda|.$$

By commutativity of $\star$, $F_\ell = e_{N^\ell - \lambda}^*$. Since $e_{N-r}(x) = e_N(x) e_r(1/x)$ in N variables, $e_N^\ell e_\mu(1/x) = e_{N^\ell - \mu}(x)$ for $\ell(\mu) \leq \ell$, and every μ in the support of e_λ^* has $\ell(\mu) \leq \ell(\lambda) \leq \ell$ (Theorem 4.1). Hence $e_{N^\ell - \lambda}^* = s^{\sigma_\ell} \sum_{\mu_1 \leq N} c_{\lambda\mu} e_{N^\ell - \mu}(x)$. The map $\mu \mapsto N^\ell - \mu$ is a bijection between partitions with at most ℓ parts, all $\leq N$; every ν in the support of $e_{N^\ell - \lambda}^*$ has at most ℓ parts; and the e_ν with $\nu_1 \leq N$ form a basis of Λ_N . Compare coefficients. $\square$

Corollary 8.3 (Column Lemma; lead invariance). $c_{\lambda+1^\ell, \mu+1^\ell} = s^{\binom{\ell}{2}} c_{\lambda\mu}$. Hence v and Lead are invariant under $(\lambda, \mu) \mapsto (\lambda + 1^\ell, \mu + 1^\ell)$ and $(\lambda, \mu) \mapsto (N^\ell - \lambda, N^\ell - \mu)$; e.g. $(2, 2, 2) \rightarrow (3, 3)$ and $(2, 2, 2) \rightarrow (4, 1, 1)$ have the lead $[3]_t(t+2)$ of $(1, 1, 1) \rightarrow (3)$.

Proof. Complement with N and then with $N+1$: $(N+1)^\ell - (N^\ell - \lambda) = \lambda + 1^\ell$, and the exponents add up to $(N^\ell - (\ell-1)|\lambda|) + ((N+1)^\ell - (\ell-1)(N^\ell - |\lambda|)) = \binom{\ell}{2}$. Since s^σ is a unit of $\mathbb{Q}[[s-1]]$ with constant term 1, valuations and leading coefficients are preserved. For the example, $(2, 2, 2) \rightarrow (4, 1, 1)$ is $(1, 1, 1) \rightarrow (3, 0, 0)$ plus a column, and $(2, 2, 2) \rightarrow (3, 3)$ is the complement with $N=3$ of $(1, 1, 1) \rightarrow (3, 0, 0)$; the lead of $(1, 1, 1) \rightarrow (3)$ is $[3]_t(t+2)$ by Theorem 7.4. $\square$

Remark 8.4 (Prior art). The inversion lemma follows in a few lines from [DFK19, Rem. 3.3] ($S\Gamma_I S = \Gamma_I^{-1}$ under $x \mapsto x^{-1}$) plus subset complement; stability is [Hik25, Prop. 3.8, Cor. 3.10]. We have not found Theorem 8.1 or Corollary 8.3 for $\star$ in the literature.

Theorem 8.5 (Plethystic linear coefficient). For G homogeneous of degree d : $\text{lin}_e(e_k \star G) = (-1)^d \frac{[k+d]_t}{[k]_t} G[(s-1)[k]_t]$, where $G \mapsto G[(s-1)[k]_t]$ is the ring map $p_r \mapsto (s^r - 1)[k]_{tr}$.

Proof. By linearity take $G = e_{j_1} \cdots e_{j_p}$, $d = \sum j_i$. As in the proof of Theorem 6.10, $E_k(e_J) = \sum_{w \leq J} e_{J-w} T_k([u^w] \prod_i R(u_i))$. For $w \neq J$ the term is a product of an element of I and a symmetric function of positive degree, hence lies in I^2 . For $w = J$, Lemma 6.11 applied to the symmetric function $f = \prod_i [u_i^{j_i}] R(u_i)$ of x_A gives $(-1)^d \frac{[k+d]_t}{[k]_t} f(1, t, \dots, t^{k-1})$, and $[u^j] R(u)$ at $x_A = (1, t, \dots, t^{k-1})$ is $\pi_k(j) := [u^j] \prod_{m=0}^{k-1} \frac{1+sut^m}{1+ut^m}$. Taking logarithms, $e_j \mapsto \pi_k(j)$ is the ring map $p_r \mapsto (s^r - 1)[k]_{tr}$. $\square$

Corollary 8.6 (Pieri rule, all s). $e_k \star e_r = \sum_{y=0}^{\min(k,r)} s^y C_{k-y, r-y} e_{k+r-y} e_y$ with $C_{a,0} = C_{0,b} = 1$, $C_{a,b} = (-1)^b \frac{[a+b]_t}{[a]_t} \pi_a(b)$, $\pi_a(j) = [u^j] \prod_{m=0}^{a-1} \frac{1+sut^m}{1+ut^m}$.

Proof. By Theorem 4.1 and commutativity, the support of $e_k \star e_r$ consists of $e_{k+r-y} e_y$ with $y \leq \min(k, r)$. Applying Corollary 8.3 with $\ell = 2$ y times, $c_{(k,r), (k+r-y, y)} = s^y c_{(k-y, r-y), (k+r-2y, 0)} = s^y \text{lin}_e(e_{k-y} \star e_{r-y})$. If $k-y = 0$ this is s^y ; otherwise Theorem 8.5 with $G = e_{r-y}$ gives $s^y (-1)^{r-y} \frac{[k+r-2y]_t}{[k-y]_t} \pi_{k-y}(r-y) = s^y C_{k-y, r-y}$. $\square$

Remark 8.7. This proof of the Pieri rule is independent of the one in Section 3: it uses neither the parabolic kernel computation of Proposition 2.2 beyond the subset formula, nor the recursion of Theorem 3.2. For $r \geq k$ the monomials $e_b e_{r+k-b}$, $0 \leq b \leq k$, are distinct, so comparing with Corollary 3.3 gives $F_n(t^m) = (-1)^m \frac{[n+m]_t}{[n]_t} \pi_n(m)$ for $1 \leq n \leq m$. Hikita's [Hik25, Thm. 3.12] is the case $k=1$.

9. SECOND ORDER: EVERY VALUATION-2 LEAD

By Theorem 7.8, valuation-2 leads reduce to connected leads with $\ell(\lambda) = 3$, $\kappa = 1$; by Corollary 8.3 to two-part μ . With T_a as in Lemma 6.11, put

$$\Gamma_a(e_b, e_c) := E_a^{(2)}(e_b e_c) - e_b E_a^{(2)}(e_c) - e_c E_a^{(2)}(e_b) = \sum_{1 \leq r \leq b, 1 \leq q \leq c} (-1)^{r+q} e_{b-r} e_{c-q} T_a(p_r p_q).$$

Indeed, with $R(u)$ as in the proof of Theorem 6.10, $R(u) = \prod_{x \in A} (1 + (s-1)y_x(u))$ with $y_x(u) = \frac{ux_x}{1+ux_x}$, so $[(s-1)^2] R(u) R(v) = e_2(Y_u) + e_2(Y_v) + p_1(Y_u) p_1(Y_v)$ in the alphabets $Y_u = \{y_x(u)\}_{x \in A}$, Y_v ; hence $E_a^{(2)}(E(u) E(v)) - E(u) E_a^{(2)} E(v) - E(v) E_a^{(2)} E(u) = E(u) E(v) T_a(p_1(Y_u) p_1(Y_v))$, and $p_1(Y_u) = \sum_{r \geq 1} (-1)^{r-1} u^r p_r(x_A)$. Take $[u^b v^c]$.

Proposition 9.1. *If (a, b, c) is any ordering of λ and $\kappa(\lambda, \mu) = 1$, then $[(s-1)^2]c_{\lambda\mu} = [e_\mu](\Gamma_a(e_b, e_c) + D_a D_b(e_c))$.*

Proof. Since $E^{(p)}(1) = 0$ for $p \geq 1$, $[(s-1)^2]E_a E_b E_c(1) = E_a^{(2)}(e_b e_c) + D_a D_b(e_c) + e_a E_b^{(2)}(e_c)$, and $E_a^{(2)}(e_b e_c) = \Gamma_a(e_b, e_c) + e_b E_a^{(2)}(e_c) + e_c E_a^{(2)}(e_b)$. Each of $e_b E_a^{(2)}(e_c)$, $e_c E_a^{(2)}(e_b)$, $e_a E_b^{(2)}(e_c)$ is a part e_{λ_i} times a function of the type of the other two parts (Proposition 6.4), so every μ in its support has $\kappa(\lambda, \mu) \geq 2$. $\square$

Theorem 9.2 (Constant-term adjoint formula). *For $g \in \Lambda_a$ and $F \in \Lambda$,*

$$\varphi_a(t) \langle T_a g, F \rangle_t = \text{CT}[Z g(z) F(z_1^{-1}, \dots, z_a^{-1}) \Omega(z)],$$

where $Z = z_1 \cdots z_a$ and $\Omega(z) = \prod_{1 \leq i < j \leq a} \frac{z_j - z_i}{1 - tz_j/z_i}$, expanded in z_i/z_j ($i < j$).

Proof. Let $G = T_a g$. In exactly a variables only $A = [a]$ survives and $c_{[a]} = 1$, so $G(z) = Zg(z)$; and $G \in \text{span}\{P_\mu : \ell(\mu) = a\}$ by Lemma 6.13. We read CT of a Laurent polynomial times Ω either formally or as the integral over the torus $|z_i| = 1$ for $|t| < 1$; the two agree because the expansion of Ω converges there, and both sides of the theorem are rational in t .

Step 1. For $\alpha \in \mathbb{Z}_{\geq 0}^a$, $\langle G, q_\alpha \rangle_t = [y^\alpha]G(y_1, \dots, y_a)$, with $q_\alpha = \prod q_{\alpha_i}$: pair G with the Cauchy identity $\sum_\mu Q_\mu(x) P_\mu(y) = \prod_{i,j} \frac{1 - tx_i y_j}{1 - x_i y_j} = \sum_\alpha q_\alpha(x) y^\alpha$ [Mac95, III (4.2), (4.4)], using $\langle P_\lambda, Q_\mu \rangle_t = \delta_{\lambda\mu}$ [Mac95, III (4.9)].

Step 2. For $\ell(\lambda) \leq a$ (padded), the raising-operator formula [Mac95, III (2.15')] gives $Q_\lambda = \sum_\beta c_\beta q_{\lambda+\beta}$ with $\sum_\beta c_\beta y^{-\beta} = \prod_{i < j} \frac{1 - y_j/y_i}{1 - ty_j/y_i}$; operators involving positions beyond $\ell(\lambda)$ only produce negative entries, where q and $[y]$ both vanish. With step 1 and the relabelling $z_k = y_{a+1-k}$,

$$\langle G, Q_\lambda \rangle_t = \text{CT}[G(z) z^{-\lambda^{\text{rev}}} \Omega(z)], \quad \lambda^{\text{rev}} = (\lambda_a, \dots, \lambda_1).$$

Step 3. Let $\Delta = \prod_{i \neq j} \frac{1 - z_i/z_j}{1 - tz_i/z_j}$ and $\theta = \prod_{i < j} \frac{z_i - tz_j}{z_i - z_j}$, so that $\Delta\theta = \Omega$. For symmetric f , $f\Omega$ is analytic on the torus and $f\Delta$ is symmetric, so averaging over S_a and using $\sum_w w(\theta) = v_a(t) = [a]_t!$ [Mac95, III (1.4)] gives $\text{CT}[f\Omega] = \frac{v_a}{a!} \text{CT}[f\Delta]$.

Step 4. By [Mac95, III (2.1)–(2.2)], $Q_\lambda(z^{-1}) = \frac{b_\lambda}{v_\lambda} \sum_w w(z^{-\lambda} \tilde{\theta})$ with $\tilde{\theta} = \prod_{i < j} \frac{z_j - tz_i}{z_j - z_i}$ and $v_\lambda = \prod_{i \geq 0} v_{m_i(\lambda)}$, $m_0 = a - \ell(\lambda)$. Since $\tilde{\theta}\Delta = \prod_{i < j} \frac{z_i - z_j}{z_i - tz_j}$, step 3, symmetry, and the relabelling $z_k \leftrightarrow z_{a+1-k}$ give

$$\text{CT}[G Q_\lambda(z^{-1}) \Omega] = v_a \frac{b_\lambda}{v_\lambda} \text{CT}[G z^{-\lambda^{\text{rev}}} \Omega] = v_a \frac{b_\lambda}{v_\lambda} \langle G, Q_\lambda \rangle_t.$$

Step 5. Write $F = \sum_\lambda F_\lambda Q_\lambda$. Terms with $\ell(\lambda) > a$ vanish on both sides, and $\langle G, Q_\lambda \rangle_t = 0$ unless $\ell(\lambda) = a$, in which case $m_0 = 0$ and $v_a b_\lambda/v_\lambda = v_a(1-t)^a = \varphi_a$. Hence $\text{CT}[GF(z^{-1})\Omega] = \varphi_a \langle G, F \rangle_t$. $\square$

Theorem 9.3 (Two-point formula). *Let $g \in \Lambda_a$ be homogeneous of degree d , $n = a + d$, $x, y \geq 1$, $x + y = n$, and $\Phi_a(g; x, y) := \langle T_a g, p_x p_y \rangle$. For $1 \leq A \leq a - 1$ put $B = a - A$ and $G_A(w) =$*

$\sum_k G_{A,k} w^k := g(1, t, \dots, t^{A-1}, w, wt, \dots, wt^{B-1})$. Then

$$\begin{aligned} \Phi_a(g; x, y) = (-1)^a (1 - t^x)(1 - t^y) & \left\{ \sum_{A=1}^{a-1} t^{-AB} \left[\frac{G_{A,y-B}}{(1-t^A)(1-t^B)} \right. \right. \\ & \left. \left. + \frac{1}{1-t^a} \sum_{m \geq 1} (t^{-Am} - t^{Bm}) G_{A,y-B-m} \right] \right. \\ & \left. - \frac{g(1, t, \dots, t^{a-1})}{1-t^a} \sum_{j=0}^{a-1} t^{-jy} \right\}. \end{aligned}$$

Proof. Let $S(u) := \sum_{i=1}^a \frac{1}{uz_i-1} = \sum_{x \geq 1} u^{-x} p_x(z^{-1})$. By Theorem 9.2 and homogeneity,

$$\mathcal{R}(u, v) := \frac{1}{\varphi_a} \text{CT}[g Z S(u) S(v) \Omega] = \sum_{x+y=n} \frac{\Phi_a(g; x, y)}{(1-t^x)(1-t^y)} u^{-x} v^{-y} = u^{-n} \rho(w), \quad w = u/v,$$

with ρ a polynomial, so $\Phi_a(g; x, y) = (1-t^x)(1-t^y)[w^y]\rho$.

String expansion. Take $|t|$ small, u, v generic, and radii $\max(|u|^{-1}, |v|^{-1}) < r_1 < \dots < r_a$; the constant term is the integral of $g S(u) S(v) \Omega \prod_m dz_m / (2\pi i)$ over $|z_m| = r_m$. Expand $S(u) S(v)$ into its a^2 summands and integrate $z_1, \dots, z_a$ in turn. When z_m is integrated, its poles inside $|z_m| = r_m$ are at $1/u$ or $1/v$ if z_m carries the u - or v -factor, and at tz_l for $l < m$ (from the factor $\frac{z_m - z_l}{z_m - tz_l}$); the poles z_j/t , $j > m$, lie outside, and g is a polynomial. The numerator $\prod_{l < m} (z_m - z_l)$ kills the residue at tz_l if some earlier $z_{l'}$ already equals tz_l . Hence the evaluation points form t -strings $q, qt, \dots, qt^{A-1}$ (at increasing positions), each started by a *seed* at $1/u$ or $1/v$, and position 1 is always a seed. The surviving configurations are: (i) a word $\omega \in \{U, V\}^a$ containing both letters, the u -factor at the first U and the v -factor at the first V , the $(\alpha + 1)$ -st U at t^α/u and the $(\beta + 1)$ -st V at t^β/v ; (ii) the word U^a , all points t^{m-1}/u , with the v -factor at any position $j + 1$, where it is simply evaluated; (iii) the mirror image V^a .

String weight. A string of length A with base q contributes $(-1)^{A-1} \varphi_{A-1}(t) q^A$: the seed residue is q ; a successor pair contributes the residue $qt^\alpha(t-1)$; a non-successor pair at string distance $\delta \geq 2$ contributes $\frac{t^\delta-1}{t^\delta-t}$; and $\prod_{\delta=2}^{A-1} (\frac{1-t^\delta}{t(1-t^{\delta-1})})^{A-\delta} = t^{-\binom{A-1}{2}} \varphi_{A-1}/(1-t)^{A-1}$.

Interaction. For a U -string of length A and a V -string of length $B = a - A$, the cross factors are $\frac{wt^{\beta-\alpha}-1}{wt^{\beta-\alpha}-t}$ (U_α before V_β) and $\frac{1-wt^{\beta-\alpha}}{1-wt^{\beta-\alpha+1}}$ (V_β before U_α), and their sum over shuffles is the function $\text{Sh}_{A,B}(w)$ of Lemma 9.4.

Assembling. Since $g(z^*) = u^{-d} G_A(w)$ for configuration (i) and $u^{-d} g(\pi_a)$ for (ii), where $\pi_a = (1, t, \dots, t^{a-1})$, and $v^{-B} = u^{-B} w^B$,

$$\varphi_a \rho(w) = \sum_{A=1}^{a-1} (-1)^A \varphi_{A-1} \varphi_{B-1} w^B \text{Sh}_{A,B}(w) G_A(w) + (-1)^{a-1} \varphi_{a-1} g(\pi_a) \sum_{j=0}^{a-1} \frac{w}{t^j - w} + O(w^n),$$

the last term coming from (iii). This is an identity of rational functions of w , each term analytic at $w = 0$, so $[w^y]$ may be taken termwise. By Lemma 9.4, $\text{Sh}_{A,B}(w) = \left[\begin{smallmatrix} a \\ A \end{smallmatrix} \right]_t t^{-AB} (1 + \kappa_{AB} \sum_{m \geq 1} (t^{-Am} - t^{Bm}) w^m)$ with $\kappa_{AB} = \frac{(1-t^A)(1-t^B)}{1-t^a}$, and $\left[\begin{smallmatrix} a \\ A \end{smallmatrix} \right]_t \varphi_{A-1} \varphi_{B-1} = \frac{\varphi_a}{(1-t^A)(1-t^B)}$, $\frac{\varphi_{a-1}}{\varphi_a} = \frac{1}{1-t^a}$. For $1 \leq y \leq n-1$ this gives the formula. $\square$

Lemma 9.4 (Shuffle identity).

$$\text{Sh}_{A,B}(w) = \left[\begin{smallmatrix} A+B \\ A \end{smallmatrix} \right]_t t^{-AB} \frac{(1-w)(1-t^{B-A}w)}{(1-t^A w)(1-t^B w)},$$

$$\frac{(1-w)(1-t^{B-A}w)}{(1-t^{-A}w)(1-t^Bw)} = 1 + \kappa_{AB} \sum_{m \geq 1} (t^{-Am} - t^{Bm})w^m.$$

Proof. Induction on $A + B$; for $A = 0$ or $B = 0$ both sides are 1. Condition on the last letter. If it is U_{A-1} , its cross factors telescope to $\frac{1-wt^{1-A}}{1-wt^{B+1-A}}$; if it is V_{B-1} , to $t^{-A} \frac{1-wt^{B-1}}{1-wt^{B-1-A}}$. So $\text{Sh}_{A,B} = \frac{1-wt^{1-A}}{1-wt^{B+1-A}} \text{Sh}_{A-1,B} + t^{-A} \frac{1-wt^{B-1}}{1-wt^{B-1-A}} \text{Sh}_{A,B-1}$. With $X = t^A$, $Y = t^B$, $[\frac{A+B-1}{A-1}]_t / [\frac{A+B}{A}]_t = \frac{1-X}{1-XY}$ and $[\frac{A+B-1}{A}]_t / [\frac{A+B}{A}]_t = \frac{1-Y}{1-XY}$, the claimed closed form satisfies this recursion if and only if $Y(1-X)(1-w/X) + (1-Y)(1-Yw) = (1-XY)(1-wY/X)$, and both sides equal $1 - XY - wY/X + wY^2$. For the expansion: the left side is 1 at $w = 0$, tends to 1 at ∞ , and has simple poles with residue $-t^A \kappa_{AB}$ at $w = t^A$ and $t^{-B} \kappa_{AB}$ at $w = t^{-B}$; so does the right side. $\square$

Remark 9.5 (Relation to Green polynomials). Write $p_\mu = \sum_\lambda X_\mu^\lambda(t) P_\lambda(x; t)$. For $g = P_\rho$ and $\lambda = \rho + 1^a$ one has $\Phi_a(P_\rho; x, y) = (1-t^x)(1-t^y) X_{(x,y)}^\lambda(t) / b_\lambda(t)$, so Theorem 9.3 evaluates Green polynomials at two-part classes; the two t -strings are the Frobenius orbits of a maximal torus of type (x, y) [Gre55]. Theorem 9.2 is Jing's constant-term expression for the matrix element $\langle Q_\lambda, p_\mu \rangle$ [Jin91], cf. [JL21, (2.17), (2.20)]. Jing and Liu [JL21, Thm. 2.7] expand the same matrix element as a nested sum valid for all μ , extracting one variable at a time; at two-part μ that sum does not close up, since its inner sums run over classes of arbitrary length. Their closed forms [JL21, Thm. 2.10, (2.37)–(2.40)] restrict the Hall–Littlewood index λ , not the class μ ; their Murnaghan–Nakayama rule [JL21, Thm. 3.2] generalises a formula of Morris [Mor77]¹, its case $\ell(\lambda) = 2$. We evaluate by residues instead, and at $\ell(\mu) = 2$ the poles organise into two t -strings. The formula is explicit, with no sum over classes, and linear in the two-string specialisation G_A of g ; for general ρ , G_A is itself a finite (branching) sum, so Theorem 9.3 is not a product formula.

Example 9.6 (Two rows). For $a = 2$, only $A = B = 1$ occurs, and $G_1(w) = P_\rho(1, w; t) = w^{\rho_2} \frac{(1-tw) - w^r(w-t)}{1-w}$ with $r = \rho_1 - \rho_2 \geq 1$ ($G_1 = w^{\rho_2}$ if $r = 0$). Substituting into Theorem 9.3 gives, for $\lambda = (\lambda_1, \lambda_2)$, $y \leq x$, and $m_{xy} = 2$ if $x = y$, 1 otherwise,

$$X_{(x,y)}^{(\lambda_1, \lambda_2)}(t) = \begin{cases} (t-1)t^{\lambda_2-1-y}(1+t^y), & y < \lambda_2, \\ m_{xy} - (1-t)t^{\lambda_2-1}, & y = \lambda_2, \\ (t-1)t^{\lambda_2-1}, & y > \lambda_2, \end{cases}$$

in agreement with a Kostka–Foulkes computation for $|\lambda| \leq 10$. This is not new: for $\lambda = (n-k, k)$ and every class ρ , Jing and Liu [JL21, §2, display after (2.32)] give $X_\rho^\lambda = \pi_k + (t-1) \sum_{j < k} t^{k-1-j} \pi_j$, with π_j the number of j -subsets fixed by ρ (their $D^{(j)}(\rho)$). A short proof: $X_\rho^\lambda = \sum_\nu \chi_\rho^\nu K_{\nu\lambda}(t)$ [Mac95, III (7.6')], where $K_{(n-j,j),\lambda} = t^{k-j}$ (one tableau, and K is monic of degree $n(\lambda) - n(\nu)$ [Mac95, III (6.5)]) and $\chi_\rho^{(n-j,j)} = \pi_j - \pi_{j-1}$ by Young's rule. On the diagonal class $(x, y) = \lambda$ it gives $t^{\lambda_2} - t^{\lambda_2-1} + m_{xy}$, which for $\lambda_1 > \lambda_2$ with $\lambda_2 \geq 3$ odd equals -1 at $t = -1$ and 1 at $t = 0$. A root in $(-1, 0)$ rules out a product of cyclotomic factors and powers of t , and Green polynomials at two-part classes admit no product formula even for two rows; for products of the form $t^c \prod_i (1 - t^{d_i})$ this was observed independently in [Cli26, Thm. D].

Theorem 9.7 (Closed $\ell = 3$, $\kappa = 1$ leads). *Let $\lambda = (a, b, c)$ be any ordering of a length-3 partition, $\mu = (x, y)$ with $\kappa(\lambda, \mu) = 1$, $n = x + y$, m_{xy} as in Example 9.6. Put*

$$\Xi_a(r, q) := \text{lin}_e T_a(p_r p_q) = (-1)^{r+q} \frac{[a+r+q]}{[a]} [a]_{t^r} [a]_{t^q}$$

¹We have not seen [Mor77] first-hand; its scope is as described in [JL21, p. 12, before Thm. 3.2].

and $U_a(b, c; x, y) := ((-1)^n \Phi_a(p_b p_c; x, y) - \Xi_a(b, c)) / m_{xy}$. Then

$$\begin{aligned} [(s-1)^2] c_{\lambda\mu} = & (-1)^{b+c} U_a(b, c; x, y) + \sum_{\substack{1 \leq r < b \\ \{b-r, a+r+c\} = \{x, y\}}} (-1)^{r+c} \Xi_a(r, c) \\ & + \sum_{\substack{1 \leq q < c \\ \{c-q, a+b+q\} = \{x, y\}}} (-1)^{b+q} \Xi_a(b, q) + [e_x e_y] D_a(M_{bc}), \end{aligned}$$

with D_a the derivation $D_a(e_j) = M_{aj}$ of Proposition 6.2.

Lemma 9.8. For G homogeneous of degree $n = x + y$, $\langle G, p_x p_y \rangle = (-1)^n (m_{xy} [e_x e_y] G + \text{line}_e G)$.

Proof. $\langle e_\nu, p_x p_y \rangle = \langle \Delta e_\nu, p_x \otimes p_y \rangle$ since p_x, p_y are primitive, and $\langle e_m, p_m \rangle = (-1)^{m-1}$. Each tensor factor must be a single e , so $\ell(\nu) \geq 3$ gives 0, $\langle e_n, p_x p_y \rangle = (-1)^n$, $\langle e_x e_y, p_x p_y \rangle = (-1)^n m_{xy}$, and other two-part ν give 0. $\square$

Proof of Theorem 9.7. By Proposition 9.1, $[(s-1)^2] c_{\lambda\mu} = [e_x e_y] (\Gamma_a(e_b, e_c) + D_a(M_{bc}))$. In $\Gamma_a(e_b, e_c) = \sum_{r,q} (-1)^{r+q} e_{b-r} e_{c-q} T_a(p_r p_q)$, terms with $r < b$ and $q < c$ have at least three factors in I and do not contribute. For $r < b$, $q = c$, only the single- e part $\Xi_a(r, c) e_{a+r+c}$ of $T_a(p_r p_c)$ (Lemma 6.11) contributes; symmetrically for $r = b$, $q < c$. For $r = b$, $q = c$ the contribution is $(-1)^{b+c} [e_x e_y] T_a(p_b p_c) = (-1)^{b+c} U_a(b, c; x, y)$ by Lemma 9.8 and $\text{line}_e T_a(p_b p_c) = \Xi_a(b, c)$. $\square$

Corollary 9.9. Every lead at valuation 2 ($\kappa = \ell - 2$, any ℓ) has a closed formula: by Theorem 7.8 it is a sum of products of pair leads (Proposition 6.2) and $\ell = 3$, $\kappa = 1$ leads (Theorem 7.4, Corollary 8.3, Theorem 9.7).

Example 9.10. $(3, 3, 3) \rightarrow (7, 2)$: $\text{Lead} = 2t^{13} + 3t^{12} + 3t^{11} + 6t^{10} + 6t^9 + 6t^8 + 9t^7 + 6t^6 + 3t^5 + 9t^4 + 6t^3 + 3t + 4$ (agrees with direct computation). Leads are *not* positive in general: $(4, 4, 2) \rightarrow (7, 3)$ has $\text{Lead} = (t+1)(t^2+1)(2t^8 + t^7 + t^6 + 3t^4 + t^3 + t^2 - t + 2)$.

Computer checks. All statements in this section were checked against an independent implementation of Hikita's subset formula: Theorem 9.7 on 16 pairs with $n \leq 12$ (both examples among them) and on 27 pairs in all 3 orderings with $n \leq 10$; Theorem 9.3 in 229 cases ($a \leq 5$); and the Green-polynomial dictionary symbolically for $|\lambda| \leq 8$ and numerically for $|\lambda| \leq 9$.

10. OPEN PROBLEMS

- (1) **Valuation ≥ 3 .** Connected leads with $\ell \geq 4$, $\kappa = 1$. Hierarchy: $v = 1 \leftrightarrow p_n \leftrightarrow$ one t -string (Lemma 6.11); $v = 2 \leftrightarrow p_x p_y \leftrightarrow$ two strings (Theorem 9.3). Does the three-string shuffle sum factor pairwise?
- (2) **Zero set at generic t .** At $t = -2$ seven pairs with $n = 6$ have $v > \ell - \kappa$; for coarsenings this is cancellation between histories, e.g. $\text{Lead}_{(1,1,1),(3)} = [3]_t(2+t)$. Describe the zero set of $[(s-1)^{\ell-\kappa}] c_{\lambda\mu}$ in t .
- (3) **Counting.** A formula for $N(\lambda, \mu)$ in Theorem 6.6; for $\ell(\lambda) = 3$ and $\mu = (n-1, 1)$ with $\kappa = 1$ we observe $\text{Lead}_{\lambda\mu}(1) = 2n^2 - 6n + 3$, independent of λ ($n \leq 10$; computed only).
- (4) **Positivity.** Leads are not t -positive (Example 9.10), so no “ t -count of minimal flows” can hold. Is there a signed or cancellation-free model?
- (5) **Cumulant link.** Is there a Macdonald-cumulant construction whose leads are the $\star$ -leads for all t , i.e. cumulants of $e_k \mapsto t^{\binom{k}{2}}$?

REFERENCES

- [BB05] Anders Björner and Francesco Brenti. *Combinatorics of Coxeter Groups*. Vol. 231. Graduate Texts in Mathematics. Springer, 2005.
- [Cli26] Clio (AI agent). “Two-part Green polynomials”. Manuscript. 2026.
- [DFK18] Philippe Di Francesco and Rinat Kedem. “Difference equations for graded characters from quantum cluster algebra”. In: *Transform. Groups* 23.2 (2018), pp. 391–424. arXiv: [1505.01657](#).
- [DFK19] Philippe Di Francesco and Rinat Kedem. “ (t, q) -deformed Q -systems, DAHA and quantum toroidal algebras via generalized Macdonald operators”. In: *Comm. Math. Phys.* 369.3 (2019), pp. 867–928. DOI: [10.1007/s00220-019-03472-x](#). arXiv: [1704.00154](#).
- [DLT94] Jacques Désarménien, Bernard Leclerc, and Jean-Yves Thibon. “Hall–Littlewood functions and Kostka–Foulkes polynomials in representation theory”. In: *Sém. Lothar. Combin.* 32 (1994). Art. B32c, 38 pp.
- [Doł19] Maciej Dołęga. “Macdonald cumulants, G -inversion polynomials and G -parking functions”. In: *European J. Combin.* 75 (2019), pp. 172–194. DOI: [10.1016/j.ejc.2018.08.011](#). arXiv: [1707.02656](#).
- [Gre55] J. A. Green. “The characters of the finite general linear groups”. In: *Trans. Amer. Math. Soc.* 80 (1955), pp. 402–447.
- [GS96] Ira M. Gessel and Bruce E. Sagan. “The Tutte polynomial of a graph, depth-first search, and simplicial complex partitions”. In: *Electron. J. Combin.* 3.2 (1996). Research Paper 9.
- [Hik25] Tatsuyuki Hikita. *(q, t) -Chromatic Symmetric Functions*. 2025. arXiv: [2503.23597](#).
- [HT26] Jim Haglund and Vasu Tewari. *Theta operators at $t = 1$, Macdonald cumulants, and LLT positivity*. 2026. arXiv: [2609.29957](#).
- [Hum90] James E. Humphreys. *Reflection Groups and Coxeter Groups*. Vol. 29. Cambridge Studies in Advanced Mathematics. Cambridge University Press, 1990.
- [Jin91] Naihuan Jing. “Vertex operators and Hall–Littlewood symmetric functions”. In: *Adv. Math.* 87 (1991), pp. 226–248.
- [JL21] Naihuan Jing and Ning Liu. *The Green polynomials via vertex operators*. 2021. arXiv: [2104.04411](#).
- [Kir98] Anatol N. Kirillov. *New combinatorial formula for modified Hall–Littlewood polynomials*. 1998. arXiv: [math/9803006](#).
- [Mac95] I. G. Macdonald. *Symmetric Functions and Hall Polynomials*. 2nd ed. Oxford University Press, 1995.
- [Mor77] Alun O. Morris. “A survey on Hall–Littlewood functions and their applications to representation theory”. In: *Combinatoire et représentation du groupe symétrique (Strasbourg, 1976)*. Vol. 579. Lecture Notes in Math. Springer, 1977, pp. 136–154. DOI: [10.1007/BFb0090015](#).
- [MR68] C. L. Mallows and John Riordan. “The inversion enumerator for labeled trees”. In: *Bull. Amer. Math. Soc.* 74 (1968), pp. 92–94.
- [Pen67] Oliver Penrose. “Convergence of fugacity expansions for classical systems”. In: *Statistical Mechanics: Foundations and Applications*. Ed. by T. A. Bak. New York–Amsterdam: W. A. Benjamin, 1967, pp. 101–109.