

























normally consume, such as copepods and phytoplankton, and could have led to the observation of a higher average prey weight per fish. We are uncertain what effect these diet shifts may have on the growth and survival of these fish populations in the long-term.

Gelatinous zooplankton are generally of poorer nutritional quality than other likely prey (Doyle et al. 2007), where a reliance on these prey may lead to reduced growth and reproductive output. However, laboratory studies have indicated that pelagic species such as Atlantic mackerel *Scomber scombrus* will feed on small medusae even when offered copepods, and can capture these very efficiently (close to 100% within the swimming path) due to their reduced avoidance capability (Runge et al. 1987). Moreover, these authors found that a single medusa has the same energy content as 10 small copepods due to their substantially larger size. A more direct measure of their condition would be to examine the energy density and lipid dynamics of these forage species among the different environmental conditions (Rosa et al. 2010, Heintz et al. 2013, Albo-Puigserver et al. 2017). Previous analyses in the NCC demonstrated that northern anchovy, Pacific sardine and Pacific herring all had higher lipid levels in the early summer of a normal upwelling year (2006) compared to 2005, a year of delayed upwelling and reduced productivity (Litz et al. 2010).

Overall, we observed major dietary shifts in a suite of planktivorous forage fishes in the northern California Current related to anomalously warm ocean conditions occurring during 2 recent years. In particular, these forage fishes shifted from having an interannually variable diet consisting mostly of crustacean prey to a more consistent diet with important contributions from gelatinous prey sources, with potentially important ecological and energetic implications. We suggest that it is imperative to continue monitoring the trophic ecology of these species, given the potential for increasing jellyfish blooms in the world's oceans (Richardson et al. 2009, Condon et al. 2013). With projections of increased extreme warming events and changes in productivity due to climate change forecast for the northeast Pacific Ocean (Timmermann et al. 2009, Cai et al. 2015), we may expect future range shifts of prey taxa including gelatinous zooplankton and alterations to food web components that depend upon them (Ruzicka et al. 2012, Albouy et al. 2014).

It is uncertain to what degree the Northern California Current and other eastern boundary upwelling ecosystems will be affected by increasing ocean tem-

peratures, but the expectation is that conditions will be substantially altered and probably less favorable for pelagic fish production in the coming decades (Doney et al. 2012, Bakun et al. 2015, García-Reyes et al. 2015). A better understanding of how environmental conditions affect prey availability, prey quality and the bioenergetics of forage fishes will help provide a clearer picture of the potential impacts of rising ocean temperatures on marine food web dynamics in the Northern California Current ecosystem and elsewhere.

**Acknowledgements.** We thank the staff of the Estuarine and Ocean Ecology Program from NOAA and OSU for assistance in conducting the surveys and processing the fish. We particularly thank Drew Hill and Elizabeth Daly for their help with stomach analyses and Yi Gong, Ashley Yarbrough, and Morgan Kroeger for assistance in measuring and weighing fish in the laboratory. Drew Hill provided the R scripts to run the ordination analyses. Elizabeth Daly, Brian Wells, and the anonymous journal reviewers provided helpful comments on an earlier version of the manuscript. This work was funded by the Northwest Fisheries Science Center (Internal Grant Program) and Bonneville Power Administration. This paper is contribution number 2018\_1 from the NOAA Integrated Ecosystem Assessment Program.

#### LITERATURE CITED

- Albo-Puigserver M, Muñoz A, Navarro J, Coll M, Pethybridge H, Sánchez S, Palomera I (2017) Ecological energetics of forage fish from the Mediterranean Sea: seasonal dynamics and interspecific differences. *Deep Sea Res II* 140:74–82
- Albouy C, Velez L, Coll M, Colloca F, Le Loc'h F, Mouillot D, Gravel D (2014) From projected species distribution to food-web structure under climate change. *Glob Change Biol* 20:730–741
- Alegre A, Bertrand A, Espino M, Espinoza P and others (2015) Diet diversity of jack and chub mackerels and ecosystem changes in the northern Humboldt Current system: a long-term study. *Prog Oceanogr* 137:299–313
- Arai MN, Welch DW, Dunsmuir AL, Jacobs MC, Ladouceur AR (2003) Digestion of pelagic Ctenophora and Cnidaria by fish. *Can J Fish Aquat Sci* 60:825–829
- Auth TD, Daly EA, Brodeur RD, Fisher JL (2018) Phenological and distributional shifts in ichthyoplankton associated with recent marine warming in the northeast Pacific Ocean. *Glob Change Biol* 24:259–272
- Bachiller E, Irigoien X (2015) Trophodynamics and diet overlap of small pelagic fish species in the Bay of Biscay. *Mar Ecol Prog Ser* 534:179–198
- Bakun A, Black BA, Bograd SJ, García-Reyes M, Miller AJ, Rykaczewski RR, Sydeman WJ (2015) Anticipated effects of climate change on coastal upwelling ecosystems. *Curr Clim Change Rep* 1:85–93
- Baumgartner TR, Soutar A, Ferreira-Bartrina V (1992) Reconstruction of the history of Pacific Sardine and northern anchovy populations over the past two millennia from sediments of the Santa Barbara Basin, California. *CCOFI Rep* 33:24–40

- Bond NA, Cronin MF, Freeland H, Mantua N (2015) Causes and impacts of the 2014 warm anomaly in the NE Pacific. *Geophys Res Lett* 42:3414–3420
- Brodeur RD, Pearcy WG (1992) Effects of environmental variability on trophic interactions and food web structure in a pelagic upwelling ecosystem. *Mar Ecol Prog Ser* 84: 101–119
- Brodeur RD, Lorz HV, Pearcy WG (1987) Food habits and dietary variability of pelagic nekton off Oregon and Washington, 1979–1984. NOAA Tech Rep NMFS 57, US Department of Commerce, Springfield, VA
- Brodeur RD, Fisher JP, Emmett RL, Morgan CA, Casillas E (2005) Species composition and community structure of pelagic nekton off Oregon and Washington under variable oceanographic conditions. *Mar Ecol Prog Ser* 298:41–57
- Brodeur RD, Buchanan JC, Emmett RL (2014) Pelagic and demersal fish predators on juvenile and adult forage fishes in the northern California Current: spatial and temporal variations. *CCOFI Rep* 55:96–116
- Brodeur RD, Perry RI, Boldt JL, Flostrand L and others (2018) An unusual gelatinous plankton event in the NE Pacific: the great pyrosome bloom of 2017. *PICES Press* 26:22–27
- Brosset P, Le Bourg B, Costalago D, Banaru D and others (2016) Linking small pelagic dietary shifts with ecosystem changes in the Gulf of Lions. *Mar Ecol Prog Ser* 554:157–171
- Cai W, Wang G, Santoso A, McPhaden MJ and others (2015) Increased frequency of extreme La Nina events under greenhouse warming. *Nat Clim Chang* 5:132–137
- Cardona L, De Quevedo IÁ, Borrell A, Aguilar A (2012) Massive consumption of gelatinous plankton by Mediterranean apex predators. *PLOS ONE* 7:e31329
- Cavole LM, Demko AM, Diner RE, Giddings A and others (2016) Biological impacts of the 2013–2015 warm-water anomaly in the Northeast Pacific: winners, losers, and the future. *Oceanography (Wash DC)* 29:273–285
- Chavez FP, Ryan J, Lluch-Cota SE, Niquen M (2003) From anchovies to sardines and back: multidecadal change in the Pacific Ocean. *Science* 299:217–221
- Checkley DM Jr, Barth JA (2009) Patterns and processes in the California Current System. *Prog Oceanogr* 83:49–64
- Clarke KR, Gorley RN (2015) *PRIMER v7: user manual/tutorial*. PRIMER-E: Plymouth
- Clarke RK, Somerfield PJ, Gorley RN (2008) Testing of null hypotheses in exploratory community analyses: similarity profiles and biota-environment linkage. *J Exp Mar Biol Ecol* 366:56–69
- Condon RH, Duarte CM, Pitt KA, Robinson KL and others (2013) Recurrent jellyfish blooms are a consequence of global oscillations. *Proc Natl Acad Sci USA* 110: 1000–1005
- Cury PM, Shannon LJ, Roux JP, Daskalov GM, Jarre A, Moloney CL, Pauly D (2005) Trophodynamic indicators for an ecosystem approach to fisheries. *ICES J Mar Sci* 62:430–442
- Cury PM, Boyd IL, Bonhommeau S, Anker-Nilssen T and others (2011) Global seabird response to forage fish depletion—one-third for the birds. *Science* 334:1703–1706
- Daly EA, Brodeur RD (2015) Warming ocean conditions relate to increased trophic requirements of threatened and endangered salmon. *PLOS ONE* 10:e0144066
- Davis ND, Myers KW, Ishida Y (1998) Caloric values of high-seas salmon prey organisms and simulated salmon ocean growth and prey consumption. *N Pac Anadromous Fish Comm Bull* 1:146–162
- Di Lorenzo E, Mantua N (2016) Multi-year persistence of the 2014/15 North Pacific marine heatwave. *Nat Clim Chang* 6:1042–1047
- Di Lorenzo E, Schneider N, Cobb KM, Chhak K and others (2008) North Pacific Gyre Oscillation links ocean climate and ecosystem change. *Geophys Res Lett* 35:L08607
- Doney SC, Ruckelshaus M, Duffy JE, Barry JP and others (2012) Climate change impacts on marine ecosystems. *Annu Rev Mar Sci* 4:11–37
- Doyle TK, Houghton JDR, McDevitt R, Davenport J, Hays GC (2007) The energy density of jellyfish: estimates from bomb-calorimetry and proximate composition. *J Exp Mar Biol Ecol* 343:239–252
- Dufrene M, Legendre P (1997) Species assemblages and indicator species: the need for a flexible asymmetrical approach. *Ecol Monogr* 67:345–366
- Engelhard GH, Peck MA, Rindorf A, Smout SC and others (2014) Forage fish, their fisheries, and their predators: who drives whom? *ICES J Mar Sci* 71:90–104
- Espinoza P, Bertrand A, van der Lingen CD, Garrido S, de Mendiola BR (2009) Diet of sardine (*Sardinops sagax*) in the northern Humboldt Current system and comparison with the diets of clupeoids in this and other eastern boundary upwelling systems. *Prog Oceanogr* 83:242–250
- Field DB, Baumgartner TR, Ferreira V, Gutierrez D, Lozano-Montes H, Salvatelli R, Soutar A (2009) Variability from scales in marine sediments and other historical records. In: Checkley J, Alheit J, Oozeki Y, Roy C (eds) *Climate change and small pelagic fish*. Cambridge University Press, Cambridge, p 45–63
- Fisher JL, Peterson WT, Rykaczewski RR (2015) The impact of El Niño events on the pelagic food chain in the northern California Current. *Glob Change Biol* 21:4401–4414
- Francis TB, Scheuerell MD, Brodeur RD, Levin PS, Ruzicka JJ, Tolimieri N, Peterson WT (2012) Climate shifts the interaction web of a marine plankton community. *Glob Change Biol* 18:2498–2508
- Galbraith M, Young K (2017) Zooplankton along the B.C. continental margin. In: Chandler PC, King SA, Boldt J (eds) *State of the physical, biological and selected fishery resources of Pacific Canadian marine ecosystems in 2016*. Can Tech Rep Fish Aquat Sci 3225:67–75
- García-Reyes M, Sydeman WJ, Schoeman DS, Rykaczewski RR, Black BA, Smit AJ, Bograd SJ (2015) Under pressure: climate change, upwelling, and eastern boundary upwelling ecosystems. *Front Mar Sci* 2:109
- Garrido S, van der Lingen CD (2014) Feeding biology and ecology. In: Ganas K (ed) *Biology and ecology of sardines and anchovies*. CRC Press, Boca Raton, FL, p 122–189
- Garrido S, Silva A, Pastor J, Dominguez R, Silva AV, Santos AM (2015) Trophic ecology of pelagic fish species off the Iberian coast: diet overlap, cannibalism and intraguild predation. *Mar Ecol Prog Ser* 539:271–286
- Heintz RA, Siddon EC, Farley EV Jr, Napp JM (2013) Correlation between recruitment and fall condition of age-0 pollock (*Theragra chalcogramma*) from the eastern Bering Sea under varying climate conditions. *Deep Sea Res II* 94:150–156
- Hilborn R, Amoroso RO, Bogazzia E, Jensen OP, Parma AP, Szuwalski C, Walters CJ (2017) When does fishing forage species affect their predators? *Fish Res* 191:211–221
- Hill AD, Daly EA, Brodeur RD (2015) Diet variability of forage fishes in the Northern California Current System. *J Mar Syst* 146:121–130

- Jacox MG, Hazen EL, Zaba KD, Rudnick DL, Edwards CA, Moore AM, Bograd SJ (2016) Impacts of the 2015–2016 El Niño on the California Current System: early assessment and comparison to past events. *Geophys Res Lett* 43:7072–7080
- Kaplan IC, Brown CJ, Fulton EA, Gray IA, Field JC, Smith AD (2013) Impacts of depleting forage species in the California Current. *Environ Conserv* 40:380–393
- Koehn LE, Essington TE, Marshall KN, Kaplan IC, Sydeman WJ, Szoboszlai AI, Thayer JA (2016) Developing a high taxonomic resolution food web model to assess the functional role of forage fish in the California Current ecosystem. *Ecol Modell* 335:87–100
- Koehn LE, Essington TE, Marshall KN, Sydeman WJ, Szoboszlai AI, Thayer JA (2017) Trade-offs between forage fish fisheries and their predators in the California Current. *ICES J Mar Sci* 74:2448–2458
- Lee YW, Sampson DB (2009) Dietary variations in three co-occurring rockfish species off the Pacific Northwest during anomalous oceanographic events in 1998 and 1999. *Fish Bull* 107:510–522
- Leising A, Schroeder ID, Bograd SJ, Abell J and others (2015) State of the California Current 2014–15: impacts of the warm-water ‘Blob’. *CCOFI Rep* 56:31–68
- Link JS, Ford MD (2006) Widespread and persistent increase of Ctenophora in the continental shelf ecosystem off NE USA. *Mar Ecol Prog Ser* 320:153–159
- Litz MC, Brodeur RD, Emmett RL, Heppell SS, Rasmussen RS, O’Higgins L, Morris MS (2010) Effects of variable oceanographic conditions on forage fish lipid content and fatty acid composition in the northern California Current. *Mar Ecol Prog Ser* 405:71–85
- Litz MNC, Emmett RL, Bentley PJ, Claiborne AM, Barceló C (2014) Biotic and abiotic factors influencing forage fish and pelagic nekton community in the Columbia River plume (USA) throughout the upwelling season 1999–2009. *ICES J Mar Sci* 71:5–18
- Mantua NJ, Hare SR, Zhang Y, Wallace JM, Francis RC (1997) A Pacific interdecadal climate oscillation with impacts on salmon production. *Bull Am Meteorol Soc* 78:1069–1079
- McClatchie S, Goericke R, Leising A, Auth T, and others (2016) State of the California Current 2015–16: comparisons with the 1997–98 El Niño. *CCOFI Rep* 57:1–57
- McCune B, Grace JB (2002) Analysis of ecological communities. MjM Software Design, Gleneden Beach, OR
- McCune B, Mefford MJ (1999) PC-ORD, multivariate analysis of ecological data, users guide. MjM Software Design, Gleneden Beach, OR
- Mianzan H, Mari N, Prenski B, Sanchez F (1996) Fish predation on neritic ctenophores from Argentine continental shelf: a neglected food source? *Fish Res* 27:69–79
- Mianzan H, Pajaro M, Colombo GA, Madirolas A (2001) Feeding on survival food: gelatinous zooplankton as a source food for anchovies. *Hydrobiologia* 451:45–53
- Milisenda G, Rosa S, Fuentes VL, Boero F, Guglielmo L, Purcell JE, Piraino S (2014) Jellyfish as prey: frequency of predation and selective foraging of *Boops boops* (Vertebrata, Actinopterygii) on the mauve stinger *Pelagia noctiluca* (Cnidaria, Scyphozoa). *PLOS ONE* 9:e94600
- Miller TW, Brodeur RD (2007) Diet of and trophic relationships among dominant marine nekton within the Northern California Current ecosystem. *Fish Bull* 105: 548–559
- Miller CB, Batchelder HP, Brodeur RD, Pearcy WG (1985) Response of the zooplankton and ichthyoplankton off Oregon to the El Niño event of 1983. In: Wooster WS & Fluharty DL (eds) *El Niño North: Niño effects in the Eastern Subarctic Pacific Ocean*. Washington Sea Grant Program, University of Washington, p 185–187
- Miller TW, Brodeur RD, Rau G, Omori K (2010) Prey dominance shapes trophic structure of the northern California Current pelagic food web: evidence from stable isotopes and diet analysis. *Mar Ecol Prog Ser* 420:15–26
- Oksanen J, Blanchett FG, Kindt R, Legendre P and others (2012) *Vegan: community ecology package*. R Package 2.0.3. <http://CRAN.R-project.org/package=vegan>
- Peck MA, Neuenfeldt S, Essington TE, Trenkel VM and others (2014) Forage fish interactions: a symposium on ‘Creating the tools for ecosystem-based management of marine resources’. *ICES J Mar Sci* 71:1–4
- Peterson WT, Keister JE, Feinberg LR (2002) The effects of the 1997–98 El Niño/La Niña events on hydrography and zooplankton off the central Oregon coast. *Prog Oceanogr* 54:381–398
- Peterson WT, Fisher JL, Strub T, Du X, Risien C, Peterson J, Shaw CT (2017) The pelagic ecosystem in the northern California Current off Oregon during the 2014–2016 warm anomalies within the context of the past 20 years. *J Geophys Res* 122:7267–7290
- Pikitch E, Boersma PD, Boyd IL, Conover DO and others (2012) Little fish, big impact: managing a crucial link in ocean food webs. Lenfest Ocean Program, Washington, DC
- Purcell JE, Arai MN (2001) Interactions of pelagic cnidarians and ctenophores with fish: a review. *Hydrobiologia* 451:27–44
- R Development Core Team (2014) *R: a language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna
- Richardson AJ, Bakun A, Hays GC, Gibbons MJ (2009) The jellyfish joyride: causes, consequences and management responses to a more gelatinous future. *Trends Ecol Evol* 24:312–322
- Rosa R, Gonzalez L, Broitman BR, Garrido S, Santos AMP, Nunes ML (2010) Bioenergetics of small pelagic fishes in upwelling systems: relationship between fish condition, coastal ecosystem dynamics and fisheries. *Mar Ecol Prog Ser* 410:205–218
- Runge JA, Pepin P, Silvert W (1987) Feeding behavior of the Atlantic mackerel *Scomber scombrus* on the hydro-medusae *Aglantha digitale*. *Mar Biol* 94:329–333
- Ruzicka JJ, Brodeur RD, Emmett RL, Steele JH and others (2012) Interannual variability in the Northern California Current food web structure changes in energy flow pathways and the role of forage fish, euphausiids, and jellyfish. *Prog Oceanogr* 102:19–41
- Sakuma KM, Field JC, Mantua NJ, Ralston S, Marinovic BB, Carrion CN (2016) Anomalous epipelagic micronekton assemblage patterns in the neritic waters of the California Current in Spring 2015 during a period of extreme ocean conditions. *CCOFI Rep* 57:163–183
- Schwartzlose RA, Alheit J, Bakun A, Baumgartner TR and others (1999) Worldwide large-scale fluctuations of sardine and anchovy populations. *S Afr J Mar Sci* 21: 289–347
- Sydeman WJ, Santora JA, Thompson SA, Marinovic B, Di Lorenzo E (2013) Increasing variance in North Pacific climate relates to unprecedented ecosystem variability off California. *Glob Change Biol* 19:1662–1675

- Szoboszlai AI, Thayer JA, Wood SA, Sydeman WJ, Koehn LE (2015) Forage species in predator diets: synthesis of data from the California Current. *Ecol Inform* 29:45–56
- Timmermann A, Oberhuber J, Bacher A, Esch M, Latif M, Roeckner E (1999) Increased El Niño frequency in a climate model forced by future greenhouse warming. *Nature* 398:694–696
- van der Lingen CD (2002) Diet of sardine *Sardinops sagax* in the southern Benguela upwelling ecosystem. *S Afr J Mar Sci* 24:301–316
- van der Lingen CD, Bertrand A, Bode A, Brodeur RD and others (2009) Trophic dynamics. In: Checkley J, Alheit J, Oozeki Y, Roy C (eds) *Climate change and small pelagic fish*. Cambridge University Press, Cambridge, p 112–157
- Walters C, Christensen V, Fulton B, Smith ADM, Hilborn R (2016) Predictions from simple predator-prey theory about impacts of harvesting forage fishes. *Ecol Modell* 337:272–280
- Ware DM, Thomson RE (2005) Bottom-up ecosystem trophic dynamics determine fish production in the northeast Pacific. *Science* 308:1280–1284
- Wood S (2012) Package ‘mgcv’. R Package 2.0.3
- Yamamura O, Honda S, Shida O, Hamatsu T (2002) Diets of walleye pollock *Theragra chalcogramma* in the Doto area, northern Japan: ontogenetic and seasonal variations. *Mar Ecol Prog Ser* 238:187–198

*Editorial responsibility: Arnaud Bertrand (Guest Editor), Sète, France*

*Submitted: August 14, 2017; Accepted: January 23, 2018  
Proofs received from author(s): April 3, 2018*