

A candidate proof of Circular colouring the orthogonality graph

github.com/graph-theory-AI — machine-generated candidate writeup

16 September 2026

Abstract

A geometric odd-walk lemma excludes every homomorphism from $\mathcal{O}$ to a finite locally bipartite graph, and consequently proves $\chi_c(\mathcal{O})=4$. This candidate proof was generated by GPT-6 Astra in a single model pass for the Graph Theory AI project. It was selected from the campaign because the model marked it `would_publish: true`; it has not been checked by the adversarial referee, a human mathematician, or a novelty search.

1 The problem and the claimed result

This document concerns [Circular colouring the orthogonality graph](http://www.openproblemgarden.org/op/circular_colouring_the_orthogonality_graph) from http://www.openproblemgarden.org/op/circular_colouring_the_orthogonality_graph.

1.1 Problem statement

The campaign artifact does not contain a separately extracted statement; see the catalog entry linked above.

1.2 Claimed result

Model verdict: **proved** (confidence: high).

A geometric odd-walk lemma excludes every homomorphism from $\mathcal{O}$ to a finite locally bipartite graph, and consequently proves $\chi_c(\mathcal{O})=4$.

The model's own caveat: Literature priority has not been checked; the argument below is self-contained.

2 Model writeup

The text below is the model output verbatim, apart from moving its machine-readable verdict block into the summary above and shifting Markdown heading levels for this document. Mechanical TeX defects and equation tags were normalized where needed for compilation.

2.1 Circular chromatic number of the orthogonality graph

Let $\mathcal{O}$ be the graph whose vertices are the one-dimensional subspaces of $\mathbb{R}^3$, with adjacency given by orthogonality.

We prove the following stronger statement.

Theorem. There is no homomorphism from $\mathcal{O}$ to a finite graph in which every open neighborhood induces a bipartite graph. Consequently,

$$\chi_c(\mathcal{O}) = 4.$$

The key geometric fact is that the union of the neighborhoods of three linearly independent directions always contains an odd closed walk.

2.1.1 1. The geometric odd-walk lemma

For a nonzero vector a , write

$$P_a = a^\perp, \quad J_a(x) = a \times x.$$

Thus $J_a(x)$, when nonzero, represents a line perpendicular to $[x]$ and belonging to P_a .

Lemma. Let $a, b, c \in \mathbb{R}^3$ be linearly independent. The subgraph of $\mathcal{O}$ induced by all lines contained in

$$P_a \cup P_b \cup P_c$$

is not bipartite.

Proof. First suppose that two of the normals, say a, b , are perpendicular. Then

$$[a], \quad [b], \quad [a \times b]$$

form a triangle, and all three belong to the specified union. This covers that case.

We may therefore assume that a, b, c are pairwise nonorthogonal. Normalize them to unit vectors and choose their signs and an orthonormal coordinate system so that

$$a = (0, 0, 1), \quad b = (s, 0, t), \quad c = (u, v, w),$$

where

$$s > 0, \quad 0 < t < 1, \quad s^2 + t^2 = 1.$$

Linear independence and pairwise nonorthogonality give

$$v \neq 0, \quad w \neq 0, \quad su + tw \neq 0. \quad (1)$$

We will construct walks using successive cross products. All the cross products involved are nonzero: whenever $d \cdot e \neq 0$, the restriction

$$J_d : P_e \longrightarrow P_d$$

is injective, because the kernel of J_d is $\mathbb{R}d$, which does not meet P_e nontrivially. This applies to every transition below.

Identify P_a with $\mathbb{R}^2$ using $(x, y, 0) \leftrightarrow (x, y)$. Consider the two operators on P_a

$$E = J_a^2 J_b^2, \quad F = J_a J_c J_b.$$

Direct calculation gives

$$E = \begin{pmatrix} t^2 & 0 \\ 0 & 1 \end{pmatrix}, \quad F = \begin{pmatrix} 0 & su + tw \\ -tw & sv \end{pmatrix}. \quad (2)$$

These operators have a walk interpretation:

- F is a three-edge walk, with successive planes

$$P_a \xrightarrow{J_b} P_b \xrightarrow{J_c} P_c \xrightarrow{J_a} P_a.$$

- E is a four-edge walk, with successive planes

$$P_a \xrightarrow{J_b} P_b \xrightarrow{J_b} P_b \xrightarrow{J_a} P_a \xrightarrow{J_a} P_a.$$

For $n \geq 0$,

$$E^n F = \begin{pmatrix} 0 & t^{2n}(su + tw) \\ -tw & sv \end{pmatrix}.$$

Its characteristic polynomial has discriminant

$$\Delta_n = (sv)^2 - 4t^{2n}tw(su + tw). \quad (3)$$

Because $0 < t < 1$ and $sv \neq 0$,

$$\Delta_n \longrightarrow (sv)^2 > 0.$$

Choose $n \geq 0$ with $\Delta_n > 0$. Then $E^n F$ has a real eigenvector $x_0 \neq 0$. Moreover, (1) shows that $E^n F$ is invertible, so its corresponding eigenvalue λ is nonzero:

$$E^n F(x_0) = \lambda x_0.$$

Start at the line $[x_0]$, perform the three cross-product steps defining F , and then perform n copies of the four steps defining E . Every step joins perpendicular lines, every intermediate vector is nonzero, and every line lies in $P_a \cup P_b \cup P_c$. The final line is

$$[E^n F(x_0)] = [\lambda x_0] = [x_0].$$

We have therefore obtained a closed walk of odd length

$$3 + 4n.$$

A bipartite graph cannot have an odd closed walk. This proves the lemma. $\square$

A useful reformulation is the following. For $S \subseteq V(\mathcal{O})$, let

$$N_{\mathcal{O}}(S) = \bigcup_{L \in S} N_{\mathcal{O}}(L).$$

Corollary. If $\mathcal{O}[N_{\mathcal{O}}(S)]$ is bipartite, then the lines in S span a vector subspace of dimension at most 2.

Indeed, otherwise S contains three linearly independent directions, and the lemma supplies an odd closed walk inside $N_{\mathcal{O}}(S)$.

2.1.2 2. No finite locally bipartite target

Call a graph *locally bipartite* if every open neighborhood induces a bipartite graph.

Suppose, for a contradiction, that

$$h : \mathcal{O} \longrightarrow H$$

is a homomorphism to a finite locally bipartite graph H .

For any $\alpha \in V(H)$, put

$$S_{\alpha} = h^{-1}(\alpha).$$

Every vertex of $N_{\mathcal{O}}(S_{\alpha})$ is mapped into $N_H(\alpha)$. Consequently,

$$\mathcal{O}[N_{\mathcal{O}}(S_{\alpha})] \longrightarrow H[N_H(\alpha)].$$

The graph on the right is bipartite, so the graph on the left is bipartite as well. By the corollary, every fiber S_{α} spans a subspace of dimension at most 2.

These finitely many fibers cannot cover all projective directions. Here is an explicit pigeonhole argument. If $m = |V(H)|$, consider the $2m + 1$ lines

$$L_j = [(1, j, j^2)], \quad 1 \leq j \leq 2m + 1.$$

Some three have the same image under h . But any three distinct such lines are linearly independent, since

$$\det \begin{pmatrix} 1 & i & i^2 \\ 1 & j & j^2 \\ 1 & k & k^2 \end{pmatrix} = (j - i)(k - i)(k - j) \neq 0.$$

This contradicts the dimension bound on that fiber.

Thus $\mathcal{O}$ has no homomorphism to a finite locally bipartite graph.

2.1.3 3. Circular colorings below 4 have locally bipartite targets

For positive integers p, q with $p \geq 2q$, let $K_{p/q}$ be the circular clique on $\mathbb{Z}_p$, with adjacency defined by

$$a \sim b \iff q \leq (b - a \bmod p) \leq p - q.$$

Lemma. If $p < 4q$, then $K_{p/q}$ is locally bipartite.

Proof. By translation symmetry, it is enough to consider the neighborhood of 0:

$$N(0) = \{q, q + 1, \dots, p - q\}.$$

Partition it into

$$A = N(0) \cap \{q, \dots, 2q - 1\}, \quad B = N(0) \cap \{2q, \dots, 3q - 1\}.$$

These sets cover $N(0)$, because $p - q < 3q$. Each lies in a block of q consecutive colors, and such a block is independent in $K_{p/q}$. Thus A, B are a bipartition of the neighborhood. $\square$

It follows from Section 2 that

$$\mathcal{O} \nrightarrow K_{p/q} \quad \text{whenever } p/q < 4.$$

Using the finite-palette definition of circular chromatic number, this already proves

$$\chi_c(\mathcal{O}) \geq 4. \quad (4)$$

For completeness, the conclusion is unchanged if circular chromatic number for infinite graphs is defined using real-circle colorings. Suppose a circular r -coloring exists for some $r < 4$. Choose a rational number

$$r < R < 4$$

and scale the coloring to circumference R . Every edge then has circular distance at least $R/r > 1$. Write $R = p/q$, choosing a sufficiently large common multiple of its numerator and denominator that

$$\frac{2}{q} < \frac{R}{r} - 1.$$

Rounding every color to the grid of spacing $1/q$ changes each color by less than $1/q$, so every edge still has circular distance at least 1. This gives a homomorphism to the finite graph $K_{p/q}$, a contradiction. The rounding estimate is uniform and therefore applies to infinitely many vertices.

2.1.4 4. A proper four-coloring, including boundary directions

Here is an explicit four-coloring to establish the upper bound without leaving boundary choices implicit.

For every line choose its unit representative $u = (x, y, z)$ according to the rule:

- choose $z > 0$, if possible;
- if $z = 0$, choose $y > 0$, if possible;
- if $z = y = 0$, choose $x > 0$.

Except for the line $[e_3]$, the planar projection (x, y) is nonzero. Color such a line by the half-open quadrant containing (x, y) , using the argument intervals

$$[0, \pi/2), \quad [\pi/2, \pi), \quad [\pi, 3\pi/2), \quad [3\pi/2, 2\pi).$$

Give $[e_3]$ the third color.

Two nonzero planar vectors in the same half-open quadrant have strictly positive inner product. Since all selected representatives satisfy $z \geq 0$, two non-pole lines of the same color have

$$u \cdot v = (x, y) \cdot (x', y') + zz' > 0.$$

Thus they are not perpendicular.

The selected representatives of equatorial lines have arguments in $[0, \pi)$, so no equatorial line has the third color. Every other line of the third color has $z > 0$ and hence is not perpendicular to e_3 . Therefore the coloring is proper, and

$$\chi_c(\mathcal{O}) \leq \chi(\mathcal{O}) \leq 4. \quad (5)$$

Combining (4) and (5) proves

$$\boxed{\chi_c(\mathcal{O}) = 4}.$$

2.1.5 5. Finite witnesses and the previous attempt

The obstruction above can also be made into a fully specified finite construction for each palette size.

Fix m . Start with the $2m + 1$ lines

$$[(1, j, j^2)], \quad 1 \leq j \leq 2m + 1.$$

For each triple, apply the geometric lemma, choosing the least $n \geq 0$ for which (3) is positive and an eigenline of the displayed matrix $E^n F$. Include all lines of its resulting odd closed walk. Let F_m be the orthogonality graph induced on the resulting finite set of lines.

Then F_m has no homomorphism to any locally bipartite graph with at most m vertices: three of the initial lines would receive the same color, and their associated odd walk would map into that color's bipartite neighborhood. This construction uses only arithmetic, square roots, and a terminating search for n .

In particular, for each fixed circular clique $K_{p/q}$ with $p/q < 4$, there is a finite obstruction. The obstruction depends on the palette size; the proof does **not** assert that one finite subgraph has circular chromatic number 4. It is therefore compatible with the previous attempt's finite-subgraph projection observation.

The proposed discrete-spectrum reduction is not needed here. The decisive additional ingredient is the odd-walk lemma, which turns bipartiteness of a color's neighborhood into a dimension restriction on its entire fiber. No unproved conjecture, regularity assumption on the coloring, or computational assertion is used.

A Provenance and review status

UNREFEREED MODEL OUTPUT. This document was selected solely because GPT-6 Astra labelled its own result `would_publish: true` and returned `proved` or `disproved`. It has not passed the adversarial LLM referee used for the earlier Sol campaign, has not been checked by a human mathematician, and has not been checked for novelty. Treat every mathematical and bibliographic claim as unverified.

Catalog id	<code>circular_colouring_the_orthogonality_graph</code>
Catalog entry	Circular colouring the orthogonality graph
Source corpus	<code>opg</code>
Campaign leg	<code>selected second attempts (attacks_retry)</code>
Source paper / entry	http://www.openproblemgarden.org/op/circular_colouring_the_orthogonality_graph
Model verdict	proved (confidence: high)
Selection criterion	<code>would_publish: true</code> and verdict <code>proved</code> or <code>disproved</code>
Model	<code>gpt-6-astra</code> , reasoning effort <code>max</code> , mode= <code>pro</code> , flex service tier
Original artifact	<code>attacks_retry/circular_colouring_the_orthogonality_graph/output.m</code>
Independent review	None
Previous attempt	<code>attacks_opg/circular_colouring_the_orthogonality_graph: gpt-6-astra verdict partial</code>
Model's caveats	Literature priority has not been checked; the argument below is self-contained.

The generated Markdown and LaTeX sources for this PDF are stored under `to_review_astra/src/circular_colouring_the`