

A candidate proof of Finding k -edge-outerplanar graph embeddings

github.com/graph-theory-AI — machine-generated candidate writeup

16 September 2026

Abstract

A polynomial-time SPQR dynamic program, using a two-bin scheduling recurrence at parallel nodes, computes a minimum edge-outerplanar embedding. This candidate proof was generated by GPT-6 Astra in a single model pass for the Graph Theory AI project. It was selected from the campaign because the model marked it `would_publish: true`; it has not been checked by the adversarial referee, a human mathematician, or a novelty search.

1 The problem and the claimed result

This document concerns [Finding \$k\$ -edge-outerplanar graph embeddings](http://www.openproblem-garden.org/op/finding_k_edge_outerplanar_graph_embeddings) from http://www.openproblem-garden.org/op/finding_k_edge_outerplanar_graph_embeddings.

1.1 Problem statement

Conjecture It has been shown that a k -outerplanar embedding for which k is minimal can be found in polynomial time. Does a similar result hold for k -edge-outerplanar graphs?

1.2 Catalog context

A k -outerplanar graph [Baker] with $k > 0$ is a planar graph having an embedding with at most k layers of vertices such that after removing iteratively the vertices (and their adjacent edges) lying on the outer face k times, we obtain the empty graph. A k -edge-outerplanar graph [Bentz] is defined to be a planar graph having an embedding with at most k layers of edges such that after removing iteratively the edges lying on the outer face k times, we obtain a graph with no edge. All k -edge-outerplanar graphs are k -outerplanar graphs. Given a planar graph, Bienstock and Monma have shown that a k -outerplanar embedding for which k is minimal can be found in polynomial time. Does a similar result hold for k -edge-outerplanar graphs?

1.3 Claimed result

Model verdict: **proved** (confidence: high).

A polynomial-time SPQR dynamic program, using a two-bin scheduling recurrence at parallel nodes, computes a minimum edge-outerplanar embedding.

The model's own caveat: Uses the standard SPQR decomposition theorem; novelty and current literature status have not been independently checked.

2 Model writeup

The text below is the model output verbatim, apart from moving its machine-readable verdict block into the summary above and shifting Markdown heading levels for this document. Mechanical TeX defects and equation tags were normalized where needed for compilation.

2.1 A polynomial algorithm for minimum edge-outerplanarity

I give an algorithmic proof. The main point is that the dual distance across a pertinent SPQR component is independent of its embedding. The remaining ordering problem at a parallel node has a polynomial dynamic program.

The proof also handles cutvertices. They require some care: one cannot simply take the maximum of the optimum values of the blocks, because an attachment vertex need not lie on the chosen outer face of its block.

2.1.1 1. Statement

Theorem. Given a finite planar graph G , one can compute, in polynomial time, a plane embedding of G minimizing the number of rounds of simultaneous deletion of all edges incident with the outer face.

The algorithm returns a combinatorial embedding and its outer face. An intentionally coarse bound for the implementation below is $O(N^8)$ operations on $O(\log N)$ -bit integers, where $N = |V(G)| + |E(G)|$.

For an edgeless graph, take the optimum to be 0; if the parameter is required to be positive, report 1 instead.

The only standard structural input is the SPQR decomposition theorem for biconnected planar graphs, including its description of all planar embeddings. No result about optimizing outerplanarity is used.

2.1.2 2. Edge peeling is an edge-distance objective in the dual

Let G be a plane graph, let $D = G^*$ be its geometric dual, and let r be the dual vertex corresponding to the outer face. Loops and parallel edges are retained in the dual.

Write

$$d(f) = \text{dist}_D(r, f).$$

For a primal edge e , let f_e, g_e be its two incident faces, possibly equal.

Lemma 1. The round in which e is deleted is

$$\text{layer}(e) = 1 + \min\{d(f_e), d(g_e)\}. \quad (1)$$

Proof. After round t , the original faces absorbed into the outer face are exactly those at dual distance at most t from r . This follows inductively: deleting the current outer-face edges absorbs precisely the neighboring original faces.

An edge is therefore exposed at the beginning of round t precisely when one of its incident original faces has distance at most $t - 1$. Its first such round is (1). This also covers a bridge, whose two incident faces are equal. $\square$

Thus the edge-outerplanarity of this particular embedding is

$$K(G, r) = \max_{e \in E(G)} (1 + \min\{d(f_e), d(g_e)\}). \quad (2)$$

In particular, the objective is **not quite the eccentricity of r in the dual**. For example, in a tetrahedral embedding of K_4 , the outer-face dual vertex has eccentricity 1, but edge peeling takes 2 rounds.

We will also use the following immediate consequence.

Deletion monotonicity. Deleting edges from a plane graph, while keeping the inherited outer face, cannot increase the deletion round of any remaining edge.

Indeed, deletion merges faces, so dual distances cannot increase.

2.1.3 3. A weighted problem needed for cutvertices

For a plane graph B , define

$$a_B(v) = \min\{d(f) : f \text{ is incident with } v\}. \quad (3)$$

This is the distance from the outer face to a nearest face incident with v .

Give the vertices nonnegative integer weights $w(v)$, and define

$$\Psi(B, w) = \max \left\{ K(B, r), \max_{v \in V(B)} (w(v) + a_B(v)) \right\}. \quad (4)$$

We first solve the following problem for a biconnected planar graph B :

- minimize $\Psi(B, w)$ over its embeddings and outer faces;
- optionally require a specified vertex c to be incident with the outer face.

The weights will eventually be optimum values of graphs attached at cutvertices.

A zero weight imposes no additional restriction: for every nonisolated vertex v , an edge bounding a nearest incident face has layer $a_B(v) + 1$, so $a_B(v) < K(B, r)$.

2.1.4 4. Rooted components and their profiles

Fix an edge of a biconnected graph and root its SPQR decomposition at that edge.

It is convenient to use the usual edge-substitution formulation. A pertinent component is a two-terminal primal graph H , with distinct poles s, t , together with an additional reference edge $\rho = st$. The augmented graph

$$\widehat{H} = H + \rho$$

is biconnected and planar.

For an embedding of $\widehat{H}$:

- let A, B be the faces on the two sides of ρ ;
- take the dual of $\widehat{H}$;
- delete the dual edge ρ^* .

The resulting connected two-terminal dual graph is denoted D_H , with terminals A, B . It has one edge for each real edge of H .

2.1.4.1 4.1 The terminal distance is embedding-invariant Set

$$\ell(H) = \text{dist}_{D_H}(A, B).$$

Lemma 2. The integer $\ell(H)$ is independent of the embedding choices in the rooted SPQR subtree. Moreover,

$$1 \leq \ell(H) \leq |E(H)|. \quad (5)$$

Proof. Use induction through the SPQR decomposition.

A real-edge leaf gives a single dual edge, so its value is 1.

At an internal node, replacing a primal skeleton edge by a child component replaces the corresponding dual edge by the child's two-terminal dual graph. Consequently, for distances between skeleton faces, that child may be replaced by an edge of length $\ell(H_i)$.

There are three skeleton types.

- For an S -node, the primal skeleton is a cycle. After removing the reference dual edge, its dual is a parallel composition. Hence

$$\ell(H) = \min_i \ell(H_i).$$

- For a P -node, the primal skeleton is a dipole. Its dual, after removing the reference dual edge, is a path through the children. Hence

$$\ell(H) = \sum_i \ell(H_i),$$

independently of their order.

- For an R -node, the skeleton has a unique spherical embedding up to reflection. Its terminal distance is the shortest-path distance in the fixed dual skeleton with child-edge lengths $\ell(H_i)$. Reflection does not change that distance.

This proves invariance. The bounds follow because D_H is connected and has $|E(H)|$ unit-length edges. $\square$

This invariance is what permits a polynomial number of states.

2.1.4.2 4.2 Profile definition Weights on the poles s, t are temporarily ignored. All other vertex weights are included.

For an integer d with

$$0 \leq d \leq \ell(H),$$

give the two dual terminals external distance labels $0, d$. In a particular embedding, the resulting distance label of a dual vertex z is

$$\delta(z) = \min \left\{ \text{dist}_{D_H}(A, z), d + \text{dist}_{D_H}(B, z) \right\}. \quad (6)$$

For that embedding, take the maximum of:

1. for every real dual edge xy ,

$$1 + \min\{\delta(x), \delta(y)\};$$

2. for every internal primal vertex $v \neq s, t$,

$$w(v) + \min_{f \ni v} \delta(f).$$

Let $F_H(d)$ be the minimum of this maximum over the embeddings of the pertinent component, including reflection.

Reflection exchanges the two boundary faces while preserving the primal pole labels. Therefore, if the actual terminal labels are a, b , with

$$|a - b| \leq \ell(H),$$

the optimum contribution of the component is

$$\min\{a, b\} + F_H(|a - b|). \quad (7)$$

Each profile has at most $|E(H)| + 1$ entries.

For a real-edge leaf,

$$\ell(H) = 1, \quad F_H(0) = F_H(1) = 1. \quad (8)$$

2.1.5 5. Composition over a fixed skeleton

Consider a skeleton Σ , with its reference edge, whose other edges correspond to children H_i . Write

$$\ell_i = \ell(H_i), \quad F_i = F_{H_i}.$$

Let W be the skeleton dual with the reference dual edge removed, giving its child edge i length ℓ_i . Its reference-face terminals are A, B .

For prescribed terminal labels a, b , compute

$$\delta(z) = \min\{a + \text{dist}_W(A, z), b + \text{dist}_W(B, z)\}. \quad (9)$$

If child edge i has dual endpoints x_i, y_i , its optimum contribution is

$$\min\{\delta(x_i), \delta(y_i)\} + F_i(|\delta(x_i) - \delta(y_i)|). \quad (10)$$

The profile argument is valid because

$$|\delta(x_i) - \delta(y_i)| \leq \ell_i.$$

2.1.5.1 Why child optimization is independent Distances between skeleton faces depend only on the invariant lengths ℓ_i , not on the child embeddings.

For an internal face of a child, every path from outside the child enters through one of its two boundary faces. Thus its distance is obtained from those two boundary labels exactly as in (6). Each child can consequently be optimized independently for its prescribed labels.

2.1.5.2 Vertex weights localize to the skeleton Suppose v is a skeleton vertex other than the parent poles. A new face inside a child incident with v has distance at least the minimum of the child's two boundary labels. Both boundary faces are themselves incident with v .

It follows that, after all substitutions,

$$\min_{f \ni v} \delta(f) = \min_{\substack{f \text{ a skeleton face} \\ f \ni v}} \delta(f). \quad (11)$$

Every internal primal vertex is either:

- internal to exactly one child; or
- a skeleton vertex other than the parent poles.

Thus (10), together with the skeleton-vertex terms from (11), accounts for every edge and every relevant vertex weight.

Define

$$C_{\Sigma}(a, b) = \max \left\{ \begin{array}{l} \max_i [\min\{\delta(x_i), \delta(y_i)\} + F_i(|\delta(x_i) - \delta(y_i)|)], \\ \max_{v \in V(\Sigma) \setminus \{s, t\}} \left[w(v) + \min_{f \ni v} \delta(f) \right] \end{array} \right\}. \quad (12)$$

For a fixed skeleton embedding, this is the exact optimum over its child embeddings.

2.1.5.3 5.1 R -nodes Compute $\ell(H)$ as the weighted terminal distance in W . Both skeleton reflections must be considered:

$$F_H(d) = \min\{C_{\Sigma}(0, d), C_{\Sigma}(d, 0)\}, \quad 0 \leq d \leq \ell(H). \quad (13)$$

The minimum in (13) is important: the two faces adjacent to a reference edge need not have equivalent distance behavior.

2.1.5.4 5.2 S -nodes The two skeleton faces are common terminals of all children. Every internal skeleton vertex is incident with both faces. Therefore,

$$\ell(H) = \min_i \ell_i$$

and

$$F_H(d) = \max \left\{ \max_i F_i(d), \max_{v \in V(\Sigma) \setminus \{s, t\}} w(v) \right\}. \quad (14)$$

It remains to treat P -nodes, where the children can be arbitrarily ordered.

2.1.6 6. The polynomial recurrence at a P -node

At a P -node, the dual network is a series composition of the children in an arbitrary order. There are no internal skeleton vertices, so there are no additional skeleton-vertex weights.

Put

$$\Lambda = \sum_{i=1}^r \ell_i = \ell(H), \quad p_i = F_i(\ell_i). \quad (15)$$

The quantity p_i is the optimum one-sided cost of child i . Indeed, with terminal labels $0, \ell_i$, the first terminal alone determines all distances, since

$$\text{dist}(A, z) \leq \ell_i + \text{dist}(B, z).$$

Fix a profile argument $d \in \{0, \dots, \Lambda\}$. Along a weighted series path, a skeleton vertex at distance x from the left terminal has label

$$\delta(x) = \min\{x, d + \Lambda - x\}. \quad (16)$$

The two expressions meet at

$$h = \frac{\Lambda + d}{2}.$$

Choose a child j whose interval contains h . If h is at a junction, choose either adjacent child. Call j the central child.

Let:

- X be the total length of the children before j ;
- Y be the total length of the children after j .

Then

$$X + Y + \ell_j = \Lambda.$$

The labels at the two ends of the central child are

$$\alpha = X, \quad \beta = d + Y.$$

The condition that the meeting point lies in the central child is exactly

$$|\alpha - \beta| \leq \ell_j. \quad (17)$$

Every child strictly before j is reached optimally from the left; every child strictly after j is reached optimally from the right. Their costs are consequently one-sided costs.

Thus, except for the central child, the problem is an assignment of children to two arms, followed by ordering within each arm.

2.1.6.1 6.1 Optimal ordering within an arm For child i , define

$$q_i = p_i - \ell_i. \quad (18)$$

On either arm, listed from its source toward the central child, there is an optimum order of nonincreasing q_i .

To verify this, suppose $q_i \geq q_j$, and compare consecutive orders ij and ji , starting at accumulated label A . The two costs in order ij are

$$A + \ell_i + q_i, \quad A + \ell_i + \ell_j + q_j.$$

Both are at most

$$A + \ell_i + \ell_j + q_i,$$

which is one of the costs in order ji . Swapping an inverted pair therefore cannot worsen the maximum. Later costs are unchanged.

Reordering within an arm does not change its total length, so it also leaves the central-child condition and contribution unchanged.

2.1.6.2 6.2 Assigning children to the two arms Fix the central child j , and sort the remaining children by nonincreasing q_i .

After processing some initial segment of this list, let its total length be S . Define

$$T(x)$$

to be the minimum possible maximum cost so far when the left arm has total length x , and the right arm has total length $S - x$.

Initially,

$$T(0) = 0,$$

with all other states infeasible.

For the next child i , there are two transitions.

- Put it on the left:

$$T_{\text{new}}(x + \ell_i) \leftarrow \min \{T_{\text{new}}(x + \ell_i), \max\{T(x), x + p_i\}\}. \quad (19)$$

- Put it on the right:

$$T_{\text{new}}(x) \leftarrow \min \{T_{\text{new}}(x), \max\{T(x), d + (S - x) + p_i\}\}. \quad (20)$$

Both arm orders are subsequences of the common sorted list. The actual left-to-right series order is the left list, followed by j , followed by the reverse of the right list. Hence every assignment produced by this dynamic program is realizable.

After all noncentral children have been processed, a state X has

$$Y = \Lambda - \ell_j - X.$$

If

$$|X - (d + Y)| \leq \ell_j,$$

its full cost is

$$\max \{T(X), \min\{X, d + Y\} + F_j(|X - (d + Y)|)\}. \quad (21)$$

Take the minimum of (21) over all feasible X , and then over all choices of j . The result is $F_H(d)$.

2.1.6.3 6.3 Completeness of the recurrence Every ordered series composition has a child containing the meeting point in (16). All other children are one-sided. Sorting each arm as above cannot worsen its contribution, and does not change the central contribution.

Conversely, every state satisfying (17) gives a valid ordering with precisely the terminal labels used in (21). Independently realizing the appropriate child profiles realizes the stated cost.

Thus the recurrence is exact, including:

- a meeting point at a junction;
- a meeting point at the right endpoint, when $d = \Lambda$;
- a composition with only one child.

2.1.6.4 6.4 Complexity For fixed j, d , the arm-assignment dynamic program has $O(\Lambda)$ states and $r - 1$ stages. Computing the whole profile takes

$$O(r^2 \Lambda^2) \quad (22)$$

operations, apart from lower-order sorting work.

The lengths here are not arbitrary binary-encoded numbers: $\Lambda \leq |E(H)|$. This is why the assignment dynamic program is polynomial in the graph size.

2.1.7 7. Solving the weighted biconnected problem

Let B be a biconnected planar graph with vertex weights w . Fix a real edge $e = st$, and require e to be incident with the outer face.

Root the SPQR decomposition at e , and let

$$H = B - e,$$

using e as the reference edge.

In the full dual, the reference dual edge joins the outer-face terminal to the other terminal by an edge of length 1. Thus the terminal distance labels are 0, 1.

The reference primal edge itself has layer 1. Its endpoints are incident with the outer face, so their weighted contributions are $w(s), w(t)$. Consequently, the optimum with e exposed is

$$\text{Opt}(B, w; e) = \max\{1, F_H(1), w(s), w(t)\}. \quad (23)$$

All other edges and vertex weights are included in $F_H(1)$.

It follows that

$$\text{Opt}(B, w) = \min_{e \in E(B)} \text{Opt}(B, w; e). \quad (24)$$

If a specified vertex c must be incident with the outer face, use

$$\text{Opt}(B, w; c) = \min_{\substack{e \in E(B) \\ e \ni c}} \text{Opt}(B, w; e). \quad (25)$$

Every outer-face incidence at c includes an incident edge, so (25) loses no embedding.

Equations (8), (13), (14), and (19)–(21) provide a bottom-up polynomial computation of all required profiles. The standard completeness property of the SPQR decomposition ensures that all embeddings are covered.

2.1.8 8. Cutvertices

The following gluing lemma supplies the reduction to weighted blocks.

2.1.8.1 8.1 Gluing lemma Suppose a connected graph consists of a block B , together with connected branches $H_{v,i}$ attached at vertices $v \in V(B)$. Each branch meets B only at v ; different branches are otherwise disjoint. As in a rooted block-cut tree, each nontrivial branch minus its attachment vertex is connected.

Let

$$h_{v,i}$$

be the minimum edge-outerplanarity of $H_{v,i}$ among embeddings in which v is incident with its outer face. Set

$$w(v) = \max_i h_{v,i}, \quad (26)$$

with $w(v) = 0$ when there is no branch.

Lemma 3. Among embeddings of the whole graph whose outer face is incident with an edge of B , the optimum equals

$$\min_{\text{embeddings and outer faces of } B} \Psi(B, w). \quad (27)$$

The analogous assertion holds with a required outer-face incidence in B .

Proof: upper bound. Choose an embedding of B . For each v , choose an incident face f_v satisfying

$$d(f_v) = a_B(v).$$

Embed every $H_{v,i}$ in a small region of f_v adjoining v , using an optimum embedding with v on its outer face. These regions can be chosen pairwise disjoint, except for their prescribed attachment vertices.

In the dual, each branch dual is joined to the dual of B at the single vertex f_v . Distances between faces of B do not change. All edge layers in the branch increase by exactly $a_B(v)$.

The resulting value is therefore

$$\max \left\{ K(B, r), \max_{v,i} (a_B(v) + h_{v,i}) \right\} = \Psi(B, w).$$

Proof: lower bound. Consider any embedding of the whole graph whose outer face is incident with an edge of B . Restrict it to B , obtaining a fixed embedding of B .

For one branch $H_{v,i}$, delete all other branches. By deletion monotonicity, this cannot increase any remaining layer.

The graph $H_{v,i} - v$ lies in a single face f of B , incident with v . Because an edge of B is on the outer face, the branch does not enclose B . In its induced standalone embedding, v is therefore incident with the outer face.

The dual of $B \cup H_{v,i}$ is obtained by identifying the outer-face vertex of $H_{v,i}^*$ with the vertex f of B^* . Hence the largest layer among the branch edges is at least

$$d_B(f) + h_{v,i} \geq a_B(v) + h_{v,i}.$$

Also, restricting to B shows that the whole graph has value at least $K(B, r)$. Taking the maximum over branches proves the lower bound. $\square$

Notice that this argument does not assume that an arbitrary embedding has nonnested branches. Nesting is handled in the lower bound by deleting all other branches. The upper bound constructs a nonnested embedding attaining the weighted objective.

2.1.8.2 8.2 Dynamic programming on the block-cut tree Choose a root block B_0 and orient the block-cut tree away from it.

For a nonroot block B , let c be its parent cutvertex. Its subtree contains no other block incident with c . Thus, in the subtree, requiring c to be on the outer face requires an outer-face incidence in B itself.

Process the block-cut tree bottom-up.

- At each vertex v of a block, use as $w(v)$ the maximum of the already computed values of the child branches attached there.
- For an ordinary biconnected block B , compute its message toward its parent by the constrained weighted algorithm (25).
- A bridge block has one face, so its message is simply

$$\max\{1, \max_{v \in V(B)} w(v)\}. \quad (28)$$

- At the root block B_0 , use the unconstrained weighted algorithm (24), or (28) if it is a bridge.

By Lemma 3 and induction, this computes the optimum among embeddings whose outer face is incident with B_0 .

Finally, try every block as B_0 . Every embedding of a connected graph with an edge has some block incident with its outer face. Thus the minimum over root blocks is the unrestricted optimum.

For a disconnected graph, solve each component separately and place the resulting embeddings side by side. The answer is the maximum of the component answers. Restriction to a component gives the matching lower bound.

2.1.9 9. Polynomial running time and reconstruction

Let $N = |V(G)| + |E(G)|$.

For one rooted SPQR computation:

- there are $O(N)$ nodes and total skeleton size $O(N)$;
- every profile has $O(N)$ entries;
- an R -node uses weighted shortest paths and table lookups;
- an S -node uses entrywise maxima;
- a P -node uses at most $O(N^4)$ operations by (22).

Thus $O(N^5)$ operations is a safe coarse bound for one rooted computation.

Trying root edges contributes a factor at most N . Computing messages for all blocks for a fixed root block contributes another factor at most N , and trying all root blocks contributes one more. This gives the advertised coarse bound

$$O(N^8).$$

All relevant values have $O(\log N)$ bits. In particular, a branch's optimum is at most its number of edges. The profile and scheduling values are also bounded by a constant multiple of N .

For reconstruction, retain minimizing choices:

- the selected root block and root edge;
- each R -node reflection;
- child profile arguments and orientations;
- each P -node's central child and arm assignment;

- a nearest incident face at every cutvertex.

The SPQR choices produce the block embeddings. Lemma 3 then specifies how to insert the branch embeddings. Hence the algorithm returns an embedding, not merely the optimum value.

For multigraphs, parallel edges are already covered by the dipole skeletons. A loop can be treated as an elementary one-edge block, with the same value formula as (28). Isolated vertices have no effect.

2.1.10 10. Conclusion and dependencies

The algorithm is exact. Its two essential ingredients are:

1. **Invariant interface lengths:** a pertinent component's shortest dual distance between its two boundary faces does not depend on its embedding.
2. **A polynomial ordering step:** at a parallel node, at most one child receives genuinely two-sided distances; all other children lie on two one-sided arms, each optimally sorted by $F_i(\ell_i) - \ell_i$.

The first makes child optimization independent at fixed skeletons. The second removes the exponential permutation choice at parallel skeletons. Vertex weights and Lemma 3 handle all cutvertices.

This proves the polynomial-time claim. The external structural dependency is the standard SPQR decomposition theorem and its completeness for planar embeddings. I have not independently established whether this particular algorithm, or an equivalent one, is already present in the literature; the argument above is an independent proof rather than a literature-status assertion.

A Provenance and review status

UNREFEREED MODEL OUTPUT. This document was selected solely because GPT-6 Astra labelled its own result `would_publish: true` and returned `proved` or `disproved`. It has not passed the adversarial LLM referee used for the earlier Sol campaign, has not been checked by a human mathematician, and has not been checked for novelty. Treat every mathematical and bibliographic claim as unverified.

Catalog id	finding_k_edge_outerplanar_graph_embeddings
Catalog entry	Finding k-edge-outerplanar graph embeddings
Source corpus	opg
Campaign leg	OpenProblemGarden first pass (attacks_opg)
Source paper / entry	http://www.openproblemgarden.org/op/finding_k_edge_outerplanar_graph_embeddings
Model verdict	proved (confidence: high)
Selection criterion	would_publish: true and verdict proved or disproved
Model	gpt-6-astra, reasoning effort max, mode=pro, flex service tier
Original artifact	attacks_opg/finding_k_edge_outerplanar_graph_embeddings/output.md
Independent review	None
Model's caveats	Uses the standard SPQR decomposition theorem; novelty and current literature status have not been independently checked.

The generated Markdown and LaTeX sources for this PDF are stored under `to_review_astra/src/finding_k_edge_outerp`