

Recasting activities at LH2017

L. Perrozzi¹, Fabio Maltoni, Sabine Kraml, Gabriel Facini, David Grellscheid, Sezen Sekmen, Johnatan Butterworth, Nishita Desai, Andy Buckley, Benjamin Fuks, Eric Conte, Peter Richardson, Olivier Mattelaer, Pasquale Musella, Alexandra Oliveira Carvalho, Ursula Laa, Kristin Lohwasser, Thrynova, Efe Yazgan, Philippe Gras, Sylvain

¹IPA at ETH Zurich, Switzerland

² ...

Abstract

Recasting activities at Les Houches 2017.

1. INTRODUCTION

Searches for new physics constitute a basic ingredient of the LHC physics program. Their variety and large number pose severe challenges to both the experimental and theory communities. In fact, hundreds of searches are performed by different collaborations, a wide variety of final states is used, new ideas on how to probe new models and non-trivial signatures and improve the sensitivity of existing searches constantly emerge. The ultimate goal of this effort is to discover new physics if such exists within the range of the LHC, and to test the widest possible range of hypothetical new physics models.

A typical analysis defines quantities that aid in classifying the event as signal or background: for example the properties of analysis objects such as jets, electrons, muons, etc., or global event variables such as object multiplicities, transverse momenta, transverse masses, etc. An analysis can be very complex and feature many intricate definitions of object and event variables, some of which cannot be expressed in closed algebraic form and must be defined algorithmically. This complexity renders the task of visualizing, understanding, developing and interpreting analyses increasingly challenging.

One obvious way to cope with the complexity is to devise ways to enforce absolute clarity in the description of analyses. A discussion was started in the Les Houches PhysTeV workshop in 2011, and continued thereafter within a wider group of LHC physicists, in order to determine what information is crucial for describing an analysis. The outcome of this discussion was reported in the Recommendations for Presentation of LHC Results[524,525], and has been embraced by many LHC physicists.

The current practice in our community is to write an analysis in non-public computer codes, which often rely on event objects specific to the experimental collaboration in question, and then make public a description of the analysis via journal publications or other documents.

These efforts, which merit our great appreciation, have significantly increased the scientific value of many important experimental results.

There is significant precedent for the effectiveness of such community standards. Several accords have been established to standardize the communication of physics modeling information, notably the Les Houches Event Accord (LHE) [529, 530] and the SUSY Les Houches Accord (SLHA) [190, 531]. These, respectively, standardize the description of hard-process particles in simulated collision events, and the details of all the parameters that define a BSM model point. Both accords are widely used in high-energy physics and have greatly helped to simplify and make more efficient the communication between physicists. In this report, we underscore the need for a standardized format an analysis description accord capable of describing the contents of an analysis in an unambiguous way, which can be fully exploited by the whole particle physics community. The accord must be capable of describing all object and event selections, as well as quantities such as efficiencies, analytic and algorithmic observables, and advanced multivariate selections. In the Les Houches PhysTeV workshop in 2015, a dedicated discussion has been initiated on how such an accord can be realized. The important motivations and use-cases for a standard analysis description accord were envisaged as:

- Analysis preservation
- Analysis design
- Analysis review and communication
- Interpretation studies and analysis reimplementation
- Easier comparison of analyses

In addition, there are several desirable features which would further improve the utility of the accord, which, however, may be nontrivial to simultaneously fulfill. Therefore, here we list all the desirable features, leaving it to the community discussion (and to individual prototypes) to decide which of these desiderata should inform an eventual accord. We start by listing the features we believe essential to the success of such an accord.

BASIC REQUIREMENTS:

- Public availability
- Completeness
- Longevity
- Correctness and validatability

DESIRABLE FEATURES:

- Human readability and writeability
- Self-contained
- Language independence
- Framework independence
- Support for combination of analyses

Several discussions and progresses have been made, but the proposal is not yet final and has not been widely adopted yet.

The scope of this report, therefore, is to document the advancements obtained so far on the recasting activities and attempt a first benchmark to compare different tools to reproduce several ATLAS and CMS analyses' results.

2. General Activities

- Feasibility study of the implementation/portability of complicated MVA techniques (BDT, NN,) into the analyses
- Improvement of results and recastability: how to provide correlations signal systematics, possibility of providing a few key observables unfolded.
- Comparison of between DELPHES results and simple object smearing.
- Trying out the use of particle-level measurements to constrain model models

3. Formats

Object efficiency tables : which format (HEPDATA?)

4. Benchmarking/Comparisons

- Implementation of analyses of increasing complexity in the Analysis Description Format (LHADA Proposal) and in (BSM) Rivet and their comparison.
- Choose an analysis of ATLAS or CMS which has cutflow and detector effects provided in some form, and possibly is already been implemented in the recasting codes CheckMate/MadAnalysis/Rivet/ATOM.
- Implement the same analysis in LHADA and then use the dedicated parsers to provide the analysis for the recasting codes.
- Reproduce the NP interpretation of the original paper (=validation implementation).




Fig. 1: Search reach for the $\mu\gamma E_T$ signal (as defined in the text) for 300 fb^{-1} integrated luminosity at the LHC.

- Recast the analysis for an other new physics model and compare the results.
- Go to point one and choose a more complicated analysis

it would be interesting to see how Delphes performance looks without analysis-specific cards, since a lot of people (outside the big recasting groups) are using it that way.

5. How to validate the analyses

6. The analysis frameworks and tools

In this section we describe the analysis frameworks and tools used for the comparison and benchmarking

6.1 ATOM

Brief description of ATOM

6.2 CheckMate

Brief description of CheckMate

6.3 MadAnalysis

Brief description of MadAnalysis

6.4 Rivet

Brief description of Rivet

6.5 Generic Analysis Description Proposal

Brief description of the Generic Analysis Description Proposal

7. Analyses benchmarking, comparisons and results

7.1 arxiv:1605.03814 - Jets+MET - ATLAS - 13 TeV

Brief description of the ATLAS analysis Jets+MET at 13 TeV (arxiv:1605.03814). Results are reported in table 1.

	Rivet			MadAnalysis5			CheckMATE
Description	#evt	tot.eff	rel.eff	#evt	tot.eff	rel.eff	tot.eff
2jl cut-flow	31250	1	-	31250	1	-	
Pre-sel+MET+pT1	28592	0.91	0.91	28626	0.92	0.92	
Njet	28592	0.91	1	28625	0.92	1	
Dphi_min(j,MET)	17297	0.55	0.6	17301	0.55	0.6	
pT2	17067	0.55	0.99	17042	0.55	0.99	
MET/sqrtHT	8900	0.28	0.52	8898	0.28	0.52	
m_eff(incl)	8896	0.28	1	8897	0.28	1	
2jm cut-flow	31250	1	-	31250	1	-	1
Pre-sel+MET+pT1	28472	0.91	0.91	28478	0.91	0.91	0.91
Njet	28472	0.91	1	28477	0.91	1	0.91
Dphi_min(j,MET)	22950	0.73	0.81	22889	0.73	0.8	0.73
pT2	22950	0.73	1	22889	0.73	1	0.73
MET/sqrtHT	10730	0.34	0.47	10710	0.34	0.47	0.33
m_eff(incl)	10630	0.34	0.99	10609	0.34	0.99	0.32
2jt cut-flow	31250	1	-	31250	1	-	
Pre-sel+MET+pT1	28592	0.91	0.91	28626	0.92	0.92	
Njet	28592	0.91	1	28625	0.92	1	
Dphi_min(j,MET)	17297	0.55	0.6	17301	0.55	0.6	
pT2	17067	0.55	0.99	17042	0.55	0.99	
MET/sqrtHT	5083	0.16	0.3	5098	0.16	0.3	
Pass m_eff(incl)	4861	0.16	0.96	4889	0.16	0.96	
4jt cut-flow	31250	1	-	31250	1	-	1
Pre-sel+MET+pT1	28592	0.91	0.91	28626	0.92	0.92	0.91
Njet	27322	0.87	0.96	27128	0.87	0.95	0.87
Dphi_min(j,MET)	18929	0.61	0.69	18829	0.6	0.69	0.6
pT2	18715	0.6	0.99	18825	0.6	1	-
pT4	16610	0.53	0.89	16430	0.53	0.87	0.52
Aplanarity	11849	0.38	0.71	11395	0.36	0.69	0.36
MET/m_eff(Nj)	8334	0.27	0.7	7971	0.26	0.7	0.25
m_eff(incl)	7201	0.23	0.86	6972	0.22	0.87	0.21
5j cut-flow	31250	1	-	31250	1	-	1
Pre-sel+MET+pT1	28592	0.91	0.91	28626	0.92	0.92	0.91
Njet	21234	0.68	0.74	21185	0.68	0.74	0.68
Dphi_min(j,MET)	14294	0.46	0.67	14292	0.46	0.67	0.45
pT2	14146	0.45	0.99	14289	0.46	1	-
pT4	13229	0.42	0.94	13228	0.42	0.93	0.42
Aplanarity	9836	0.31	0.74	9576	0.31	0.72	0.3
MET/m_eff(Nj)	4643	0.15	0.47	4506	0.14	0.47	0.13
m_eff(incl)	4620	0.15	1	4476	0.14	0.99	0.13
6jm cut-flow	31250	1	-	31250	1	-	1
Pre-sel+MET+pT1	28592	0.91	0.91	28626	0.92	0.92	0.91
Njet	13235	0.42	0.46	13236	0.42	0.46	0.41
Dphi_min(j,MET)	8520	0.27	0.64	8553	0.27	0.65	0.26
pT2	8436	0.27	0.99	8551	0.27	1	-
pT4	8135	0.26	0.96	8217	0.26	0.96	0.25
Aplanarity	6365	0.2	0.78	6307	0.2	0.77	0.19
MET/m_eff(Nj)	2675	0.09	0.42	2665	0.09	0.42	0.08
m_eff(incl)	2670	0.09	1	2656	0.08	1	0.08
6jt cut-flow	31250	1	-	31250	1	-	
Pre-sel+MET+pT1	28592	0.91	0.91	28626	0.92	0.92	
Njet	13235	0.42	0.46	13236	0.42	0.46	
Dphi_min(j,MET)	8520	0.27	0.64	8553	0.27	0.65	
pT2	8436	0.27	0.99	8551	0.27	1	
pT4	8135	0.26	0.96	8217	0.26	0.96	
Aplanarity	6365	0.2	0.78	6307	0.2	0.77	
MET/m_eff(Nj)	3900	0.12	0.61	3839	0.12	0.61	
m_eff(incl)	3715	0.12	0.95	3672	0.12	0.96	

Table 1: 1605.03814 cut flow

7.2 arxiv:1704.03848 - Monophoton - ATLAS - 13 TeV

Brief description of the ATLAS analysis Monophoton at 13 TeV (arxiv:1704.03848).

7.3 CMS-SUS-16-039 - 3 leptons + MET - CMS - 13 TeV

Brief description of the CMS analysis 3 leptons + MET at 13 TeV (CMS-SUS-16-039).

7.4 arxiv:1706.04402 - 1 lepton + MET + Jets ($\geq 1b$) - CMS - 13 TeV

(topness variable?) Brief description of the CMS analysis 1 lepton + MET + Jets ($\geq 1b$) (topness variable) at 13 TeV (arxiv:1706.04402).

CONCLUSIONS

We are cool.

ACKNOWLEDGEMENTS

We acknowledge the acknowledgements.

References