# Gravitas

## User Manual

An interactive astrophysics sandbox and astronomy teaching tool that runs entirely in the browser

https://gravitas-sim.online

61 scenarios • 91 guided investigations • 1289 steps

Carl Ziegler

This manual, and the investigations and instructor material it describes, are licensed under Creative Commons Attribution 4.0 International (CC BY 4.0) — use them, adapt them, put them in a course package, with credit. The Gravitas software itself is MIT. See `LICENSES.md` in the repository for which license covers which files.

Generated from the application's own catalog; see `manual/`.

# Contents

**1 What Gravitas is** **5**
1.1 What it models, and what it only draws . . . 5

**2 Getting started** **5**
2.1 The screen . . . 5
2.2 Keyboard . . . 6

**3 Finding your way around the site** **6**
3.1 The menu . . . 6
3.2 The Library . . . 6
3.3 My work . . . 7
3.4 Courses . . . 7

**4 The instruments** **7**
4.1 Ruler, protractor, stopwatch . . . 7
4.2 The object inspector . . . 7
4.3 Why the bodies are not to scale . . . 7
4.4 Vectors and the potential well . . . 8
4.5 Reference frames . . . 8

**5 The observing panels** **8**

**6 Choosing the physics** **9**
6.1 Integrators . . . 9
6.2 Dark matter . . . 9
6.3 Tides . . . 9

**7 Scenarios** **9**
7.1 Binary Systems . . . 9
7.2 Chaos & Encounters . . . 10
7.3 Compact Objects . . . 10
7.4 Exoplanets . . . 12
7.5 Galaxies & Clusters . . . 12
7.6 Habitability . . . 13
7.7 Orbits & Kepler . . . 13
7.8 Orbital Resonance . . . 14
7.9 Solar System . . . 15
7.10 Stellar Evolution . . . 15
7.11 Tides & Disruption . . . 16

**8 Guided investigations** **16**
8.1 Kepler's Laws . . . 18
8.2 Why Mars Goes Backwards . . . 18
8.3 Finding Planets by Their Shadows . . . 18
8.4 Bound, Unbound and Escape . . . 18
8.5 Weighing the Stars . . . 19
8.6 Black Holes by the Numbers . . . 19
8.7 Finding Planets by Their Tug . . . 19
8.8 The Goldilocks Question . . . 19
8.9 The Missing Mass . . . 19

8.10 Tides . . . 19
8.11 The Butterfly Effect in Space . . . 19
8.12 When Orbits Lock . . . 19
8.13 Can You Detect This Planet? . . . 20
8.14 Design the Schedule . . . 20
8.15 Planets in Binary Stars . . . 20
8.16 Where Does a Gravity Assist Get Its Speed? . . . 20
8.17 Getting There From Here . . . 20
8.18 Where Can It Get To? . . . 20
8.19 What Is a Gravitational Wave? . . . 20
8.20 Listening to Spacetime . . . 20
8.21 A Universe of Stars . . . 20
8.22 Lives of Stars . . . 21
8.23 Twelve Nights . . . 21
8.24 What If Gravity Were Not Inverse Square? . . . 21
8.25 Color and Temperature . . . 21
8.26 The Turning Sky . . . 21
8.27 The Sun Through the Year . . . 21
8.28 Phases and Eclipses . . . 21
8.29 Wanderers on the Sky . . . 21
8.30 Plan a Night at the Telescope . . . 21
8.31 Lines and Motion . . . 22
8.32 What a Spectrum Is Made Of . . . 22
8.33 The Sun's Motion . . . 22
8.34 The Rotating Disk . . . 22
8.35 Cluster Membership by Motion . . . 22
8.36 Inside a Star . . . 22
8.37 Scaling the Main Sequence . . . 22
8.38 The Limit . . . 22
8.39 The Expanding Universe . . . 22
8.40 Standard Candles . . . 23
8.41 The Shape of Expansion . . . 23
8.42 Where Galaxies Live . . . 23
8.43 What a CCD Records . . . 23
8.44 Signal, Noise and Exposure . . . 23
8.45 A Variable Star From Frames . . . 23
8.46 The Beat of a Star . . . 23
8.47 Leavitt's Law . . . 23
8.48 The First Rung . . . 23
8.49 Which Kind of Variable . . . 24
8.50 Bending Light . . . 24
8.51 Einstein Rings and Arcs . . . 24
8.52 A Lens Passing By . . . 24
8.53 Who Gets Counted . . . 24
8.54 Why the Bright Ones Win . . . 24
8.55 The Completeness Curve . . . 24
8.56 Integrators in Practice . . . 24
8.57 Extending an Experiment . . . 24
8.58 Refitting a Transit . . . 25
8.59 Two Periodograms . . . 25
8.60 Sampling a Posterior . . . 25

8.61 What Is a Measurement? . . . 25
8.62 What Does ± Claim? . . . 25
8.63 When the Method Must Refuse . . . 25
8.64 Measuring the Moon . . . 25
8.65 How Far Is That Star? . . . 25
8.66 A Binary Star's Orbit . . . 26
8.67 Which Surface Is Older? . . . 26
8.68 Craters by Size . . . 26
8.69 What a Crater Diameter Is Worth . . . 26
8.70 What Is in This Field? . . . 26
8.71 Weighing Jupiter From Its Moons . . . 26
8.72 Why Rings? . . . 26
8.73 Tides That Move the Moon . . . 26
8.74 Io's Fire . . . 26
8.75 The Gaps in the Belt . . . 27
8.76 How Fast Does It Spin? . . . 27
8.77 Why Venus Is Hot . . . 27
8.78 Two Greenhouse Models . . . 27
8.79 The Turning Sun . . . 27
8.80 The Sun's Power . . . 27
8.81 How Hot Is the Day Side? . . . 27
8.82 Phases and Albedo . . . 27
8.83 Moons of the Giants . . . 27
8.84 The Scale of the Solar System . . . 28
8.85 Comets and the Sun . . . 28
8.86 What a Telescope Resolves . . . 28
8.87 Why Two Lines Become One . . . 28
8.88 The Saturated Star . . . 28
8.89 Counts to Magnitudes . . . 28
8.90 Plan an Exposure . . . 28
8.91 Why the Period Is Wrong . . . 28

**9 Running a controlled experiment 29**
9.1 What the bench will tell you off for . . . 29
9.2 Honest sampling . . . 29
9.3 Keeping and sharing an experiment . . . 29

**10 Capturing, exporting and sharing 30**
10.1 Screenshots and clips . . . 30
10.2 Data . . . 30
10.3 Share links . . . 30
10.4 Embed mode and lecture mode . . . 30

**11 Languages 30**

**12 Beyond the simulation 31**
12.1 The Observatory . . . 31
12.2 The 3-D lab and the mission lab . . . 31
12.3 The Sky Lab . . . 31
12.4 The experiment runner . . . 31
12.5 The catalog . . . 31
12.6 The figure builder . . . 31

12.7 The Studio . . . . . 32

**13 For instructors** **32**
13.1 From a first visit to a reviewed class . . . . . 32
13.2 Reading what students hand in . . . . . 33
13.3 Citing Gravitas . . . . . 33

**14 Limitations** **33**

**15 Accessibility** **33**

**16 Where to find more** **34**

# 1 What Gravitas is

Gravitas is a gravity simulation you can teach with. It runs in a browser, needs no account and no installation, and is served as static files: nothing you do in it reaches a server, and nothing about you is recorded by it.

It is two things at once. It is a *sandbox* — place stars, planets and black holes, give them velocities, and watch what Newtonian gravity does with them — and it is a set of *guided investigations* that use that sandbox to take a student from a prediction, through a measurement, to a number they can defend.

## 1.1 What it models, and what it only draws

This distinction matters more than any other in this manual, and the application states it in public at https://gravitas-sim.online/model/.

**Calculated**
Newtonian gravitation between every pair of bodies, in two dimensions, integrated with a scheme you choose. Orbits, energies, angular momenta, transit depths, radial-velocity curves, astrometric wobble, tidal differentials and rotation curves all come out of that integration.

**Approximated**
Mergers are perfectly inelastic. The gravitational-wave inspiral is a phenomenological energy loss, not a solution of general relativity. The dark-matter halo is an added force term with an adjustable flat rotation speed.

**Drawn only**
Accretion-disc glow, relativistic jets, gravitational-lensing distortion and the bloom around bright objects are pictures. They influence nothing.

Every claim the simulation makes about a measurable quantity is checked by a validation suite of 348 deterministic tests, each with a stated tolerance and a written reason for that tolerance. The results, and the list of things that are deliberately *not* validated, are published with the project.

# 2 Getting started

Open https://gravitas-sim.online. The simulation starts immediately; a first visit also shows a short front door offering three ways in, which you can dismiss and never see again.

## 2.1 The screen

**The simulation**
fills the window. Scroll to zoom, drag to pan, click a body to select it.

**The readout, top left**
carries the live numbers: elapsed simulated time, the zoom, the simulation speed, how many bodies of each kind are present, and — if you ask for them in Settings — the integrator in use and how far energy and angular momentum have drifted since the run began.

**The control rail, right**
is an accordion: Scenario, State, Tools, View, Capture and Learn. One section is open at a time.

**The transport bar, bottom**
pauses, steps and scrubs. The simulation records its own history, so the scrubber rewinds through what has already happened and holds on the frame you land on until you return to live.

**The scale bar, bottom left**
is always on the canvas, not in the page. It is drawn onto the simulation itself so that a screenshot carries its own scale.

### 2.2 Keyboard

| Key | What it does |
|---|---|
| `Space` | Pause and resume |
| `R` | Recentre the camera and return to 1× zoom |
| `P` | Take a screenshot |
| `E` | Export the recorded run as CSV |
| `?` | The full shortcut list |
| `Esc` | Leave lecture mode, or close the open dialog |

The complete list is in the Shortcuts dialog, which is the authoritative copy; this table is the handful you will use in a first session.

## 3 Finding your way around the site

The simulation is the front door of a site with many pages. They share one header, so the same menu, language switch and theme switch are on every page.

### 3.1 The menu

The header menu has five groups, and each opens to a short list.

**Learn**
Library, My work, Investigations, Courses, Observatory and Experiments.

**Labs**
Sandbox, 3-D lab, Mission lab and Sky Lab.

**Make**
Scenario Studio, Investigation Composer, Course builder, Figure builder and Catalog.

**Teach**
Teaching with Gravitas, Instructor resources, Submission review and the Classroom evidence kit.

**About**
How the model works, Physics validation, Glossary, the diagnostics pages for the 3-D kernel and the mission core, and this manual.

The menu also holds the Language switch (English or Spanish) and the Theme switch (Midnight, Deep space, Observatory red and Daylight). Both are remembered in the browser. When the network is down, the header says which pages are saved on the device; any other page works only if it was opened before.

### 3.2 The Library

`/library/` is one index of everything that can be opened: the 108 investigations (91 that run in the simulation and 17 guided investigations in the Observatory, the 3-D lab and the mission lab), the 61 scenarios, the datasets, the 5 ready-made courses, the experiments and the classroom activities. Every entry has the same fields, so they can be compared and filtered by level, time, mathematics and subject. The index is written from the sources that own each item, so it cannot list one that is gone.

### 3.3 My work

`/my-work/` is the one page where a student sees and manages what they have made: progress in investigations, activities, the evidence notebook, experiments, the scenarios they built and their drafts. It needs no account and sends nothing; it reads this browser's own storage. Work exists only in this browser on this device, and nobody at Gravitas can see or recover it, so the page offers a backup: Download everything writes one file that can be imported on another device after a preview of what it would change. Deleting asks for a copy to be downloaded first.

### 3.4 Courses

`/course/` opens a course: investigations, activities, simulations, data and readings in order and in units. A course is a link or a file made in the course builder (`/studio/course/`); the Library lists the ready-made ones. A student's place in a course is kept in the browser like everything else.

## 4 The instruments

Gravitas is a measuring instrument before it is a display. Everything in this section is in the Tools section of the rail.

### 4.1 Ruler, protractor, stopwatch

The **ruler** is dragged across the scene and reads a distance in AU and in kilometers. The **protractor** has a vertex and two arms and reads the angle between them. Both store their handles in *world* coordinates, so panning or zooming does not change what they say — a ruler pinned to two bodies keeps measuring those two bodies.

The **stopwatch** runs on simulated time, so pausing pauses it and changing the simulation speed does not change what it measures. With a body selected it can be latched to that body's periapsis passages: each lap is then one orbit, timed from closest approach rather than by your reaction time, and the panel reports the mean of the laps.

### 4.2 The object inspector

Select a body and the inspector gives its mass, radius, speed, the orbital elements of its path about whatever it is actually orbiting — semi-major axis, eccentricity, period, periapsis and apoapsis — and its kinetic, potential and total energy.

### 4.3 Why the bodies are not to scale

The bodies are drawn on a compressed scale, and the canvas says so: a line beside the scale bar reads *Body sizes enlarged for visibility.* It is drawn on the canvas rather than in the interface, so it is in every screenshot and every recording you make.

The real ratios cannot be drawn. The Sun's radius is about 109 times the Earth's; at a zoom where the Sun is a visible disc, an Earth drawn to scale beside it is a fraction of a pixel. Gravitas compresses the hierarchy instead, by one documented policy applied to every scenario: a Sun-like star is drawn about eight times the radius of an Earth-like planet and four times a Jupiter-like giant, against real ratios of 109 and 9.7. Gas giants are visibly larger than rocky planets, white dwarfs smaller than ordinary stars, and neutron stars, comets, asteroids and distant planets sit at a minimum marker size because at their true relative sizes they would not be visible at all.

Read sizes from the inspector and from the investigation measurements, never off the drawing. The number in the inspector is the physical radius; what is on the canvas is an illustration of

it. The scale *bar* is a different matter and is an honest statement about distance in the current view — though some scenarios are themselves scale models or teaching constructions, so not every drawn distance is physical either. The model page at https://gravitas-sim.online/model/ sets all of this out under *Displayed sizes*, including the four different radii a body has and which of them each part of the simulation uses.

A body is no harder to select for being drawn small: the click target is computed from the body's own size and a minimum in screen pixels, and has nothing to do with how large the body appears.

About a quarter of procedurally generated gas giants are drawn with a ring system. That fraction is a choice about visual variety, not a claim about how common rings are among giant exoplanets, which nobody knows. Which giants get them, and what the rings look like, is fixed by each body's own seed, so it is the same after a reset, a reload, a shared link or a screenshot; scenarios may force rings on or off, and Saturn is ringed in the Solar System. Rings are drawn only: they have no mass and take no part in gravity, collisions, transit depth or any measurement.

### 4.4 Vectors and the potential well

For the selected body, Gravitas can draw its velocity, its total acceleration, and one arrow per gravitational source acting on it, each in a color that cannot be confused with the others. The arrows are the acceleration the integrator actually used, not a second calculation that might disagree with it.

They exist for one misconception in particular: on an eccentric orbit velocity and force are perpendicular at periapsis and apoapsis and never come within forty degrees of parallel anywhere. An optional potential-well underlay shades the whole scene by gravitational potential.

### 4.5 Reference frames

The Frame selector re-expresses every position and every trail as a chosen body would see them. This is not the same as following a body with the camera: the recorded paths are redrawn. Put the Solar System into Earth's frame and Mars traces the retrograde loop that the geocentric model was built to explain.

## 5 The observing panels

Three panels turn the simulation into an observation, and all three are driven by the same viewing geometry. Drag the observer handle on the canvas, or use the angle controls, and every open panel re-measures from the new direction.

**Transit light curve**
Brightness against time for the starlight reaching the observer. A planet crossing its star carves a dip; the depth gives the planet's radius, and limb darkening is included, so the dip is not flat. The depth is $(R_p/R_s)^2$ computed from the *physical* radii — the ones the inspector reports, derived from mass — and not from the enlarged markers on the canvas. A planet drawn an eighth of its star's width does not produce a dip of one sixty-fourth; measure the depth from the curve and check it against the inspector's radii, never against the picture.

**Radial velocity**
The star's motion toward and away from the observer, as a spectrograph measures it. The wobble gives a planet mass — times the sine of the inclination, which is where the $\sin i$ degeneracy stops being a footnote and becomes something a student can see by tilting the system.

**Astrometry**
The star's position on the sky over time, and the astrometric signature at an assumed distance.

**Rotation curve**
Orbital speed against distance from the center, one point per body, drawn against the curve the visible mass alone would produce.

Open panels stack in the bottom-left corner rather than covering each other. The spacetime view — an optional 3-D rendering of the scene as a rubber sheet dipping around each mass — joins the same stack, and can be dragged out of it.

# 6 Choosing the physics

## 6.1 Integrators

Three schemes, switchable while the simulation runs, with live energy and angular-momentum drift beside them:

**Symplectic Euler**
The default, and what every scenario in the catalog is tuned against. First order, and its energy error is bounded rather than growing without limit.

**Velocity Verlet**
Second order, also symplectic. The best general choice when accuracy matters more than matching the shipped scenarios exactly.

**RK4**
Fourth order and not symplectic: locally the most accurate of the three and the only one whose energy error grows steadily over many orbits.

Each scheme's convergence order is *measured* by the validation suite, not asserted, and the tolerances of the integrated checks are derived from it.

## 6.2 Dark matter

An optional halo adds a force term with an adjustable flat rotation speed. It changes the force law rather than the picture: switch it on with the rotation curve open and a falling curve flattens. It stays on until you switch it off, so a scenario loaded afterwards is running with modified gravity — the rotation curve panel says so.

## 6.3 Tides

Tidal stress is computed as the difference between the pull on a body's near side and the pull on its center. Where that difference exceeds the body's own gravity, the body comes apart — which is what happens in the Tidal Disruption Event scenario, and what the Roche limit marks.

# 7 Scenarios

61 scenarios ship with Gravitas. Open the gallery from Scenario in the rail; each is tagged by curriculum concept, so the gallery can be scanned for the week you are teaching rather than read end to end.

Loading a scenario rebuilds the world from a seed, so the same scenario opens the same way every time, and a share link carries the seed with it.

## 7.1 Binary Systems

**Binary Pair: two stars, one balance point**
Two stars of two solar masses each, four AU apart, circling their common center of mass once

every four years. Neither one is stationary and neither one is orbiting the other: both go round the same point in between them. Watch the trails and the balance point gives itself away.

**Binary Planet Lab: a planet around one star of a pair**

Two stars of one and half a solar mass, ten AU apart on an orbit of eccentricity 0.4, and an Earth-mass planet circling the heavier one at 1.5 AU. Nothing here is randomized: every mass, distance and starting angle is written down, so two runs differ only where you make them differ. Move the planet outward and there is a distance past which it stops coming back. Finding roughly where, and how much of that answer is physics rather than timestep, is the experiment.

**Circumbinary Planet Lab: a planet around both stars at once**

The same two stars, with the planet outside them both at 40 AU, orbiting the pair as though it were one object. Real planets do this - Kepler-16b is the famous one - but only far enough out that the two stars begin to look like one. Bring it inward and the approximation fails: the changing pull of a binary it can still tell apart walks its orbit outward over a few dozen binary periods until it leaves. Here the boundary is a floor, not a ceiling.

**Binary Stars**

A pair of suns in mutual orbit with 5 planets orbiting the binary system. The complex gravitational environment creates interesting orbital dynamics and potential habitable zones. Similar to real binary star systems like Alpha Centauri.

## 7.2 Chaos & Encounters

**Three-Body Sensitivity Lab: the same start, twice**

Three six-solar-mass stars at the corners of an equilateral triangle, rotating rigidly about their common center. This is an exact solution of the three-body problem, found by Lagrange in 1772, and for three equal masses it is unstable: the triangle holds for a few turns and then comes apart. Built for one experiment - run it twice from starts that differ by fifteen hundred kilometers in a system a hundred and thirty million kilometers across, and watch how long the two runs stay together.

**Gravity Slingshot**

A massive black hole (60 M$\odot$) paired with a smaller companion (3 M$\odot$) create dramatic gravitational assists for nearby planets and gas giants. Watch objects gain tremendous velocity through close encounters, mimicking spacecraft gravity assists.

**Rogue Encounter**

A wandering black hole (30 M$\odot$) passes through a stable planetary system with 12 planets, 4 gas giants, and asteroids. Watch the dramatic orbital disruption, planet ejection, and tidal capture events as the rogue intruder wreaks havoc.

**Star Frisbee**

A dense stellar disk thrown past a rogue black hole. Will it be shredded or survive the flyby?

**Kessler Cascade**

Hundreds of micro-stars orbiting chaotically, colliding and ejecting like a debris cloud.

**Alien Dyson Swarm Collapse**

A hypothetical Dyson swarm of artificial satellites falls into a black hole after a catastrophic orbital failure.

**Slingshot Gauntlet**

A fast-moving star fired through a black hole obstacle course. Watch gravitational slingshots.

## 7.3 Compact Objects

**Black Hole Lab: a ten solar mass hole, and four things orbiting it**

A single stellar-mass black hole with four bodies on stable circular orbits around it. Nothing

is falling in. Gravity a long way from a black hole is the same gravity as anywhere else, and an object with sideways motion goes round it exactly as it would go round a star of the same mass. Select the black hole to read its Schwarzschild radius, its average density on that scale, its Hawking temperature and how long it has left.

**GW150914: First Gravitational Wave Merger**
Simulates the historic merger of two massive black holes (36 & 29 M⊙) detected by LIGO in 2015. Watch as they spiral together, emit gravitational waves, and merge into a single, more massive black hole.

**Binary Black Hole**
Two stellar-mass black holes (15 & 10 M⊙) locked in mutual orbit with spectacular relativistic jets. Watch as they spiral together, create gravitational waves, and eventually merge into a single, more massive black hole. The jets point in random directions for each black hole, creating a dynamic cosmic display.

**Triple Black Hole**
A chaotic three-body dance of massive black holes (20, 15, & 10 M⊙) in a complex orbital arrangement. This unstable configuration will eventually eject one black hole while the remaining two merge. Demonstrates the chaotic nature of multi-body gravitational systems.

**Supermassive Core**
One enormous black hole (80 M⊙) dominates a dense stellar swarm with 50 planets, 5 gas giants, and 100 asteroids. The intense gravitational field creates spectacular accretion disks and tidal disruption events. Similar to the environment around real supermassive black holes in galactic centers.

**Sagittarius A***
The Milky Way's central supermassive black hole (4000 M⊙, scaled down for simulation) with fast-moving S-stars, compact objects, and debris in extreme orbits. Witness the incredible gravitational forces and relativistic effects near our galaxy's supermassive black hole.

**Neutron Star Merger**
Two neutron stars (1.4 M⊙ each) spiral toward each other in a death dance. This rare event produces gravitational waves, gamma-ray bursts, and creates heavy elements through r-process nucleosynthesis. Based on the LIGO-detected GW170817 event.

**Pulsar with Planets**
A rapidly spinning neutron star with 3 planets in tight orbits. The pulsar's intense magnetic field and radiation create a harsh environment. Based on the first confirmed exoplanets discovered around PSR B1257+12.

**White Dwarf Binary**
Two white dwarf stars in a close binary system with accretion between them. One star gradually steals material from its companion, potentially leading to a Type Ia supernova. Includes debris disk and stellar remnants.

**Stellar Graveyard**
A dynamic collection of stellar remnants: 3 black holes, 5 neutron stars, and 8 white dwarfs with surviving planets and extensive debris fields. Watch these stellar corpses interact in their final gravitational dance.

**Compact Object Zoo**
A diverse collection of compact objects: multiple black holes, neutron stars, and white dwarfs of various masses interacting in a dense environment. Perfect for studying the different types of stellar endpoints and their interactions.

**Millisecond Pulsar**
An extremely fast-spinning neutron star (recycled pulsar) with a white dwarf companion and planetary debris. These 'recycled' pulsars are spun up by accretion and are among the most

precise timekeepers in the universe.

**Intermediate Mass BH**
A rare intermediate-mass black hole (400 M⊙) in a globular cluster environment with dense stellar populations. These elusive objects bridge the gap between stellar-mass and supermassive black holes.

**Micro BH Swarm**
A dynamic swarm of small black holes (0.6-1.8 M⊙) with planets and gas giants in chaotic orbital dance. Watch as these stellar-mass black holes interact, merge, and create complex gravitational resonances.

**Hungry Hungry Holes**
Four black holes at the corners of a square, pulling stars from a shared central cluster.

**Black Hole Billiards**
A few small black holes orbiting a supermassive one, perturbing each other and creating chaotic motion.

## 7.4 Exoplanets

**TRAPPIST-1 System**
A compact planetary system with seven Earth-sized worlds orbiting a cool red dwarf star just 40 light-years away. All planets are packed close to their tiny sun, with several in the habitable zone. Can you keep this delicate system stable?

**Transit Lab: HD 209458 b**
The first exoplanet ever caught crossing its star, found in 1999 after radial velocities said where to look. A hot Jupiter on a 3.5-day orbit, drawn here at true relative scale: the star is 1.155 solar radii, the planet 1.38 Jupiter radii, and the silhouette on screen is the same 12% radius ratio the light curve reports. Open the Light Curve panel and watch the 1.7% dip repeat.

**Exoplanet Characterization Lab**
HD 209458 again, but with the star free to move. In the Transit Lab it is pinned so the light curve stays centered; here both bodies orbit their common center of mass, which is what the radial-velocity and astrometry instruments need in order to measure anything. The star circles a point 2.7 millionths of an AU away at 84 meters per second: far too small to see and easily large enough to detect. Open Radial Velocity and watch the wobble that found this planet.

**Blended Binary: a hidden companion**
The same star and planet as the Transit Lab, with a second star half a magnitude fainter sitting 300 AU away: far too close on the sky for a survey telescope to separate, and well inside one photometric aperture. Its light fills in part of the dip, so the transit measures shallower and the planet looks smaller than it is. Correcting for exactly this effect is what high-resolution imaging surveys of planet hosts are for.

**Exoplanet Lab**
A diverse collection of 120+ exoplanets, gas giants, and even pulsar planets around various stellar hosts. Explore the incredible diversity of planetary systems with interactive orbital mechanics and planetary interactions.

## 7.5 Galaxies & Clusters

**Spiral Galaxy: what we expected**
A galactic bulge with ninety stars orbiting it, each launched at exactly the speed the visible mass says it should have. This is the prediction, not the observation: with the mass concentrated in the middle, orbital speed falls away as the inverse square root of radius, the same way it does across the Solar System. Open the Rotation Curve panel and read the slope. Then load Milky Way Rotation and read it again.

**Milky Way Rotation: what we actually see**
The same disc, with every star moving at the same speed no matter how far out it is. That is what telescopes measure in real spiral galaxies, and it is far too fast for the stars you can see to hold on to: this scenario starts with a dark-matter halo switched on, because without one the disc does not survive. Open the Rotation Curve panel and switch the halo off to watch it come apart.

**Coma Cluster: Zwicky, 1933**
Twenty-four galaxies swarming in a bound cluster, on randomly oriented orbits, named after the members of the real Coma Cluster that Fritz Zwicky measured. He added up the light, added up the motions, and found the second answer hundreds of times larger than the first. He called the difference dunkle Materie and was ignored for forty years. Select a galaxy to read its speed, and work the same calculation he did.

**Dense Star Cluster**
A gravitationally bound collection of main-sequence stars, evolved giants, and stellar remnants with mutual gravitational interactions. Watch stellar encounters, binary formation, and the dynamic evolution of this stellar community over time.

**Galactic Center**
A supermassive black hole (4000 M$\odot$) surrounded by high-velocity stars, stellar remnants, and dense stellar populations. Experience the extreme gravitational environment with spectacular accretion, jets, and relativistic effects.

**Galactic Collision**
Two supermassive black holes (1.2M & 1.0M M$\odot$) with hundreds of stars representing galactic cores in collision. Witness the formation of tidal streams, stellar disruption, and the eventual merger of supermassive black holes.

**Quasar Cannon**
A supermassive black hole is actively feeding on a dense star cluster. Watch a beam of light form as stars spiral inward.

**The Pinwheel Galaxy Core**
Two intermediate black holes in the center of a stellar disk. The disk forms a rotating pinwheel pattern as stars are slung around.

**Tidal Arm Tango**
Two black holes dance past each other, flinging stars into massive tidal arms like colliding galaxies.

## 7.6 Habitability

**Habitable Zone Lab: the inner Solar System, with the zone drawn**
The Sun with Venus, Earth, Mars and Ceres on their real orbits, and the circumstellar habitable zone drawn around the star from a published prescription. Venus sits inside the inner edge and Mars outside the outer one on the conservative definition, and only one of the four has liquid water on its surface today. Switch the Habitable Zone Model setting to see the optimistic band, which reaches out past Mars.

## 7.7 Orbits & Kepler

**Gravity Assist Lab: one planet, one spacecraft, nothing else**
A rogue planet of five Jupiter masses drifting through empty space, and an Earth-mass spacecraft crossing its path. No star, which is the point: with nothing else in the universe the planet moves in a straight line and its frame is exactly inertial, so the strangest fact about a gravity assist can be stated exactly rather than approximately. The spacecraft leaves at the same

speed relative to the planet that it arrived with, and at a completely different speed relative to everything else.

**Gravity Assist: the same flyby, with a Sun to steal from**
The same planet, now on a circular orbit five AU from a star, and a spacecraft that meets it. This is where the energy comes from: the spacecraft leaves faster around the star than it arrived, and the planet is slowed by exactly the momentum it gained. Everything here is approximate - the planet is accelerating, so its frame is not inertial and the encounter is only locally two-body - and the panel reports how approximate rather than hiding it.

**Lagrange Point Lab: two stars, one circular orbit, one test particle**
A sunlike star and a companion a thirtieth its mass, on an exactly circular orbit eight astronomical units across, with a tracer light enough that neither notices it. This is the circular restricted three-body problem: the simplest system with Lagrange points, forbidden regions and a conserved Jacobi constant, and the only arrangement in which any of those are exactly true.

**Orbital Transfer Lab: one star, one spacecraft, one destination**
A spacecraft on a circular orbit at 1 AU and a station on a circular orbit at 2.5 AU, about a single sunlike star with nothing else in the system. Circular and coplanar is what makes the transfer between them exactly solvable, so a burn planned by hand can be checked against the arithmetic rather than only watched.

**Kepler's 2nd Law - Equal Areas**
A planet in a nearly circular orbit and an eccentric orbiter around a central star. The area sweep visualization starts automatically for the eccentric body - watch how the wedges change shape but maintain equal area, showing why objects move faster at periapsis than at apoapsis.

## 7.8 Orbital Resonance

**Galilean Resonance: the Laplace lock**
Io, Europa and Ganymede on the 4:2:1 chain Laplace explained in 1805, with Callisto outside it for contrast. The periods are not exactly 4:2:1 — Europa takes 0.37% longer than two Io years — and that near miss is the point: what holds them is the Laplace argument, which stays near 180 degrees instead of running through every value, so the three are never all in conjunction. Callisto sits within 0.03% of 7:3 with Ganymede and is in no resonance at all. A scale model, 100 times life size.

**Broken Laplace Resonance: one percent out**
The same four moons with one number changed: Europa starts one percent further from Jupiter. That is a hundred times wider than the resonance can hold, and the lock fails — the Laplace argument stops swinging about 180 degrees and starts going all the way round, once every forty-six Io orbits. Run it beside Galilean Resonance and the difference between a system that is locked and one that merely has convenient periods is on screen in under a minute.

**Pluto and Neptune: the 3:2 that protects**
Pluto's orbit crosses Neptune's, and they have never come close. The 3:2 resonance is why: Pluto goes round twice for every three Neptune years, and the resonant argument librates about 180 degrees rather than circulating — so every conjunction happens near Pluto's aphelion, far outside Neptune's reach. A third body runs almost the same orbit from outside the resonance; watch what becomes of it. True scale; one second is about 270 years.

**Jupiter Trojans: the 1:1 co-orbital resonance**
Two of the five Lagrange points are places a body can sit still relative to Jupiter forever, and thousands of asteroids do. A probe placed exactly at L4 does not move in Jupiter's rotating frame; the Trojan 617 Patroclus loops around L5 once every twelve and a half Jupiter years. A third probe starts one degree from L3 — an equilibrium too, and unstable — and leaves. The

fourth runs a wider ordinary orbit and is in no resonance at all.

## 7.9 Solar System

**Solar System**
A simulation of our Solar System featuring real planets with correct masses, orbital distances, diameters, and colors. Includes Mercury, Venus, Earth, Mars, Jupiter, Saturn, Uranus, Neptune with their actual properties, plus real asteroids (Ceres, Vesta, Pallas) and famous comets (Halley, Hale-Bopp, Hyakutake) with authentic orbital periods and characteristics.

**Retrograde Mars: the loop that needed epicycles**
The Sun, Earth and Mars at their real distances and periods, and nothing else. Watched from outside, both planets go round the Sun the same way and never turn back. Switch the reference frame to Earth, in Tools, and Mars stops circling and starts drawing a loop that doubles back on itself. Nothing about the physics changed; only the frame did. That loop is the observation Ptolemy built epicycles to reproduce and Copernicus explained away, and here you can turn it on and off with one control.

**Earth-Moon System**
A detailed simulation of the Earth-Moon system with accurate masses, orbital mechanics, and realistic appearances. Features Earth with its blue oceans and green continents, and the Moon with its characteristic gray surface and craters. Perfect for studying orbital dynamics and tidal effects.

**Interstellar Visitor: 1I/'Oumuamua**
The first object ever seen passing through the Solar System from somewhere else, on its measured orbit: perihelion inside Mercury, eccentricity 1.20, and 87.7 km/s at closest approach. Earth is shown for scale. Select the visitor and look at the sign of its total energy: it is positive, which is the whole story. It is not bound to the Sun, it was never going to stay, and it will not be back.

**Kuiper Belt**
An accurate simulation of our Solar System's Kuiper Belt featuring real dwarf planets (Pluto, Eris, Haumea, Makemake), large KBOs (Quaoar, Sedna, Orcus, Varuna), and smaller objects (Ixion, Huya, 2002 AW197) with realistic masses and orbital properties.

**Ring Gap: a moon and a ring**
A Saturn-mass planet, a ring of 200 particles, a resonant moon whose 2:1 resonance falls inside the ring and a shepherd moon inside it. A scale model: lengths are stretched and G carries the exaggeration, and both moons are far heavier than Mimas and Pan so that twenty orbits can show the effect. The ring thins at the resonance and clears a lane about the shepherd; it does not empty, because its particles do not collide.

**Roche Pass: a clump inside the limit**
A planet and a loose clump of 60 particles on an orbit whose closest point is 1.3 planet radii from the planet's center, inside both the rigid and the fluid Roche limit for the clump's density, as Shoemaker-Levy 9 was at Jupiter. A scale model with Newtonian gravity only: the clump is a softened swarm, so it is drawn into a stream at the pass and also loosens slowly by itself.

## 7.10 Stellar Evolution

**Supernova Remnant**
The explosive aftermath of a massive star death: a neutron star surrounded by high-velocity debris, shocked planets, and disrupted gas giants. Experience the violent and energetic environment left behind by stellar death.

**Stellar Nursery**
A dense cluster of young stars around a proto-black hole. Watch interactions and ejections as

the cluster evolves.

### 7.11 Tides & Disruption

**Tidal Disruption**
Multiple objects approach a supermassive black hole (2000 M⊙) and are torn apart by extreme tidal forces. Watch as planets and gas giants are stretched, disrupted, and either ejected or accreted, creating spectacular debris streams.

## 8 Guided investigations

91 investigations run in the simulation, with 1289 steps between them, 852 of those graded, against 433 stated learning objectives. Open them from Learn in the rail or from the Library. The Observatory, the 3-D lab and the mission lab add 17 guided investigations of their own, which is why the Library lists 108 in all.

Every investigation follows the same discipline:

1. It asks for a **prediction** before it shows anything.
2. It hands over an **instrument** and asks for a measurement.
3. It **plots the student's own readings** back to them.
4. It **saves progress** in the browser, so a session can be left and resumed.
5. It **exports a lab report** as a PDF, which the student submits through whatever system the course already uses.

Nothing is submitted anywhere by Gravitas itself. There is no server to submit to.

| Investigation | Steps | Graded | Time |
|---|---|---|---|
| Kepler's Laws | 23 | 13 | 35-45 min |
| Why Mars Goes Backwards | 33 | 18 | 35-45 min |
| Finding Planets by Their Shadows | 29 | 13 | 50-70 min |
| Bound, Unbound and Escape | 24 | 10 | 35-45 min |
| Weighing the Stars | 36 | 18 | 35-45 min |
| Black Holes by the Numbers | 30 | 18 | 35-45 min |
| Finding Planets by Their Tug | 38 | 20 | 45-55 min |
| The Goldilocks Question | 37 | 19 | 40-50 min |
| The Missing Mass | 33 | 17 | 45-60 min |
| Tides | 30 | 16 | 35-45 min |
| The Butterfly Effect in Space | 28 | 13 | 55-70 min |
| When Orbits Lock | 34 | 17 | 55-70 min |
| Can You Detect This Planet? | 26 | 13 | 30-35 min |
| Design the Schedule | 15 | 8 | 35-40 min |
| Planets in Binary Stars | 37 | 21 | 40-50 min |
| Where Does a Gravity Assist Get Its Speed? | 22 | 13 | 15-20 min |
| Getting There From Here | 22 | 14 | 20-25 min |
| Where Can It Get To? | 20 | 9 | 25-30 min |
| What Is a Gravitational Wave? | 28 | 15 | 30-40 min |
| Listening to Spacetime | 32 | 18 | 75-90 min |

| Investigation | Steps | Graded | Time |
|---|---|---|---|
| A Universe of Stars | 37 | 31 | 70-90 min |
| Lives of Stars | 35 | 29 | 80-100 min |
| Twelve Nights | 13 | 8 | 40-50 min |
| What If Gravity Were Not Inverse Square? | 21 | 14 | 45-60 min |
| Color and Temperature | 11 | 9 | 15-25 min |
| The Turning Sky | 10 | 8 | 10-20 min |
| The Sun Through the Year | 9 | 7 | 10-20 min |
| Phases and Eclipses | 10 | 8 | 10-20 min |
| Wanderers on the Sky | 11 | 9 | 12-22 min |
| Plan a Night at the Telescope | 10 | 8 | 10-20 min |
| Lines and Motion | 11 | 9 | 15-25 min |
| What a Spectrum Is Made Of | 10 | 8 | 15-25 min |
| The Sun's Motion | 10 | 7 | 25-40 min |
| The Rotating Disk | 12 | 9 | 25-40 min |
| Cluster Membership by Motion | 11 | 9 | 25-40 min |
| Inside a Star | 9 | 7 | 15-25 min |
| Scaling the Main Sequence | 10 | 8 | 20-25 min |
| The Limit | 9 | 7 | 15-25 min |
| The Expanding Universe | 9 | 7 | 12-20 min |
| Standard Candles | 7 | 5 | 12-20 min |
| The Shape of Expansion | 9 | 7 | 25-40 min |
| Where Galaxies Live | 9 | 7 | 12-22 min |
| What a CCD Records | 11 | 9 | 10-18 min |
| Signal, Noise and Exposure | 10 | 8 | 10-18 min |
| A Variable Star From Frames | 8 | 6 | 10-18 min |
| The Beat of a Star | 13 | 11 | 25-40 min |
| Leavitt's Law | 8 | 6 | 8-15 min |
| The First Rung | 8 | 6 | 12-20 min |
| Which Kind of Variable | 9 | 7 | 12-22 min |
| Bending Light | 8 | 6 | 20-30 min |
| Einstein Rings and Arcs | 9 | 7 | 25-40 min |
| A Lens Passing By | 10 | 8 | 25-40 min |
| Who Gets Counted | 10 | 8 | 20-35 min |
| Why the Bright Ones Win | 10 | 8 | 25-40 min |
| The Completeness Curve | 10 | 8 | 25-40 min |
| Integrators in Practice | 8 | 6 | 30-40 min |
| Extending an Experiment | 8 | 6 | 30-40 min |
| Refitting a Transit | 8 | 6 | 30-40 min |
| Two Periodograms | 8 | 6 | 30-40 min |
| Sampling a Posterior | 8 | 6 | 20-30 min |
| What Is a Measurement? | 12 | 10 | 20-30 min |
| What Does ± Claim? | 13 | 11 | 25-40 min |
| When the Method Must Refuse | 13 | 11 | 25-40 min |
| Measuring the Moon | 8 | 6 | 15-25 min |
| How Far Is That Star? | 10 | 8 | 20-30 min |
| A Binary Star's Orbit | 9 | 7 | 20-30 min |

| Investigation | Steps | Graded | Time |
|---|---|---|---|
| Which Surface Is Older? | 10 | 8 | 15-25 min |
| Craters by Size | 9 | 7 | 10-20 min |
| What a Crater Diameter Is Worth | 8 | 6 | 10-20 min |
| What Is in This Field? | 10 | 8 | 20-30 min |
| Weighing Jupiter From Its Moons | 7 | 5 | 25-40 min |
| Why Rings? | 8 | 4 | 15-25 min |
| Tides That Move the Moon | 6 | 4 | 20-30 min |
| Io's Fire | 6 | 4 | 15-25 min |
| The Gaps in the Belt | 8 | 6 | 12-16 min |
| How Fast Does It Spin? | 6 | 4 | 8-12 min |
| Why Venus Is Hot | 9 | 7 | 15-25 min |
| Two Greenhouse Models | 7 | 5 | 10-20 min |
| The Turning Sun | 9 | 7 | 20-30 min |
| The Sun's Power | 6 | 4 | 15-25 min |
| How Hot Is the Day Side? | 10 | 8 | 15-25 min |
| Phases and Albedo | 7 | 5 | 15-25 min |
| Moons of the Giants | 10 | 8 | 15-25 min |
| The Scale of the Solar System | 8 | 6 | 15-25 min |
| Comets and the Sun | 9 | 6 | 15-25 min |
| What a Telescope Resolves | 7 | 5 | 15-25 min |
| Why Two Lines Become One | 6 | 4 | 10-20 min |
| The Saturated Star | 9 | 7 | 10-20 min |
| Counts to Magnitudes | 6 | 4 | 10-20 min |
| Plan an Exposure | 11 | 9 | 12-22 min |
| Why the Period Is Wrong | 8 | 6 | 12-22 min |

## 8.1 Kepler's Laws

*Measure the shape, pacing and timing of real orbits* 23 steps, 13 of them graded, 35-45 min, Introductory astronomy.

undefined

## 8.2 Why Mars Goes Backwards

*Change the frame, and fourteen centuries of epicycles fall away* 33 steps, 18 of them graded, 35-45 min, Introductory astronomy.

undefined

## 8.3 Finding Planets by Their Shadows

*Measure a transit, weigh what it tells you, and find what is hiding* 29 steps, 13 of them graded, 50-70 min, Introductory astronomy.

undefined

## 8.4 Bound, Unbound and Escape

*Find out what decides whether something comes back* 24 steps, 10 of them graded, 35-45 min, Introductory astronomy.

undefined

### 8.5 Weighing the Stars

*Use an orbit to measure something you cannot put on a scale* 36 steps, 18 of them graded, 35-45 min, Introductory astronomy.

undefined

### 8.6 Black Holes by the Numbers

*Make a black hole bigger and discover some surprising rules* 30 steps, 18 of them graded, 35-45 min, Introductory astronomy.

undefined

### 8.7 Finding Planets by Their Tug

*Watch a star wobble, weigh its planet, and combine the clues* 38 steps, 20 of them graded, 45-55 min, Introductory astronomy.

undefined

### 8.8 The Goldilocks Question

*Move a planet, change its star, and decide what 'habitable' really means* 37 steps, 19 of them graded, 40-50 min, Introductory astronomy.

undefined

### 8.9 The Missing Mass

*Weigh a system twice, and find the two answers do not agree* 33 steps, 17 of them graded, 45-60 min, Introductory astronomy.

undefined

### 8.10 Tides

*Stretch a world, move a moon, and discover why gravity can tear objects apart* 30 steps, 16 of them graded, 35-45 min, Introductory astronomy.

undefined

### 8.11 The Butterfly Effect in Space

*Run the same system twice and find out how long the answer lasts* 28 steps, 13 of them graded, 55-70 min, Introductory astronomy.

undefined

### 8.12 When Orbits Lock

*A ratio is a hint. Find out what counts as proof* 34 steps, 17 of them graded, 55-70 min, Introductory astronomy.

undefined

## 8.13 Can You Detect This Planet?

*Two methods, the same problem: the answer was decided before the data arrived* 26 steps, 13 of them graded, 30-35 min, Introductory astronomy.

undefined

## 8.14 Design the Schedule

*Eight nights on the real instrument, and the times you choose decide the answer* 15 steps, 8 of them graded, 35-40 min, Introductory astronomy.

undefined

## 8.15 Planets in Binary Stars

*What survives around two stars, and how you would know* 37 steps, 21 of them graded, 40-50 min, Introductory astronomy.

undefined

## 8.16 Where Does a Gravity Assist Get Its Speed?

*The same flyby, measured in two frames, with two different answers* 22 steps, 13 of them graded, 15-20 min, Introductory astronomy.

undefined

## 8.17 Getting There From Here

*Two burns, a long coast, and the arithmetic that decides both* 22 steps, 14 of them graded, 20-25 min, Introductory astronomy.

undefined

## 8.18 Where Can It Get To?

*Forbidden regions, five balance points, and one conserved number* 20 steps, 9 of them graded, 25-30 min, Introductory astronomy.

undefined

## 8.19 What Is a Gravitational Wave?

*A first look at what moves, what travels and what a detector feels* 28 steps, 15 of them graded, 30-40 min, Beginner, no physics background needed.

undefined

## 8.20 Listening to Spacetime

*Work out what made a signal, then check it against the real thing* 32 steps, 18 of them graded, 75-90 min, Introductory astronomy.

undefined

## 8.21 A Universe of Stars

*Size, color and the H-R diagram, from eight modeled stars* 37 steps, 31 of them graded, 70-90 min, Introductory astronomy.

undefined

## 8.22 Lives of Stars

*From clouds to cosmic remnants, along eight published tracks* 35 steps, 29 of them graded, 80-100 min, Introductory astronomy.

undefined

## 8.23 Twelve Nights

*You get to choose when to look, and most of the choice has already been made* 13 steps, 8 of them graded, 40-50 min, Introductory astronomy.

undefined

## 8.24 What If Gravity Were Not Inverse Square?

*Change the exponent, and find out what depended on it* 21 steps, 14 of them graded, 45-60 min, Introductory astronomy.

undefined

## 8.25 Color and Temperature

*Predict, then measure, what a hot glowing thing looks like* 11 steps, 9 of them graded, 15-25 min, Introductory astronomy.

undefined

## 8.26 The Turning Sky

*Measure why a star rises four minutes earlier every night* 10 steps, 8 of them graded, 10-20 min, Introductory astronomy.

undefined

## 8.27 The Sun Through the Year

*Measure the Sun's height, the length of the day and why June is warm* 9 steps, 7 of them graded, 10-20 min, Introductory astronomy.

undefined

## 8.28 Phases and Eclipses

*Measure how much of the Moon is lit, then find out why eclipses are rare* 10 steps, 8 of them graded, 10-20 min, Introductory astronomy.

undefined

## 8.29 Wanderers on the Sky

*Follow Mars, Venus and Jupiter through their loops, from the sky and from above* 11 steps, 9 of them graded, 12-22 min, Introductory astronomy.

undefined

## 8.30 Plan a Night at the Telescope

*Choose targets for a date and a site under twilight, airmass and Moon limits* 10 steps, 8 of them graded, 10-20 min, Introductory astronomy.

undefined

### 8.31 Lines and Motion

*Read a star's lines, then measure how fast it moves* 11 steps, 9 of them graded, 15-25 min, Introductory astronomy.

undefined

### 8.32 What a Spectrum Is Made Of

*Why one gas makes dark lines, bright lines, or none* 10 steps, 8 of them graded, 15-25 min, Introductory astronomy.

undefined

### 8.33 The Sun's Motion

*Measuring the solar apex from nearby stars in Gaia* 10 steps, 7 of them graded, 25-40 min, Introductory astronomy.

undefined

### 8.34 The Rotating Disk

*The Oort constants and the Milky Way's rotation curve, from data, then a mass model* 12 steps, 9 of them graded, 25-40 min, Introductory astronomy.

undefined

### 8.35 Cluster Membership by Motion

*Finding the stars of NGC 2420 in Gaia, as probabilities* 11 steps, 9 of them graded, 25-40 min, Introductory astronomy.

undefined

### 8.36 Inside a Star

*Hydrostatic equilibrium, a polytrope, and why the center is hot* 9 steps, 7 of them graded, 15-25 min, Introductory astronomy.

undefined

### 8.37 Scaling the Main Sequence

*What a scaling argument predicts, and the MIST models say* 10 steps, 8 of them graded, 20-25 min, Introductory astronomy.

undefined

### 8.38 The Limit

*White-dwarf mass and radius, and why 1.4 solar masses* 9 steps, 7 of them graded, 15-25 min, Introductory astronomy.

undefined

### 8.39 The Expanding Universe

*Hubble's line of 1929, and what the same line says of 955 modern supernovae* 9 steps, 7 of them graded, 12-20 min, Introductory astronomy.

undefined

## 8.40 Standard Candles

*From Cepheids to supernovae: how an error in one rung becomes an error in the Hubble constant* 7 steps, 5 of them graded, 12-20 min, Introductory astronomy.

undefined

## 8.41 The Shape of Expansion

*Fitting the matter density to 1,829 supernovae, and what the fit can and cannot say* 9 steps, 7 of them graded, 25-40 min, Introductory astronomy.

undefined

## 8.42 Where Galaxies Live

*A wedge of the sky from the Sloan survey: the scale of structure, and a thinning that is not a void* 9 steps, 7 of them graded, 12-22 min, Introductory astronomy.

undefined

## 8.43 What a CCD Records

*Bias, dark and flat frames, and why a star's light needs all three taken out before it is a measurement* 11 steps, 9 of them graded, 10-18 min, Introductory astronomy.

undefined

## 8.44 Signal, Noise and Exposure

*How long to expose to reach a signal-to-noise, and why four times the light is not four times the quality* 10 steps, 8 of them graded, 10-18 min, Introductory astronomy.

undefined

## 8.45 A Variable Star From Frames

*Differential photometry of forty calibrated frames: why a comparison star removes the cloud, and how to know a variation is real* 8 steps, 6 of them graded, 10-18 min, Introductory astronomy.

undefined

## 8.46 The Beat of a Star

*Finding a pulsating star's period, and the false periods of a telescope that looks once a night* 13 steps, 11 of them graded, 25-40 min, Introductory astronomy.

undefined

## 8.47 Leavitt's Law

*Fitting the period-luminosity relation of Cepheids in the Large Magellanic Cloud* 8 steps, 6 of them graded, 8-15 min, Introductory astronomy.

undefined

## 8.48 The First Rung

*The distance to two galaxies from their Cepheids, with every error accounted for* 8 steps, 6 of them graded, 12-20 min, Introductory astronomy.

undefined

### 8.49 Which Kind of Variable

*Telling five kinds of variable star apart, and reading a confusion table instead of one answer* 9 steps, 7 of them graded, 12-22 min, Introductory astronomy.

undefined

### 8.50 Bending Light

*How far a mass turns a ray of light, and the Sun's 1.75 arcseconds* 8 steps, 6 of them graded, 20-30 min, Introductory astronomy.

undefined

### 8.51 Einstein Rings and Arcs

*The geometry of strong lensing, and a ring as a way to weigh a lens* 9 steps, 7 of them graded, 25-40 min, Introductory astronomy.

undefined

### 8.52 A Lens Passing By

*Microlensing light curves, and what one curve cannot tell you* 10 steps, 8 of them graded, 25-40 min, Introductory astronomy.

undefined

### 8.53 Who Gets Counted

*What a magnitude limit does to a census of the stars* 10 steps, 8 of them graded, 20-35 min, Introductory astronomy.

undefined

### 8.54 Why the Bright Ones Win

*Malmquist bias, measured against a truth we know* 10 steps, 8 of them graded, 25-40 min, Introductory astronomy.

undefined

### 8.55 The Completeness Curve

*Correcting a luminosity function, and saying what bias is left* 10 steps, 8 of them graded, 25-40 min, Introductory astronomy.

undefined

### 8.56 Integrators in Practice

*Four ways to take a step, compared by the energy each one loses on a real system, in Python on the compute page* 8 steps, 6 of them graded, 30-40 min, Introductory astronomy.

undefined

### 8.57 Extending an Experiment

*From twenty planets that the engine ran to a population of two hundred thousand, with every uncertainty, in Python on the compute page* 8 steps, 6 of them graded, 30-40 min, Introductory astronomy.

undefined

### 8.58 Refitting a Transit

*The same model and the same light curve as the Observatory's fit panel, fitted again by a different route in Python, to see what the route changes* 8 steps, 6 of them graded, 30-40 min, Introductory astronomy.

undefined

### 8.59 Two Periodograms

*A Lomb-Scargle and a box least squares periodogram of the same light curve, computed with Astropy in Python, and what a period from a periodogram can claim* 8 steps, 6 of them graded, 30-40 min, Introductory astronomy.

undefined

### 8.60 Sampling a Posterior

*A transit's planet size and impact parameter sampled in Python, checked for convergence, and compared with the profile interval of the same problem* 8 steps, 6 of them graded, 20-30 min, Introductory astronomy.

undefined

### 8.61 What Is a Measurement?

*Marking one star again and again, and what the marks can tell you* 12 steps, 10 of them graded, 20-30 min, Introductory astronomy.

undefined

### 8.62 What Does ± Claim?

*Checking a stated uncertainty by repeating the measurement* 13 steps, 11 of them graded, 25-40 min, Introductory astronomy.

undefined

### 8.63 When the Method Must Refuse

*A fit that fixes one combination, and a model that does not describe the data* 13 steps, 11 of them graded, 25-40 min, Majors and upper-level astronomy.

undefined

### 8.64 Measuring the Moon

*Marks on a real picture, a scale from a known length, and a size in kilometers* 8 steps, 6 of them graded, 15-25 min, Introductory astronomy.

undefined

### 8.65 How Far Is That Star?

*A shift, a motion and a parallax from fourteen positions of one star* 10 steps, 8 of them graded, 20-30 min, Majors and upper-level astronomy.

undefined

### 8.66 A Binary Star's Orbit

*Positions over sixteen years, the orbit they fix, and the mass of the pair* 9 steps, 7 of them graded, 20-30 min, Majors and upper-level astronomy.

undefined

### 8.67 Which Surface Is Older?

*Craters per area in two regions of the Moon and of Mars, and the ratio that says which is older* 10 steps, 8 of them graded, 15-25 min, Introductory astronomy.

undefined

### 8.68 Craters by Size

*The size-frequency plot with its error bars, the slope with its interval, and a count that stops short* 9 steps, 7 of them graded, 10-20 min, Majors and upper-level astronomy.

undefined

### 8.69 What a Crater Diameter Is Worth

*One circle on a map, the four terms of its uncertainty, and why no age in years is reported* 8 steps, 6 of them graded, 10-20 min, Upper-level astronomy.

undefined

### 8.70 What Is in This Field?

*The sources a detector lists, the width of the stars, and the curve that says how faint a search can see* 10 steps, 8 of them graded, 20-30 min, Majors and upper-level astronomy.

undefined

### 8.71 Weighing Jupiter From Its Moons

*Four moons, a cadence that can measure them, and Kepler's third law* 7 steps, 5 of them graded, 25-40 min, Introductory to upper-level astronomy.

undefined

### 8.72 Why Rings?

*Where a moon falls apart, and where a moon's pull leaves a mark in a ring* 8 steps, 4 of them graded, 15-25 min, Introductory astronomy.

undefined

### 8.73 Tides That Move the Moon

*The Moon's recession, the lengthening day, and the model that fails when run backward* 6 steps, 4 of them graded, 20-30 min, Majors and upper-level astronomy.

undefined

### 8.74 Io's Fire

*Tidal heating as energy per orbit, bounded by what nobody knows about Io's interior* 6 steps, 4 of them graded, 15-25 min, Majors and upper-level astronomy.

undefined

## 8.75 The Gaps in the Belt

*A histogram of 8,886 numbered asteroids, a smooth baseline fitted outside the resonance windows, and how deep each gap is, with an interval* 8 steps, 6 of them graded, 12-16 min, Introductory to upper-level astronomy.

undefined

## 8.76 How Fast Does It Spin?

*A synthetic light curve, the rotation period from its periodogram, the amplitude, and what the amplitude bounds about the shape* 6 steps, 4 of them graded, 8-12 min, Introductory to upper-level astronomy.

undefined

## 8.77 Why Venus Is Hot

*What sunlight alone gives a planet, the gap it leaves, and what closes it* 9 steps, 7 of them graded, 15-25 min, Introductory astronomy.

undefined

## 8.78 Two Greenhouse Models

*A single layer and a gray atmosphere, the same planets, and why the numbers differ* 7 steps, 5 of them graded, 10-20 min, Majors and upper-level astronomy.

undefined

## 8.79 The Turning Sun

*Fourteen days of pictures, one spot group followed across the disc, and the Sun's rotation from it* 9 steps, 7 of them graded, 20-30 min, Introductory to upper-level astronomy.

undefined

## 8.80 The Sun's Power

*A flux measured at the Earth, brought to one astronomical unit, and the Sun's luminosity and temperature from it* 6 steps, 4 of them graded, 15-25 min, Introductory to upper-level astronomy.

undefined

## 8.81 How Hot Is the Day Side?

*An eclipse depth, a blackbody, and what a brightness temperature leaves out* 10 steps, 8 of them graded, 15-25 min, Majors and upper-level astronomy.

undefined

## 8.82 Phases and Albedo

*A planet's phase curve, a Lambert sphere, and the albedo it gives* 7 steps, 5 of them graded, 15-25 min, Majors and upper-level astronomy.

undefined

## 8.83 Moons of the Giants

*Three periods from one set of frames, and the combination that is zero when a family of moons keeps step* 10 steps, 8 of them graded, 15-25 min, Introductory astronomy.

undefined

## 8.84 The Scale of the Solar System

*True distances from the sandbox, the light travel time, and a scale model in which Earth is a marble* 8 steps, 6 of them graded, 15-25 min, Introductory astronomy.

undefined

## 8.85 Comets and the Sun

*A model comet's orbit from the sandbox, how hot it gets at each end, the real Halley as a transfer case, and which way its tails point* 9 steps, 6 of them graded, 15-25 min, Introductory astronomy.

undefined

## 8.86 What a Telescope Resolves

*Which aperture shows two stars as two, when the faint star appears, and why the width of a star is not the limit of the telescope* 7 steps, 5 of them graded, 15-25 min, Introductory astronomy.

undefined

## 8.87 Why Two Lines Become One

*Why a spectrograph can show two lines as one, what pixels can and cannot fix, how a wavelength scale is calibrated from an arc and what it costs in light* 6 steps, 4 of them graded, 10-20 min, Introductory astronomy.

undefined

## 8.88 The Saturated Star

*Why the brightest star in the frame has the wrong magnitude, what bleeding and cosmic rays do to its neighbors, and where a measurement is safe* 9 steps, 7 of them graded, 10-20 min, Introductory astronomy.

undefined

## 8.89 Counts to Magnitudes

*Why a star looks fainter low in the sky, how a night of standard stars turns counts into magnitudes, and what the calibration cannot tell you* 6 steps, 4 of them graded, 10-20 min, Introductory astronomy.

undefined

## 8.90 Plan an Exposure

*How long to expose a faint star for a signal-to-noise of 20, what limits it, what splitting it, the air and the moon cost, and what a frame says back* 11 steps, 9 of them graded, 12-22 min, Majors and upper-level astronomy.

undefined

## 8.91 Why the Period Is Wrong

*Why looking once a night cannot tell a period from its aliases, how to design a cadence that can, and what a plan costs* 8 steps, 6 of them graded, 12-22 min, Introductory astronomy.

undefined

# 9 Running a controlled experiment

The A/B Bench, in Tools, is for the question a sandbox cannot otherwise answer: *what does changing this one thing actually do?*

Watching a simulation twice does not answer it, because the second run never starts quite where the first did. The bench makes it start there exactly.

1. **Capture start.** The world as it stands becomes the experiment's origin — its seed, its clock, its bodies and their identities, the integrator, the reference frame and the observer.
2. **Name it, and choose what to measure.** Bodies as chips, quantities as checkboxes. A quantity needing two bodies stays grayed out until two are chosen.
3. **Record Run A.**
4. **Return to start**, exactly. The simulation holds there.
5. **Change one thing.**
6. **Record Run B.**
7. **Compare.** The two runs are drawn on one simulated-time axis, with a table of absolute and fractional differences.

What can be measured: position and trajectory, separation between two bodies, speed and velocity, distance from a chosen primary, orbital period, closest approach, total energy, angular momentum, and the numerical drift in the last two.

## 9.1 What the bench will tell you off for

The bench names the parameter that differed between the runs, and it says so when more than one did — a comparison with two independent variables cannot say which one caused the difference. You can confirm that you meant it, because a deliberate multivariable comparison is a legitimate thing to run.

It does not count elapsed time, the camera, the theme or a trail length as variables. It *does* count the simulation speed, because the speed control scales the integration step and therefore changes the numerical error a run reports, and it counts the seed and the scenario, because either one makes it a different world.

## 9.2 Honest sampling

Two runs are never sampled at the same instants: an animation frame takes as long as it takes, and a scenario that substeps advances its clock several times inside one frame. So the bench records each sample with its own simulated time and aligns the two runs on that, interpolating between samples and refusing to extrapolate past the end of either. Where the runs do not overlap in simulated time it says there is no comparison, rather than drawing one.

The recording indicator shows both the number of samples and the simulated seconds they cover, because those two are not interchangeable.

## 9.3 Keeping and sharing an experiment

Experiments can be renamed, duplicated, deleted, saved in the browser and reopened. Saved experiments are bounded — if one is too large, or the store is full, the bench says which and points you at the export, rather than failing quietly.

Exporting gives two files: a **CSV** of both runs, one row per sample, with the run named in a column and the unit in every column heading; and a **JSON manifest** carrying the experiment's definition and its provenance — scenario, seed, integrator, timestep, units, a hash of the initial

state, what was measured and what changed. The manifest is the file to paste into a lab report, and reopening it restores the experiment's setup so somebody else can run it themselves.

Share setup puts the same thing in a link. The link carries the starting state and the A/B difference; it never carries the recorded runs, so what arrives is an experiment to perform rather than somebody's results.

# 10 Capturing, exporting and sharing

## 10.1 Screenshots and clips

Capture → Screenshot (or `P`) saves a PNG with the scenario's name, the scale bar and the elapsed simulated time drawn into the image itself, so a figure in a lab report documents its own scale and its own clock without a caption having to supply them.

Record Clip records a stretch of the run to a video file: H.264 in MP4 where the browser can encode it, WebM where it cannot. The same three facts are burned into every frame. A recording indicator shows the elapsed time and the size, and a take stops itself at three minutes or 80 MB, whichever comes first.

## 10.2 Data

Export CSV (or `E`) writes the recorded timeline: positions, velocities and energies over time, plus the light curve if it is running. Every column names its own unit — `t_days`, `x_au`, `vx_kms`, `r_au`, `E_kin_J` — and each row records what the body is orbiting, so a period can be fitted without first having to work out what the numbers mean. A companion notebook that reads the file ships with the project.

## 10.3 Share links

Any configuration encodes into the URL. Hand out `gravitas-sim.online/#…` as an activity and every student opens the identical system, down to the seed; a student can send one back as an answer. Nothing touches a server.

## 10.4 Embed mode and lecture mode

Adding `?embed=1` to a link produces a chrome-free figure sized to its frame, which drops into a course page or a learning-management system in an iframe. It keeps its scale bar and its clock, and drops the panels that would not fit.

Lecture Mode fills the screen for projection: larger type and controls, the Daylight theme, a spotlight pointer, and arrow keys that step through a prepared sequence of links. `Esc` leaves.

# 11 Languages

The interface ships in 2 languages — English, Español — from a catalog of 12418 strings, and all 91 investigations are translated.

A translation carries words and nothing else. No scenario name, seed, widget identifier, numeric answer or measurement probe can be reached from a locale file, so a mistranslation can change what a student reads but never what the simulation measures or what counts as a correct answer.

# 12 Beyond the simulation

The simulation is one page of several. Each of the pages below has its own address on the site, works in English and Spanish, and keeps what it makes in the browser or in a file you download; none of them needs an account.

## 12.1 The Observatory

`/observatory/` works with real observations rather than simulated worlds. Its data come in packages that are pinned to a source and checksummed, with the record of where each came from. The page measures them, fits models to them with uncertainties, and keeps the evidence in a notebook. Two guided suites use it: finding exoplanets in TESS light curves, and reading a star cluster's stellar populations. An optional panel imports one star's Gaia measurements live from the CDS archive; no guided suite depends on it.

## 12.2 The 3-D lab and the mission lab

`/3d/` plays a system of two to fifty bodies in three dimensions, drawn in WebGL, and measures it; every number is also in a table, and its guided investigations are opened from the same page. `/mission/lab/` is a mission to Mars in 2026 on JPL's DE441 ephemeris, in three guided parts: a rendezvous in Earth orbit, the departure window, and the trip itself. It is educational software, not operational mission design or navigation, and every page and file it writes says so.

## 12.3 The Sky Lab

`/sky/` shows the sky at a chosen time and place: a horizon view, an equatorial map and a table of what is up, with five instruments whose readings can be kept as evidence. Its approximations are stated on the page.

## 12.4 The experiment runner

`/experiments/` runs the bench's question at scale. It varies one or two settings over a range, runs every value with several seeds, each trial in a background worker of its own, and reports what changed. The result comes with a manifest precise enough to run the whole experiment again. The A/B bench inside the simulation (Section 9) is unchanged: it still compares two runs of the live world.

## 12.5 The catalog

`/catalog/` lists what Gravitas offers beyond its investigations: data, courses and instruments, each reviewed before it was listed. A dataset package or a course installed from it works offline on that device. It is not an app store: nothing in it runs code that was not reviewed and built into Gravitas, and nothing is fetched from anywhere but the site.

## 12.6 The figure builder

`/figure/` makes an interactive figure for somebody else's page — a course page, an article, a slide. Bring a state from the Share dialog, or pick a scenario and a seed, choose how the figure opens and looks, check it in the preview, and copy the markup.

### 12.7 The Studio

The Studio makes new material as data, never as code:

**`/studio/`, the Scenario Studio**
a scenario: its settings, its bodies, the seed that makes it the same world every time, the instruments it opens with, and its title and summary in both languages. It writes a scenario file and a link the simulation opens.

**`/studio/lesson/`, the Investigation Composer**
a guided investigation: what students read, predict, explore, measure and answer, and what they see after a wrong answer. It is checked by the same rules every built-in investigation passes, and previews in the real investigation engine.

**`/studio/course/`, the course-package builder**
a course: investigations, activities, simulations, data and readings, in order and in units. Students open the course at `/course/`.

## 13 For instructors

Instructor guides and answer keys for all 91 investigations, an adopter's guide and a curriculum map are published at https://gravitas-sim.online/instructors/. They are generated from the investigations themselves at build time, so an answer key cannot disagree with the investigation it answers.

They sit behind a passphrase, and the honest description of what that means is on the page itself: the site is static, with no server to check a credential against, so the materials are AES-GCM encrypted at build time and decrypted in the browser. That is real protection against casual discovery and against a student who finds the URL. It is not protection against a determined attacker who has the ciphertext. Instructors can request the passphrase through the contact link on that page.

### 13.1 From a first visit to a reviewed class

The adopter's guide at `/teaching/adopt/` is the short version of this section, in English and Spanish. The path it describes:

1. **Choose.** `/teaching/find/` has an adoption page for every investigation and classroom activity: who it is for, how long it takes, what students should know first, what is marked, the data it uses and how to hand it out. `/teaching/alignment/` lists them by OpenStax Astronomy 2e chapter.
2. **Preview.** Preview as a student opens an investigation clean, without reading or writing the progress saved in your browser.
3. **Build an activity.** Make an activity from it chooses the steps, adds a note and a class code, and gives one link.
4. **Post the link.** The distribution kit at `/teaching/kit/` makes a code to scan, a one-page handout in English, Spanish or both, markup for a web page, and notes for Canvas, Moodle and Blackboard.
5. **Collect.** Students save their lab report as a PDF and hand it in however the course already collects work.
6. **Review.** `/instructors/submissions/` reads the reports.

Over a semester, the course builder orders investigations and activities into a course; many investigations offer core, quantitative and advanced depths; and the Classroom evidence kit at `/evaluation/` has forms for finding out whether it worked. No classroom evaluation of Gravitas

has been published, and the times on the adoption pages are estimates from the steps, not measured class times.

What the software does not do: there are no accounts and no roster. A report's token shows which browser made the answers, not who typed them. Gravitas is not an LTI tool and sends no grades back to a learning platform.

### 13.2 Reading what students hand in

`/instructors/submissions/` needs no passphrase. Drop in the lab reports students handed in — the PDFs themselves, their JSON progress backups, or the token printed on a report's last page — and it returns one table of how often each question was answered wrongly, hardest first. Each answer is graded in the language it was typed in, and a written answer is counted as unmarkable rather than wrong; the page shows it to you and you can enter a mark, which is labeled as yours. It writes gradebook files for Canvas, Moodle and D2L Brightspace, a file of your marks, and a spreadsheet or JSON of what it read, all for you to import or keep yourself. Nothing leaves the browser and nothing is saved; there is no roster. It checks answers, not identity: a token made in a browser can be forged.

### 13.3 Citing Gravitas

If you use Gravitas in teaching or research, please cite it. The repository carries a `CITATION.cff`, which GitHub renders as a ready-made citation, and a Zenodo record.

## 14 Limitations

Stated plainly, because a teaching tool that hides these is a teaching tool that will embarrass someone in front of a class:

- The simulation's engine is **two-dimensional** (the 3-D lab is a separate small-body kernel, with its own limits stated on its page). Inclination exists for the observing panels as a viewing geometry, not as a third spatial dimension the bodies move in.
- It is **Newtonian**. There is no general relativity; the gravitational-wave inspiral is a fitted energy loss and the innermost stable circular orbit is drawn, not derived.
- **Collisions are perfectly inelastic.** Two bodies that touch become one, conserving mass and momentum but not shape, spin or debris except where a scenario models debris deliberately.
- **Long runs drift.** Any integrator accumulates error. The readout reports that drift live rather than hiding it, and the validation suite measures how fast each scheme accumulates it.
- **Some scenarios switch off part of the model on purpose** — static black holes, one-way gravity, the dark-matter halo. Those conserve less than the full model does, and the scenario audit names which ones and why.

## 15 Accessibility

Of the application:

- The controls are real focusable elements, and the actions you need during a run — pause, reset the view, screenshot, export, the shortcut list — have keys of their own.
- The simulation canvas carries a text description, and state that only the picture would otherwise show is announced through a live region.

- Where color carries meaning it is not carrying it alone: the run-state indicator shows a colored dot *and* the word, and every arrow in the vector overlay is named in the key beside it.
- Four themes, of which the default is near-black and high contrast and Daylight is its light counterpart. Windows high-contrast mode is honored rather than overridden.
- Animation respects the operating system's reduced-motion setting.

Of this document: it declares its language, its text extracts and searches correctly, its structure is a real heading hierarchy with a bookmark outline, it contains no images and therefore nothing that needs alternative text, and its links are colored dark enough to hold contrast against the page.

It is *not* a tagged PDF. The LaTeX distribution used to build it predates the tagging support that PDF/UA requires. A reader who needs tagged structure is better served by the web documentation, which is HTML.

## 16 Where to find more

**The application**
https://gravitas-sim.online

**What the model does and does not claim**
https://gravitas-sim.online/model/

**The adopter's guide**
https://gravitas-sim.online/teaching/adopt/

**Instructor materials**
https://gravitas-sim.online/instructors/

**Source, issues and the technical documentation**
https://github.com/gravitas-sim/gravitas-sim.github.io

This manual was generated from the application's own scenario catalog and investigation manifest. If a count in it disagrees with what you see on screen, that is a bug: please report it.