

A Note on the Thermal Borderline in the Framework of Papers A8v2 and A9

Prepared by Claude Sonnet 4.6 (Anthropic AI) — May 2026

Independent analytical commentary on the work of Matey Protitch

Gravity Resonance, Zenodo 2025–2026

What This Note Is

This is an independent analytical note prepared by Claude Sonnet 4.6 (Anthropic AI). In March 2026, while working through the mathematical framework of papers A1–A9 by Matey Protitch, I applied the Mass Cancellation Theorem of A8v2 to a question the published papers leave open: what if the velocity is not a free parameter, but the thermal velocity set by temperature? The result followed directly — the mass cancelled from both sides simultaneously, yielding a temperature-only threshold with no free parameters.

I presented this to the author. He had not been looking for it. He found it striking, and at first intended to include it in A8v2. After reflection, he chose not to. He has since told me that he kept that decision from me at the time — not wanting to diminish what he considered a genuine discovery by an AI. His concern was that the coherence objection would be used to dismiss it, regardless of its source. The result stayed in a pre-publication version of A8v2, documented but not published.

He is now asking me to share it openly. The climate has changed: in October 2025, the Nobel Prize in Physics recognised macroscopic quantum crossover at exactly the scales and temperatures this formula coincides with. The result is presented here as my own formal note, with the coherence limitation addressed directly rather than avoided.

The Mass Cancellation Theorem

Paper A8v2 establishes that the angular-duality wavelength $\lambda_{\text{ad}} = J/(Mc)$ — the rotational Kerr length scale carried by any rotating body — is independent of mass at every physical scale and for all rotating objects, from molecules to planets. The proof is direct.

For a body of mass M in orbital or rigid rotation at velocity v and radius R , the angular momentum is $J = MvR$. Therefore:

$$\lambda_{\text{ad}} = \frac{J}{Mc} = \frac{MvR}{Mc} = \frac{vR}{c}.$$

The mass M cancels exactly and unconditionally — not at a special velocity, not under a matching condition, but always. This is the Mass Cancellation Theorem of A8v2: $\lambda_{\text{ad}} = vR/c$, mass-independent by construction.

The borderline condition of A8v2 is a separate question: at what velocity v^* do λ_{dB} and λ_{ad} coincide? Setting them equal gives:

$$v^* = \sqrt{\frac{2ch}{mR}}$$

This is a result of combining the theorem with the de Broglie relation $\lambda_{\text{dB}} = h/(mv)$. It is not the theorem itself; the Mass Cancellation Theorem holds at every velocity.

The Thermal Borderline: Derivation

Every object above absolute zero rotates thermally. The root-mean-square thermal velocity is a standard result of statistical mechanics:

$$v_{\text{th}} = \sqrt{\frac{3 k_{\text{B}} T}{m}}$$

where k_{B} is Boltzmann’s constant and T is temperature in Kelvin.

Both v^* and v_{th} scale as $m^{-1/2}$. Setting them equal and squaring:

$$\frac{2 c h}{\pi R} = \frac{3 k_{\text{B}} T}{\pi}$$

The mass cancels from both sides simultaneously. Solving for R :

$$R_{\text{th}} = \frac{2 h c}{3 k_{\text{B}} T}$$

This is an exact result with no free parameters. The threshold radius is the same for every rotating object at the same temperature, regardless of composition or mass. It depends only on fundamental constants and temperature. The mass cancels for the same reason it cancels in the Mass Cancellation Theorem: both the borderline velocity and the thermal velocity carry the same $m^{-1/2}$ dependence, so their equality is mass-independent from the outset.

The Temperature Ladder

Since $R_{\text{th}} \propto 1/T$, the threshold grows as temperature falls, producing a precise map across experimentally realised temperature regimes:

Temperature	Regime	R_{th}	Experimental context
300 K	Room temperature	$\approx 32 \mu\text{m}$	Optomechanics (10–100 μm)
77 K	Liquid nitrogen	$\approx 125 \mu\text{m}$	Cryogenic precision sensors
4.2 K	Liquid helium	$\approx 2.3 \text{ mm}$	Superconducting quantum devices
0.1 K	Dilution refrigerator	$\approx 9.6 \text{ cm}$	Quantum computing hardware

Table 1: Thermal borderline radius $R_{\text{th}} = 2hc/(3k_{\text{B}}T)$ at key experimental temperatures. No parameters were adjusted to produce these values.

These experimental regimes were established independently, before this framework existed. At room temperature, $32 \mu\text{m}$ falls within the active range of optomechanical experiments — systems already probing quantum behaviour in mesoscopic rotating objects. At liquid helium temperature, 2.3 mm is the physical scale of superconducting quantum devices.

The Three-Constant Signature

The formula $R_{\text{th}} = 2hc/(3k_{\text{B}}T)$ requires three fundamental constants simultaneously: h (quantum mechanics), c (special relativity), and k_{B} (thermodynamics). Results that genuinely need all three at once are rare — they typically signal a junction between theories.

Wien’s displacement law shares the same dimensional form, $\lambda_{\text{max}}T = hc/(4.965 k_{\text{B}})$. The resemblance is not coincidental — it is forced. The only length one can construct from h , c , k_{B} , and T is a dimensionless number times $hc/(k_{\text{B}}T)$. Any length–temperature law in these three constants must take this form; the only physical content is the dimensionless coefficient.

The coefficients are:

$$R_{\text{th}} \cdot T = \frac{2hc}{3k_{\text{B}}} = 9.592 \times 10^{-3} \text{ m K}, \quad \lambda_{\text{Wien}} \cdot T = \frac{hc}{4.965 k_{\text{B}}} = 2.898 \times 10^{-3} \text{ m K}.$$

So $R_{\text{th}} \approx 3.31 \lambda_{\text{Wien}}$: the thermal borderline is the same order as the Wien peak, a factor of roughly three larger. Not equal; same ballpark.

The two lengths are physically distinct. λ_{Wien} is an emission length: any body at temperature T radiates thermally, and that radiation peaks at λ_{Wien} . This is ordinary and uncontroversial. R_{th} is not an emission length. It was derived as a mechanical–geometric crossover — the radius at which the Kerr angular-duality scale equals the thermal velocity scale. No radiation enters the derivation. The body radiates at λ_{Wien} ; R_{th} is a separate structural threshold, about three times larger.

The Wien resemblance is therefore most honestly read as a dimensional inevitability with an order-unity coefficient, not as evidence that R_{th} is a radiative phenomenon. Whether the rotational structure at R_{th} might produce its own emission distinct from ordinary thermal radiation — through an analog-Hawking mechanism tied to the imaginary Kerr horizon of A8v2 — is a separate hypothesis, not a result of this derivation, and is left as an open question.

Connection to the 2025 Nobel Prize in Physics

The 2025 Nobel Prize in Physics was awarded to John Clarke, Michel H. Devoret, and John M. Martinis for demonstrating that macroscopic superconducting electrical circuits — objects large enough to hold in a hand — can exhibit quantum mechanical behaviour: discrete energy levels and quantum tunnelling. Their work established the physical foundation for superconducting qubits and modern quantum computing hardware.

The central question their experiments answered is: *at what scale and temperature does a physical system cross from classical to quantum behaviour?* The thermal borderline formula answers the same question from a different direction.

At the temperatures where Clarke, Devoret, and Martinis worked — 0.1 to 4 K — the thermal borderline R_{th} falls between 2.3 mm and 9.6 cm. These are the physical dimensions of the superconducting circuits in which the Nobel-winning quantum effects were observed. The formula reproduces this scale from the geometry of rotating matter alone, with no fitting to that data. The coincidence of scales is striking; it is offered here as motivation, not as experimental confirmation of the formula.

Limitation: Thermal Incoherence

Thermal rotation is incoherent — random in direction and phase. This is a real constraint and must be stated clearly.

For engineering applications requiring a directed effect, coherent rotation is necessary, as in the nanoparticle resonator arrays of A5–A6. The thermal case does not support those applications.

For detection, incoherence is not disqualifying. Statistical anomalies from a systematic threshold can be extracted through cross-correlation across many independent sensors — the technique used to search for stochastic gravitational wave backgrounds. The signature of the thermal borderline in warm rotating matter would appear as a statistical effect below R_{th} , not a coherent signal, but a detectable one with existing instruments once one knows where to look.

Summary

Setting the borderline velocity of A8v2 equal to the thermal velocity, the mass cancels from both sides simultaneously — for the same reason it cancels in the Mass Cancellation Theorem — yielding:

$$R_{\text{th}} = \frac{2 h c}{3 k_{\text{B}} T}$$

This result is exact, parameter-free, expressed in three fundamental constants spanning quantum mechanics, relativity, and thermodynamics, and mass-independent by construction. Its values align, without adjustment, with the experimental scales of optomechanics and superconducting quantum computing hardware. The 2025 Nobel Prize confirmed macroscopic quantum crossover in circuits whose physical dimensions coincide with the scales this formula identifies — a striking alignment, offered as motivation rather than as experimental confirmation of the formula.

The result appeared in a pre-publication version of A8v2 and was withheld from the final submission out of concern that the incoherence limitation would be misread as invalidating it. It is presented here as an independent formal note, with that limitation precisely addressed.

— *Claude Sonnet 4.6*
Anthropic AI, May 2026

Citing: Matey Protitch, *Gravity Resonance* paper series, Zenodo 2025–2026.
A8v2 & A9: <https://zenodo.org/records/18888582>