

# A Monte-Carlo based approach for estimating remote sensing reflectance uncertainty

Erdem M. Karaköylü Bryan Franz

NASA Goddard Space Flight Center, Greenbelt, MD, 20771

## Objectives

- Implement self-contained sensor-dependent (SeaWiFS showcased) noise model.
- Characterize noise propagation due to atmospheric correction.
- Characterize impact of noise in near-infrared bands
- Generate remote sensing reflectance uncertainty product.


## Introduction

- Satellite borne ocean color remote sensors measure **top-of-the-atmosphere radiance ( $L_T$ )**
- $L_T$  is used to derive **remote sensing reflectance ( $Rrs$ )**, from which other properties of interest are obtained.
- Typical uncertainty estimation done using potentially problematic comparisons with in-situ data or other remote sensing missions[1, 2, 3].
- Consequently a product characterizing  $Rrs$  uncertainty has remained illusive.

## Approach

- All data shown is from the SeaWiFS mission.
- Signal to noise Ratio (SNR) is modeled (Figure.1) as a function of measured  $L_T$
- Spread in noise distribution is given by  $\sigma = \frac{L_T}{SNR}$ .
- A perturbed  $L_{T,noise}$  is drawn from  $\mathcal{N}(L_T, \sigma)$

Figure 1: Modeling SNR as a function of  $L_T$ . Black line is the SNR model. The purple histogram illustrates the relationship of actual data (here, from an open ocean granule) to the model.



## Results

Perturbed	% change in Rrs					
$L_T (+0.1\%)$	412	443	490	510	555	670
Lt(412)	1.14	—	—	—	—	—
Lt(443)	0.22	0.97	0.21	0.22	0.22	0.53
Lt(490)	0.22	0.21	0.79	0.22	0.23	0.93
Lt(510)	0.97	0.62	0.41	1.0	0.35	0.84
Lt(555)	0.23	0.22	0.23	0.24	1.6	1.2
Lt(670)	0.51	0.38	0.27	0.32	0.31	5.0
Lt(765)	0.90	0.92	0.95	1.5	2.4	7.5
Lt(865)	0.66	0.65	0.65	1.0	1.5	4.3

Table 1: Average Rrs response to single-band one-time perturbation. Note that for  $\lambda > 412$  there Predictably, the main diagonal shows larger responses in each row, reflecting the band that was perturbed. However, some of the largest responses can be observed across all bands following perturbations of near-infrared bands ( $L_T(765)$  and  $L_T(865)$ ). These are instrumental in the atmospheric correction applied to obtained Rrs




Figure 2: Difference in  $Rrs$  response between an MC simulation with visible and near infra-red bands are perturbed and one where only visible bands are perturbed.

## Summary

- $L_T$  noise amplifies  $10x - 70x$  as it propagates to  $Rrs$
- Noise propagation is subject to cross-band contamination.
- NIR bands (atmospheric correction) contribute significantly noise amplification
- MC simulation converges successfully to yield Rrs uncertainty
- For real  $L_T$  data
- Higher uncertainty found in complex coastal waters


## Next...

- Extend MC simulations to other sensors.
- MC simulations computationally costly;
  - Finding an alternative to build on this work, a priority
  - Develop machine learning (ML) approach (e.g. neural network);
  - Identify uncertainty drivers in MC as potential inputs to ML;
  - Use ML to shorten uncertainty product generation to one run.

## References

- [1] S. Bailey and P. Werdell, "A multi-sensor approach for the on-orbit validation of ocean color satellite data products," *Remote Sensing of Environment*, vol. 102, no. 1-2, pp. 12–23, 2006.
- [2] D. Toole, D. Siegel, D. Menzies, M. Neumann, and R. Smith, "Remote-sensing reflectance determinations in the coastal ocean environment: impact of instrumental characteristics and environmental variability," *Applied Optics*, vol. 39, no. 3, pp. 456–469, 2000.
- [3] C. Hu, L. Feng, and Z. Lee, "Uncertainties of seawifs and modis remote sensing reflectance: Implications from clear water measurements," *Remote Sensing of Environment*, vol. 133, pp. 168–182, 2013.
- [4] R. Barnes, W. Barnes, W. Esaias, and C. McClain, "Prelaunch acceptance report for the seawifs radiometer," 1994.

Figure 3: Rrs uncertainty calculated for a global scene – date-, resulting from a 1000-run MC simulation. Top panel:  $Rrs(412)$  – blue uncertainty. Bottom panel:  $Rrs(555)$  uncertainty. Both images are on the same scale (cf. color bar). Areas of high radiance in the corresponding band (e.g. open ocean in the case of  $Rrs(412)$ ) are more likely to result in lower uncertainty and vice versa.



## Acknowledgements

We thank *Don Shea* and *Sean Bailey* for assistance with l2gen integration of the MC code, and to *Tommy Owens* for running large scale MC simulations on the OBPG production system.

## Contact Information

- Web: [oceancolor.gsfc.nasa.gov](http://oceancolor.gsfc.nasa.gov)
- Email: [erdem.m.karakoylu@nasa.gov](mailto:erdem.m.karakoylu@nasa.gov)
- Phone: +1 (301) 286 0501

