Formal Languages Non-Regular Languages

Non-regular languages

$$\{a^n b^n : n \ge 0\}$$

 $\{vv^R : v \in \{a,b\}^*\}$

Regular languages * 1.

$$a*b$$
 $b*c+a$ $b+c(a+b)*$ $etc...$

How can we prove that a language L is not regular?

Prove that there is no DFA that accepts $\,L\,$


Problem: this is not easy to prove

Solution: the Pumping Lemma!!!





The Pigeonhole Principle


4 pigeons




3 pigeonholes










A pigeonhole must contain at least two pigeons







n pigeons




The Pigeonhole Principle


n pigeons

m pigeonholes


n > m

There is a pigeonhole with at least 2 pigeons










The Pigeonhole Principle

and

DFAs

DFA with 4 states




In walks of strings: a


aa

aab

no state is repeated




In walks of strings: aabb a state bbaa is repeated abbabb abbabbabb...



If string w has length $|w| \ge 4$:

Then the transitions of string w are more than the states of the DFA

Thus, a state must be repeated




In general, for any DFA:

String w has length \geq number of states



A state q must be repeated in the walk of w





In other words for a string w:

 \xrightarrow{a} transitions are pigeons



(q) states are pigeonholes






The Pumping Lemma

Take an infinite regular language L

There exists a DFA that accepts L



Take string w with $w \in L$


There is a walk with label W:

$$\longrightarrow$$
 walk W


If string w has length $|w| \ge m$ (number of states of DFA)

then, from the pigeonhole principle:


a state is repeated in the walk w



Let q be the first state repeated in the walk of w




Write w = x y z




Observations:


 $| length \ | \ x \ y \ | \le m \ number$ of states $| length \ | \ y \ | \ge 1$ of DFA




Observation: The string xz is accepted



Observation: The string x y y z is accepted




Observation: The string x y y y z is accepted



In General:


The string $xy^{l}z$ is accepted i=0,1,2,...



In General:
$$x y^i z \in L$$

 $i = 0, 1, 2, \dots$

Language accepted by the DFA



In other words, we described:







The Pumping Lemma!!!





The Pumping Lemma:

- \cdot Given an infinite regular languageL
- there exists an integer m
- for any string $w \in L$ with length $|w| \ge m$
- we can write w = x y z
- with $|xy| \le m$ and $|y| \ge 1$
- such that: $x y^l z \in L$ i = 0, 1, 2, ...

Applications

of

the Pumping Lemma

Theorem: The language
$$L = \{a^nb^n : n \ge 0\}$$
 is not regular

Proof: Use the Pumping Lemma

$$L = \{a^n b^n : n \ge 0\}$$

Assume for contradiction that L is a regular language

Since L is infinite we can apply the Pumping Lemma

$$L = \{a^n b^n : n \ge 0\}$$

Let m be the integer in the Pumping Lemma

Pick a string w such that: $w \in L$

length $|w| \ge m$

We pick
$$w = a^m b^m$$

Write:
$$a^m b^m = x y z$$

From the Pumping Lemma it must be that length $|x y| \le m$, $|y| \ge 1$

$$xyz = a^m b^m = \underbrace{a...aa...aa...ab...b}_{m}$$

Thus:
$$y = a^k$$
, $k \ge 1$

$$x y z = a^m b^m$$

$$y = a^k, \quad k \ge 1$$

$$x y^{i} z \in L$$
 $i = 0, 1, 2, \dots$

Thus:
$$x y^2 z \in L$$

$$x y z = a^m b^m \qquad y = a^k, \quad k \ge 1$$

From the Pumping Lemma: $x y^2 z \in L$

$$xy^{2}z = \underbrace{a...aa...aa...aa...ab...b}_{m+k} \in L$$

Thus:
$$a^{m+k}b^m \in L$$

$$a^{m+k}b^m \in L$$

$$k \ge 1$$

BUT:
$$L = \{a^n b^n : n \ge 0\}$$



$$a^{m+k}b^m \notin L$$

CONTRADICTION!!!

Therefore: Our assumption that L is a regular language is not true

Conclusion: L is not a regular language

Non-regular languages $\{a^nb^n: n \ge 0\}$

