

Exact Exchange in Density Functional Theory

S. Sharma, J. K. Dewhurst and C. Ambrosch-Draxl

Institut für Physik
Karl-Franzens-Universität Graz
Universitätsplatz 5
A-8010 Graz, Austria

19th January 2005

Density functional theory

Many-body theory involves solving for functions in $3N$ coordinates:
 $\Psi(\mathbf{r}_1, \mathbf{r}_2, \dots, \mathbf{r}_N)$

Density functional theory (DFT) requires solving for functions in 3 coordinates: $n(\mathbf{r})$

Density functional theory

Many-body theory involves solving for functions in $3N$ coordinates:
 $\Psi(\mathbf{r}_1, \mathbf{r}_2, \dots, \mathbf{r}_N)$

Density functional theory (DFT) requires solving for functions in 3 coordinates: $n(\mathbf{r})$

Density functional theory

Hohenberg-Kohn Theorems (1964)

- 1 External potential $v(\mathbf{r})$ is uniquely determined by $n(\mathbf{r})$
- 2 The variational principle holds

$$E_0 = E_{v_0}[n_0] < E_{v_0}[n]$$

- 3 $E_{v_0}[n] = F[n] + \int d\mathbf{r} v_0(\mathbf{r})n(\mathbf{r})$

Density functional theory

Hohenberg-Kohn Theorems (1964)

- 1 External potential $v(\mathbf{r})$ is uniquely determined by $n(\mathbf{r})$
- 2 The variational principle holds

$$E_0 = E_{v_0}[n_0] < E_{v_0}[n]$$

- 3 $E_{v_0}[n] = F[n] + \int d\mathbf{r} v_0(\mathbf{r})n(\mathbf{r})$

Density functional theory

Hohenberg-Kohn Theorems (1964)

- 1 External potential $v(\mathbf{r})$ is uniquely determined by $n(\mathbf{r})$
- 2 The variational principle holds

$$E_0 = E_{v_0}[n_0] < E_{v_0}[n]$$

- 3 $E_{v_0}[n] = F[n] + \int d\mathbf{r} v_0(\mathbf{r})n(\mathbf{r})$

Density functional theory

DFT is an exact theory for interacting systems in the ground state

Kohn-Sham equations

Find set of auxilliary single particle orbitals such that

$$\left[-\frac{1}{2} \nabla^2 + v_s(\mathbf{r}) \right] \phi_i(\mathbf{r}) = \epsilon_i \phi_i(\mathbf{r})$$

and

$$n(\mathbf{r}) = \sum_i^{\text{occ}} |\phi_i(\mathbf{r})|^2.$$

If

$$F[n] = T_s[n] + \frac{1}{2} \int d\mathbf{r} d\mathbf{r}' \frac{n(\mathbf{r})n(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|} + E_{\text{xc}}[n]$$

Kohn-Sham equations

Find set of auxilliary single particle orbitals such that

$$\left[-\frac{1}{2} \nabla^2 + v_s(\mathbf{r}) \right] \phi_i(\mathbf{r}) = \epsilon_i \phi_i(\mathbf{r})$$

and

$$n(\mathbf{r}) = \sum_i^{\text{occ}} |\phi_i(\mathbf{r})|^2.$$

If

$$F[n] = T_s[n] + \frac{1}{2} \int d\mathbf{r} d\mathbf{r}' \frac{n(\mathbf{r})n(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|} + E_{\text{xc}}[n]$$

Kohn-Sham equations

Then

$$v_s[n](\mathbf{r}) = v(\mathbf{r}) + \int d\mathbf{r}' \frac{n(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|} + v_{xc}[n](\mathbf{r})$$

where

$$v_{xc}[n](\mathbf{r}) = \frac{\delta E_{xc}[n]}{\delta n(\mathbf{r})}.$$

Many sins hidden in $E_{xc}[n]$!

Kohn-Sham equations

Then

$$v_s[n](\mathbf{r}) = v(\mathbf{r}) + \int d\mathbf{r}' \frac{n(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|} + v_{xc}[n](\mathbf{r})$$

where

$$v_{xc}[n](\mathbf{r}) = \frac{\delta E_{xc}[n]}{\delta n(\mathbf{r})}.$$

Many sins hidden in $E_{xc}[n]$!

Exchange-correlation functionals

First generation : Local density approximation (LDA)

$$E_{\text{xc}}^{\text{LDA}}[n] = \int d\mathbf{r} n(\mathbf{r}) e_{\text{xc}}^{\text{unif}}(n(\mathbf{r}))$$

Second generation : Generalised gradient approximations

$$E_{\text{xc}}^{\text{GGA}}[n] = \int d\mathbf{r} f(n(\mathbf{r}), \nabla n(\mathbf{r}))$$

$$E_{\text{xc}}^{\text{Meta-GGA}}[n] = \int d\mathbf{r} g(n(\mathbf{r}), \nabla n(\mathbf{r}), \tau(\mathbf{r}))$$

Exchange-correlation functionals

First generation : Local density approximation (LDA)

$$E_{\text{xc}}^{\text{LDA}}[n] = \int d\mathbf{r} n(\mathbf{r}) e_{\text{xc}}^{\text{unif}}(n(\mathbf{r}))$$

Second generation : Generalised gradient approximations

$$E_{\text{xc}}^{\text{GGA}}[n] = \int d\mathbf{r} f(n(\mathbf{r}), \nabla n(\mathbf{r}))$$

$$E_{\text{xc}}^{\text{Meta-GGA}}[n] = \int d\mathbf{r} g(n(\mathbf{r}), \nabla n(\mathbf{r}), \tau(\mathbf{r}))$$

Exchange-correlation functionals

First generation : Local density approximation (LDA)

$$E_{\text{xc}}^{\text{LDA}}[n] = \int d\mathbf{r} n(\mathbf{r}) e_{\text{xc}}^{\text{unif}}(n(\mathbf{r}))$$

Second generation : Generalised gradient approximations

$$E_{\text{xc}}^{\text{GGA}}[n] = \int d\mathbf{r} f(n(\mathbf{r}), \nabla n(\mathbf{r}))$$

$$E_{\text{xc}}^{\text{Meta-GGA}}[n] = \int d\mathbf{r} g(n(\mathbf{r}), \nabla n(\mathbf{r}), \tau(\mathbf{r}))$$

Exchange-correlation functionals

Development of new functionals leads not only to improved accuracy but also correct qualitative features

Exchange-correlation functionals

Third generation: Exact exchange (EXX)

Neglect correlation and use the Hartree-Fock exchange energy

$$E_x[n] = -\frac{1}{2} \sum_{i,j}^{\text{occ}} \int d\mathbf{r} d\mathbf{r}' \frac{\phi_i^*(\mathbf{r})\phi_j^*(\mathbf{r}')\phi_j(\mathbf{r})\phi_i(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|}$$

To solve the Kohn-Sham system we require

$$v_x[n](\mathbf{r}) = \frac{\delta E_x[n]}{\delta n(\mathbf{r})}$$

Exchange-correlation functionals

Third generation: Exact exchange (EXX)

Neglect correlation and use the Hartree-Fock exchange energy

$$E_x[n] = -\frac{1}{2} \sum_{i,j}^{\text{occ}} \int d\mathbf{r} d\mathbf{r}' \frac{\phi_i^*(\mathbf{r})\phi_j^*(\mathbf{r}')\phi_j(\mathbf{r})\phi_i(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|}$$

To solve the Kohn-Sham system we require

$$v_x[n](\mathbf{r}) = \frac{\delta E_x[n]}{\delta n(\mathbf{r})}$$

Exchange-correlation functionals

Using the functional derivative chain rule:

$$v_x[n](\mathbf{r}) = \sum_i^{\text{occ}} \int d\mathbf{r} d\mathbf{r}'' \left[\frac{\delta E_x}{\delta \phi_i(\mathbf{r}'')} \frac{\delta \phi_i(\mathbf{r}'')}{\delta v_s(\mathbf{r}')} + \frac{\delta E_x}{\delta \phi_i^*(\mathbf{r}'')} \frac{\delta \phi_i^*(\mathbf{r}'')}{\delta v_s(\mathbf{r}')} \right] \frac{\delta v_s(\mathbf{r}')}{\delta n(\mathbf{r})}$$

$$= \int d\mathbf{r}' \left[\sum_i^{\text{occ}} \sum_j^{\text{unocc}} \langle \phi_i | \hat{v}_x^{\text{NL}} | \phi_j \rangle \frac{\phi_j^*(\mathbf{r}') \phi_i(\mathbf{r}')}{\epsilon_i - \epsilon_j} + \text{c.c.} \right] \frac{\delta v_s(\mathbf{r}')}{\delta n(\mathbf{r})},$$

where

$$\langle \phi_{i\mathbf{k}} | \hat{v}_x^{\text{NL}} | \phi_{j\mathbf{k}} \rangle = \sum_{l\mathbf{k}'}^{\text{occ}} w_{\mathbf{k}'} \int d\mathbf{r} d\mathbf{r}' \frac{\phi_{i\mathbf{k}}^*(\mathbf{r}) \phi_{l\mathbf{k}'}(\mathbf{r}) \phi_{l\mathbf{k}'}^*(\mathbf{r}') \phi_{j\mathbf{k}}(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|}.$$

Exchange-correlation functionals

Using the linear-response operator

$$\begin{aligned}\chi(\mathbf{r}, \mathbf{r}') &\equiv \frac{\delta n(\mathbf{r})}{\delta v_s(\mathbf{r}')} \\ &= \sum_i^{\text{occ}} \sum_j^{\text{unocc}} \frac{\phi_i^*(\mathbf{r}) \phi_j(\mathbf{r}) \phi_j^*(\mathbf{r}') \phi_i(\mathbf{r}')}{\epsilon_i - \epsilon_j} + \text{c.c.}\end{aligned}$$

we have

$$\frac{\delta v_s(\mathbf{r}')}{\delta n(\mathbf{r})} = \tilde{\chi}^{-1}(\mathbf{r}, \mathbf{r}').$$

basis

Exchange-correlation functionals

Exact exchange is **NOT**

- Hartree-Fock
- a mean-field theory
- an attempt to solve the quasi-particle equation
- a parameterised correction to LDA
- easy to implement

Exchange-correlation functionals

Exact exchange is **NOT**

- Hartree-Fock
- a mean-field theory
- an attempt to solve the quasi-particle equation
- a parameterised correction to LDA
- easy to implement

Exchange-correlation functionals

Exact exchange is **NOT**

- Hartree-Fock
- a mean-field theory
- an attempt to solve the quasi-particle equation
- a parameterised correction to LDA
- easy to implement

Exchange-correlation functionals

Exact exchange is **NOT**

- Hartree-Fock
- a mean-field theory
- an attempt to solve the quasi-particle equation
- a parameterised correction to LDA
- easy to implement

Exchange-correlation functionals

Exact exchange is **NOT**

- Hartree-Fock
- a mean-field theory
- an attempt to solve the quasi-particle equation
- a parameterised correction to LDA
- easy to implement

Practicalities

Pseudopotentials (PP)

- atomic core is frozen and represented by a non-local potential
- does not react properly to the solid state environment: no relaxation of core states
- planewaves are used as the basis

Practicalities

Pseudopotentials (PP)

- atomic core is frozen and represented by a non-local potential
- does not react properly to the solid state environment: no relaxation of core states
- planewaves are used as the basis

Practicalities

Pseudopotentials (PP)

- atomic core is frozen and represented by a non-local potential
- does not react properly to the solid state environment: no relaxation of core states
- planewaves are used as the basis

Practicalities

Atomic sphere approximation (ASA)

- includes core orbitals
- potential is spherically symmetric
- muffin-tins are space-filling

Practicalities

Atomic sphere approximation (ASA)

- includes core orbitals
- potential is spherically symmetric
- muffin-tins are space-filling

Practicalities


Atomic sphere approximation (ASA)

- includes core orbitals
- potential is spherically symmetric
- muffin-tins are space-filling

Practicalities

Full-potential linearised augmented planewaves (FP-LAPW)


- includes core orbitals
- potential is fully described
- space divided into muffin-tin and interstitial regions
- most precise method available



Practicalities

Full-potential linearised augmented planewaves (FP-LAPW)


- includes core orbitals
- potential is fully described
- space divided into muffin-tin and interstitial regions
- most precise method available



Practicalities

Full-potential linearised augmented planewaves (FP-LAPW)


- includes core orbitals
- potential is fully described
- space divided into **muffin-tin** and interstitial regions
- most precise method available



Practicalities

Full-potential linearised augmented planewaves (FP-LAPW)


- includes core orbitals
- potential is fully described
- space divided into muffin-tin and **interstitial** regions
- most precise method available



Practicalities

Full-potential linearised augmented planewaves (FP-LAPW)

- includes core orbitals
- potential is fully described
- space divided into muffin-tin and interstitial regions
- most precise method available



Practicalities

Basis set

- An efficient basis set is required so that the response

$$\chi(\mathbf{r}, \mathbf{r}') \equiv \frac{\delta n(\mathbf{r})}{\delta v_s(\mathbf{r}')}$$

can be inverted

- Basis set should not contain constant functions

response

Practicalities

Basis set

- An efficient basis set is required so that the response

$$\chi(\mathbf{r}, \mathbf{r}') \equiv \frac{\delta n(\mathbf{r})}{\delta v_s(\mathbf{r}')}$$

can be inverted

- Basis set should not contain constant functions

response

Practicalities

Choose the overlap densities

$$\rho_{\alpha}(\mathbf{r}) \equiv \phi_{i\mathbf{k}}^*(\mathbf{r})\phi_{j\mathbf{k}}(\mathbf{r}),$$

and complex conjugates, where $\alpha \equiv (i\mathbf{k}, j\mathbf{k})$.

Diagonalise

$$O_{\alpha\beta} \equiv \int d\mathbf{r} \rho_{\alpha}^*(\mathbf{r})\rho_{\beta}(\mathbf{r}),$$

and eliminate eigenvectors with small eigenvalues.

Practicalities

Choose the overlap densities

$$\rho_{\alpha}(\mathbf{r}) \equiv \phi_{i\mathbf{k}}^*(\mathbf{r})\phi_{j\mathbf{k}}(\mathbf{r}),$$

and complex conjugates, where $\alpha \equiv (i\mathbf{k}, j\mathbf{k})$.

Diagonalise

$$O_{\alpha\beta} \equiv \int d\mathbf{r} \rho_{\alpha}^*(\mathbf{r})\rho_{\beta}(\mathbf{r}),$$

and eliminate eigenvectors with small eigenvalues.

Practicalities

Find transformation matrix C such that if

$$\tilde{\rho}_{\alpha}(\mathbf{r}) = \sum_{\beta} C_{\beta}^{\alpha} \sum_{\gamma} v_{\gamma}^{\beta} \rho_{\gamma}(\mathbf{r})$$

then

$$\int d\mathbf{r} \tilde{\rho}_{\alpha}^{*}(\mathbf{r}) \tilde{\rho}_{\beta}(\mathbf{r}) = \delta_{\alpha\beta}.$$

The matrix equation

$$CC^{\dagger} = \left(v^{\dagger} O v \right)^{-1}$$

is solved by Cholesky decomposition.

Practicalities

Find transformation matrix C such that if

$$\tilde{\rho}_{\alpha}(\mathbf{r}) = \sum_{\beta} C_{\beta}^{\alpha} \sum_{\gamma} v_{\gamma}^{\beta} \rho_{\gamma}(\mathbf{r})$$

then

$$\int d\mathbf{r} \tilde{\rho}_{\alpha}^{*}(\mathbf{r}) \tilde{\rho}_{\beta}(\mathbf{r}) = \delta_{\alpha\beta}.$$

The matrix equation

$$CC^{\dagger} = \left(v^{\dagger} O v\right)^{-1}$$

is solved by Cholesky decomposition.

Practicalities

By construction $\int d\mathbf{r} \tilde{\rho}_\alpha(\mathbf{r}) = 0$, so $\{\tilde{\rho}_\alpha\}$ form an ideal basis for inversion of χ

Practicalities

NL Lastly, given $\mathbf{q} \equiv \mathbf{k} - \mathbf{k}'$, the long range coulomb term of the NL matrix elements

$$\langle \phi_{i\mathbf{k}} | \hat{v}_x^{\text{NL}} | \phi_{j\mathbf{k}} \rangle_{\text{LR}} = \sum_{l\mathbf{q}}^{\text{occ}} w_{\mathbf{q}} \frac{4\pi\Omega}{q^2} \rho_{il}^*(\mathbf{q}) \rho_{lj}(\mathbf{q}),$$

where $\rho_{il}(\mathbf{q})$ and $\rho_{lj}(\mathbf{q})$ are the pseudo-charge densities.

Poor convergence with respect to the number of q -points.

Approximate it by an integral over a sphere of volume equivalent to that of the BZ

$$\langle \phi_{i\mathbf{k}} | \hat{v}_x^{\text{NL}} | \phi_{j\mathbf{k}} \rangle_{\text{LR}} \simeq 2 \left(\frac{6\Omega^5}{\pi} \right)^{1/3} \sum_{l\mathbf{q}}^{\text{occ}} w_{\mathbf{q}} \rho_{il}^*(\mathbf{q}) \rho_{lj}(\mathbf{q}).$$

Practicalities

NL Lastly, given $\mathbf{q} \equiv \mathbf{k} - \mathbf{k}'$, the long range coulomb term of the NL matrix elements

$$\langle \phi_{i\mathbf{k}} | \hat{v}_x^{\text{NL}} | \phi_{j\mathbf{k}} \rangle_{\text{LR}} = \sum_{l\mathbf{q}}^{\text{occ}} w_{\mathbf{q}} \frac{4\pi\Omega}{q^2} \rho_{il}^*(\mathbf{q}) \rho_{lj}(\mathbf{q}),$$

where $\rho_{il}(\mathbf{q})$ and $\rho_{lj}(\mathbf{q})$ are the pseudo-charge densities.

Poor convergence with respect to the number of q -points.

Approximate it by an integral over a sphere of volume equivalent to that of the BZ

$$\langle \phi_{i\mathbf{k}} | \hat{v}_x^{\text{NL}} | \phi_{j\mathbf{k}} \rangle_{\text{LR}} \simeq 2 \left(\frac{6\Omega^5}{\pi} \right)^{1/3} \sum_{l\mathbf{q}}^{\text{occ}} w_{\mathbf{q}} \rho_{il}^*(\mathbf{q}) \rho_{lj}(\mathbf{q}).$$

Practicalities

NL Lastly, given $\mathbf{q} \equiv \mathbf{k} - \mathbf{k}'$, the long range coulomb term of the NL matrix elements

$$\langle \phi_{i\mathbf{k}} | \hat{v}_x^{\text{NL}} | \phi_{j\mathbf{k}} \rangle_{\text{LR}} = \sum_{l\mathbf{q}}^{\text{occ}} w_{\mathbf{q}} \frac{4\pi\Omega}{q^2} \rho_{il}^*(\mathbf{q}) \rho_{lj}(\mathbf{q}),$$

where $\rho_{il}(\mathbf{q})$ and $\rho_{lj}(\mathbf{q})$ are the pseudo-charge densities.

Poor convergence with respect to the number of q -points.

Approximate it by an integral over a sphere of volume equivalent to that of the BZ

$$\langle \phi_{i\mathbf{k}} | \hat{v}_x^{\text{NL}} | \phi_{j\mathbf{k}} \rangle_{\text{LR}} \simeq 2 \left(\frac{6\Omega^5}{\pi} \right)^{1/3} \sum_{l\mathbf{q}}^{\text{occ}} w_{\mathbf{q}} \rho_{il}^*(\mathbf{q}) \rho_{lj}(\mathbf{q}).$$

Applications

EXX Applied to

- 1 Magnetic metals
- 2 Semiconductors and insulators

Magnetic metals: introduction to the problem

Magnetic moment in Bohr magneton

Compound	FP-LDA	Experiment
FeAl	0.71	0.0

① P. Mohn et al. Phys. Rev. Lett. 87 196401 (2001): LDA+ U

② Petukhov et al. Phys. Rev. B 67 153106 (2003): LDA+ U +DMFT


Magnetic metals: introduction to the problem

Magnetic moment in Bohr magneton


Compound	FP-LDA	Experiment
FeAl	0.71	0.0

- 1 P. Mohn et al. Phys. Rev. Lett. 87 196401 (2001): LDA+ U
- 2 Petukhov et al. Phys. Rev. B 67 153106 (2003): LDA+ U +DMFT


Magnetic metals: FeAl




Magnetic metals: FeAl




Magnetic metals: FeAl



Magnetic metals: FeAl



Magnetic metals: FeAl



Magnetic metals: stringent tests for EXX

Magnetic moment in Bohr magneton

Compound	FP-LDA	FP-EXX	Experiment
FeAl	0.71	0.0	0.0
Ni ₃ Ga	0.79	?	0.0
Ni ₃ Al	0.70	?	0.23

<http://arxiv.org/abs/cond-mat/0501258>

Magnetic metals: FeAl, Ni₃Ga and Ni₃Al


Magnetic moment in Bohr magneton

Compound	FP-LDA	FP-EXX	Experiment
FeAl	0.71	0.0	0.0
Ni ₃ Ga	0.79	0.0	0.0
Ni ₃ Al	0.70	0.20	0.23


<http://arxiv.org/abs/cond-mat/0501258>

Semiconductors and insulators: band-gaps

PP-EXX : Städele et al. PRB 59 10031



PP-EXX: Magyar et al. PRB 69 045111



The fundamental band-gap for materials:

$$E_g = A - I = (E_{N+1} - E_N) - (E_N - E_{N-1})$$

$$E_g = \frac{\delta E}{\delta n^+} - \frac{\delta E}{\delta n^-}$$

$$E[n] = T[n] + U_s[n] + E_{xc}[n]$$

$$E_g = \left(\frac{\delta T}{\delta n^+} - \frac{\delta T}{\delta n^-} \right) + \left(\frac{\delta v_{xc}}{\delta n^+} - \frac{\delta v_{xc}}{\delta n^-} \right)$$

$$E_g = E_g^{\text{KS}} + \Delta_{xc}$$

The fundamental band-gap for materials:

$$E_g = A - I = (E_{N+1} - E_N) - (E_N - E_{N-1})$$

$$E_g = \frac{\delta E}{\delta n^+} - \frac{\delta E}{\delta n^-}$$

$$E[n] = T[n] + U_s[n] + E_{xc}[n]$$

$$E_g = \left(\frac{\delta T}{\delta n^+} - \frac{\delta T}{\delta n^-} \right) + \left(\frac{\delta v_{xc}}{\delta n^+} - \frac{\delta v_{xc}}{\delta n^-} \right)$$

$$E_g = E_g^{\text{KS}} + \Delta_{xc}$$

The fundamental band-gap for materials:

$$E_g = A - I = (E_{N+1} - E_N) - (E_N - E_{N-1})$$

$$E_g = \frac{\delta E}{\delta n^+} - \frac{\delta E}{\delta n^-}$$

$$E[n] = T[n] + U_s[n] + E_{xc}[n]$$

$$E_g = \left(\frac{\delta T}{\delta n^+} - \frac{\delta T}{\delta n^-} \right) + \left(\frac{\delta v_{xc}}{\delta n^+} - \frac{\delta v_{xc}}{\delta n^-} \right)$$

$$E_g = E_g^{KS} + \Delta_{xc}$$

The fundamental band-gap for materials:

$$E_g = A - I = (E_{N+1} - E_N) - (E_N - E_{N-1})$$

$$E_g = \frac{\delta E}{\delta n^+} - \frac{\delta E}{\delta n^-}$$

$$E[n] = T[n] + U_s[n] + E_{xc}[n]$$

$$E_g = \left(\frac{\delta T}{\delta n^+} - \frac{\delta T}{\delta n^-} \right) + \left(\frac{\delta v_{xc}}{\delta n^+} - \frac{\delta v_{xc}}{\delta n^-} \right)$$

$$E_g = E_g^{KS} + \Delta_{xc}$$

The fundamental band-gap for materials:

$$E_g = A - I = (E_{N+1} - E_N) - (E_N - E_{N-1})$$


$$E_g = \frac{\delta E}{\delta n^+} - \frac{\delta E}{\delta n^-}$$

$$E[n] = T[n] + U_s[n] + E_{xc}[n]$$


$$E_g = \left(\frac{\delta T}{\delta n^+} - \frac{\delta T}{\delta n^-} \right) + \left(\frac{\delta v_{xc}}{\delta n^+} - \frac{\delta v_{xc}}{\delta n^-} \right)$$

$$E_g = E_g^{\text{KS}} + \Delta_{xc}$$

Semiconductors and insulators: band-gaps




Semiconductors and insulators: band-gaps




Semiconductors and insulators: band-gaps


<http://arxiv.org/abs/cond-mat/0501353>



Semiconductors and insulators: *d*-band position




Semiconductors and insulators: *d*-band position



GW LMTO-ASA: Aryasetiawan et al. PRB 54 17564 FPLMTO: Kotani et al. SSC 121

Semiconductors and insulators: *d*-band position



<http://arxiv.org/abs/cond-mat/0501353>

Conclusions

- 1 EXX within an all electron full potential method is implemented within **EXC!TING** code. Right now this is the only FP code to be able to do EXX.
- 2 A new and one of the most optimal basis is proposed for calculating and inverting the response. This basis may be useful for future TD-DFT and GW calculations.
- 3 Magnetic metals: **Asymmetry** in exchange potential is very important to get the correct ground-state.
- 4 Semiconductors and insulators: **Core-valence interaction** is crucial for correct treatment of the EXX.
- 5 Lack of **Asymmetry** and/or **core-valence interaction** could lead to spurious agreement with experiments.

Conclusions

- 1 EXX within an all electron full potential method is implemented within **EXC!TING** code. Right now this is the only FP code to be able to do EXX.
- 2 A new and one of the most optimal basis is proposed for calculating and inverting the response. This basis may be useful for future TD-DFT and GW calculations.
- 3 Magnetic metals: **Asymmetry** in exchange potential is very important to get the correct ground-state.
- 4 Semiconductors and insulators: **Core-valence interaction** is crucial for correct treatment of the EXX.
- 5 Lack of **Asymmetry** and/or **core-valence interaction** could lead to spurious agreement with experiments.

Conclusions

- 1 EXX within an all electron full potential method is implemented within **EXC!TING** code. Right now this is the only FP code to be able to do EXX.
- 2 A new and one of the most optimal basis is proposed for calculating and inverting the response. This basis may be useful for future TD-DFT and GW calculations.
- 3 Magnetic metals: **Asymmetry** in exchange potential is very important to get the correct ground-state.
- 4 Semiconductors and insulators: **Core-valence interaction** is crucial for correct treatment of the EXX.
- 5 Lack of **Asymmetry** and/or **core-valence interaction** could lead to spurious agreement with experiments.

Conclusions

- 1 EXX within an all electron full potential method is implemented within **EXC!TING** code. Right now this is the only FP code to be able to do EXX.
- 2 A new and one of the most optimal basis is proposed for calculating and inverting the response. This basis may be useful for future TD-DFT and GW calculations.
- 3 Magnetic metals: **Asymmetry** in exchange potential is very important to get the correct ground-state.
- 4 Semiconductors and insulators: **Core-valence interaction** is crucial for correct treatment of the EXX.
- 5 Lack of **Asymmetry** and/or **core-valence interaction** could lead to spurious agreement with experiments.

Conclusions

- 1 EXX within an all electron full potential method is implemented within **EXC!TING** code. Right now this is the only FP code to be able to do EXX.
- 2 A new and one of the most optimal basis is proposed for calculating and inverting the response. This basis may be useful for future TD-DFT and GW calculations.
- 3 Magnetic metals: **Asymmetry** in exchange potential is very important to get the correct ground-state.
- 4 Semiconductors and insulators: **Core-valence interaction** is crucial for correct treatment of the EXX.
- 5 Lack of **Asymmetry** and/or **core-valence interaction** could lead to spurious agreement with experiments.

Outlook

- 1 EXX can be generalised to handle non-collinear magnetism (derivatives w.r.t. $n_{\sigma\sigma'}(\mathbf{r})$).
- 2 Future inclusion of exact correlation may be possible (multiconfiguration approach or adiabatic fluctuation dissipation)

Outlook

- 1 EXX can be generalised to handle non-collinear magnetism (derivatives w.r.t. $n_{\sigma\sigma'}(\mathbf{r})$).
- 2 Future inclusion of exact correlation may be possible (multiconfiguration approach or adiabatic fluctuation dissipation)

Acknowledgements

Prof. P. Mohn

Magnetic metals work done in collaboration with Dr. C. Persson.

Austrian Science Fund (project P16227)

EXCITING network funded by the EU (HPRN-CT-2002-00317)

Code available at:

http://physik.kfunigraz.ac.at/~kde/secret_garden/exciting.html