

Linear Algebra

Theorem 2

Let $(V; F; +; \cdot)$ be a vector space over the field F . Let $n \in \mathbb{N}^*$ and $U = \{\vec{u}_1, \vec{u}_2, \dots, \vec{u}_n\}$ be a set of vectors $\in V$. Then $\text{span}(U)$ is a subspace of V .

Proof. Let $\text{span}(U)$ be the set of vectors which can be written as a linear combination of the vectors in U

$$\bullet \vec{0} \in \text{span}(U) \text{ because } \vec{0} = \overbrace{\lambda_1}^0 \cdot \vec{u}_1 + \overbrace{\lambda_2}^0 \cdot \vec{u}_2 + \overbrace{\lambda_n}^0 \cdot \vec{u}_n$$

- Let $\vec{u}, \vec{w} \in \text{span}(U)$ and let $\mu \in F$
We want to show that $\vec{u} + \mu \cdot \vec{w} \in \text{span}(U)$

$$\text{Let } \vec{u} = \alpha_1 \vec{u}_1 + \dots + \alpha_n \vec{u}_n \quad \vec{w} = \beta_1 \vec{u}_1 + \dots + \beta_n \vec{u}_n$$

$$\begin{aligned} \vec{u} + \mu \vec{w} &= \alpha_1 \vec{u}_1 + \dots + \alpha_n \vec{u}_n + \mu (\beta_1 \vec{u}_1 + \dots + \beta_n \vec{u}_n) = \\ &= \underbrace{(\alpha_1 + \mu \beta_1)}_{\in F} \cdot \vec{u}_1 + \dots + \underbrace{(\alpha_n + \mu \beta_n)}_{\in F} \cdot \vec{u}_n \in \text{span}(U) \Rightarrow \text{span}(U) \text{ is a subset of } V. \end{aligned}$$

□

Theorem 3

Uniqueness of representation. Let $(V; F; +; \cdot)$ be a vector space over the field F . Let $B = \{\vec{b}_1, \vec{b}_2, \dots, \vec{b}_n\}$ be a set of vectors $\in V$. Then B is a basis of V if and only if every vector $\vec{v} \in V$ can be written uniquely as a linear combination of the basis vectors of B .

Proof. Let $B = \{\vec{b}_1, \vec{b}_2, \dots, \vec{b}_n\}$ be a set of vectors in V .

→ Let B be a basis of V .

We want to show that every vector $\vec{v} \in V$ can be written uniquely as a linear combination of the vectors of B .

Assume that \vec{v} can't be written uniquely as a linear combination of the vectors of B :

Since B is a basis of V , then \vec{v} can be written as a linear combination of the vectors of B .

$$\vec{v} = \lambda_1 \cdot \vec{b}_1 + \dots + \lambda_n \cdot \vec{b}_n \text{ and } \vec{v} = \alpha_1 \cdot \vec{b}_1 + \dots + \alpha_n \cdot \vec{b}_n$$

$$\vec{v} - \vec{v} = (\lambda_1 \cdot \vec{b}_1 + \dots + \lambda_n \cdot \vec{b}_n) - (\alpha_1 \cdot \vec{b}_1 + \dots + \alpha_n \cdot \vec{b}_n) = \vec{0}$$

$\iff \vec{0} = (\lambda_1 - \alpha_1) \cdot \vec{b}_1 + \dots + (\lambda_n - \alpha_n) \cdot \vec{b}_n$ // Since B is a basis of V , then $\vec{b}_1, \dots, \vec{b}_n$ are linearly independent which implies:

$$\lambda_1 - \alpha_1 = 0, \dots, \lambda_n - \alpha_n = 0 \iff \lambda_1 = \alpha_1, \dots, \lambda_n = \alpha_n$$

which is a contradiction, since it proves that any vector $\vec{v} \in V$ is written uniquely as a linear combination of the basis vectors of B .

← Assume that every vector $\vec{v} \in V$ can be written uniquely as a linear combination of the vectors of B .

We want to show that B is a basis of V

Let $\vec{v} \in \text{span}(B)$: $\vec{v} = \alpha_1 \cdot \vec{b}_1 + \dots + \alpha_n \cdot \vec{b}_n \in V$ since V is a vector space and closed under vector addition and scalar multiplication. Hence $\text{span}(B) = V$ since every vector $\vec{v} \in V$ can be written as a linear combination of the vectors of B by hypothesis.

$$\begin{cases} \vec{0} = \lambda_1 \cdot \vec{b}_1 + \dots + \lambda_n \cdot \vec{b}_n \\ \vec{0} = 0 \cdot \vec{b}_1 + \dots + 0 \cdot \vec{b}_n \end{cases}$$

Since we assumed that there is only one way to write every vector $\vec{v} \in V$, then $\lambda_1 = 0, \dots, \lambda_n = 0$. It proves that the vectors of the set B are linearly independent. $\Rightarrow B$ is a basis of V . □

Theorem 5

Kernel. Let $f: V \rightarrow W$ be a linear map. Then $\ker(f)$ is a subspace of V .

Proof. Let $f: V \rightarrow W$ be a linear map.

$$f(\vec{u} + \lambda \vec{v}) = f(\vec{u}) + \lambda \cdot f(\vec{v}) \quad \forall \vec{u}, \vec{v} \in V$$

Since V is a vector space, then $V \neq \emptyset$ by definition.

$\Rightarrow \exists \vec{u} \in V$ and since V is a vector space, there is closure under scalar multiplication, ie $\lambda \cdot \vec{u} \in V$
 $\forall \lambda \in$

$$\Rightarrow 0 \cdot \vec{u} \in V \iff \vec{0} \in V$$

1) We want to show that $\ker(f) \neq \emptyset$ by showing that $\vec{0} \in \ker(f)$

$$f(\underbrace{\vec{0}}_{\in V}) = f(0 \cdot \vec{u}) \xrightarrow{f = \text{lin. map}} 0 \cdot f(\vec{u}) = \underbrace{\vec{0}}_{\in W} \Rightarrow \vec{0} \in \ker(f)$$

2) Let $\vec{u}, \vec{v} \in \ker(f), \lambda \in \mathbb{R}$

Since $\vec{u}, \vec{v} \in \ker(f) \Rightarrow f(\vec{u}) = \vec{0}$ and $f(\vec{v}) = \vec{0}$

$$f(\vec{u} + \lambda \vec{v}) \xrightarrow{f = \text{lin. map}} f(\vec{u}) + \lambda \cdot f(\vec{v}) = \vec{0} + \lambda \vec{0} = \vec{0} \Rightarrow \vec{u} + \lambda \vec{v} \in \ker(f)$$

Since $\vec{0} \in \ker(f)$ and $\vec{u} + \lambda \vec{v} \in \ker(f) \quad \forall \vec{u}, \vec{v} \in \ker(f)$

$\Rightarrow \ker(f)$ is a subspace of V

□

Theorem 6

Range. Let $f: V \rightarrow W$ be a linear map. Then $\text{Im}(f)$ is a subspace of W .

Proof.

1) We have show that $f(\vec{0}) = \underbrace{\vec{0}}_{\in W}$

$\vec{0} \in \text{Im}(f)$ since $\exists \vec{0} \in V$ such that $f(\vec{0}) = \vec{0}$

2) Let $\vec{w}_1, \vec{w}_2 \in \text{Im}(f)$

$\exists \vec{u}_1, \vec{u}_2 \in V$ such that $f(\vec{u}_1) = \vec{w}_1$ and $f(\vec{u}_2) = \vec{w}_2$

Let $\lambda \in \mathbb{R}$: We want to show that $\vec{w}_1 + \lambda \vec{w}_2 \in \text{Im}(f)$

$$\vec{w}_1 + \lambda \cdot \vec{w}_2 = f(\vec{u}_1) + \lambda \cdot f(\vec{u}_2) \xrightarrow{f = \text{lin. map}} \underbrace{f(\vec{u}_1 + \lambda \cdot \vec{u}_2)}_{\in W (\text{since } V \text{ is a vec. space})}$$

Since $\exists \vec{u}_1 + \lambda \vec{u}_2 \in V$ such that

$f(\vec{u}_1 + \lambda \vec{u}_2) = \vec{w}_1 + \lambda \vec{w}_2$, then $\vec{w}_1 + \lambda \vec{w}_2 \in \text{Im}(f)$

$\Rightarrow \text{Im}(f)$ is a subspace of W

□