

Math Review Notes—Asymptotics and Convergence

Gregory Faletto

Contents

1	Asymptotics and Convergence	3
1.1	Preliminaries (5.9 and 7.1, Grimmett and Stirzaker)	3
1.2	Inequalities (8.6 of Pesaran)	4
1.3	Modes of Convergence (7.2 of Grimmett and Stirzaker, 8.2 and 8.4 of Pesaran)	9
1.4	More on convergence (7.2 of Grimmett and Stirzaker)	13
1.4.1	Slutsky's Convergence Theorems (8.4.1 of Pesaran, 7.3 of Grimmett and Stirzaker) . .	17
1.5	Stochastic orders $\mathcal{O}_p(\cdot)$ and $o_p(\cdot)$ (Pesaran 8.5)	18
1.6	Laws of Large Numbers and Central Limit Theorems (Pesaran 8.6; Grimmett and Stirzaker 7.4, 7.5)	19
1.7	The case of dependent and heterogeneously distributed observations (Pesaran 8.8)	21
1.8	Worked Examples from Math 505A Midterm 2	22
1.9	Estimators and Central Limit Theorems (DSO 607)	27

Last updated July 11, 2019

1 Asymptotics and Convergence

These notes are based on my notes from chapter 8 of *Time Series and Panel Data Econometrics* (1st edition) by M. Hashem Pesaran [Pesaran, 2015] and coursework for Economics 613: Economic and Financial Time Series I at USC, as well as Math 505A and Math 541A at USC and chapter 7 from *Probability and Random Processes* (Grimmett and Stirzaker) 3rd edition [Grimmett and Stirzaker, 2001].

1.1 Preliminaries (5.9 and 7.1, Grimmett and Stirzaker)

Definition 1.1. Definition 7.1.4, Grimmett and Stirzaker. If for all $x \in [0, 1]$ the sequence $\{f_n(x)\}$ of real numbers satisfies $f_n(x) \rightarrow f(x)$ as $n \rightarrow \infty$ then we say $f_n \rightarrow f$ **pointwise**.

Remark. In practice pointwise convergence is often not useful for functions because a sequence of functions may be continuous while its limit is not. For instance, consider $\{f_n : f_n = x^n \forall x \in [0, 1]\}$. Then f_n is continuous for all n but

$$\lim_{n \rightarrow \infty} f_n = \begin{cases} 0 & x < 1 \\ 1 & x = 1 \end{cases}$$

Instead, the following definition is often more useful.

Definition 1.2. (from class notes.) We say that f_n **uniformly converges to f on $[a, b]$** if for every $\epsilon > 0$ there exists N such that for every $n > N$,

$$\forall x \in [a, b] |f_n(x) - f(x)| < \epsilon$$

Definition 1.3. (Definition 7.1.5, Grimmett and Stirzaker.) Let V be a collection of functions mapping $[0, 1]$ into \mathbb{R} and assume V is endowed with a function $\|\cdot\| : V \rightarrow \mathbb{R}$ satisfying

- (a) $\|f\| \geq 0$ for all $f \in V$
- (b) $\|f\| = 0$ if and only if f is the zero function (or equivalent to it)
- (c) $\|af\| = |a| \cdot \|f\|$ for all $a \in \mathbb{R}$, $f \in V$
- (d) $\|f + g\| \leq \|f\| + \|g\|$ (Triangle Inequality)

The function $\|\cdot\|$ is called a **norm**. If $\{f_n\}$ is a sequence of members of V then we say that $f_n \rightarrow f$ **with respect to the norm $\|\cdot\|$** if $\|f_n - f\| \rightarrow 0$ as $n \rightarrow \infty$.

Definition 1.4. (Definition 7.16, Grimmett and Stirzaker.) Let $\epsilon > 0$ be prescribed, and define the distance between two functions $g, h : [0, 1] \rightarrow \mathbb{R}$ by

$$d_\epsilon(g, h) = \int_E dx$$

where $E = \{u \in [0, 1] : |g(u) - h(u)| > \epsilon\}$. We say that $f_n \rightarrow f$ **in measure** if

$$d_\epsilon(f_n, f) \rightarrow 0 \text{ as } n \rightarrow \infty \text{ for all } \epsilon > 0$$

Theorem 1. Inversion Theorem (Theorem 5.9.2, Grimmett and Stirzaker). Let X have distribution function F and characteristic function ϕ . Define $\bar{F} : \mathbb{R} \rightarrow [0, 1]$ by

$$\bar{F}(x) = \frac{1}{2} [F(x) + \lim_{y \rightarrow x^-} F(y)]$$

Then

$$\bar{F}(b) - \bar{F}(a) = \lim_{N \rightarrow \infty} \int_{-N}^N \frac{\exp(-iat) - \exp(-ibt)}{2\pi it} \cdot \phi(t) dt$$

Proof. See [Kingman and Taylor \[1966\]](#). □

Corollary 1.1. Corollary 5.9.3. Random variables X and Y have the same characteristic function if and only if they have the same distribution function.

Proof. Available in Grimmett and Stirzaker section 5.9, pp. 189 - 190. □

Definition 1.5. (Definition 5.9.4, Grimmett and Stirzaker.) We say that the sequence F_1, F_2, \dots of distribution functions **converges** to the distribution function F (written $F_n \rightarrow F$) if $F(x) = \lim_{n \rightarrow \infty} F_n(x)$ at each point x where F is continuous.

Theorem 2. Continuity theorem (Theorem 5.9.5; in notes from Friday 10/26, Lecture 28).

Suppose that F_1, F_2, \dots is a sequence of distribution functions with corresponding characteristic functions ϕ_1, ϕ_2, \dots

- (a) If $F_n(x) \rightarrow F(x)$ for some distribution function F with characteristic function ϕ (at x where F is continuous), then $\phi_n(t) \rightarrow \phi(t)$ for all t .
- (b) Conversely, if $\phi(t) = \lim_{n \rightarrow \infty} \phi_n(t)$ exists and $\phi(t)$ is continuous at $t = 0$, then ϕ is the characteristic function of some distribution function F , and $F_n \rightarrow F$.

Proof. See [Kingman and Taylor \[1966\]](#). □

1.2 Inequalities (8.6 of Pesaran)

Inequalities

- Probabilities

—

Lemma 3. Markov's Inequality (Grimmett and Stirzaker p. 311, 319) : Let $X : \Omega \rightarrow [-\infty, \infty]$ be a random variable. Then for all $a > 0$,

$$\Pr(|X| \geq a) \leq \frac{\mathbb{E}(|X|)}{a}$$

Proof. Note $t \cdot \mathbf{1}_{\{|X| \geq t\}} \leq |X|$, where $\mathbf{1}$ is the indicator function. Dividing both sides by t and taking expectations, we have

$$\mathbb{E}(\mathbf{1}_{\{|X| \geq t\}}) \leq \frac{\mathbb{E}|X|}{t} \iff \Pr(|X| \geq t) \leq \frac{\mathbb{E}|X|}{t}, \quad \forall t > 0.$$

□

Corollary 3.1. If n is a positive integer, then

$$\Pr(|X| \geq t) \leq \frac{\mathbb{E}(|X|^n)}{t^n} \quad \forall t > 0$$

Proof. By Markov's Inequality (Theorem 3),

$$\Pr(|X| \geq t) = \Pr(|X|^n \geq t^n) \leq \frac{\mathbb{E}(|X|^n)}{t^n}$$

□

—

Theorem 4. Chebyshev's Inequality: (probability p. 319) Let $X : \Omega \rightarrow [-\infty, \infty]$ be an (integrable) random variable with $\mathbb{E}(X^2) < \infty$. Then for any real number $k > 0$

$$\Pr(|X - \mathbb{E}(X)| \geq k\sqrt{\text{Var}(X)}) \leq \frac{1}{k^2}$$

This can also be written as

$$\Pr(|X - \mathbb{E}(X)| \geq k) \leq \frac{\text{Var}(X)}{k^2}$$

(Can be used to demonstrate consistency of estimators: if we can show that as $T \rightarrow \infty$ $\text{Var}(X) = \sigma^2 \rightarrow 0$, then this implies $\Pr(|X - \mu| \geq k\sigma) \rightarrow 0$ as $T \rightarrow \infty$, showing consistency.)

—

Theorem 5. Chernoff For $x \geq 0$, $a > 0$, $\forall t > 0$,

$$\Pr(X \geq a) = \Pr(e^{tX} \geq e^{ta}) \leq \frac{\mathbb{E}(e^{tX})}{e^{ta}}$$

• Moments

—

Theorem 6 (Cauchy-Schwarz (and Bunyakovsky)). If X and Y are random variables with finite variance then

$$\mathbb{E}(XY)^2 \leq \mathbb{E}(X^2)\mathbb{E}(Y^2)$$

Note that this can be a corollary of Theorem 9 with $p = q = 2$. We can also prove this theorem on its own in a different one. We first prove a useful result.

Lemma 7. If $\text{Var}(X) = 0$ then X is almost surely constant; that is, $\Pr(X = a) = 1$ for some $a \in \mathbb{R}$.

Proof. Note that because $\text{Var}(X) = 0 < \infty$, we know that $\mathbb{E}(X)$ and $\mathbb{E}(X^2)$ exist. We have

$$\text{Var}(X) = \mathbb{E}[(X - \mathbb{E}(X))^2] = 0$$

Let $Y = (X - \mathbb{E}(X))^2$. Note that $Y = (X - \mathbb{E}(X))^2 \geq 0$ and that $\mathbb{E}(Y) = \text{Var}(X) = 0$. Therefore $\Pr(Y = 0) = 1$, so $\Pr(Y \neq 0) = 0$. To see why, in the case that X is discrete,

$$\mathbb{E}(Y) = \sum_{k=0}^{\infty} k \cdot \Pr(Y = k) = \text{Var}(X) = 0$$

which is true if and only if $\Pr(Y = k) = 0$ for all $k > 0$. Since we already showed that $\Pr(Y < 0) = 0$, it follows that $\Pr(Y = 0) = 1$. In the continuous case,

$$\mathbb{E}(Y) = \int_0^{\infty} y \cdot f_Y(y) dy = \text{Var}(X) = 0$$

which implies that $f_Y(x) = 0$ for all $x > 0$. Again, since $\Pr(Y < 0) = 0$, we have $\Pr(Y \neq 0) = 0$. But $Y = 0 \iff X = \mathbb{E}(X)$ so we have $\Pr(X = \mathbb{E}(X)) = 1$. \square

Remark. Note that Lemma 7 along with Proposition ?? imply that X has variance 0 if and only if it is (almost surely) constant.

We are now ready to prove the Cauchy-Schwarz Inequality.

Proof. if $\mathbb{E}(X^2) = 0$ or $\mathbb{E}(Y^2) = 0$, the Cauchy-Schwarz Inequality follows immediately. To see why, suppose without loss of generality that $\mathbb{E}(X^2) = 0$. Then the right side is 0. Also, $0 \leq \text{Var}(X) = \mathbb{E}(X^2) - \mathbb{E}(X)^2 = -\mathbb{E}(X)^2$. Since $\mathbb{E}(X)^2 \geq 0$, we must have $\mathbb{E}(X)^2 = 0$ and therefore $\text{Var}(X) = 0$. Therefore by Lemma 7, X is almost surely constant, which means that $\text{Cov}(X, Y) = 0$.

In the case that $\mathbb{E}(X^2) > 0$ and $\mathbb{E}(Y^2) > 0$, for $a, b \in \mathbb{R}$, let $Z = aX - bY$. Then

$$0 \leq \mathbb{E}(Z^2) = a^2\mathbb{E}(X^2) - 2ab\mathbb{E}(XY) + b^2\mathbb{E}(Y^2) \quad (1)$$

The right side of (1) is quadratic in a . Because it is greater than or equal to zero, it has at most one real root, which means its discriminant must be non-positive. That is, if $b \neq 0$,

$$(-2b\mathbb{E}(XY))^2 - 4b^2\mathbb{E}(X^2)\mathbb{E}(Y^2) \leq 0 \iff \mathbb{E}(XY)^2 - \mathbb{E}(X^2)\mathbb{E}(Y^2) \leq 0$$

which yields the result. Note that equality holds if and only if $\Pr(aX = bY) = 1$ because the discriminant is zero if and only if the quadratic has a real root, which occurs if and only if

$$\mathbb{E}[(aX - bY)^2] = 0$$

which is true if and only if $\Pr(aX = bY) = 1$ by Lemma 7 and Proposition ??. \square

– **Krylov**

–

Definition 1.6. Let $\phi : \mathbb{R} \rightarrow \mathbb{R}$. We say that ϕ is **convex** if for any $x, y \in \mathbb{R}$ and for any $t \in [0, 1]$, we have

$$\phi(tx + (1 - t)y) \leq t\phi(x) + (1 - t)\phi(y)$$

Theorem 8 (Jensen's Inequality, from Math 541A. Also Grimmett and Stirzaker p.181, 349). Let $X : \Omega \rightarrow [-\infty, \infty]$ be a random variable. Let $\phi : \mathbb{R} \rightarrow \mathbb{R}$ be convex. If $\mathbb{E}|X| < \infty$ and $\mathbb{E}|\phi(X)| < \infty$, then

$$\phi(\mathbb{E}X) \leq \mathbb{E}\phi(X).$$

(See also Theorem ??.)

For the definition of convexity, see Definition ??.

Proof. Note that from Theorem ??, for any $y \in \mathbb{R}$ there exists a constant a and a function L such that

$$a(x - y) + \phi(y) \leq \phi(x) \quad \forall x \in \mathbb{R}$$

Letting $y = \mathbb{E}(X)$ we have

$$a(X - \mathbb{E}X) + \phi(\mathbb{E}X) \leq \phi(X)$$

Since expectations preserve inequalities,

$$\mathbb{E}[a(X - \mathbb{E}X) + \phi(\mathbb{E}X)] \leq \mathbb{E}\phi(X)$$

But

$$\mathbb{E}[a(X - \mathbb{E}X) + \phi(\mathbb{E}X)] = a(\mathbb{E}X - \mathbb{E}X) + \mathbb{E}(\phi(\mathbb{E}X)) = \phi(\mathbb{E}X)$$

which yields

$$\phi(\mathbb{E}X) \leq \mathbb{E}\phi(X).$$

□

For some corollaries, see section ??.

Theorem 9. [Hölder (Grimmett and Stirzaker p. p. 143, 319; Theorem 1.99 in Math 541A lecture notes) Generalization of Cauchy-Schwarz] Let $X, Y : \Omega \rightarrow \mathbb{R}$ be random variables. For $p, q \geq 1$ satisfying $1/p + 1/q = 1$ we have

$$\mathbb{E}(|XY|) \leq (\mathbb{E}(|X|^p))^{1/p} (\mathbb{E}(|Y|^q))^{1/q} = \|X\|_p \|Y\|_q.$$

The equality case happens only if X is a constant multiple of Y with probability 1. Note that the case $p = q = 2$ recovers the Cauchy-Schwarz Inequality (Theorem 6).

Proof. Assume without loss of generality that $\|X\|_p = \|Y\|_q = 1$. Also, the case $p = 1, q = \infty$ follows from the triangle inequality, so we assume $1 < p < \infty$. From concavity of the log function, we have

$$\begin{aligned} \log((x^p)^{1/p} (y^q)^{1/q}) &= (1/p) \log(x^p) + (1/q) \log(y^q) \\ &\leq \log\left(\frac{1}{p} x^p + \frac{1}{q} y^q\right) \end{aligned}$$

$$\implies (x^p)^{1/p}(y^q)^{1/q} \leq \frac{1}{p}x^p + \frac{1}{q}y^q$$

Fixing an $\omega \in \Omega$, we have

$$|X(\omega)Y(\omega)| = (|X(\omega)|^p)^{1/p}(|Y(\omega)|^q)^{1/q} \leq \frac{1}{p}|X(\omega)|^p + \frac{1}{q}|Y(\omega)|^q$$

Integrating we have...

□

Theorem 10 (Hölder (vector form)). For any $u, v \in \mathbb{R}^n$,

$$|u^T v| \leq \|u\|_p \|v\|_q$$

for any $p, q \in [0, \infty]$ satisfying $1/p + 1/q = 1$.

—

Theorem 11. Minkowski (Grimmett and Stirzaker p. p. 143) For $p \geq 1$,

$$[\mathbb{E}(|X + Y|^p)]^{1/p} \leq (\mathbb{E}|X|^p)^{1/p} + (\mathbb{E}|Y|^p)^{1/p}$$

– Useful for showing lower order moments are finite (e.g. finite variance implies finite mean).

Lemma 12. Lyapunov's Inequality (Grimmett and Stirzaker p. 143). For $0 < r \leq s < \infty$,

$$\mathbb{E}(|X|^r)^{1/r} \leq \mathbb{E}(|X|^s)^{1/s}$$

—

Theorem 13. Triangle Inequality: Let $X, Y : \Omega \rightarrow \mathbb{R}$ be random variables. Let $1 \leq p \leq \infty$. Then

$$\|X + Y\|_p \leq \|X\|_p + \|Y\|_p, 1 \leq p \leq \infty$$

Proof. The case $p = \infty$ follows from the scalar triangle inequality, so assume $1 \leq p < \infty$. By scaling, we may assume $\|X\|_p = 1 - t, \|Y\|_p = t$, for some $t \in (0, 1)$ (zeroes and infinities being trivial). Define $V := X/(1 - t), W := Y/t$. Then by convexity of $x \rightarrow |x|^p$ on \mathbb{R} ,

$$|(1 - t)V(\omega) + t(W(\omega))|^p \leq (1 - t)|V(\omega)|^p + t|W(\omega)|^p$$

Take expectation of both sides:

$$\mathbb{E}|X + Y|^p \leq (1 - t)^{1-p} \mathbb{E}|X|^p + t^{1-p} \mathbb{E}|Y|^p$$

Since $\|X\|_p = t, \|Y\|_p = 1 - t$, we have that the right side is $1 - t + t = 1$. (Note: $\|Y\|_p = t, \mathbb{E}|Y|^p = t^p, \|X\|_p = 1 - t$ Therefore

$$(\mathbb{E}|X + Y|^p)^{1/p} = \|X + Y\|_p \leq 1$$

□

Remark. See also Theorem ?? and Corollary ??.

Theorem 14 (Chernoff Bound). Let X be a random variable and let $r > 0$. Define $M_X(t) := \mathbb{E}e^{tX}$ for any $t \in \mathbb{R}$. Then for any $t > 0$,

$$\mathbb{P}(X > r) \leq e^{-tr} M_X(t).$$

Proof. Using Markov's Inequality (Theorem 3) on e^{tX} , we have

$$\Pr(X \geq r) = \Pr(e^{tX} \geq e^{tr}) \leq \frac{\mathbb{E}e^{tX}}{e^{tr}} = e^{-tr} M_X(t), \quad \forall t > 0.$$

□

Remark. Consequently, if X_1, \dots, X_n are independent random variables with the same CDF, and if $r, t > 0$,

$$\mathbb{P}\left(\frac{1}{n} \sum_{i=1}^n X_i > r\right) \leq e^{-trn} (M_{X_1}(t))^n.$$

For example, if X_1, \dots, X_n are independent Bernoulli random variables with parameter $0 < p < 1$, and if $r, t > 0$,

$$\mathbb{P}\left(\frac{X_1 + \dots + X_n}{n} - p > r\right) \leq e^{-trn} (e^{-tp}[pe^t + (1-p)])^n.$$

And if we choose t appropriately, then the quantity $\mathbb{P}\left(\frac{1}{n} \sum_{i=1}^n (X_i - p) > r\right)$ becomes exponentially small as either n or r become large. That is, $\frac{1}{n} \sum_{i=1}^n X_i$ becomes very close to its mean. Importantly, the Chernoff bound is much stronger than either Markov's or Cheyshev's inequality, since they only respectively imply that

$$\mathbb{P}\left(\left|\frac{X_1 + \dots + X_n}{n} - p\right| > r\right) \leq \frac{2p(1-p)}{r}, \quad \mathbb{P}\left(\left|\frac{X_1 + \dots + X_n}{n} - p\right| > r\right) \leq \frac{p(1-p)}{nr^2}.$$

Monotone convergence theorem.

Dominated Convergence Theorem (Theorem ??).

1.3 Modes of Convergence (7.2 of Grimmett and Stirzaker, 8.2 and 8.4 of Pesaran)

Let $\{X_n\} = \{X_1, X_2, \dots\}$ and X be random variables defined on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$.

Definition 1.7. Convergence in probability. $\{X_n\}$ is said to **converge in probability** to X if

- Grimmett and Strizaker definition:

$$\lim_{n \rightarrow \infty} \Pr(|X_n - X| > \epsilon) = 0, \text{ for every } \epsilon > 0$$

- Pesaran definition:

$$\lim_{n \rightarrow \infty} \Pr(|X_n - X| < \epsilon) = 1, \text{ for every } \epsilon > 0$$

- More formal (from Math 541A):

$$\forall \epsilon > 0, \lim_{n \rightarrow \infty} \Pr(\{\omega \in \Omega : |X_n(\omega) - X(\omega)| > \epsilon\}) = 0$$

Remark. This mode of convergence is also often denoted by $X_n \xrightarrow{p} X$ and when X is a fixed constant it is referred to as the **probability limit of X_n** , written as $\text{Plim}(X_n) = x$, as $n \rightarrow \infty$.

The above concept is readily extended to multivariate cases where $\{\mathbf{X}_n, n = 1, 2, \dots\}$ denote m -dimensional vectors of random variables. Then the condition is

$$\lim_{n \rightarrow \infty} \Pr(\|\mathbf{X}_n - \mathbf{X}\| < \epsilon) = 1, \text{ for every } \epsilon > 0$$

where $\|\cdot\|$ denotes an appropriate norm (say ℓ_2). Convergence in probability is often referred to as "weak convergence" (in contrast to convergence with probability 1, below).

Definition 1.8. Convergence with probability 1 or almost surely. The sequence of random variables $\{X_n\}$ is said to **converge with probability 1** (or **almost surely**) to X if

- (505A class notes definition)

$$\Pr(\{\omega \in \Omega : \lim_{n \rightarrow \infty} X_n(\omega) = X(\omega)\}) = 1$$

(Note: pointwise convergence can hardly ever be shown here and is not useful.)

- Grimmett and Stirzaker textbook definition:

$$\Pr(\{\omega \in \Omega : X_n(\omega) \rightarrow X(\omega) \text{ as } n \rightarrow \infty\}) = 1$$

- Pesaran textbook definition:

$$\Pr\left(\lim_{n \rightarrow \infty} X_n = X\right) = 1$$

Remark. This is often written as $X_n \xrightarrow{w.p.1} X$ or $X_n \xrightarrow{a.s.} X$. An equivalent condition for convergence with probability 1 is given by

$$\lim_{n \rightarrow \infty} \Pr(|X_m - X| < \epsilon, \text{ for all } m \geq n) = 1, \text{ for every } \epsilon > 0$$

which shows that convergence in probability is a special case of convergence with probability 1 (obtained by setting $m = n$). Convergence with probability 1 is stronger than convergence in probability and is often referred to as "strong convergence."

Definition 1.9. Convergence in r -th mean or convergence in ℓ_p . $X_n \rightarrow X$ in r th mean (or in ℓ_p) where $r \geq 1$ (or $0 < p \leq \infty$) if $\mathbb{E}|X_n^r| < \infty$ for all n and

$$\lim_{n \rightarrow \infty} \mathbb{E}(|X_n - X|^r) = 0$$

or if $\|X\|_p < \infty$ and

$$\lim_{n \rightarrow \infty} \|X_n - X\|_p = 0$$

Remark. Recall that $\|X\|_p := (\mathbb{E}(X)^p)^{1/p}$ if $0 < p < \infty$ and $\|X\|_\infty := \inf\{c > 0 : \Pr(|X| \leq c) = 1\}$. Note that if $p < 1$, $\|\cdot\|_p$ is no longer a norm because it does not satisfy the Triangle Inequality (Corollary ?? and Theorem 13), but this property still holds. Convergence in r th mean is often written $X_n \xrightarrow{r} X$.

Definition 1.10. Convergence in Distribution. Let X_1, X_2, \dots have distribution functions $F_1(\cdot), F_2(\cdot), \dots$ respectively. Then X_n is said to **converge in distribution to X** if

$$\lim_{n \rightarrow \infty} \Pr(X_n \leq u) = \Pr(X \leq u)$$

for all u at which $F_X(x) = \Pr(X \leq x)$ is continuous. This can also be written

$$\lim_{n \rightarrow \infty} F_n(u) = F(u)$$

for all u at which F is continuous.

Remark. Convergence in distribution is usually denoted by $X_n \xrightarrow{d} X$, $X_n \xrightarrow{L} X$, or $F_n \implies F$. By the Continuity Theorem (Theorem 2, section 1.1), this is equivalent to

$$\lim_{n \rightarrow \infty} \phi_{X_n}(t) = \phi_X(t), \quad t \in \mathbb{R}.$$

Note that the random variables are allowed to have different domains.

Definition 1.11. (Convergence in distribution for vector-valued random variables.) We say that random variables $Y^{(1)}, \dots, : \Omega \rightarrow \mathbb{R}^d$ **converge in distribution** to $Y : \Omega \rightarrow \mathbb{R}^d$ if for all $v \in \mathbb{R}^d$, $\langle v, Y^{(1)} \rangle, \langle v, Y^{(2)} \rangle, \dots$ converges in distribution to $\langle v, Y \rangle$.

Theorem 15 ((Theorem 7.2.3, Grimmett and Stirzaker.)). The following implications hold:

- (a) $(X_n \xrightarrow{a.s.} X) \implies (X_n \xrightarrow{p} X)$
- (b) $(X_n \xrightarrow{r} X) \implies (X_n \xrightarrow{p} X)$ for any $r \geq 1$
- (c) $(X_n \xrightarrow{p} X) \implies (X_n \xrightarrow{d} X)$

Also, if $r > s \geq 1$, then $(X_n \xrightarrow{r} X) \implies (X_n \xrightarrow{s} X)$. No other implications hold in general.

Proof. (a) By Markov's Inequality (Lemma 3),

$$\Pr(|X_n - X| > \epsilon) \leq \frac{\mathbb{E}|X_n - X|}{\epsilon} \quad \text{for all } \epsilon > 0$$

Therefore if $X_n \xrightarrow{1} X$; that is, $\lim_{n \rightarrow \infty} \mathbb{E}(|X_n - X|) = 0$, then $\lim_{n \rightarrow \infty} \Pr(|X_n - X| > \epsilon) = 0$ for every $\epsilon > 0$, so $X_n \xrightarrow{p} X$.

(b) Let $\epsilon > 0$ and let $0 < r < \infty$. From Markov's Inequality,

$$\Pr(|X_n - X| > \epsilon) = \Pr(|X_n - X|^r > \epsilon^r) \leq \frac{\mathbb{E}|X_n - X|^r}{\epsilon^r}$$

The right side converges to zero by assumption. Therefore X_1, X_2, \dots converges to X in probability. The case $r = \infty$ follows from any case $r < \infty$, since for example

$$(\mathbb{E}(X_n) - X)^2 \leq \mathbb{E}(|X_n - X|^2) \leq \|X_n - X\|_\infty$$

which shows that L_∞ convergence implies L_2 convergence. So since L_2 convergence implies convergence in probability, L_∞ convergence does too. We can also see this by simply examining the definition of the L_∞ norm:

$$\|X_n - X\|_\infty := \inf\{c > 0 : \Pr(|X_n - X| \leq c) = 1\}.$$

Clearly if this number goes to 0, then $X_n \xrightarrow{p} X$.

(c)

□

Remark. Here are counterexamples showing that the converses are not always true:

(a) To see the converse fails, define an independent sequence $\{X_n\}$ by

$$X_n = \begin{cases} n^3 & \text{with probability } n^{-2} \\ 0 & \text{with probability } 1 - n^{-2} \end{cases}$$

Then $\Pr(|X_n| > \epsilon) = n^{-2}$ for all large n , and so $X_n \xrightarrow{p} 0$. However, $\mathbb{E}|X_n| = n \rightarrow \infty$.

(b) To see why the converse is false, fix $0 < r < \infty$, let $\Omega := [0, 1]$ with \mathbb{P} uniform on Ω and consider $X_n := n^{1/r} \mathbf{1}_{[0, 1/n]}$. Then X_1, X_2, \dots converges in probability to 0 since if $1 > \epsilon > 0$, then $\mathbb{P}(|X_n - 0| > \epsilon) \leq \mathbb{P}([0, 1/n]) = 1/n \rightarrow 0$ as $n \rightarrow \infty$. However, X_1, X_2, \dots does not converge in L_r to 0 since $\mathbb{E}|X_n - 0|^r = n/n = 1$ for all $n \geq 1$.

(c)

Theorem 16. Some exceptions (Theorem 7.2.4).

(a) If $X_n \xrightarrow{d} c$ where c is constant, then $X_n \xrightarrow{p} c$.

(b) If $X_n \xrightarrow{p} X$ and $\Pr(|X_n| \leq k) = 0$ for all n and some k , then $X_n \xrightarrow{r} X$ for all $r \geq 1$.

(c) If $P_n(\epsilon) = \Pr(|X_n - X| > \epsilon)$ satisfies $\sum_n P_n(\epsilon) < \infty$ for all $\epsilon > 0$, then $X_n \xrightarrow{a.s.} X$.

Proof. (Part (c).) Let $A_n(\epsilon) = \{|X_n - X| > \epsilon\}$ (so that $P_n(\epsilon) = \Pr[A_n(\epsilon)]$), and let $B_m(\epsilon) = \bigcup_{n \geq m} A_n(\epsilon)$. Then

$$\Pr(B_m(\epsilon)) \leq \sum_{n=m}^{\infty} \Pr(A_n(\epsilon))$$

so $\lim_{m \rightarrow \infty} \Pr(B_m(\epsilon)) = 0$ whenever $\sum_n \Pr(A_n(\epsilon)) < \infty$. See also Lemma 18 part (b). □

Lemma 17. (Lemma 7.2.6 from Grimmett and Stirzaker)

- (a) If $r > s \geq 1$ and $X_n \xrightarrow{r} X$, then $X_n \xrightarrow{s} X$.
- (b) If $X_n \xrightarrow{1} X$ then $X_n \xrightarrow{p} X$.

The converse assertions fail in general.

Proof. (a) Using Lyapunov's Inequality (Lemma 12), if $r > s \geq 1$

$$[\mathbb{E}(|X_n - X|^s)]^{1/s} \leq [\mathbb{E}(|X_n - X|^r)]^{1/r}$$

Therefore if $X_n \xrightarrow{r} X$ (meaning $\lim_{n \rightarrow \infty} \mathbb{E}(|X_n - X|^r) = 0$), (then $\lim_{n \rightarrow \infty} \mathbb{E}(|X_n - X|^s) = 0$, so $X_n \xrightarrow{s} X$. We show the converse fails by counterexample:

$$X_n = \begin{cases} n & \text{with probability } n^{-(1/2)(r+s)} \\ 0 & \text{with probability } 1 - n^{-(1/2)(r+s)} \end{cases}$$

Then $\mathbb{E}|X_n^s| = n^{(1/2)(s-r)} \rightarrow 0$ and $\mathbb{E}|X_n^r| = n^{(1/2)(r-s)} \rightarrow \infty$.

- (b) See proof of Theorem 15(a).

□

1.4 More on convergence (7.2 of Grimmett and Stirzaker)

Other theorems to include: Fatou's Lemma, Fubini's Theorem, Kolmogorov's Maximal Inequality, Kolmogorov Three-Series Test, Lindeberg Feller Central Limit Theorem, **this and more at beginning of Mike's 505A qual solutions.**

Definition 1.12. Cauchy Convergence. We say that the sequence $\{X_n : n \geq 1\}$ of random variables on the probability space $(\Omega, \mathcal{F}, \mathbb{P})$ is **almost surely Cauchy convergent** if

$$\Pr(\{\omega \in \Omega : X_m(\omega) - X_n(\omega) \rightarrow 0 \text{ as } m, n \rightarrow \infty\}) = 1$$

That is, the set of points ω of the sample space for which the real sequence $\{X_n(\omega) : n \geq 1\}$ is Cauchy convergent is an event having probability 1.

Lemma 18. (Lemma 7.2.10, Grimmett and Stirzaker.) Let $A_n(\epsilon) = \{|X_n - X| > \epsilon\}$ and $B_m(\epsilon) = \cup_{n \geq m} A_n(\epsilon)$. Then:

- (a) $X_n \xrightarrow{a.s.} X$ if and only if $\Pr(B_m(\epsilon)) \rightarrow 0$ as $m \rightarrow \infty$ for all $\epsilon > 0$.
- (b) $X_n \xrightarrow{a.s.} X$ if $\sum_n \Pr(A_n(\epsilon)) < \infty$ for all $\epsilon > 0$.
- (c) If $X_n \xrightarrow{a.s.} X$ then $X_n \xrightarrow{p} X$, but the converse fails in general.

Proof. (a)

(b) As for Theorem 16 part (c).

(c) To see the converse fails, define an independent sequence $\{X_n\}$ by

$$X_n = \begin{cases} 1 & \text{with probability } n^{-1} \\ 0 & \text{with probability } 1 - n^{-1} \end{cases}$$

Clearly $X_n \xrightarrow{p} 0$. However, if $0 < \epsilon < 1$,

$$\Pr(B_m(\epsilon)) = 1 - \lim_{r \rightarrow \infty} \Pr(X_n = 0 \text{ for all } n \text{ such that } m \leq n \leq r) \text{ (by Lemma 1.3.5)}$$

$$= 1 - \left(1 - \frac{1}{m}\right) \left(1 - \frac{1}{m+1}\right) \cdots \text{ (by independence)}$$

$$= 1 - \lim_{M \rightarrow \infty} \left(\frac{m-1}{m} \cdot \frac{m}{m+1} \cdot \frac{m+1}{m+2} \cdots \frac{M}{M+1} \right)$$

$$= 1 - \lim_{M \rightarrow \infty} \frac{m-1}{M+1} = 1$$

and so $\{X_n\}$ does not converge almost surely. □

Lemma 19. (Lemma 7.2.12, Grimmett and Stirzaker.) There exist sequences which

(a) converge almost surely but not in mean,

(b) converge in mean but not almost surely.

Proof. (a) As for Lemma 17 part (b). □

Theorem 20. (Theorem 7.2.13, Grimmett and Stirzaker.) If $X_n \xrightarrow{p} X$, there exists a non-random increasing sequence of integers n_1, n_2, \dots such that $X_{n_i} \xrightarrow{a.s.} X$ as $i \rightarrow \infty$.

Theorem 21. Skorokhod's representation theorem (Theorem 7.2.14, Grimmett and Stirzaker.) If $\{X_n\}$ and X with distribution functions $\{F_n\}$ and F are such that $X_n \xrightarrow{d} X$ (or equivalently, $F_n \rightarrow F$) as $n \rightarrow \infty$, then there exists a probability space $(\Omega', \mathcal{F}', \mathbb{P}')$ and random variables $\{Y_n\}$ and Y mapping Ω' into \mathbb{R} such that

(a) $\{Y_n\}$ and Y have distribution functions $\{F_n\}$ and F

(b) $Y_n \xrightarrow{a.s.} Y$ as $n \rightarrow \infty$

Therefore, although X_n may fail to converge to X in any mode other than in distribution, there exists a sequence $\{Y_n\}$ such that Y_n is distributed identically to X_n for every n , which converges almost surely to a copy of X .

Theorem 22. (Theorem 7.2.19, Grimmett and Stirzaker; same as Portmanteau Theorem?)

The following three statements are equivalent:

- (a) $X_n \xrightarrow{d} X$
- (b) $\mathbb{E}[g(X_n)] \rightarrow \mathbb{E}[g(X)]$ for all bounded continuous functions g .
- (c) $\mathbb{E}[g(X_n)] \rightarrow \mathbb{E}[g(X)]$ for all functions g of the form $g(x) = f(x)\mathbf{1}_{[a,b]}(x)$ where f is continuous on $[a, b]$ and a and b are points of continuity of the distribution function of the random variable X .

Theorem 23. (Grimmett and Stirzaker Theorem 7.3.9.)

- (a) If $X_n \xrightarrow{a.s.} X$ and $Y_n \xrightarrow{a.s.} Y$ then $X_n + Y_n \xrightarrow{a.s.} X + Y$.
- (b) If $X_n \xrightarrow{r} X$ and $Y_n \xrightarrow{r} Y$ then $X_n + Y_n \xrightarrow{r} X + Y$.
- (c) If $X_n \xrightarrow{p} X$ and $Y_n \xrightarrow{p} Y$ then $X_n + Y_n \xrightarrow{p} X + Y$.
- (d) It is not in general true that $X_n + Y_n \xrightarrow{d} X + Y$ whenever $X_n \xrightarrow{d} X$ and $Y_n \xrightarrow{d} Y$.

Proposition 24. Almost sure convergence does not imply convergence in L_2 , and convergence in L_2 does not imply almost sure convergence. That is, find random variables that converge in L_2 but not almost surely. Then, find random variables that converge almost surely but not in L_2 .

Proof. (a) **Almost sure convergence does not imply ℓ_2 convergence:** Let $Z_n := n\mathbf{1}_{[0,1/n]}$. Then Z_1, Z_2, \dots converges almost surely to 0 since $\lim_{n \rightarrow \infty} Z_n = 0$ for all $\omega \in (0, 1]$, but it does not converge in L_2 since $\mathbb{E}(Z_n - 0)^2 = \mathbb{E}Z_n^2 = n^2\mathbb{E}\mathbf{1}_{[0,1/n]} = n \rightarrow \infty$ as $n \rightarrow \infty$.

(b) **ℓ_2 convergence does not imply almost sure convergence:** Define an independent sequence $\{X_n\}$ by

$$X_n = \begin{cases} 1 & \text{with probability } n^{-1} \\ 0 & \text{with probability } 1 - n^{-1} \end{cases}$$

and let $B_m(\epsilon) = \cup_{n \geq m} \{|X_n - X| > \epsilon\}$. $X_n \xrightarrow{2} 0$ because

$$\lim_{n \rightarrow \infty} \|X_n - 0\|_2 = 0$$

However, if $0 < \epsilon < 1$,

$$\Pr(B_m(\epsilon)) = 1 - \lim_{r \rightarrow \infty} \Pr(X_n = 0 \text{ for all } n \text{ such that } m \leq n \leq r)$$

$$= 1 - \left(1 - \frac{1}{m}\right) \left(1 - \frac{1}{m+1}\right) \cdots \text{ (by independence)}$$

$$= 1 - \lim_{M \rightarrow \infty} \left(\frac{m-1}{m} \cdot \frac{m}{m+1} \cdot \frac{m+1}{m+2} \cdots \frac{M}{M+1} \right)$$

$$= 1 - \lim_{M \rightarrow \infty} \frac{m-1}{M+1} = 1$$

and so $\{X_n\}$ does not converge almost surely.

□

Theorem 25. Borel-Cantelli lemmas (Grimmett and Stirzaker Theorem 7.3.10.) Let $\{A_n\}$ be an infinite sequence of events from some probability space $(\Omega, \mathcal{F}, \mathbb{P})$. Let $A = \bigcap_n \bigcup_{m=n}^{\infty} A_m = \limsup_{n \rightarrow \infty} A_n = \{A_n \text{ i.o.}\}$ be the event that infinitely many of the A_n occur. Then:

- (a) $\Pr(A) = 0$ if $\sum_n \Pr(A_n) < \infty$
- (b) $\Pr(A) = 1$ if $\sum_n \Pr(A_n) = \infty$ and A_1, A_2, \dots are independent events.

Proof. (a) We have that $A \subseteq \bigcup_{m=n}^{\infty} A_m$ for all n , so

$$\Pr(A) \leq \sum_{m=n}^{\infty} \Pr(A_m) \rightarrow 0 \text{ as } n \rightarrow \infty$$

whenever $\sum_n \Pr(A_n) < \infty$.

- (b) One can confirm that

$$A^c = \bigcup_n \bigcap_{m=n}^{\infty} A_m^c$$

But

$$\begin{aligned} \Pr\left(\bigcap_{m=n}^{\infty} A_m^c\right) &= \lim_{r \rightarrow \infty} \Pr\left(\bigcap_{m=n}^r A_m^c\right) = \prod_{m=n}^{\infty} [1 - \Pr(A_m)] \text{ (by independence)} \leq \prod_{m=n}^{\infty} \exp(-\Pr(A_m)) \\ &= \exp\left(-\sum_{m=n}^{\infty} \Pr(A_m)\right) = 0 \end{aligned}$$

whenever $\sum_n \Pr(A_n) = \infty$, where the fourth step follows since $1 - x \leq e^{-x}$ if $x \geq 0$. Thus

$$\Pr(A^c) = \lim_{n \rightarrow \infty} \Pr\left(\bigcap_{m=n}^{\infty} A_m^c\right) = 0$$

so $\Pr(A) = 1$. □

Theorem 26. Kolmogorov's Two-Series Theorem. Let X_1, X_2, \dots be independent random variables with $\mathbb{E}(X_n) = \mu_n$ and $\text{Var}(X_n) = \sigma_n^2$ such that $\sum_{n=1}^{\infty} \mu_n < \infty$ and $\sum_{n=1}^{\infty} \sigma_n^2 < \infty$. Then $\sum_{n=1}^{\infty} X_n$ converges in \mathbb{R} almost surely.

Proof. Available on wikipedia, https://en.wikipedia.org/wiki/Kolmogorov%27s_two-series_theorem. □

1.4.1 Slutsky's Convergence Theorems (8.4.1 of Pesaran, 7.3 of Grimmett and Stirzaker)

Theorem 27. Theorem 6 of Pesaran, Section 8.4.1, p. 173. Let $\{x_t, y_t\}, t = 1, 2, \dots$ be a sequence of pairs of random variables with $y_t \xrightarrow{d} y$ and $|y_t - x_t| \xrightarrow{p} 0$. Then $x_t \xrightarrow{d} y$.

Theorem 28. Theorem 7 in Pesaran, on p.318 (section 7.3) of Grimmett and Stirzaker. (Section 8.4.1, p. 174) If $x_t \xrightarrow{d} x$ and $y_t \xrightarrow{p} c$ where c is a finite constant, then

- (i) $x_t + y_t \xrightarrow{d} x + c$
- (ii) $y_t x_t \xrightarrow{d} cx$
- (iii) $x_t/y_t \xrightarrow{d} x/c$, if $c \neq 0$.

Theorem 29. on p.318 (section 7.3) of Grimmett and Stirzaker. Suppose that $X_n \xrightarrow{d} 0$ and $Y_n \xrightarrow{p} Y$, and let $g : \mathbb{R}^2 \rightarrow \mathbb{R}$ be such that $g(x, y)$ is a continuous function of y for all x , and $g(x, y)$ is continuous at $x = 0$ for all y . Then $g(X_n, Y_n) \xrightarrow{p} g(0, Y)$.

Theorem 30 (Continuous Mapping Theorem (Theorem 9 of Pesaran, Section 8.4.1, p. 176: convergence properties of transformed sequences.)). Suppose $\{\mathbf{x}_j\}, \{\mathbf{y}_j\}, \mathbf{x}$, and \mathbf{y} are $k \times 1$ vectors of random variables on a probability space, and let $\mathbf{g}(\cdot)$ be a continuous vector-valued function. (Alternatively, suppose \mathbf{g} has the set of discontinuity points D_g such that $\Pr(X \in D_g) = 0$.) Then

- (i) $\mathbf{x}_j \xrightarrow{a.s.} x \implies \mathbf{g}(\mathbf{x}_j) \xrightarrow{a.s.} \mathbf{g}(x)$
- (ii) $\mathbf{x}_j \xrightarrow{p} x \implies \mathbf{g}(\mathbf{x}_j) \xrightarrow{p} \mathbf{g}(x)$
- (iii) $\mathbf{x}_j \xrightarrow{d} x \implies \mathbf{g}(\mathbf{x}_j) \xrightarrow{d} \mathbf{g}(x)$
- (iv) $\mathbf{x}_j - \mathbf{y}_j \xrightarrow{p} \mathbf{0}$ and $\mathbf{y}_j \xrightarrow{d} \mathbf{y} \implies \mathbf{g}(\mathbf{x}_j) - \mathbf{g}(\mathbf{y}_j) \xrightarrow{d} \mathbf{0}(x)$

where $x = (c_1, \dots, c_k) \in \mathbb{R}^k$.

Proof (part (b), continuous case, one-dimensional codomain). Let $\mathbf{x}_j = (M_{j,1}, \dots, M_{j,k})$. We have that

$$\begin{aligned} \forall \epsilon_j > 0, \lim_{n \rightarrow \infty} \Pr(\{\omega \in \Omega : |M_{j,n}(\omega) - c_j| > \epsilon_j\}) &= 0, \quad \forall j \in \{1, \dots, k\}. \\ \iff \forall \epsilon_j > 0, \lim_{n \rightarrow \infty} \Pr(\{\omega \in \Omega : |M_{j,n}(\omega) - c_j| < \epsilon_j\}) &= 1, \quad \forall j \in \{1, \dots, k\}. \end{aligned} \quad (2)$$

Because g is continuous, we have that for every $\epsilon^* > 0$ there exists a $\delta^* > 0$ such that

$$0 < \|(M_{1,n}(\omega), \dots, M_{j,n}(\omega))\|_2 < \delta^* \implies |g(M_{1,n}, \dots, M_{j,n}) - g(c_1, \dots, c_j)| < \epsilon^*. \quad (3)$$

Note that since in \mathbb{R} the L_2 and L_1 norms are equivalent,

$$\begin{aligned}
|M_{j,n}(\omega) - c_j| < \epsilon_j &\iff \|M_{j,n}(\omega) - c_j\|_2 < \epsilon_j \implies \sum_{j=1}^k \|M_{j,n}(\omega) - c_j\|_2 < \sum_{j=1}^k \epsilon_j \\
&\implies \|(M_{1,n}(\omega), \dots, M_{j,n}(\omega))\|_2 < \sum_{j=1}^k \epsilon_j
\end{aligned}$$

where the last step follows by the Triangle Inequality. Therefore letting $\delta^* = \sum_{j=1}^k \epsilon_j$, we have

$$\begin{aligned}
\Pr(\{\omega \in \Omega : |M_{j,n}(\omega) - c_j| < \epsilon_j\}) &\leq \Pr(0 < \|(M_{1,n}(\omega), \dots, M_{j,n}(\omega))\|_2 < \delta^*) \\
&\leq \Pr(\{\omega \in \Omega : |g(M_{1,n}(\omega), \dots, M_{j,n}(\omega)) - g(c_1, \dots, c_j)| < \epsilon^*\})
\end{aligned}$$

where the last step follows from (3). So

$$\Pr(\{\omega \in \Omega : |M_{j,n}(\omega) - c_j| < \epsilon_j\}) \leq \Pr(\{\omega \in \Omega : |g(M_{1,n}(\omega), \dots, M_{j,n}(\omega)) - g(c_1, \dots, c_j)| < \epsilon^*\}). \quad (4)$$

Taking limits of (4) and substituting in (2), we have

$$\begin{aligned}
&\forall \epsilon^* > 0, \lim_{n \rightarrow \infty} \Pr(\{\omega \in \Omega : |g(M_{1,n}(\omega), \dots, M_{j,n}(\omega)) - g(c_1, \dots, c_j)| < \epsilon^*\}) \geq 1 \\
&\iff \forall \epsilon^* > 0, \lim_{n \rightarrow \infty} \Pr(\{\omega \in \Omega : |g(M_{1,n}(\omega), \dots, M_{j,n}(\omega)) - g(c_1, \dots, c_j)| > \epsilon^*\}) = 0 \\
&\iff g(M_{1,n}, \dots, M_{j,n}) \xrightarrow{p} g(c_1, \dots, c_j).
\end{aligned}$$

For remaining parts, see [Serfling \[1980\]](#) or [Rao \[1973\]](#). □

See also:

Theorem 31. (Theorem 7.2.18, Grimmett and Stirzaker.) If $X_n \xrightarrow{d} X$ and $g : \mathbb{R} \rightarrow \mathbb{R}$ is continuous, then $g(X_n) \xrightarrow{d} g(X)$.

1.5 Stochastic orders $\mathcal{O}_p(\cdot)$ and $o_p(\cdot)$ (Pesaran 8.5)

Definition 1.13 (Pesaran 8.5 Definition 6.). Let $\{a_t\}$ be a sequence of positive numbers and $\{x_t\}$ be a sequence of random variables. Then

- (i) $x_t = \mathcal{O}_p(a_t)$, or x_t/a_t is bounded in probability, if for every $\epsilon > 0$ there exist real numbers M_ϵ and N_ϵ such that

$$\Pr \left(\frac{|x_t|}{a_t} > M_\epsilon \right) < \epsilon, \quad \text{for } t > N_\epsilon$$

- (ii) $x_t = o_p(a_t)$ if

$$\frac{x_t}{a_t} \xrightarrow{p} 0$$

Definition 1.14 (Ross ISE 620 Definition). We say that $f(x)$ is $o(h)$ if $\lim_{h \rightarrow 0} f(h)/h = 0$.

1.6 Laws of Large Numbers and Central Limit Theorems (Pesaran 8.6; Grimmett and Stirzaker 7.4, 7.5)

Theorem 32. Weak Law of Large Numbers (Khinchine) (Pesaran 8.6 Theorem 10, Grimmett and Stirzaker Theorem 7.4.7, 541A notes Theorem 2.10). Suppose that $\{X_k\}$ is a sequence of (i) independent (ii) identically distributed random variables with (iii) constant means, i.e., $\mathbb{E}(X_k) = \mu < \infty$. Then

$$\bar{X}_k = \frac{1}{n} \sum_{k=1}^n X_k \xrightarrow{p} \mu$$

Theorem 33. Weak Law of Large Numbers (Chebyshev) (Pesaran Section 8.6, p. 178, Theorem 11.) Let $\{X_k\}$ be a sequence of random variables. If (i) $\mathbb{E}(X_k) = \mu_k$, (ii) $\text{Var}(X_k) = \sigma_k^2$, and (iii) $\text{Cov}(X_k, X_j) = 0$, $k \neq j$, and (iv)

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n \sigma_k^2 < \infty$$

then we have $\bar{X}_n - \bar{\mu}_n \xrightarrow{p} 0$, where $\bar{\mu}_n = n^{-1} \sum_{k=1}^n \mu_k$.

Theorem 34. Strong Law of Large Numbers (Grimmett and Stirzaker Theorem 7.4.3). Let $\{X_k\}$ be a sequence of (i) independent (ii) identically distributed random variables with (iii) $\mathbb{E}(X_k) = \mu$ and (iv) $\mathbb{E}(X_k^2) < \infty$. Then

$$\frac{1}{n} \sum_{k=1}^n X_k \rightarrow \mu \text{ almost surely and in mean square.}$$

Theorem 35 (Strong Law of Large Numbers (Grimmett and Stirzaker Theorem 7.5.1, 541A notes Theorem 2.11)). Let $\{X_k\}$ be a sequence of (i) independent (ii) identically distributed random variables. Then if and only if (iii) $\mathbb{E}|X_k| < \infty$,

$$\frac{1}{n} \sum_{k=1}^n X_k \xrightarrow{a.s.} \mu$$

Theorem 36. Strong Law of Large Numbers 1 (Kolmogorov) (Pesaran 8.8 Theorem 12). Let $\{X_k\}$ be a sequence of (i) independent random variables with (ii) $\mathbb{E}(X_k) = \mu_k < \infty$ and (iii) $\text{Var}(X_k) = \sigma_k^2$ such that (iii)

$$\sum_{k=1}^{\infty} \frac{\sigma_k^2}{k^2} < \infty$$

Then $\bar{X}_n - \bar{\mu}_n \xrightarrow{wp1} 0$. If the independence assumption (i) is replaced by a lack of correlation (i.e. $\text{Cov}(X_k, X_j) = 0, k \neq j$), the convergence of $\bar{X}_n - \bar{\mu}_n$ with probability one requires the stronger condition

$$\sum_{k=1}^{\infty} \frac{\sigma_k^2 (\log k)^2}{k^2} < \infty$$

Theorem 37. Strong Law of Large Numbers 2 (Pesaran 8.8 Theorem 13) Suppose that X_1, X_2, \dots are (i) independent random variables, and that (ii) $\mathbb{E}(X_k) = 0$, (iii) $\mathbb{E}(X_k^4) \leq M \forall k$ where M is an arbitrary positive constant. Then

$$\bar{X}_n = \frac{1}{n} \sum_{k=1}^n X_k \xrightarrow{a.s.} 0$$

Theorem 38. Central Limit Theorem (Grimmett and Stirzaker theorem 5.10.4.) Let X_1, X_2, \dots be a sequence of independent identically distributed random variables with finite mean μ and finite non-zero variance σ^2 , and let $S_n = \sum_{i=1}^n X_i$. Then

$$\frac{S_n - n\mu}{\sqrt{n\sigma^2}} \xrightarrow{d} \mathcal{N}(0, 1)$$

Theorem 39. (Berry-Esseen Central Limit Theorem.) There exists $c > 0$ such that the following holds. Let X_1, X_2, \dots be i.i.d. real-valued random variables with mean zero, variance 1, and $\mathbb{E}(|X_1|^3) < \infty$. Let Z be a standard Gaussian random variable. Then for any $n \geq 1$,

$$\sup_{t \in \mathbb{R}} \left| \Pr \left(\frac{X_1 + \dots + X_n}{\sqrt{n}} \leq t \right) - \Pr(Z \leq t) \right| \leq c \cdot \frac{\mathbb{E}(|X_1|^3)}{\sqrt{n}}$$

Remark. You can look up what the c is; Heilman doesn't think it's any bigger than around 10.

Theorem 40. (Central Limit Theorem in \mathbb{R}^d , Heilman notes Theorem 2.33.) Let $X^{(1)}, X^{(2)}, \dots$ be a sequence of independent identically distributed \mathbb{R}^d -valued random variables. (Notation: we write $X^{(1)} = (X_1^{(1)}, \dots, X_d^{(1)})$.) Assume $\mathbb{E}(X^{(n)}) = \boldsymbol{\mu}$ for all $n \geq 1$ and for any $1 \leq i < j \leq d$, all of the covariances

$$a_{ij} = \mathbb{E}[(X_i^{(1)} - \mathbb{E}(X_i^{(1)}))(X_j^{(1)} - \mathbb{E}(X_j^{(1)}))]$$

are finite. Let $\mathbf{S}_n = \sum_{i=1}^n X^{(i)}$. Then as $n \rightarrow \infty$,

$$\frac{\mathbf{S}_n - n\boldsymbol{\mu}}{\sqrt{n\sigma^2}} \xrightarrow{d} \mathcal{N}(\boldsymbol{\mu}, [a_{ij}])$$

Theorem 41. (Grimmett and Stirzaker theorem 5.10.5.) Let X_1, X_2, \dots be independent random variables satisfying $\mathbb{E}(X_j) = 0$, $\text{Var}(X_j) = \sigma_j^2$, $\mathbb{E}|X_j^3| < \infty$ such that

$$\lim_{n \rightarrow \infty} \frac{1}{\sigma(n)^3} \sum_{j=1}^n \mathbb{E}|X_j^3| = 0$$

where $\sigma(n)^2 = \text{Var}(\sum_{j=1}^n X_j) = \sum_{j=1}^n \sigma_j^2$. Then

$$\frac{1}{\sigma(n)} \sum_{j=1}^n X_j \xrightarrow{d} \mathcal{N}(0, 1)$$

Proof. See Loeve [1977, p. 287] and Grimmett and Stirzaker Problem 5.12.40. □

Lemma 42. Lindeberg's Condition: Let $\{X_k\}$ be a sequence of independent (not necessarily identically distributed) random variables with expectations μ_k and finite variances σ_k^2 . Let $s_n^2 = \sum_{k=1}^n \sigma_k^2$. If such a sequence of independent random variables X_k satisfies the condition

$$\lim_{n \rightarrow \infty} \frac{1}{s_n^2} \sum_{k=1}^n \mathbb{E}[(X_k - \mu_k)^2 \cdot \mathbf{1}_{\{|X_k - \mu_k| > \epsilon s_n\}}] = 0$$

for all $\epsilon > 0$ then the central limit theorem holds; that is, the random variables

$$Z_n = \frac{1}{s_n} \sum_{k=1}^n (X_k - \mu_k)$$

converge in distribution to $\mathcal{N}(0, 1)$ as $n \rightarrow \infty$.

1.7 The case of dependent and heterogeneously distributed observations (Pesaran 8.8)

Theorem 43. Central limit theorem for martingale difference sequences (Pesaran 8.8 Theorem 28). Let $\{x_t\}$ be a martingale difference sequence with respect to the information set Ω_t . Let $\bar{\sigma}_T^2 = \text{Var}(\sqrt{T}\bar{x}_T) = T^{-1} \sum_{t=1}^T \sigma_t^2$. If $\mathbb{E}(|x_t|^r) < K < \infty$ for any $r > 2$ and for all t , and

$$\frac{1}{T} \sum_{t=1}^T x_t^2 - \bar{\sigma}_T^2 \xrightarrow{p} 0$$

then $\sqrt{T}\bar{x}_T / \bar{\sigma}_T \xrightarrow{d} \mathcal{N}(0, 1)$.

1.8 Worked Examples from Math 505A Midterm 2

- (1) (a) **Fall 2010 Problem 1.** Let X_k , $k \geq 1$, be i.i.d. random variables with mean 1 and variance 1. Show that the limit

$$\lim_{n \rightarrow \infty} \frac{\sum_{k=1}^n X_k}{\sum_{k=1}^n X_k^2}$$

exists in an appropriate sense, and identify the limit.

- (b) **Not included on midterm or final.** Let $(X_j)_{j \geq 1}$ be i.i.d. uniform on $(-1, 1)$. Let

$$Y_n = \frac{\sum_{j=1}^n X_j}{\sum_{j=1}^n X_j^2 + \sum_{j=1}^n X_j^3}$$

Prove that $\lim_{n \rightarrow \infty} \sqrt{n}Y_n$ exists in an appropriate sense, and identify the limit.

Solution.

- (a)

$$\lim_{n \rightarrow \infty} \frac{\sum_{k=1}^n X_k}{\sum_{k=1}^n X_k^2} = \lim_{n \rightarrow \infty} \frac{n^{-1} \sum_{k=1}^n X_k}{n^{-1} \sum_{k=1}^n X_k^2}$$

Since X_1, X_2, \dots are i.i.d., $E(X_1^2) = \text{Var}(X_1) + (\mathbb{E}(X_1))^2 = 2 < \infty$, we have

$$n^{-1} \sum_{k=1}^n X_k \xrightarrow{a.s.} \mathbb{E}(X_1) = 1 \text{ as } n \rightarrow \infty$$

by Theorem 34 (Strong Law of Large Numbers). Also, X_1^2, X_2^2, \dots are clearly identically distributed, and are independent by Theorem 4.2.3 (“If X and Y are independent, then so are $g(X)$ and $g(Y)$.”). It is clear also that $\mathbb{E}(|X_1^2|) = \mathbb{E}(X_1^2) = \text{Var}(X_1) + \mathbb{E}(X_1)^2 = 1 + 1 = 2 < \infty$. Therefore by Theorem 35 (Strong Law of Large Numbers),

$$n^{-1} \sum_{k=1}^n X_k^2 \xrightarrow{a.s.} \mathbb{E}(X_1^2) = 2 \text{ as } n \rightarrow \infty$$

(From here I had two different ways of finishing the problem.)

- Because we have almost sure convergence in the numerator and denominator, by the Continuous Mapping Theorem (Theorem 30),

$$\lim_{n \rightarrow \infty} \frac{n^{-1} \sum_{k=1}^n X_k}{n^{-1} \sum_{k=1}^n X_k^2} = \frac{\lim_{n \rightarrow \infty} n^{-1} \sum_{k=1}^n X_k}{\lim_{n \rightarrow \infty} n^{-1} \sum_{k=1}^n X_k^2} \xrightarrow{a.s.} \boxed{\frac{1}{2}}$$

- Then, using one of Slutsky’s convergence theorems (Theorem 28: “If $x_t \xrightarrow{d} x$ and $y_t \xrightarrow{p} c$ where c is a finite constant, then $x_t/y_t \xrightarrow{d} x/c$, if $c \neq 0$.”), we have

$$\frac{n^{-1} \sum_{k=1}^n X_k}{n^{-1} \sum_{k=1}^n X_k^2} \xrightarrow{d} \frac{\mathbb{E}(X_1)}{\mathbb{E}(X_1^2)} = \frac{\mathbb{E}(X_1)}{\text{Var}(X_1) + \mathbb{E}(X_1)^2} = \frac{1}{1+1} = \frac{1}{2}$$

But then, by Theorem 16 (Theorem 7.2.4(a) in Grimmett and Stirzaker: “If $X_n \xrightarrow{d} c$ where c is constant, then $X_n \xrightarrow{p} c$.”), we have $\frac{n^{-1} \sum_{k=1}^n X_k}{n^{-1} \sum_{k=1}^n X_k^2} \xrightarrow{p} 1/2$.

(b) (Not included on midterm or final.)

$$Y_n = \frac{\sum_{j=1}^n X_j}{\sum_{j=1}^n X_j^2 + \sum_{j=1}^n X_j^3} = \frac{n^{-1} \sum_{j=1}^n X_j}{n^{-1} \sum_{j=1}^n X_j^2 + n^{-1} \sum_{j=1}^n X_j^3}$$

Note that $\mathbb{E}(X_1) = 0, \mathbb{E}(X_1^2) = \text{Var}(X_1) + \mathbb{E}(X_1)^2 = (1 - (-1))^2/12 + 0^2 = 1/3, \mathbb{E}(X_1^3) = (1/2) \int_{-1}^1 x^3 dx = 0$. (We derived the formulae for the first three moments of a uniform distribution on Homework 4 problem 2(2).)

$$\Rightarrow \sqrt{n}Y_n = \frac{\sqrt{1/3}(\sum_{j=1}^n X_j - n\mathbb{E}(X_1))/\sqrt{n \cdot 1/3}}{n^{-1} \sum_{j=1}^n X_j^2 + n^{-1} \sum_{j=1}^n X_j^3}$$

By the Central Limit Theorem (Theorem 38),

$$\frac{\sum_{j=1}^n X_j - n\mathbb{E}(X_1)}{\sqrt{n \cdot 1/3}} \xrightarrow{d} \mathcal{N}(0, 1)$$

By the Law of Large Numbers (Theorem 35), since $\mathbb{E}(|X_1^2|) = \mathbb{E}(X_1^2) = 1/3 < \infty$,

$$\frac{1}{n} \sum_{j=1}^n X_j^2 \xrightarrow{a.s.} \mathbb{E}(X_1^2) = 1/3$$

By the Law of Large Numbers (Theorem 35), since $\mathbb{E}(|X_1^3|) = (1/2) \int_{-1}^1 |x^3| dx = \int_0^1 x^3 dx = 1/4 < \infty$,

$$\frac{1}{n} \sum_{j=1}^n X_j^3 \xrightarrow{a.s.} \mathbb{E}(X_1^3) = 0$$

In the denominator, since we have almost sure convergence, the regular rules of calculus/real analysis apply. That is, using the above results,

$$n^{-1} \sum_{j=1}^n X_j^2 + n^{-1} \sum_{j=1}^n X_j^3 \xrightarrow{a.s.} 1/3$$

Therefore

$$\sqrt{n}Y_n = \frac{\sqrt{1/3}(\sum_{j=1}^n X_j - n\mathbb{E}(X_1))/\sqrt{n \cdot 1/3}}{n^{-1} \sum_{j=1}^n X_j^2 + n^{-1} \sum_{j=1}^n X_j^3} \xrightarrow{d} \frac{\sqrt{1/3}}{1/3} \mathcal{N}(0, 1) = \boxed{\mathcal{N}(0, 3)}$$

(2) **Fall 2010 Problem 2.** Fix $p \in (0, 1)$ and consider independent Poisson random variables $X_k, k \geq 1$ with

$$\mathbb{E}X_k = \frac{p^k}{k}$$

Verify that the sum $\sum_{k=1}^{\infty} kX_k$ converges with probability one and determine the distribution of the random variable $Y = \sum_{k=1}^{\infty} kX_k$.

Solution. Melike's solution (use for midterm): We have $\mathbb{E}[kX_k] = p^k$ and $\sum_{k=1}^{\infty} p^k = p/(1-p) < \infty$, and $\text{Var}(kX_k) = kp^k$ and

$$\sum_{k=1}^{\infty} kp^k = p \sum_{k=1}^{\infty} kp^{k-1} = p \frac{d}{dp} \sum_{k=1}^{\infty} p^k = p \frac{d}{dp} \frac{p}{1-p} = p \cdot \frac{(1-p) - p(-1)}{(1-p)^2} = \frac{p}{(1-p)^2} < \infty$$

Since the sequence $\{Y_k\}_{k \geq 1}$ is independent, by Kolmogorov's Two Series Theorem (Theorem 26: "Let X_1, X_2, \dots be independent random variables with $\mathbb{E}(X_n) = \mu_n$ and $\text{Var}(X_n) = \sigma_n^2$ such that $\sum_{n=1}^{\infty} \mu_n < \infty$ and $\sum_{n=1}^{\infty} \sigma_n^2 < \infty$. Then $\sum_{n=1}^{\infty} X_n$ converges in \mathbb{R} almost surely."), we conclude that $\sum_{k=1}^{\infty} kX_k$ converges almost surely.

To find the distribution of Y , let X be a Poisson random variable and consider its probability generating function:

$$G_X(s) = \mathbb{E}(s^X) = \sum_{k=0}^{\infty} s^k e^{-\lambda} \frac{\lambda^k}{k!} = e^{-\lambda} \sum_{k=0}^{\infty} \frac{(\lambda s)^k}{k!} = e^{-\lambda} e^{\lambda s} = e^{\lambda(s-1)}$$

So $\mathbb{E}(s^{X_k}) = \exp\left(\frac{p^k}{k}(s-1)\right)$ and $\mathbb{E}(s^{kX_k}) = \mathbb{E}[(s^k)^{X_k}] = \exp\left(\frac{p^k}{k}(s^k-1)\right)$. Then define $Y_n = \sum_{k=1}^n kX_k$ and consider

$$\begin{aligned} G_{Y_n}(s) &= \mathbb{E}(s^{Y_n}) = \mathbb{E}\left(\prod_{k=1}^n s^{kX_k}\right) = \prod_{k=1}^n \mathbb{E}(s^{kX_k}) = \prod_{k=1}^n \exp\left(\frac{p^k}{k}(s^k-1)\right) = \exp\left(\sum_{k=1}^n \frac{p^k}{k}(s^k-1)\right) \\ &= \exp\left(\sum_{k=1}^n \frac{(ps)^k}{k} - \sum_{k=1}^n \frac{p^k}{k}\right) \end{aligned}$$

Now, by taking limits as $n \rightarrow \infty$ (since we are allowed to take limit inside of expectation here), we get

$$\begin{aligned} G_Y(s) &= \mathbb{E}(s^Y) = \exp\left(\sum_{k=1}^{\infty} \frac{(ps)^k}{k} - \sum_{k=1}^{\infty} \frac{p^k}{k}\right) = \exp\left(\int \sum_{k=1}^{\infty} (ps)^{k-1} dp - \int \sum_{k=1}^{\infty} p^{k-1} dp\right) \\ &= \exp\left(\int \frac{1}{1-ps} dp - \int \frac{1}{1-p} dp\right) = \exp(-\log(1-ps) + \log(1-p)), \quad -1 \leq ps < 1 \text{ and } -1 \leq p < 1 \\ &= \frac{1-p}{1-ps}, \quad -1 \leq ps < 1 \end{aligned}$$

Since we know $\Pr(X = k) = \frac{G_X^{(k)}(0)}{k!}$, we have

$$G_Y(s) = \frac{1-p}{1-sp}, \quad G'(s) = \frac{p(1-p)}{(1-sp)^2}, \quad G''(s) = \frac{2p^2(1-p)}{(1-sp)^3}, \quad G^{(3)}(s) = \frac{3 \cdot 2p^3(1-p)}{(1-sp)^3}, \dots$$

$$G^{(k)}(s) = \frac{k!p^k(1-p)}{(1-sp)^k} \text{ for } k = 0, 1, 2, \dots$$

So we have

$$\Pr(Y = k) = (1-p)p^k, \quad k = 0, 1, 2, \dots$$

$$= \Pr(G_1(1-p) = k+1) = \Pr(G_1(1-p) - 1 = k)$$

which means $Y \sim G_1(1-p) - 1$.

(3) Spring 2017 Problem 3.

- (a) Consider the sequence $\{X_k, k \geq 1\}$ of random variables such that X_1 is uniform on $(0, 1)$ and, given X_k , the distribution of X_{k+1} is uniform on $(0, CX_k)$, where $\sqrt{3} < C < 2$.
- (i) For $n \geq 1$, compute the conditional expectation $\mathbb{E}(X_{n+1}^r | X_n)$.
 - (ii) For $n \geq 1$, compute $\mathbb{E}(X_n^r)$.
 - (iii) Show that $\lim_{n \rightarrow \infty} X_n = 0$ in ℓ_1 and with probability one, but not in ℓ_2 .
 - (iv) Investigate the same questions for all other values of $C > 0$.
- (b) Let $a > 0$, let $X_n, n \geq 1$ be i.i.d. random variables that are uniform on $(0, a)$, and let $Y_n = \prod_{k=1}^n X_k$. Determine, with a proof, all values of a for which $\lim_{n \rightarrow \infty} Y_n = 0$ with probability one.

Solution.

- (a) (i) We have that $X_{n+1} | X_n \sim U(0, CX_n)$. Therefore

$$\begin{aligned} \mathbb{E}(X_{n+1}^r | X_n) &= \frac{1}{CX_n} \int_0^{CX_n} x^r dx = \frac{1}{CX_n} \cdot \frac{x^{r+1}}{r+1} \Big|_0^{CX_n} = \frac{C^r X_n^r}{r+1} \\ \implies \mathbb{E}(X_{n+1}^r) &= \mathbb{E}[\mathbb{E}(X_{n+1}^r | X_n)] = \frac{C^r}{r+1} \cdot \mathbb{E}(X_n^r) \\ \implies \boxed{\mathbb{E}(X_{n+1}^r | X_n) &= \frac{C^r}{r+1} X_n^r} \end{aligned}$$

- (ii) Note that $\mathbb{E}(X_1^r) = \int_0^1 x^r dx = 1/(r+1)$. Therefore

$$\mathbb{E}(X_{n+1}^r) = \frac{C^r}{r+1} \cdot \mathbb{E}(X_n^r) = \left(\frac{C^r}{r+1} \right)^n \cdot \mathbb{E}(X_1^r) = \boxed{\left(\frac{C^r}{r+1} \right)^n \cdot \frac{1}{r+1}}$$

- (iii) We would like to show that $X_n \xrightarrow{w.p.1} 0$ and that $X_n \xrightarrow{1} 0$, but that the same result does not follow for the ℓ_2 norm.

- **Convergence with probability one:** We seek to show that $\Pr(\{\omega \in \Omega : \lim_{n \rightarrow \infty} X_n(\omega) = 0\}) = 1$. By Markov's Inequality (Lemma 3), we have

$$\Pr(|X_n| \geq a) \leq \frac{\mathbb{E}(X_n)}{a} \quad \forall a > 0$$

$$\iff \Pr(|X_n| \geq a) \leq \left(\frac{C^1}{1+1} \right)^{n-1} \cdot \frac{1}{1+1} \cdot \frac{1}{a} = \left(\frac{C}{2} \right)^{n-1} \cdot \frac{1}{2a} \quad \forall a > 0$$

Since $\sqrt{3} < C < 2$, $\sqrt{3}/2 < C/2 < 1$. Since $X_n \in [0, CX_{n-1}]$, $X_n \geq 0$, so $|X_n| = X_n$. Therefore we have

$$\Pr(\lim_{n \rightarrow \infty} |X_n| \geq a) = \Pr(\lim_{n \rightarrow \infty} X_n \geq a) \leq \lim_{n \rightarrow \infty} \left(\frac{C}{2} \right)^{n-1} \cdot \frac{1}{2a} = 0 \quad \forall a > 0$$

Since $|X_n| \geq 0$, this implies that $\Pr(\lim_{n \rightarrow \infty} X_n = 0) = \Pr(\{\omega \in \Omega : \lim_{n \rightarrow \infty} X_n(\omega) = 0\}) = 1$, so by the Borel-Cantelli Lemma (Theorem 25), X_n converges to 0 with probability 1.

- **Convergence in ℓ_1 norm:** We seek to show that $\lim_{n \rightarrow \infty} \mathbb{E}(|X_n|) = 0$. Since $X_n \in [0, CX_{n-1}]$, $X_n \geq 0$, so $|X_n| = X_n$. Therefore

$$\lim_{n \rightarrow \infty} \mathbb{E}(|X_n|) = \lim_{n \rightarrow \infty} \mathbb{E}(X_n) = \lim_{n \rightarrow \infty} \left(\frac{C}{2}\right)^{n-1} \cdot \frac{1}{2}$$

Since $\sqrt{3} < C < 2$, $\sqrt{3}/2 < C/2 < 1$, so $C/2 < 1$. Therefore we have

$$\lim_{n \rightarrow \infty} \mathbb{E}(|X_n|) = \lim_{n \rightarrow \infty} \left(\frac{C}{2}\right)^{n-1} \cdot \frac{1}{2} = 0$$

so X_n converges to 0 in 1st mean.

- **Convergence in ℓ_2 norm:** We seek to show that $\lim_{n \rightarrow \infty} \mathbb{E}(|X_n|^2) \neq 0$. We have

$$\lim_{n \rightarrow \infty} \mathbb{E}(|X_n|^2) = \lim_{n \rightarrow \infty} \mathbb{E}(X_n^2) = \lim_{n \rightarrow \infty} \left(\frac{C^2}{3}\right)^{n-1} \cdot \frac{1}{3}$$

Since $\sqrt{3} < C < 2$, $3/3 < C^2/3 < 4/3$, so $C^2/3 > 1$. Therefore we have

$$\lim_{n \rightarrow \infty} \mathbb{E}(|X_n|^2) = \lim_{n \rightarrow \infty} \left(\frac{C^2}{3}\right)^{n-1} \cdot \frac{1}{3} = \infty \neq 0$$

so X_n does not converge to 0 in 2nd mean.

- (iv) From the above, it is clear that for convergence with probability one or in 1st mean we require $0 < C/2 < 1$ and for convergence in second mean we require $0 < C^2/3 < 1$. For $0 < C < \sqrt{3}$, we see that X_n would converge to zero in 2nd mean since this would imply that $0 < C^2/3 < 1$. It would also still converge to 0 in 1st mean (and with probability 1) since we would have $(0 < C/2 < \sqrt{3}/2 < 1)$. For $C = \sqrt{3}$, X_n would still converge to 0 with probability one and in 1st mean for the same reasons. However, it would not converge in 2nd mean because we would have

$$\lim_{n \rightarrow \infty} \mathbb{E}(|X_n|^2) = \lim_{n \rightarrow \infty} \left(\frac{\sqrt{3}^2}{3}\right)^{n-1} \cdot \frac{1}{3} = \frac{1}{3} \neq 0$$

For $C \geq 2$, it would diverge in all three cases, since in this case $C/2 \geq 2/2 = 1$ and $C^2/3 \geq 4/3 > 1$.

- (b) **Probably won't be on midterm.** Note that

$$\lim_{n \rightarrow \infty} Y_n = \lim_{n \rightarrow \infty} \prod_{k=1}^n X_k = 0 \iff \log(Y_n) = \log\left(\prod_{k=1}^n X_k\right) = \sum_{k=1}^n \log(X_k) \rightarrow -\infty$$

Note that

$$\begin{aligned} \mathbb{E}[\log(Y_n)] &= \mathbb{E}\left(\sum_{k=1}^n \log(X_k)\right) = \sum_{k=1}^n \mathbb{E}[\log(X_k)] = \sum_{k=1}^n \mathbb{E}[\log(X_1)] = \sum_{k=1}^n \int_0^a (\log(x)/a) dx \\ &= \sum_{k=1}^n \frac{1}{a} [x \log x - x]_0^a = \sum_{k=1}^n \frac{a \log a - a}{a} = \sum_{k=1}^n (\log(a) - 1) = n(\log(a) - 1) \end{aligned}$$

As $n \rightarrow \infty$ we have

$$\mathbb{E}[\log(Y_n)] = \begin{cases} -\infty & a < e \\ 0 & a = e \\ \infty & a > e \end{cases}$$

Since $\mathbb{E}[\log(Y_n)] \rightarrow \infty$ for $a < e$, we have $\lim_{n \rightarrow \infty} Y_n = 0$ for $a < 3$. Therefore

$$\lim_{n \rightarrow \infty} Y_n = \lim_{n \rightarrow \infty} \prod_{k=1}^n X_k = 0 \iff a < e.$$

1.9 Estimators and Central Limit Theorems (DSO 607)

Lyapunov (?) condition: can prove central limit theorem if we check 3rd moment. Lindeberg's Condition (Lemma 42)

References

- G. Grimmett and D. Stirzaker. *Probability and random processes*, volume 80. Oxford university press, 2001. URL http://scholar.google.com/scholar.bib?q=info:xzStZXK20NkJ:scholar.google.com/&output=citation&hl=en&as_sdt=0,5&ct=citation&cd=0.
- J. Kingman and S. Taylor. *Introduction to Measure and Probability*. Cambridge University Press, 1966. URL <https://books.google.com/books?id=JbtEAAAAIAAJ>.
- M. Loeve. *Probability Theory I*. Comprehensive Manuals of Surgical Specialties. Springer, 1977. ISBN 9780387902104. URL https://books.google.com/books?id=_9xWBlvUEuIC.
- M. H. Pesaran. *Time Series and Panel Data Econometrics*. Number 9780198759980 in OUP Catalogue. Oxford University Press, 2015. ISBN ARRAY(0x3bdaaf68). URL <https://ideas.repec.org/b/oxp/obooks/9780198759980.html>.
- C. Rao. *Linear statistical inference and its applications*. Wiley series in probability and mathematical statistics: Probability and mathematical statistics. Wiley, 1973. ISBN 9780471708230. URL <https://books.google.com/books?id=lPhQAAAAMAAJ>.
- R. Serfling. *Approximation theorems of mathematical statistics*. Wiley series in probability and mathematical statistics : Probability and mathematical statistics. Wiley, New York, NY [u.a.], [nachdr.] edition, 1980. ISBN 0471024031. URL http://gso.gbv.de/DB=2.1/CMD?ACT=SRCHA&SRT=YOP&IKT=1016&TRM=ppn+024353353&sourceid=fbw_bibsonomy.