

Aromatic interactions in model systems

Marcey L Waters

A thorough knowledge of noncovalent interactions is crucial to the understanding of biological complexity. One of the less well understood but significant weak interactions in nature is the aromatic interaction. Recent studies have provided new insight into the driving force, stability and selectivity of these interactions. The contribution of solvophobic and electrostatic interactions have been shown to be inextricably linked. Moreover, the influence of electrostatic and solvophobic components on the selectivity of aromatic interactions has been demonstrated.

Addresses

Department of Chemistry, University of North Carolina, Chapel Hill, NC 27599-3290, USA; e-mail: mlwaters@email.unc.edu

Current Opinion in Chemical Biology 2002, 6:736–741

1367-5931/02/\$ — see front matter

© 2002 Elsevier Science Ltd. All rights reserved.

Published online 18 October 2002

Abbreviation

Cha cyclohexylalanine

Introduction

Aromatic interactions are ubiquitous in nature. They are believed to provide stability to duplex DNA [1]; they have been proposed to contribute to the unique properties of thermophilic proteins [2•]; they may play a role in aggregation of amyloid β in Alzheimer's disease [3]; and they are common motifs in biomolecular recognition. Extensive work has focused on these interactions to determine their importance as a recognition element, including statistical analyses of the Protein Data Bank [4,5•,6–8], mutation studies in proteins [9] and protein–nucleic-acid complexes [10,11], and theoretical studies [12••]. Model systems have also proven to be an informative method of investigating the nature and significance of aromatic interactions as molecular recognition elements in biological and non-biological systems. This review presents a general background on aromatic interactions and recent studies of aromatic interactions in solution-phase synthetic and biological model systems that have advanced the understanding of this biologically important noncovalent interaction.

The nature and geometry of aromatic interactions

Aromatic interactions have been proposed to consist of van der Waals, hydrophobic and electrostatic forces [12••]. The relative contribution and magnitude of each of these components is still under investigation. This is complicated by the fact that aromatic groups interact in one of several geometries, depending on the nature of the rings involved. Nonetheless, aromatic interactions are intriguing molecular recognition elements because they are expected to be strong in water because of the hydrophobic component of


the interaction yet, at the same time, the interaction should be selective if the electrostatic component is significant, thus providing the best features of both hydrophobic interactions and hydrogen bonding.

Several geometries are attractive, and have been proposed on the basis of the electrostatic component of the interaction (Figure 1) [13]. The electrostatic component has been proposed to arise from interactions of the quadrupole moments of the aromatic rings. Although benzene has no net dipole, it has an uneven distribution of charge, with greater electron-density on the face of the ring and reduced electron-density on the edge, which gives rise to the quadrupole moment. The edge-face geometry (Figure 1a), which can be considered a CH– π interaction, is found in benzene in the solid state, and is commonly observed between aromatic residues in proteins. The offset stacked orientation (Figure 1b) is also commonly found in proteins and is the geometry of base stacking in DNA. In this geometry, more surface area is buried, and the van der Waals and hydrophobic interactions are increased. This orientation appears to be more common when the electron density on the face of one or both rings is reduced. A third possible geometry is the face-to-face stacked orientation (Figure 1c). This is commonly observed with donor–acceptor pairs and compounds that have opposite quadrupole moments, such that the interaction between the faces of the rings is attractive. The benzene–perfluorobenzene interaction is an excellent example of this type of aromatic interaction, and has been calculated to provide $-15.5 \text{ kJ mol}^{-1}$ in stability [14]. In an elegant study, Cozzi and Siegel [15] have demonstrated the electrostatic contribution to aromatic interactions in the face-to-face stacked conformation. However, questions still exist regarding the importance of electrostatics relative to dispersion and hydrophobic forces in the edge-face and offset-stacked geometries, which are the geometries commonly found in nature.

Solvent effects on aromatic interactions

Several groups have recently published solvent studies on aromatic interactions to address the importance of electrostatic and solvophobic components in aromatic interactions. A solvent study by Cammers-Goodwin [16] and co-workers suggests that water interacts preferentially with the hydrogens of an aromatic ring, influencing the propensity for stacking. Moore and co-workers [17,18••] have designed both cyclic and linear systems made up of *m*-phenylene ethynylene units that spontaneously associate in polar solvents (Figure 2). The driving force for association is believed to be stacking of the aromatic rings. The fact that association occurs preferentially in polar solvents suggests that solvophobic forces influence this system significantly. However, folding is highly dependent on the nature of the substituents on the rings, suggesting an electronic tuning of the interactions

Figure 1




Geometries of aromatic interactions. (a) edge-face; (b) offset stacked; (c) face-to-face stacked.

through minimization of repulsive quadrupole–quadrupole interactions [19]. Intriguingly, these molecules were also found to fold in aromatic solvents, which was unexpected. The same phenomenon has also been observed in *m*-diethynylbenzene macrocycles [20].

Gellman and co-workers [21] have probed the solvent effect on a simple aromatic interaction in the context of a secondary amide model system (Figure 2, R = H; Ar₁ = phenyl). In this system, the *E* to *Z* ratio of amide rotamers is influenced by the Ar₂-group and the solvent. When Ar₂ = H, the *Z*-isomer is favored as a result of reduced steric interactions between side chains relative to the *E*-isomer and the *E*:*Z* ratio is independent of solvent (water or chloroform). By contrast, when Ar₂ = naphthyl or biphenyl, the amount of *E*-isomer is fourfold higher in water than in chloroform. In addition, the amount of *E*-isomer increases with increasing temperature. Both the solvent dependence of the rotamer populations and the temperature dependence of the equilibrium constant are consistent with a hydrophobically driven association between the aromatics groups.

Iverson and co-workers [22**] have performed an elegant solvent study to address the nature of aromatic interactions. Three different aromatic pairs were shown to have varying degrees of sensitivity to solvent polarity, depending on the nature of the electrostatic interaction between the two rings (Figure 4). For the donor–acceptor pair, which interact in the face-to-face stacked orientation, a strong solvent effect was observed, in which the interaction was stronger in more polar solvents. This suggests that dispersion and/or hydrophobic interactions dominate. In the case of the *self*-association of the electron-poor aromatic (1, Figure 4), although there is the possibility of greater burial of surface area, these molecules are found to stack in the offset

Figure 2



Cyclic *m*-phenylene ethynylenes.


orientation in the solid state, as predicted by the electrostatic model. Moreover, the solvent effect was an order of magnitude smaller for this system, presumably because less surface area is buried. The *self*-association of the electron-rich aromatic ring (2), which was the weakest of the three interactions and is believed to interact in the edge–face orientation, showed almost no solvent dependence. Thus, it appears that the electrostatic component influences the geometry of the interaction, which in turn influences the magnitude of the solvent effect on the interaction. The range of strengths of these interactions in water were about -6.7 to -18.8 kJ mol⁻¹. It is also worth noting that this study demonstrates the potential for selective aromatic interactions.

Substituent effects on aromatic interactions

Hunter and co-workers [23] have designed a system for measurement of both edge–face and stacking interactions through the use of a chemical double mutant cycle (Figure 5). With this system, they have found that there is an electronic effect on edge–face interactions in chloroform. Changes in the substituents, X and Y, gave a linear free-energy relationship with interaction energies from $+1.2$ kJ mol⁻¹ to -4.6 kJ mol⁻¹, with the unsubstituted edge–face interaction worth -1.4 kJ mol⁻¹. This is in contrast to previous work by Wilcox and co-workers [24,25], in which little electronic effect was observed and dispersion forces were suggested to be the most important component in edge–face interactions.

There have been some questions about the design of Hunter's system for double mutant cycles, including the rigidity of the 1,1-diphenylcyclohexyl subunit [26].

Figure 5



Hunter's double mutant cycle for measuring edge-face interactions. Deletion of one or the other aromatic ring (A to B or A to C) results in loss of the aromatic interaction, but may also have other secondary effects on the

strengths of the hydrogen bonds and other contributing interactions. The double mutant, D, accounts for these secondary changes, such that the sum $\Delta\Delta G(\pi-\pi)$ represents only the energy of the edge-face interaction.

Quadrupole-quadrupole interactions were also proposed to be important for electron-poor rings such as 4-nitroindole. Rosemeyer and Seela [35] have also shown that the polarizability of the dangling base correlates with stability of the duplex better than hydrophobicity as measured from RP-HPLC retention times, and have interpreted this as indicating that polarizability, rather than hydrophobicity, is the most important factor in stabilizing duplex DNA. A general difficulty with interpreting the results from dangling-base studies is that the geometry of interaction in these systems typically is not known. For example, Richert and co-workers [36[•]] found that duplex DNA with a 5'-quinolone cap was significantly stabilized relative to the uncapped sequence. However, detailed structural analysis of this system indicates that the quinolone does not cap the terminal T:A base pair, but in fact disrupts the terminal pair in favor of quinolone-adenine stacking against the penultimate G:C base pair.

Gellman and co-workers [37[•]] have studied the role of heteroatoms in aromatic interactions in their amide model system (Figure 3, $R = CH_2CO_2X$). In this system, the number and location of nitrogen atoms in two aromatic rings, Ar_1 and Ar_2 , were varied, and the effect on amide rotamer population was determined in water. The *E*-rotamer places Ar_1 and Ar_2 in close proximity, and a crystal structure indicates that Ar_1 and Ar_2 interact in an offset-stacked geometry in this rotamer. In this system, Gellman has found that, in general, an increase in the number of heteroatoms in either or both Ar_1 and Ar_2 resulted in an increase in the *E*-rotamer, presumably due to a favorable aromatic interaction. This argues for a significant polar component to the interaction. Nonetheless, when the experiments were performed in chloroform, very little effect was observed, indicating that aqueous solvent is required for significant interaction.

Conclusions

Recent studies of aromatic interactions have provided new insight into the nature of these biologically important non-covalent interactions in terms of their driving force, stability and selectivity. It is clear that the hydrophobic effect plays a role in these interactions, but it is not the sole factor involved in these attractive interactions. Moreover, its contribution depends on the geometry of the interaction, which is influenced by electrostatics. Recent work has demonstrated that aromatic interactions can provide *selectivity* as well as stability. Selective association between different aromatics demonstrates the importance of electrostatics to the magnitude of the interaction. Moreover, the observed selectivity between aromatic and aliphatic groups in water demonstrates the fundamental differences between aromatic and purely hydrophobic interactions. The advances in understanding of aromatic interactions should prove useful in their application in molecular and biomolecular recognition.

References and recommended reading

Papers of particular interest, published within the annual period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Kool ET: **Hydrogen bonding, base stacking, and steric effects in DNA replication.** *Annu Rev Biophys Biomol Struct* 2001, 30:1-22.
2. Kannan N, Vishveshwara S: **Aromatic clusters: a determinant of thermal stability of thermophilic proteins.** *Protein Eng* 2000, 13:753-761.
- Aromatic interactions are proposed to contribute to the thermal stability of thermophilic proteins in this article.
3. Gazit E: **A possible role for pi-stacking in the self-assembly of amyloid fibrils.** *FASEB J* 2002, 16:77-83.
4. Bhattacharyya R, Samanta U, Chakrabarti P: **Aromatic-aromatic interactions in and around alpha-helices.** *Protein Eng* 2002, 15:91-100.
5. Thomas A, Meurisse, R, Charleatoux B, Brasseur R: **Aromatic side chain interactions in proteins. I. Main structural features.** *Proteins* 2002, 48:628-634.
- These papers (see also [6]) describe where aromatic interactions are favored in protein structures.
6. Thomas A, Meurisse R, Brasseur R: **Aromatic side-chain interactions in proteins. II. Near- and far-sequence Phe-X pairs.** *Proteins*, 2002, 48:635-644.
7. McGaughey GB, Gagne M, Rappe AK: **Pi-stacking interactions — alive and well in proteins.** *J Biol Chem* 1998, 273:15458-15463.
8. Burley SK, Petsko GA: **Aromatic-aromatic interaction — a mechanism of protein-structure stabilization.** *Science* 1985, 229:23-28.
9. Serrano L, Bycroft M, Fersht AR: **Aromatic aromatic interactions and protein stability — investigation by double-mutant cycles.** *J Mol Biol* 1991, 218:465-475.
10. Shiels JC, Tuite JB, Nolan SJ, Baranger AM: **Investigation of a conserved stacking interaction in target site recognition by the U1A protein.** *Nucleic Acids Res* 2002, 30:550-558.
11. Nolan SJ, Shiels JC, Tuite JB, Cecere KL, Baranger AM: **Recognition of an essential adenine at a protein-RNA interface: comparison of the contributions of hydrogen bonds and a stacking interaction.** *J Am Chem Soc* 1999, 121:8951-8952.
12. Hunter CA, Lawson KR, Perkins J, Urch CJ: **Aromatic interactions.** •• *J Chem Soc Perkin Trans* 2001:651-669.
This is an excellent recent review of aromatic interactions.
13. Hunter CA, Sanders JKM: **The nature of pi-pi interactions.** *J Am Chem Soc* 1990, 112:5525-5534.
14. West J, Mecozzi S, Dougherty DA: **Theoretical studies of the supramolecular synthon benzene-hexafluorobenzene.** *J Phys Org Chem* 1997, 10:347-350.
15. Cozzi F, Siegel JS: **Interaction between stacked aryl groups in 1,8-diarylnaphthalenes — dominance of polar/pi over charge-transfer effects.** *Pure Appl Chem* 1995, 67:683-689.
16. Sindkhedkar MD, Mulla HR, Cammers-Goodwin A: **Three-state, conformational probe for hydrophobic, pi-stacking interactions in aqueous and mixed aqueous solvent systems: anisotropic solvation of aromatic rings.** *J Am Chem Soc* 2000, 122:9271-9277.
17. Lahiri S, Thompson JL, Moore JS: **Solvophobically driven pi-stacking of phenylene ethynylene macrocycles and oligomers.** *J Am Chem Soc* 2000, 122:11315-11319.
18. Hill DJ, Moore JS: **Helicogenicity of solvents in the conformational equilibrium of oligo(m-phenylene ethynylene)s: implications for foldamer research.** *Proc Natl Acad Sci USA* 2002, 99:5053-5057.
This is a thorough solvent study of aromatic interactions.
19. Shetty AS, Zhang JS, Moore JS: **Aromatic pi-stacking in solution as revealed through the aggregation of phenylacetylene macrocycles.** *J Am Chem Soc* 1996, 118:1019-1027.
20. Tobe Y, Utsumi N, Kawabata K, Nagano A, Adachi K, Araki S, Sonoda M, Hirose K, Naemura K: **m-Diethynylbenzene macrocycles: syntheses and self-association behavior in solution.** *J Am Chem Soc* 2002, 124:5350-5364.
21. Gardner RR, McKay SL, Gellman SH: **Solvent-dependent stabilization of the E configuration of propargylic secondary amides.** *Org Lett* 2000, 2:2335-2338.
22. Cubberley MS, Iverson BL: **H-1 NMR investigation of solvent effects in aromatic stacking interactions.** *J Am Chem Soc* 2001, 123:7560-7563.
This paper provides a clear example of the contribution of both solvophobic and electrostatic contributions to aromatic interactions.
23. Carver FJ, Hunter CA, Seward EM: **Structure-activity relationship for quantifying aromatic interactions.** *Chem Commun* 1998:775-776.
24. Paliwal S, Geib S, Wilcox CS: **Chemistry of synthetic receptors and functional-group arrays. 24. Molecular torsion balance for weak molecular recognition forces — effects of tilted-t edge-to-face aromatic interactions on conformational selection and solid-state structure.** *J Am Chem Soc* 1994, 116:4497-4498.
25. Kim E, Paliwal S, Wilcox CS: **Measurements of molecular electrostatic field effects in edge-to-face aromatic interactions and CH-pi interactions with implications for protein folding and molecular recognition.** *J Am Chem Soc* 1998, 120:11192-11193.
26. Martinez AG, Barcina JO, Cerezo AD: **Influence of highly preorganised 7,7-diphenylbornane in the free energy of edge-to-face aromatic interactions.** *Chem Eur J* 2001, 7:1171-1175.
27. Carver FJ, Hunter CA, Jones PS, Livingstone DJ, McCabe JF, Seward EM, Tiger P, Spey SE: **Quantitative measurements of edge-to-face aromatic interactions by using chemical double-mutant cycles.** *Chem Eur J* 2001, 7:4854-4862.
28. Adams H, Hunter CA, Lawson KR, Perkins J, Spey SE, Urch CJ, Sanderson JM: **A supramolecular system for quantifying aromatic stacking interactions.** *Chem Eur J* 2001, 7:4863-4877.
29. Adams H, Blanco LJ, Chessari G, Hunter CA, Low CMR, Sanderson JM, Vinter JG: **Quantitative determination of intermolecular interactions with fluorinated aromatic rings.** *Chem Eur J* 2001, 7:3494-3503.
30. Goodman AJ, Breinlinger EC, McIntosh CM, Grimaldi LN, Rotello VM: **Model systems for flavoenzyme activity. Control of flavin recognition via specific electrostatic interactions.** *Org Lett* 2001, 3:1531-1534.
31. Rashkin MJ, Waters ML: **Unexpected substituent effects in offset pi-pi stacked interactions in water.** *J Am Chem Soc* 2002, 124:1860-1861.
32. Butterfield SM, Patel PR, Waters ML: **Contribution of aromatic interactions to alpha-helix stability.** *J Am Chem Soc* 2002, 124:9751-9755.
33. Tatko CD, Waters ML: **Selective aromatic interactions in beta-hairpin peptides.** • *J Am Chem Soc* 2002, 124:9372-9373.
This paper clearly demonstrates the selectivity of aromatic interactions relative to aliphatic interactions in a peptide model system.

34. Guckian KM, Schweitzer BA, Ren RXF, Sheils CJ, Tahmassebi DC, Kool ET: **Factors contributing to aromatic stacking in water: evaluation in the context of DNA.** *J Am Chem Soc* 2000, **122**:2213-2222.

This is a thorough paper investigating the correlation between a dangling aromatic group and duplex DNA stability.

35. Rosemeyer H, Seela F: **Modified purine nucleosides as dangling ends of DNA duplexes: the effect of the nucleobase polarizability on stacking interactions.** *J Chem Soc Perkin Trans* 2002:746-750.

36. Tuma J, Connors WH, Stitelman DH, Richert C: **On the effect of covalently appended quinolones on termini of DNA duplexes.** *J Am Chem Soc* 2002, **124**:4236-4246.

This paper provides a detailed structural investigation of a dangling aromatic group and demonstrates that not all dangling aromatics stack at the terminus of the duplex.

37. McKay SL, Haptonstall B, Gellman SH: **Beyond the hydrophobic effect: attractions involving heteroaromatic rings in aqueous solution.** *J Am Chem Soc* 2001, **123**:1244-1245.

This paper provides the only example of the stacking of heteroaromatic rings in a context other than nucleic acids.