

“Wreath products of groups acting with bounded orbits” answer to referee’s comments

Paul-Henry Leemann, Grégoire Schneeberger

March 1, 2022

We thank the anonymous reviewer for their careful reading of our manuscript and their insightful comments and suggestions, that we have taken into account.

While we took in account all the referee’s suggestions and remarks, we will not comment below the ones concerning grammatical issues and obvious typos.

Several comments (25, 26, 27, 29) of the reviewer asked for rewriting some statements and proofs with a more axiomatic and general approach. While we agree with the referee suggestion, the implementation of this change was a little more heavy than the referee suggested. In order to keep the article readable by non-specialists in category theory, we choose to add more contexts and examples before the statements of the axioms. In practice, this means that the material on pages 11 to 13 (starting at the § before Definition 2.11 and ending just before Subsection 2.4) is either new, or has been considerably rewritten. We also added at the beginning of Section 3 a table recapitulating which axioms are satisfied by the categories under consideration. The proof of Lemma 3.9 (new notation) has also been slightly modified to work in this general context, but the changes are small. And a small discussion has been added after the proof.

2. *Theorem B: The result is rather elementary, maybe a proposition instead of a theorem?*

We followed the referee’s suggestion.

3. *Page 4, Median graphs: It could be interesting to mention explicitly the connection with $CAT(0)$ cube complexes (Gerasimov, Chepoi, Roller).*

We added the following “The class of median graphs was introduced by Nebeský in 1971 [Neb71] and Roller [Rol98] and Chepoi [Che00] realized independently that this class can be naturally identified to the class of $CAT(0)$ cube complexes.”

4. *Page 5, line -4: It is not clear that the sequence is increasing. I guess the sequence $(H_n)_n$ should be replaced with a well-chosen increasing subsequence $(H_{r_n})_n$.*

Indeed, the sequence $(H_n)_n$ is only decreasing. We added “Since they are proper subgroups and $H_n \leq H_{n+1}$, we can extract an increasing subsequence $(H_{r_n})_n$ that still satisfies $G = \bigcup_n H_{r_n}$.”

5. *Page 6, line -15: I do not find that the example about graphs is convincing: describing graphs as discrete metric spaces, isometries are automatically automorphisms. But the same example makes sense for cube complexes.*

We changed the example from graphs to cube complexes. We also added a reference to Example 2.8.

6. *Page 7, line 4: Isn't it already the case for Bergman's original example $\text{Sym}(N)$ (much before Cornulier)?*

Indeed. The reference was changed accordingly.

7. *Page 7, Proposition 2.5: In my copy of the article, there are TeX problems with the arrows. Is $FH \implies FR$ missing?*

Added the arrow, the sentence “The implication $[FH \implies FR]$ follows from the fact that real trees are median metric space, and that such spaces can be embedded into L^1 -spaces (see for instance [Ver93, Theorem V.2.4]).” and the proof that the converse implication does not holds (similar to properties FA versus FW).

8. *Page 7, line -8: A triangle Coxeter group would be a more explicit example of a group with FA but not FW.*

Such an example as been added after the proof of Proposition 2.5. We choose to not use this example in the proof, as it use the wall structure and not the median graph structure.

12. *Page 9, line -9: This is not true: the infinite dihedral group does not surject onto Z since it is generated by elements of order two. However, looking at the subcategory Z with orientation-preserving isometries, the property BS amounts to not having Z as a quotient.*

Indeed. This has been corrected following the referee's suggestion.

13. *Page 9, line -6: No need of Bass-Serre theory: Z is a tree, so the implication is obvious.*

This has been corrected following the referee's suggestion.

14. *Page 10, Lemma 2.10: There is a more direct argument: Let X be a metric space. Let $G(X)$ denotes the graph obtained from a vertex-set X by applying the following process: for any two $x, y \in X$ add a path of length $[d(x, y)] + 1$ between x and y . $G(X)$ is connected and the obvious inclusion $\iota: X \rightarrow G(X)$ is a quasi-isometric embedding. Moreover, the construction is canonical, so every group action on X extends to a group action on $G(X)$, making ι equivariant. So if a group satisfying the bounded orbit property on connected graphs acts on a metric space X , then its induced action on $G(X)$ has bounded orbits, which implies that its orbits in X are bounded.*

We followed the referee's suggestion. Nevertheless, we kept the reference to Cornulier's characterization of Bergmann's property after the proof.

15. *Page 10, Definition 2.11: Is it a good terminology? The subcategory of bounded metric spaces has unbounded Cartesian products...*

It is not! We changed it to “bornological cartesian product”.

16. *Page 10, Definition 2.11, item 2: Not clear. Does it mean that the obvious image of $\text{Aut}(X)^n \rtimes \text{Sym}(n)$ in $\text{Bij}(X^n)$ lies in $\text{Aut}_{\mathbf{S}}(X^n)$ or that $\text{Aut}_{\mathbf{S}}(X^n)$ contains a subgroup isomorphic to $\text{Aut}(X)^n \rtimes \text{Sym}(n)$? The meaning is clear from the proofs, but it should be precise already in the definition.*

This has been clarified to “the canonical image of $\text{Aut}_{\mathbf{S}}(X)^n \rtimes \text{Sym}(n)$ in $\text{Bij}(X^n)$ lies in $\text{Aut}_{\mathbf{S}}(X^n)$ ”.

17. *Page 12, line 6: Roller proved the general bounded orbit property much before Cornulier.*

The reference has been updated to [Roll98].

18. *Page 12, second paragraph: It could be worth mentioning that one recovers a global fixed point in the category of $\text{CAT}(0)$ cube complexes.*

Added the precision “(i.e. complete $\text{CAT}(0)$ spaces and in particular $\text{CAT}(0)$ cube complexes which are either finite dimensional or locally finite)”. If a CCC is neither finite dimensional nor locally finite, then it is not complete and Proposition 2.12 does not apply. We do not know if it is possible to recover a global fixed point in the whole category of $\text{CAT}(0)$ cube complexes, but will be happy for any reference on this fact.

19. *Page 13, Lemma 3.1: The first sentence of the proof is not used, so it can be removed.*

This has been done.

20. *Page 14, line 19: The notation $H \cong \{giN\}$ is not clear.*

Changed to “ $H \cong \{giN \mid i \in I\}$ with the quotient multiplication”.

22. *Page 15, Lemma 3.7: The group G is not defined.*

The beginning of the Lemma is now “Let G and $(G_x)_{x \in X}$ be non-trivial groups.”

23. *Page 15, lines 13-14: Each G should be replaced with G_x .*

Done.

24. *Page 15, line 15: An enumeration of Y has to be fixed.*

This is now “So let us fix an enumeration of Y and let $K := \bigoplus_{i \geq 1} G_i$ ”. Observe the use of the new variable K to not conflict with the G of the second part of the lemma.

25. *Page 15, Lemma 3.7: I find weird that what is proved is rather different from what is claimed in the lemma. The proof is really about uncountable cofinality and not about BS. A suggestion: write your lemma as a corollary of a new lemma stating that, if a semidirect product $A \rtimes B$ has uncountable cofinality, then there exists a finite subset $S \subset A$ such that A is generated by $\bigcup_{b \in B} bSb^{-1}$ (From the characterization of uncountable cofinality in terms of subgroups, this is pretty obvious.)*

The general result is now Corollary 3.8, which follows from Lemma 3.7 (now written only for uncountable cofinality). However, we kept the old (geometric) proof as it is most likely to be generalized to other categories. Nevertheless, we added as an in-text remark that the algebraic proof is easier and more direct.

26. *Page 16, Lemma 3.8: Same remark as before: it would be more natural to state the lemma about FW and to mention the result as a corollary.*

See next item.

27. Page 16, Lemma 3.8: *It could be interesting to formulate the lemma in greater generality. If G has a non-trivial action on Z , then $G \wr_X H$ naturally acts on $\bigoplus_X Z$ with unbounded orbits if X is infinite. So you just need your category to have a well-defined infinite product operation and the property that every non-trivial group admits a non-trivial action on one space in the category. (For FW, G acts non-trivially on the tree whose vertices are G , $\{g\}$ ($g \in G$) and whose edges are given by inclusion.).*

We followed the referee's suggestion to give a more axiomatic proof. However, the axioms they proposed are not enough, as demonstrated by the case of ultrametric spaces and uncountable cofinality. The solution is to keep the non-trivial actions axiom and to require the existence of infinite Cartesian powers that are in some rough sense able to detect the number of coordinates on which two elements of the product differs. In particular, this excludes the sup distance on the Cartesian product, and rules out ultrametric spaces.

28. Page 17: Lemma 3.11 can be seen as a direct corollary of Lemma 3.10(2). *Indeed, if H' is the kernel of the action of H on X , the pre-image of H' under the quotient map $G \wr_X H \rightarrow H$ is a finite-index subgroup that splits as $H' \oplus \bigoplus_X G$. The latter must have BS, but G is a quotient of this group.*

The referee's remark is correct and, depending on personal tastes, one might argue that their proof is shorter and easier. However, we purposely tried to give geometric proofs when possible as we think that, in this particular context, they are more enlightening than more abstract algebraic proofs. Nevertheless, we added the following after the proof of Lemma 3.11.

“Observe that it is also possible to derive Lemma ?? directly from Lemma ?? (2), with a more algebraic proof. Indeed, using the notation and hypothesis of Lemma ??, let H' be the kernel of the action of H on X and $\pi: G \wr_X H \rightarrow H$ be the canonical projection. Then $\pi^{-1}(H') \cong \bigoplus_X G \oplus H'$ is a finite index subgroup of $G \wr_X H$ and hence has property BS. Since G is a quotient of $\bigoplus_X G \oplus H'$ we conclude that it also has property BS.”

29. Page 17, Theorem 3.12: *The implication $BS \implies FW$ is a rather artificial assumption. I think you should list the few axioms you are actually using. It seems to me that there are only two: (1) an axiom about products, with possibly infinitely many factors, which could generalise the definition of unbounded Cartesian powers; (2) every non-trivial group G admits a non-trivial action on a space in \mathcal{S} . If $G \wr_X H$ has BS, then X must be finite by Lemma 3.8 (deduced from (1) and (2)), H must have BS because it is a quotient, and G must have BS because of Lemma 3.11 (which is a corollary of Lemma 3.10, itself obtained from (1)). Conversely, if X is finite and G, H have BS, then $G \wr_X H$ virtually splits as a direct sum of finitely many copies of G and a finite-index subgroup of H . But stability of BS under direct sum is given by (1).*

See Item 27. More precisely, we wrote axioms (1) as the union of two axioms. The first one asserts the existence of finite Cartesian powers satisfying some compatibility condition with the bornology (diagonal(unbounded subset) is unbounded), the second one asserts the existence of infinite Cartesian powers satisfying another compatibility condition with the bornol-

ogy (given two elements $x \neq y \in X$, the set $\{f \in X^I \mid f(i) \in \{x, y\}, f(i) = x \text{ for all but finitely many } i\}$ is unbounded).

30. *Page 19, Lemma 3.13: The assumption of having finitely many orbits can be removed.*

The assertion has been removed and the proof was extended accordingly. The new proof:

“The desired result then follows from $(G \wr_X H)^{\text{ab}} \cong \bigoplus_{X/H}(G^{\text{ab}}) \times H^{\text{ab}}$ and the claim that a direct sum $\bigoplus_{y \in Y} K_y$ has a quotient isomorphic to \mathbf{Z} if and only if at least one of the factor has a quotient isomorphic to \mathbf{Z} . Indeed, one direction of the claim is trivial. For the other direction, remind that K does not project onto \mathbf{Z} if and only if any action of K by orientation preserving isomorphisms on Z the 2-regular tree has bounded orbits. But the only possibility for such an action to have bounded orbits is to be trivial. If none of the K_y projects onto \mathbf{Z} , all their actions on Z are trivial and so is any action of $\bigoplus_{y \in Y} K_y$, which can therefore not project onto \mathbf{Z} .”

31. *Page 19: As written, Theorem C is only proved for FA.*

The statement of Thm 1.2 and Proposition 3.14 have been rewritten to take in account property **FR**. Observe that the proofs remain unchanged (Theorem 1.2 is proved in [Cornulier-Kar] both for FA and **FR**).

32. *Proofs of Lemma 3.7 and Theorem B: You should mention that the proofs using the characterisation in terms of subgroups are actually much shorter. In fact, they are almost exercises.*

The introduction to the proof of Lemma 3.7 now reads “It is of course possible to prove Lemma ?? using the characterization of uncountable cofinality in terms of subgroups, *in which case the proof is a short exercise let to the reader.* However, we find enlightening to prove it using the characterization in terms of actions on ultrametric spaces.” (the italicized part is new). A similar statement is made before the proof of Thm B.

33. *In all the article: “an S-space” instead of “a S-space”?*

We followed the referee’s suggestion.

- 34.