

Property FW and wreath products of groups: a simple approach using Schreier graphs

Paul-Henry Leemann, Grégoire Schneeberger

Friday 22nd January, 2021 11:31

Abstract

The group property FW stands in-between the celebrated Kazhdan's property (T) and Serre's property FA. Among many characterizations, it might be defined, for finitely generated groups, as having all Schreier graphs one-ended. Using this, we show that a finitely generated wreath product $G \wr_X H$ has property FW if and only if both G and H have property FW and X is finite.

1 Introduction

Property FW is a group property that is (for discrete groups) a weakening of the celebrated Kazhdan's property (T). It was introduced by Barnhill and Chatterji in [1]. It is a fixed point property for actions on wall spaces or, equivalently, on CAT(0) cube complexes. So it stands between property FH (fixed points on Hilbert spaces, equivalent to property (T) for discrete groups) and property FA (fixed points on *arbres*¹). It is known that all these properties are different, see [4] for examples of groups which distinguish them.

When working with group properties, it is natural to ask if they are stable under “natural” group operations. For example, one may wonder when a property is stable by subgroups, quotients, direct products or semi-direct products.

A slightly less common operation, but still of great use in geometric group theory, is the wreath product, see Definition 2.4.

In the context of properties defined by fixed points of actions, the first result concerning wreath products was due to Cherix, Martin and Valette and latter refined by Neuhauser and concerns property (T).

Theorem 1.1 ([3, 12]). *Let G, H be two discrete groups with G non-trivial and X a set on which H acts. The wreath product $G \wr_X H$ has property (T) if and only if G and H have property (T) and X is finite.*

The aim of this note is to give an elementary proof of analogous of Theorem 1.1 for property FW:

Theorem 1.2. *Let G, H be two finitely generated groups with G non-trivial and X a set on which H acts with finitely many orbits. The wreath product $G \wr_X H$ has property FW if and only if G and H have property FW and X is finite.*

¹ *Arbres* is the french word for trees.

Est ce que
c'est écrit
quelques
part les
preuves
pour cela
? Si oui
citer si non
les écrire
pour être
complet

Our proof will rely on a characterization of property FW via ends of Schreier graphs, see Subsection 2.1 for the relevant definitions. Note that Cornulier also proved in [4] that if $G \wr_X H$ has property FW (and G is non trivial), then X is finite.

Cornulier and Kar studied the question of stability of this property for property FA in [5]. They proved a similar result for actions of H on X with finitely many orbits and without fixed point, but the condition that G has property FA replaced by G has finite abelianization and can not be expressed as an union of properly increasing sequence of subgroups.

At this point, the curious reader might have two questions. First, is it possible to extend Theorem 1.2 beyond the realms of finitely generated groups and of actions with finitely many orbits? And secondly, is there a link between Theorems 1.1 and 1.2? In both cases, the answer is yes. This is the subject of the forthcoming and more technical paper [10], which gives an unified proof of Theorems 1.1 and 1.2 as well as of similar results for the Bergman's property and more.

Peut être
mettre une
phrase qui
explique
en quoi
c'est dif-
férent de
FA

Organization of the paper The next section contains all the definitions as well as some examples, while Section 3 is devoted to the proof of Theorem 1.2 and some related results.

Acknowledgment The authors are thankful to A. Genevois, T. Nagnibeda and A. Valette for helpful comments on a previous version of this note, and to the Swiss National Fund for Scientific Research for its support.

2 Definitions and examples

This section contains all the definitions, some standard but usefull preliminary results as well as some examples.

2.1 Ends of Schreier graphs and property FW

In what follows, we will always assume that generating sets of groups are *symmetric*, that is we will look at $S \subset G$ such that $s \in S$ if and only if $s^{-1} \in S$. Our graphs will be undirected and we will sometimes identify a graph with its vertex set.

Definition 2.1. Let G be a group, H a subgroup of G and S a symmetric generating set. The (left) *Schreier graph* $\text{Sch}(G, H; S)$ is the graph with vertices the left cosets $gH = \{gh \mid h \in H\}$ and for every $\{s, s^{-1}\} \subset S$ an edge between gH and sgH .

Observe that with the above definition, a Schreier graph may have loops as well as multiple edges. Moreover, in $\text{Sch}(G, H; S)$ every vertex has degree $|S|$.

If X is a set with a left action $G \curvearrowright X$ and S is a symmetric generating set for G , the corresponding (left) *orbital graph* $\text{Sch}_\mathcal{O}(G, X; S)$ is the graph with vertex set X and with an edge between x and y for every $s \in S$ such that $s.x = y$. As the notation suggest, these two definitions are the two faces of the same coin. Indeed, we have $\text{Sch}(G, H; S) = \text{Sch}_\mathcal{O}(G, G/H; S)$ for the

action by left multiplication of G on G/H , while for every $x \in X$, the graph $\text{Sch}(G, \text{Stab}_G(x); S)$ is equal to the connected component of x in $\text{Sch}_\mathcal{O}(G, X; S)$.

Schreier graphs are generalizations of the well-known *Cayley graphs*, with $\text{Cayl}(G; S) = \text{Sch}(G, \{1\}; S)$, see Figures 1 and 2 for some examples.

If S and T are two generating sets of G , the graphs $\text{Sch}(G, H; S)$ and $\text{Sch}(G, H; T)$ need not to be isomorphic. However, if both S and T are finite, then $\text{Sch}(G, H; S)$ and $\text{Sch}(G, H; T)$ are quasi-isometric, see [6, IV.B.21.iii] for a proof and Figure 1 for an example. The only fact we will use about quasi-isometries is that they preserve “large-scale properties” of the graph, as for example the number of ends. Observe that the requirement that both S and T are finite is crucial for the existence of a quasi-isometry between the corresponding Schreier graphs. Indeed, for every group $\text{Cayl}(G; G \setminus \{1\}) = \text{Sch}(G, \{1\}; G \setminus \{1\})$ is always a complete graph on $|G|$ vertices, in particular $\text{Cayl}(\mathbf{Z}; \{1\})$ and $\text{Cayl}(\mathbf{Z}; \mathbf{Z} \setminus \{0\})$ are not quasi-isometric.




Figure 1: Fragments of two Cayley graphs of \mathbf{Z} (2 ends), for the standard generating set $\{\pm 1\}$ and for the generating set $\{\pm 2, \pm 3\}$.




Figure 2: Fragments of the Cayley graphs of \mathbf{Z}^2 (1 end) on the left and of F_2 (infinitely many ends) on the right; with standard generating sets.




Figure 3: A fragment of a Schreier graph (4 ends) of the free group $F_2 = \langle x^{\pm 1}, y^{\pm 1} \rangle$.

Let Γ be a graph and K a finite subset of vertices. The graph $\Gamma \setminus K$ is the

subgraph of Γ obtained by deleting all vertices in K and all edges adjacent to them. This graph is not necessarily connected.

Definition 2.2. Let Γ be a graph. The *number of ends* of Γ is the supremum, taken over all finite K , of the number of infinite connected components of $\Gamma \setminus K$.

There exists other characterization of the number of ends in graphs, see [7] and the references therein, but Definition 2.2 is the one that best suits our purpose. A locally finite graph (i.e. such that every vertex has finite degree) is finite if and only if it has 0 end.

An important fact about the number of ends of a graph, is that it is an invariant of quasi-isometry, see [2]. In particular, if G is a finitely generated group it is possible to speak about the number of ends of the Schreier graph $\text{Sch}(G, H; S)$ without specifying a particular finite generating set S . By a celebrated result of Hopf [9], the number of ends of a Cayley graph of a finitely generated group can only be 0, 1, 2 or infinite (in which case it is uncountable), see Figures 1 and 2 for some examples. On the other hand, Schreier graphs may have any number of ends in $\mathbf{N} \cup \{\infty\}$, see Figure 3 for an example of a graph with 4 ends. In fact, every regular graph of even degree is isomorphic to a Schreier graph, [8, 11].

We are now finally able to introduce property FW. Instead of giving the original definition in terms of actions on wall spaces, we will use an equivalent one for finitely generated group due to Cornuier [4].

Definition 2.3. A finitely generated group G has *property FW* if all its Schreier graphs have at most one end.

It directly follows from the definition that all finite groups have property FW, but that \mathbf{Z} does not have it. In fact, if G is a finitely generated group with an homomorphism onto \mathbf{Z} , then it does not have FW. Indeed, in this case $G \cong H \rtimes \mathbf{Z}$ for some H and the Schreier graph $\text{Sch}(G, H; S)$ will be isomorphic to a Cayley graph of $\mathbf{Z} \cong G/H$ and hence has 2 ends.

Property FW admits many distinct characterizations that allow to define it for groups that are non-necessarily finitely generated and even for topological groups. We refer the reader to [4] for a survey of these characterizations.

2.2 Wreath products

Let X be a set and G a group. We view $\bigoplus_X G$ as the set of functions from X to G with finite support:

$$\bigoplus_X G = \{\varphi: X \rightarrow G \mid \varphi(x) = 1 \text{ for all but finitely many } x\}.$$

This is naturally a group, where multiplication is taken componentwise.

If H is a group acting on X , then it naturally acts on $\bigoplus_X G$ by $(h \cdot \varphi)(x) = \varphi(h^{-1} \cdot x)$. This leads to the following standard definition

Definition 2.4. Let G and H be groups and X be a set on which H acts. The (restricted) wreath product $G \wr_X H$ is the group $(\bigoplus_X G) \rtimes H$.

A prominent source of examples of wreath products are the ones of the form $G \wr_H H$, where H acts on itself by left multiplication. In particular, the group

$(\mathbf{Z}/2\mathbf{Z}) \wr_{\mathbf{Z}} \mathbf{Z}$ has become well-known under the name of the *lamplighter group*. Another (trivial) examples of wreath products are direct products $G \times H$ which corresponds to wreath products over a singleton $G \wr_{\{*\}} H$.

Let S be a generating set of G and T a generating set of H . Let $\{x_i\}_{i \in I}$ be a choice of a representative in each H -orbit. Finally, let δ_x^s be the element of $\bigoplus_X G$ defined by $\delta_x^s(x) = s$ and $\delta_x^s(y) = 1_G$ if $y \neq x$ and let $\mathbf{1}$ be the constant function with value 1_G . It is then standard that

$$\{(\delta_{x_i}^s, 1_H) \mid s \in S, i \in I\} \cup \{(\mathbf{1}, t) \mid t \in T\}$$

is a generating set for $G \wr_X H$.

On the other hand, if $\{(\varphi_i, h_i) \mid i \in I\}$ is a generating set of $G \wr_X H$, then $\{h_i \mid i \in I\}$ is a generating set of H while $\{\varphi_i(x) \mid i \in I, x \in X\}$ is a generating set of G . Observe that since the φ_i take only finitely many values, if I is finite, so is $\{\varphi_i(x) \mid i \in I, x \in X\}$. We hence recover the following characterization of the finite generation of $G \wr_X H$.

Lemma 2.5. *The group $G \wr_X H$ is finitely generated if and only if both G and H are finitely generated and H acts on X with finitely many orbits.*

Proof. If $G \wr_X H$ is finitely generated, so is its abelianization $(G \wr_X H)^{\text{ab}} \cong (\bigoplus_{X/H} G^{\text{ab}}) \times H^{\text{ab}}$, which implies that the orbit set X/H is finite. The other implications directly follows from the above discussion on generating sets. \square

Using the above lemma, we could reformulate Theorem 1.2 in the following way:

Proposition 2.6. *Let G, H be two groups with G non-trivial and X a set on which H acts. Suppose that all three of G, H and $G \wr_X H$ are finitely generated. Then the wreath product $G \wr_X H$ has property FW if and only if G and H have property FW and X is finite.*

3 Proof of the main result

This section is devoted to the proof of Theorem 1.2. This proof is split into two parts: Lemma 3.1 and its Corollary 3.3 and Lemma 3.4.

We begin by a result on semi-direct products.

Lemma 3.1. *Let N and H be two finitely generated groups and $N \rtimes H$ a semi-direct product. Then*

1. *If $N \rtimes H$ has property FW, then so does H ,*
2. *If both N and H have property FW, then $N \rtimes H$ also has property FW.*

Proof. Let S , respectively T , denotes a finite generating set of N , respectively H . Then the group $G := N \rtimes H$ is finitely generated by $U = (S \times \{1\}) \cup (\{1\} \times T)$.

Suppose that H does not have property FW. Then there exists a Schreier graph $\Gamma = \text{Sch}(H, K; T)$ of H with more than one end. The group G acts on the vertices Γ via $(n, h).x := h.x$. Since H acts transitively on Γ , so does G . In fact, the graph of the action G on the vertices of Γ is isomorphic to the graph Γ with some additional loops for generators in $S \times \{1\}$. As adding loops does

not change the number of ends, this Schreier graph has more than one end and therefore G does not have property FW.

Suppose now that both N and H have property FW. We want to show that every Schreier graphs of G has at most one end. If they are all finite, then there is nothing to prove (and G is finite). So let Γ be an infinite Schreier graph of G with respect to the generating set U . The groups N and H acts on the vertices of Γ by restriction of the action of G . That is, $n.x = (n, 1).x$ and $h.x = (1, h).x$. For each vertex x we define Γ_x^H (and respectively Γ_x^N) as the Schreier graph obtained from the action of H (respectively N) on the H -orbit (respectively N -orbit) of x . These are subgraphs of Γ . As N and H have property FW, the graphs Γ_x^H and Γ_x^N are either finite or one-ended. We want to prove that in this case Γ has exactly one end.

Let K be a finite set of vertices of Γ . If x is in K and Γ_x^H is finite, add all vertices of Γ_x^H to K . By doing so for every x in K , we obtain a new finite set $K \subset K'$ of vertices of Γ . We will show that $\Gamma \setminus K'$ has only one infinite connected component. By definition of K' , if x is not in K' , then either Γ_x^H has one end or Γ_x^H does not contains vertices of K' .

Let x and y be two vertices, each of them lying in some infinite connected component of $\Gamma \setminus K'$. We will construct a path from x to y in $\Gamma \setminus K'$ as a concatenation of three smaller paths, see Figure 4, as follows. First, a path in $\Gamma_x^H \setminus K'$ from x to some z , then a path in $\Gamma_z^N \setminus K'$ from z to some $z' \in (\Gamma_z^N \cap \Gamma_y^H) \setminus K'$, and finally a path in $\Gamma_y^H \setminus K'$ from z' to y . In order to finish the proof, it remains to exhibit elements z and z' and the three desired paths.




Figure 4: The path between x and y .

The action of G on Γ being transitive, there exists an element (n_0, h_0) of $N \rtimes H$ such that $(n_0, h_0).x = y$. Since K' is finite, the set $\Gamma_x^H \setminus K'$ is infinite. Moreover, there is infinitely many z in $\Gamma_x^H \setminus K'$ such that either Γ_z^N is one-ended or Γ_z^N does not intersect K' . For such a z there exists h such that $(1, h).x = z$. Now, the vertex $z' := (hh_0^{-1}.n_0, h).x$ is both equal to $(hh_0^{-1}.n_0, 1)(1, h_0).x = (hh_0^{-1}.n_0, 1).z$ and to $(1, hh_0^{-1})(n_0, h_0).x = (1, hh_0^{-1}).y$. That is, z' is in $\Gamma_z^N \cap \Gamma_y^H$. A simple computation show us that the map $z \mapsto z'$ is injective: $z'_1 = z'_2$ if and only if $z_1 = z_2$. Since K' is finite, there is only finitely many z' in K' and hence there is infinitely many $z \in \Gamma_x^H$ such that both z and z' are not in K' and either

Γ_z^N is one-ended or Γ_z^N does not intersect K' .

In order to finish the proof, observe that the three graphs Γ_x^H , Γ_y^H and Γ_z^N are all either one-ended or do not intersect K' . Therefore, there is a path in $\Gamma_x^H \setminus K'$ from x to z as desired, and similarly for the paths from z to z' and z' to y . We just have proved that for any finite K the graph $\Gamma \setminus K$ has only one infinite connected component and therefore that Γ is one-ended. \square

As a direct corollary, we have

Corollary 3.2. *Let G and H be two finitely generated groups. Then $G \times H$ has property FW if and only if both G and H have property FW.*

By iterating Lemma 3.1, we obtain

Corollary 3.3. *Let G and H be two finitely generated groups and X a set on which H acts with finitely many orbits. Then,*

1. *If $G \wr_X H$ has property FW, then so does H ,*
2. *If both G and H have property FW and X is finite, then $G \wr_X H$ has property FW.*

The following Lemma finishes the proof of Theorem 1.2.

Lemma 3.4. *Let G and H be two finitely generated groups with G non-trivial and such that H acts on some set X with finitely many orbits. If $G \wr_X H$ has property FW, then*

1. *G has property FW,*
2. *X is finite.*

Proof. We will prove the contrapositives. The idea is to construct, for each of the two cases, a Schreier graph of $G \wr_X H$ with more than one end by exhibiting some well-chosen action of $G \wr_X H$. We fix some finite generating sets S and T of G and H and let

$$U := \{(\delta_x^s, 1_H) \mid s \in S\} \cup \{(1, t) \mid t \in T\}$$

be the standard generating set of $G \wr_X H$.

Suppose that X is an infinite set. Since H acts on X with finitely many orbits, there exists an infinite orbit X' . Let x_0 be an arbitrary vertex of X' . The group G acting on itself by left multiplications, we have the so-called *imprimitive action* of the wreath-product $G \wr_X H$ on $Y := G \times X'$:

$$(\varphi, h).(g, x) := (\varphi(h.x)g, h.x).$$

Since both $G \curvearrowright G$ and $H \curvearrowright X'$ are transitive, the action $G \wr_X H \curvearrowright Y$ is also transitive. Therefore, the orbital Schreier graph of $G \wr_X H \curvearrowright Y$ is isomorphic to the Schreier graph $\Gamma := \text{Sch}(G \wr_X H, \text{Stab}((1_G, x_0)), U)$. We decompose this graph into leaves of the form $Y_g = \{g\} \times X'$. There are two types of edges in Γ , which are coming from the two sets of generators, see Figure 5. The first ones,

of the form $(\mathbf{1}, t)$, give us on each leaf a copy of the orbital Schreier graph of $H \curvearrowright X'$. Indeed,

$$(\mathbf{1}, t).(g, x) = (g, t.x).$$

The second ones, of the form $(\delta_{x_0}^s, 1)$, give us loops everywhere except on vertices of the form (g, x_0) . By direct computation, we see that the vertices (g, x_0) and (sg, x_0) connect the leaves Y_g and Y_{sg} ,

$$(\delta_{x_0}^s, 1).(g, x) = \begin{cases} (g, x) & \text{if } x \neq x_0, \\ (sg, x) & \text{if } x = x_0. \end{cases}$$




Figure 5: The leaf structure of the orbital Schreier graph of $G \wr_X H \curvearrowright Y$. Plain edges correspond to generators of the form $(\mathbf{1}, t)$ while dotted edges correspond to generators of the form $(\delta_{x_0}^s, 1)$.

If we remove a vertex (g, x_0) we disconnect the leaf Y_g from the rest of Γ . As X' is infinite each leaf is infinite and the number of ends of Γ is then at least $|G| \geq 2$. We just proved that if X is infinite the group $G \wr_X H$ does not have property FW.

Suppose now that G does not have property FW. Then there exists a subgroup K of G such that $\text{Sch}(G, K, S)$ has more than one end. Let x_0 be any point of X and X' be its orbit under the action of H . We have the imprimitive action of $G \wr_X H$ on $G/K \times X$, which restricts to an action on $G/K \times X'$:

$$(\varphi, h).(gK, x) = (\varphi(h.x)gK, h.x).$$

As above, the action is transitive and the orbital Schreier graph of this action is isomorphic to a Schreier graph Γ of $G \wr_X H$. We decompose this graph into leaves in the same way. Now we look at the subgraph made up of vertices

(g, x_0) and edges $(\delta_{x_0}^s, 1)$ and we remark that it is isomorphic to the Schreier graph $\text{Sch}(G, K, S)$ which has more than one end. Then Γ has also more than one end, which finishes the proof. \square

References

- [1] Angela Barnhill and Indira Chatterji. Property (T) versus property FW. *Enseign. Math. (2)*, 54(1-2):16–18, 2008.
- [2] Stephen G Brick. Quasi-isometries and ends of groups. *J. Pure Appl. Algebr.*, 86(1):23–33, 1993.
- [3] Pierre-Alain Cherix, Florian Martin, and Alain Valette. Spaces with measured walls, the Haagerup property and property (T). *Ergodic Theory Dynam. Systems*, 24(6):1895–1908, 2004.
- [4] Yves Cornuier. Group actions with commensurated subsets, wallings and cubings. *arXiv e-prints*, page arXiv:1302.5982, February 2013.
- [5] Yves Cornuier and Aditi Kar. On property (FA) for wreath products. *J. Group Theory*, 14(1):165–174, 2011.
- [6] Pierre de la Harpe. *Topics in geometric group theory*. Chicago Lectures in Mathematics. University of Chicago Press, Chicago, IL, 2000.
- [7] Reinhard Diestel and Daniela Kühn. Graph-theoretical versus topological ends of graphs. *J. Combin. Theory Ser. B*, 87(1):197–206, 2003. Dedicated to Crispin St. J. A. Nash-Williams.
- [8] Jonathan L. Gross. Every connected regular graph of even degree is a Schreier coset graph. *J. Combinatorial Theory Ser. B*, 22(3):227–232, 1977.
- [9] Heinz Hopf. Enden offener Räume und unendliche diskontinuierliche Gruppen. *Comment. Math. Helv.*, 16:81–100, 1944.
- [10] Paul-Henry Leemann and Grégoire Schneeberger. Groups acting with bounded orbits and wreath products. In preparation.
- [11] Alexander Lubotzky. Cayley graphs: eigenvalues, expanders and random walks. In *Surveys in combinatorics, 1995 (Stirling)*, volume 218 of *London Math. Soc. Lecture Note Ser.*, pages 155–189. Cambridge Univ. Press, Cambridge, 1995.
- [12] Markus Neuhauser. Relative property (T) and related properties of wreath products. *Math. Z.*, 251(1):167–177, 2005.