



# Property FW and wreath products of groups: A simple approach using Schreier graphs

Paul-Henry Leemann<sup>a,1</sup>, Grégoire Schneeberger<sup>b,\*,2</sup>

<sup>a</sup> Department of Pure Mathematics, Xi'an Jiaotong-Liverpool University, 111 Ren'ai Road, 215123 Suzhou, P. R. China

<sup>b</sup> Section de mathématiques, Université de Genève, rue du Conseil Général 7-9, 1205 Genève, Switzerland

Received 21 June 2021; received in revised form 7 December 2021; accepted 12 July 2022

## Abstract

The group property FW stands in-between the celebrated Kazhdan's property (T) and Serre's property FA. Among many characterizations, it might be defined, for finitely generated groups, as having all Schreier graphs one-ended.

It follows from the work of Y. Cornulier that a finitely generated wreath product  $G \wr_X H$  has property FW if and only if both  $G$  and  $H$  have property FW and  $X$  is finite. The aim of this paper is to give an elementary, direct and explicit proof of this fact using Schreier graphs.

© 2022 The Author(s). Published by Elsevier GmbH. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

MSC: primary 20E22; secondary 20F65; 05C25

Keywords: Wreath products; Property FW; Schreier graphs; Ends

## 1. Introduction

Property FW is a group property that is (for discrete groups) a weakening of the celebrated Kazhdan's property (T). It was introduced by Barnhill and Chatterji in [1]. It is a fixed point property for actions on wall spaces or, equivalently, on CAT(0) cube

\* Corresponding author.

E-mail addresses: [PaulHenry.Leemann@xjtlu.edu.cn](mailto:PaulHenry.Leemann@xjtlu.edu.cn) (P.-H. Leemann), [gregoire.schneeberger@unige.ch](mailto:gregoire.schneeberger@unige.ch) (G. Schneeberger).

<sup>1</sup> Supported by grant 200021.188578 of the Swiss NSF.

<sup>2</sup> Supported by grants 200020-178828 and 200020-200400 of the Swiss NSF.

<https://doi.org/10.1016/j.exmath.2022.07.001>

0723-0869/© 2022 The Author(s). Published by Elsevier GmbH. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Please cite this article as: P.-H. Leemann and G. Schneeberger, Property FW and wreath products of groups: A simple approach using Schreier graphs, Expositiones Mathematicae (2022), <https://doi.org/10.1016/j.exmath.2022.07.001>.

complexes. Therefore it stands between property FH (fixed points on Hilbert spaces, equivalent to property (T) for discrete groups) and property FA (fixed points on *arbres*<sup>3</sup>) It is known that all these properties are different, see [4] for examples of groups that distinguish them.

When working with group properties, it is natural to ask if they are stable under “natural” group operations. One such operation, of great use in geometric group theory, is the wreath product, that generalizes the direct product of two groups, see Definition 2.4.

In the context of properties defined by fixed points of actions, the first result concerning wreath products is due to Cherix, Martin and Valette and later refined by Neuhauser and concerns property (T).

**Theorem 1.1** ([3,11]). *Let  $G, H$  be two discrete groups with  $G$  non-trivial and  $X$  a set on which  $H$  acts. The wreath product  $G \wr_X H$  has property (T) if and only if  $G$  and  $H$  have property (T) and  $X$  is finite.*

In [4] Cornuier studied property FW using *cardinal definite functions* and while not explicitly stated in [4], the following result can be extracted from his work.

**Theorem 1.2.** *Let  $G, H$  be two finitely generated groups with  $G$  non-trivial and  $X$  a set on which  $H$  acts with finitely many orbits. The wreath product  $G \wr_X H$  has property FW if and only if  $G$  and  $H$  have property FW and  $X$  is finite.*

The aim of this note is to give an explicit and elementary proof of this fact using a characterization of property FW via the number of ends of Schreier graphs, see Section 2.1 for the relevant definitions.

At this point, the curious reader might have two questions. First, is it possible to extend Theorem 1.2 beyond the realms of finitely generated groups and of actions with finitely many orbits? And secondly, is there a link between Theorems 1.1 and 1.2? In both cases, the answer is yes.

This is the subject of the more technical paper [9], which gives a unified proof of Theorems 1.1 and 1.2 as well as of similar results for the Bergman’s property and more.


*Organization of the paper.* The next section contains all the definitions as well as some examples, while Section 3 is devoted to the proof of Theorem 1.2 and some related results.

## 2. Definitions and examples


This section contains all the definitions, some standard but useful preliminary results as well as some examples.

### 2.1. Ends of Schreier graphs and property FW

In what follows, we will always assume that generating sets of groups are *symmetric*, that is we will look at  $S \subset G$  such that  $s \in S$  if and only if  $s^{-1} \in S$ . Our graphs will be undirected and we will sometimes identify a graph with its vertex set.



**Fig. 1.** Fragments of two Cayley graphs of  $\mathbb{Z}$  (2 ends), for the standard generating set  $\{\pm 1\}$  and for the generating set  $\{\pm 2, \pm 3\}$ .



**Fig. 2.** Fragments of the Cayley graphs of  $\mathbb{Z}^2$  (1 end) on the left and of  $F_2$  (infinitely many ends) on the right; with standard generating sets.

Let  $G$  be a group with symmetric generating set  $S$ , and let  $X$  be a non-empty set endowed with a left action  $G \curvearrowright X$ . The corresponding (left) *orbital graph*  $\text{Sch}_O(G, X; S)$  is the graph with vertex set  $X$  and with an edge between  $x$  and  $y$  for every  $s$  in  $S$  such that  $s.x = y$ .


**Definition 2.1.** Let  $G$  be a group,  $H$  a subgroup of  $G$  and  $S$  a symmetric generating set. The (left) *Schreier graph*  $\text{Sch}(G, H; S)$  is the graph with vertices the left cosets  $gH = \{gh \mid h \in H\}$  and for every  $\{s, s^{-1}\} \subset S$  an edge between  $gH$  and  $sgH$ .

As the notation suggests, these two definitions are two faces of the same coin. More precisely, Schreier graphs of  $G = \langle S \rangle$  are exactly the orbital graphs for transitive actions (equivalently: the connected components of orbital graphs) of  $G = \langle S \rangle$ . The correspondence is given by  $H \mapsto G/H$  and  $x_0 \in X \mapsto \text{Stab}_G(x_0)$ , where  $x_0$  is an arbitrary element of  $X$ .

Schreier graphs are generalizations of the well-known *Cayley graphs*, with  $\text{Cayl}(G; S) = \text{Sch}(G, \{1\}; S)$ , see Figs. 1 and 2 for some examples.

If  $S$  and  $T$  are two generating sets of  $G$ , the graphs  $\text{Sch}(G, H; S)$  and  $\text{Sch}(G, H; T)$  does not need to be isomorphic. However, if both  $S$  and  $T$  are finite, then  $\text{Sch}(G, H; S)$  and  $\text{Sch}(G, H; T)$  are quasi-isometric, see [7, IV.B.21.iii] for a proof and Fig. 1 for an

<sup>3</sup> *Arbres* is the french word for trees.



**Fig. 3.** A fragment of a Schreier graph (with 4 ends) of the free group  $F_2 = \langle x^{\pm 1}, y^{\pm 1} \rangle$  for the subgroup  $H = \{x^2, y^n xy^{-n}, xy^n xy^{-n} x^{-1} \mid n \in \mathbb{Z} \setminus \{0\}\}$ .

example. The only fact we will use about quasi-isometries is that they preserve “large-scale properties” of the graph, as for example the number of ends. Observe that the requirement that both  $S$  and  $T$  are finite is crucial for the existence of a quasi-isometry between the corresponding Schreier graphs. Indeed, for every group  $\text{Cayl}(G; G \setminus \{1\}) = \text{Sch}(G, \{1\}; G \setminus \{1\})$  is always a complete graph on  $|G|$  vertices, in particular  $\text{Cayl}(\mathbb{Z}; \{1\})$  and  $\text{Cayl}(\mathbb{Z}; \mathbb{Z} \setminus \{0\})$  are not quasi-isometric.

Let  $\Gamma$  be a graph and  $K$  a finite subset of vertices. The graph  $\Gamma \setminus K$  is the subgraph of  $\Gamma$  obtained by deleting all vertices in  $K$  and all edges containing them. This graph is not necessarily connected.

**Definition 2.2.** Let  $\Gamma$  be a graph. The *number of ends* of  $\Gamma$  is the supremum, taken over all finite  $K$ , of the number of infinite connected components of  $\Gamma \setminus K$ .

There exist other characterizations of the number of ends in graphs, see [5] and the references therein, but Definition 2.2 is the one that best suits our purpose. A locally finite graph (i.e. such that every vertex has finite degree) is finite if and only if it has 0 end.

An important fact about the number of ends of a graph, is that it is an invariant of quasi-isometry, see [2]. In particular, if  $G$  is a finitely generated group it is possible to speak about the number of ends of the Schreier graph  $\text{Sch}(G, H; S)$  without specifying a particular finite generating set  $S$ . By a celebrated result of Hopf [8], the number of ends of a Cayley graph of a finitely generated group can only be 0, 1, 2 or infinite (in which case it is uncountable), see Figs. 1 and 2 for some examples. On the other hand, Schreier graphs may have any number of ends in  $\mathbb{N} \cup \{\infty\}$ , see Fig. 3 for an example of a graph with 4 ends. In fact, every regular graph of even degree is isomorphic to a Schreier graph, [6,10].

We are now finally able to introduce property FW. Instead of giving the original definition in terms of actions on wall spaces, we will use an equivalent one for finitely generated groups, which essentially follows from [12], see [4] for a direct proof.

**Definition 2.3.** A finitely generated group  $G$  has *property FW* if all its Schreier graphs have at most one end.

It directly follows from the definition that all finite groups have property FW, but that  $\mathbb{Z}$  does not have it. In fact, if  $G$  is a finitely generated group with a homomorphism onto  $\mathbb{Z}$ , then it does not have FW. Indeed, in this case  $G \cong H \rtimes \mathbb{Z}$  for some  $H$  and the Schreier graph  $\text{Sch}(G, H; S)$  is isomorphic to a Cayley graph of  $\mathbb{Z} \cong G/H$  and hence has 2 ends.

Property FW admits many distinct characterizations that allow to define it for groups that are non-necessarily finitely generated and even for topological groups. We refer the reader to [4] for a survey of these characterizations.

## 2.2. Wreath products

Let  $X$  be a set and  $G$  a group. We view  $\bigoplus_X G$  as the set of functions from  $X$  to  $G$  with finite support:

$$\bigoplus_X G = \{\varphi: X \rightarrow G \mid \varphi(x) = 1 \text{ for all but finitely many } x\}.$$

This is naturally a group, where multiplication is taken pointwise.

If  $H$  is a group acting on  $X$ , then it naturally acts on  $\bigoplus_X G$  by  $(h.\varphi)(x) = \varphi(h^{-1}.x)$ . This leads to the following standard definition.

**Definition 2.4.** Let  $G$  and  $H$  be groups and  $X$  be a set on which  $H$  acts. The (retracted) wreath product  $G \wr_X H$  is the group  $(\bigoplus_X G) \rtimes H$ .

A prominent source of examples of wreath products is the ones of the form  $G \wr_H H$ , where  $H$  acts on itself by left multiplication. In particular, the group  $(\mathbb{Z}/2\mathbb{Z}) \wr_{\mathbb{Z}} \mathbb{Z}$  is well-known under the name of the *lamplighter group*. The direct product  $G \times H$  corresponds to wreath product over a singleton  $G \wr_{\{*\}} H$ .

Let  $S$  be a generating set of  $G$  and  $T$  a generating set of  $H$ . Let  $\{x_i\}_{i \in I}$  be a choice of a representative in each  $H$ -orbit. Finally, let  $\delta_x^s$  be the element of  $\bigoplus_X G$  defined by  $\delta_x^s(x) = s$  and  $\delta_x^s(y) = 1_G$  if  $y \neq x$  and let  $\mathbf{1}$  be the constant function with value  $1_G$ . It is a direct computation that

$$\{(\delta_{x_i}^s, 1_H) \mid s \in S, i \in I\} \cup \{(\mathbf{1}, t) \mid t \in T\}$$

is a generating set for  $G \wr_X H$ .

On the other hand, if  $\{(\varphi_i, h_i) \mid i \in I\}$  is a generating set of  $G \wr_X H$ , then  $\{h_i \mid i \in I\}$  is a generating set of  $H$  while  $\{\varphi_i(x) \mid i \in I, x \in X\}$  is a generating set of  $G$ . Observe that since the  $\varphi_i$  take only finitely many values, if  $I$  is finite, so is  $\{\varphi_i(x) \mid i \in I, x \in X\}$ . We hence recover the following characterization of the finite generation of  $G \wr_X H$ .

**Lemma 2.5.** The group  $G \wr_X H$  is finitely generated if and only if both  $G$  and  $H$  are finitely generated and  $H$  acts on  $X$  with finitely many orbits.

**Proof.** If  $G \wr_X H$  is finitely generated, so is its abelianization  $(G \wr_X H)^{\text{ab}} \cong (\bigoplus_{X/H} G^{\text{ab}}) \times H^{\text{ab}}$ , which implies that the orbit set  $X/H$  is finite. The other implications directly follow from the above discussion on generating sets.  $\square$

Using the above lemma, we could reformulate Theorem 1.2 in the following way: Let  $G, H$  be two groups with  $G$  non-trivial and  $X$  a set on which  $H$  acts. Suppose that all three of  $G, H$  and  $G \wr_X H$  are finitely generated. Then the wreath product  $G \wr_X H$  has property FW if and only if  $G$  and  $H$  have property FW and  $X$  is finite. While this formulation is formally equivalent to Theorem 1.2, it hints the fact that the

finite generation hypothesis on  $G$ ,  $H$  and  $G \wr_X H$  are not necessary, but artefacts of using Schreier graphs in the proof. Indeed, the result remains true without these hypothesis, see [4, Propositions 5.B.3 & 5.B.4] and [9, Theorem A].

### 3. Proof of the main result

This section is devoted to the proof of [Theorem 1.2](#). This proof is split into two parts: [Lemma 3.2](#) and its [Corollary 3.4](#), and [Lemma 3.5](#).

We begin by an easy result on quotients.

**Lemma 3.1.** *Let  $G$  be a finitely generated group and  $H$  a quotient of  $G$ . If  $G$  has FW, then so does  $H$ .*

**Proof.** First, remark that if  $G$  is generated by a finite symmetric set  $S$ , the group  $H$  is generated by the projection of  $S$  that we will also denote by  $S$ . Moreover, any generating set of  $H$  can be obtained in such a way.

Let  $K$  be any subgroup of  $H$  and  $L$  its preimage in  $G$ . As  $G/L \cong H/K$ , forgetting about loops and multiedges, the Schreier graphs  $\text{Sch}(G, L; S)$  and  $\text{Sch}(H, K; S)$  are isomorphic. By assumption on  $G$ , the graph  $\text{Sch}(G, L; S)$  has at most one end. As adding loops or doubling edges do not change the number of ends,  $\text{Sch}(H, K; S)$  has also at most one end, and therefore  $H$  has property FW.  $\square$

We also have the following lemma on semi-direct products:

**Lemma 3.2.** *Let  $N$  and  $H$  be two finitely generated groups and  $N \rtimes H$  a semi-direct product. Then*

1. *If  $N \rtimes H$  has property FW, then so does  $H$ ,*
2. *If both  $N$  and  $H$  have property FW, then  $N \rtimes H$  also has property FW.*

**Proof.** The first part follows from [Lemma 3.1](#).

For the second part, let  $S$ , respectively  $T$ , denote a finite generating set of  $N$ , respectively  $H$ . Then the group  $G := N \rtimes H$  is finitely generated by  $U = (S \times \{1\}) \cup (\{1\} \times T)$ .

Suppose that both  $N$  and  $H$  have property FW. We want to show that every Schreier graph of  $G$  has at most one end. If they are all finite, then there is nothing to prove (and  $G$  is finite). So let  $\Gamma$  be an infinite Schreier graph of  $G$  with respect to the generating set  $U$ . The groups  $N$  and  $H$  act on the vertices of  $\Gamma$  by restriction of the action of  $G$ . That is,  $n.x = (n, 1).x$  and  $h.x = (1, h).x$  for every vertex  $x$  of  $\Gamma$ . For each vertex  $x$  we define  $\Gamma_x^H$  (and respectively  $\Gamma_x^N$ ) as the Schreier graph obtained from the action of  $H$  (respectively  $N$ ) on the  $H$ -orbit (respectively  $N$ -orbit) of  $x$ . These are subgraphs of  $\Gamma$ . As  $N$  and  $H$  have property FW, the graphs  $\Gamma_x^H$  and  $\Gamma_x^N$  are either finite or one-ended; and we will prove that this implies that  $\Gamma$  has exactly one end.

Let  $K$  be a finite set of vertices of  $\Gamma$ . If  $x$  is in  $K$  and  $\Gamma_x^H$  is finite, add all vertices of  $\Gamma_x^H$  to  $K$ . By doing so for every  $x$  in  $K$ , we obtain a new finite set  $K' \supset K$  of vertices of  $\Gamma$ . We will show that  $\Gamma \setminus K'$  has only one infinite connected component. By definition




Fig. 4. The path between  $x$  and  $y$ .

of  $K'$ , if  $x$  is not in  $K'$ , then either  $\Gamma_x^H$  has one end or  $\Gamma_x^H$  does not contain vertices of  $K'$ .

Let  $x$  and  $y$  be two vertices, each of them lying in some infinite connected component of  $\Gamma \setminus K'$ . We will construct a path from  $x$  to  $y$  in  $\Gamma \setminus K'$  as a concatenation of three smaller paths, see Fig. 4, as follows. First, a path in  $\Gamma_x^H \setminus K'$  from  $x$  to some  $z$ , then a path in  $\Gamma_z^N \setminus K'$  from  $z$  to some  $z' \in (\Gamma_z^N \cap \Gamma_y^H) \setminus K'$ , and finally a path in  $\Gamma_y^H \setminus K'$  from  $z'$  to  $y$ . In order to finish the proof, it remains to exhibit elements  $z$  and  $z'$  and the three desired paths.

The action of  $G$  on  $\Gamma$  being transitive, there exists an element  $(n_0, h_0)$  of  $N \rtimes H$  such that  $(n_0, h_0).x = y$ . Since  $K'$  is finite, the set  $\Gamma_x^H \setminus K'$  is infinite. Moreover, there is infinitely many  $z$  in  $\Gamma_x^H \setminus K'$  such that either  $\Gamma_z^N$  is one-ended or  $\Gamma_z^N$  does not intersect  $K'$ . For such a  $z$  there exists  $h$  such that  $(1, h).x = z$ . Now, the vertex  $z' := (hh_0^{-1}.n_0, h).x$  is both equal to  $(hh_0^{-1}.n_0, 1)(1, h_0).x = (hh_0^{-1}.n_0, 1).z$  and to  $(1, hh_0^{-1})(n_0, h_0).x = (1, hh_0^{-1}).y$ . That is,  $z'$  is in  $\Gamma_z^N \cap \Gamma_y^H$ . A direct computation shows us that the map  $z \mapsto z'$  is injective:  $z'_1 = z'_2$  if and only if  $z_1 = z_2$ . Since  $K'$  is finite, there are only finitely many  $z'$  in  $K'$  and hence there are infinitely many  $z \in \Gamma_x^H$  such that both  $z$  and  $z'$  are not in  $K'$  and either  $\Gamma_z^N$  is one-ended or  $\Gamma_z^N$  does not intersect  $K'$ .

In order to finish the proof, observe that the three graphs  $\Gamma_x^H$ ,  $\Gamma_y^H$  and  $\Gamma_z^N$  are all either one-ended or do not intersect  $K'$ . Therefore, there is a path in  $\Gamma_x^H \setminus K'$  from  $x$  to  $z$  as desired, and similarly for the paths from  $z$  to  $z'$  and  $z'$  to  $y$ . We have just proved that for any finite  $K$  the graph  $\Gamma \setminus K$  has only one infinite connected component and therefore that  $\Gamma$  is one-ended.  $\square$

We have the following result on direct products that can be obtained as a corollary of Lemma 3.2. It is also possible to give a short proof of it using Lemma 3.1 and a direct argument; details are left to the reader.

**Corollary 3.3.** *Let  $G$  and  $H$  be two finitely generated groups. Then  $G \times H$  has property FW if and only if both  $G$  and  $H$  have property FW.*

By iterating Lemma 3.2, we obtain

**Corollary 3.4.** *Let  $G$  and  $H$  be two finitely generated groups and  $X$  a set on which  $H$  acts with finitely many orbits. Then,*

1. *If  $G \wr_X H$  has property FW, then so does  $H$ ,*
2. *If both  $G$  and  $H$  have property FW and  $X$  is finite, then  $G \wr_X H$  has property FW.*

The following Lemma finishes the proof of Theorem 1.2.

**Lemma 3.5.** *Let  $G$  and  $H$  be two finitely generated groups with  $G$  non-trivial and such that  $H$  acts on some set  $X$  with finitely many orbits. If  $G \wr_X H$  has property FW, then*

1.  *$G$  has property FW,*
2.  *$X$  is finite.*

**Proof.** Let us fix some finite generating sets  $S$  and  $T$  of  $G$  and  $H$  and let

$$U := \{(\delta_x^s, 1_H) \mid s \in S\} \cup \{(1, t) \mid t \in T\}$$

be the standard generating set of  $G \wr_X H$ .

*If  $G \wr_X H$  has property FW, then  $X$  is finite.* Since  $H$  acts on  $X$  with finitely many orbits, it is enough to show that each orbit is finite. So let  $X'$  be an orbit and  $x_0$  be an arbitrary vertex of  $X'$ . The group  $G$  acting on itself by left multiplications, we have the so-called *imprimitive action* of the wreath-product  $G \wr_X H$  on  $Y := G \times X'$ :

$$(\varphi, h).(g, x) := (\varphi(h.x)g, h.x).$$

Since both  $G \curvearrowright G$  and  $H \curvearrowright X'$  are transitive, the action  $G \wr_X H \curvearrowright Y$  is also transitive. Therefore, the orbital Schreier graph of  $G \wr_X H \curvearrowright Y$  is isomorphic to the Schreier graph  $\Gamma := \text{Sch}(G \wr_X H, \text{Stab}((1_G, x_0)), U)$ . We decompose this graph into leaves of the form  $Y_g = \{g\} \times X'$ . There are two types of edges in  $\Gamma$ , which are coming from the two sets of generators, see Fig. 5. The first ones, of the form  $(1, t)$ , give us on each leaf a copy of the orbital Schreier graph of  $H \curvearrowright X'$ . Indeed,

$$(1, t).(g, x) = (g, t.x).$$


The second ones, of the form  $(\delta_{x_0}^s, 1)$ , give us loops everywhere except on vertices of the form  $(g, x_0)$ . By direct computation, we see that the vertices  $(g, x_0)$  and  $(sg, x_0)$  connect the leaves  $Y_g$  and  $Y_{sg}$ ,

$$(\delta_{x_0}^s, 1).(g, x) = \begin{cases} (g, x) & \text{if } x \neq x_0, \\ (sg, x) & \text{if } x = x_0. \end{cases}$$

If we remove a vertex  $(g, x_0)$  we disconnect the leaf  $Y_g$  from the rest of  $\Gamma$ . Since  $\Gamma$  has at most one end and there is  $|G| \geq 2$  leaves, we deduce that all leaves  $Y_g$  are finite, and hence that  $X'$  itself is finite.

*The group  $G$  has property FW.* Let  $K$  be any subgroup of  $G$ . We will show that  $\text{Sch}(G, K, S)$  has at most one end. Let  $x_0$  be any point of  $X$  and  $X'$  be its orbit under





**Fig. 5.** The leaf structure of the orbital Schreier graph of  $G \wr_X H \curvearrowright Y$ . Plain edges correspond to generators of the form  $(1, t)$  while dotted edges correspond to generators of the form  $(\delta_{x_0}^s, 1)$ .

the action of  $H$ . We have the imprimitive action of  $G \wr_X H$  on  $G/K \times X$ , which restricts to an action on  $G/K \times X'$ :

$$(\varphi, h).(gK, x) = (\varphi(h.x)gK, h.x).$$

As above, the action is transitive and the orbital Schreier graph of this action is isomorphic to a Schreier graph  $\Gamma$  of  $G \wr_X H$ . We decompose this graph into leaves in the same way. Now observe that  $\text{Sch}(G, K, S)$  is isomorphic to the subgraph  $\Delta$  of  $\Gamma$  consisting of vertices  $\{(g, x_0) \mid g \in G\}$  and edges  $\{(\delta_{x_0}^s, 1) \mid s \in S\}$ . Due to the leaves structure of  $\Gamma$ , the number of ends of  $\Delta$  is bounded above by the number of ends of  $\Gamma$ , and hence is at most one. We conclude that  $\text{Sch}(G, K, S)$  too has at most one end.  $\square$

## Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

## Acknowledgements

The authors are thankful to A. Genevois, T. Nagnibeda and A. Valette for helpful comments on a previous version of this note, and to the Swiss National Science Foundation for its support. The author also thank the anonymous referee for their valuable comments.

## References

- [1] Angela Barnhill, Indira Chatterji, Property (T) versus property FW, *Enseign. Math.* (2) 54 (1–2) (2008) 16–18.
- [2] Stephen G. Brick, Quasi-isometries and ends of groups, *J. Pure Appl. Algebr.* 86 (1) (1993) 23–33.
- [3] Pierre-Alain Cherix, Florian Martin, Alain Valette, Spaces with measured walls, the Haagerup property and property (T), *Ergodic Theory Dynam. Syst.* 24 (6) (2004) 1895–1908.
- [4] Yves Cornuier, Group actions with commensurated subsets, wallings and cubings, 2013, arXiv e-prints, page arXiv:1302.5982.
- [5] Reinhard Diestel, Daniela Kühn, Graph-theoretical versus topological ends of graphs, *J. Combin. Theory Ser. B* 87 (1) (2003) 197–206.
- [6] Jonathan L. Gross, Every connected regular graph of even degree is a Schreier coset graph, *J. Combin. Theory Ser. B* 22 (3) (1977) 227–232.
- [7] Pierre de la Harpe, Topics in Geometric Group Theory, in: *Chicago Lectures in Mathematics*, University of Chicago Press, Chicago, IL, 2000.
- [8] Heinz Hopf, Enden offener Räume und unendliche diskontinuierliche Gruppen, *Comment. Math. Helv.* 16 (1944) 81–100.
- [9] Paul-Henry Leemann, Grégoire Schneeberger, Wreath products of groups acting with bounded orbits, to appear in *L'Enseignement Mathématique*.
- [10] Alexander Lubotzky, Cayley graphs: eigenvalues, expanders and random walks, in: *Surveys in combinatorics, 1995 (Stirling)*, in: *London Math. Soc. Lecture Note Ser.*, vol. 218, Cambridge Univ. Press, Cambridge, 1995, pp. 155–189.
- [11] Markus Neuhauser, Relative property (T) and related properties of wreath products, *Math. Z.* 251 (1) (2005) 167–177.
- [12] Michah Sageev, Ends of group pairs and non-positively curved cube complexes, *Proc. Lond. Math. Soc.* (3) 71 (3) (1995) 585–617.