

Confidence interval on SEIR predictions

10/04/2024

1 Proof of the method for computing confidence intervals

Assumption :

We suppose that the data of the pandemic observed follows the model h , of parameter $\theta^* \in \mathcal{R}^d$. Let $Y_i, i = 1, \dots, n$ be the number of hospitalized at each day. We suppose that : $Y_i = h_{\theta^*}(i) + \epsilon_i$, with $\epsilon_i \sim \mathcal{N}(0, \sigma^2)$, iid, and independent from all the other variables. The objective is to estimate θ^* . We use $\hat{\theta}$, the least square estimator of θ^* as an estimator of θ^* :

$$\hat{\theta} = \underset{\theta \in \mathbb{R}^d}{\operatorname{argmin}} \sum_{i=1}^n (Y_i - h_{\theta}(i))^2$$

Let :

$$Y = \begin{pmatrix} Y_1 \\ \vdots \\ Y_n \end{pmatrix}$$
$$h_{\theta} = \begin{pmatrix} h_{\theta}(1) \\ \vdots \\ h_{\theta}(n) \end{pmatrix}$$

We have :

$$\hat{\theta} = \underset{\theta \in \mathbb{R}^d}{\operatorname{argmin}} \|Y - h_{\theta}\|^2$$

Now, if θ is close enough to θ^* , we can write :

$$\forall i \in \{1, \dots, n\} : h_{\theta}(i) = h_{\theta^*}(i) + \nabla_{\theta} h_{\theta^*}(i)(\theta - \theta^*)$$

which leads to :

$$\hat{\theta} = \underset{\theta \in \mathbb{R}^d}{\operatorname{argmin}} \|Y - h_{\theta^*}(i) - \nabla_{\theta} h_{\theta^*}(i)(\theta - \theta^*)\|^2$$

Let us define :

$$\tilde{Y} = Y - h_{\theta^*}$$
$$\beta = \theta - \theta^*$$
$$\hat{\beta} = \theta - \hat{\theta}$$

and let us define the matrix $A \in \mathbb{R}^{n \times d}$ such that $\forall i \in \{1, \dots, n\}, \forall j \in \{1, \dots, d\}, A_{i,j} = \frac{dh_{\theta^*}}{d\theta_j}(i)$. The previous problem can be re-written as :

$$\hat{\beta} = \underset{\beta \in \mathbb{R}^d}{\operatorname{argmin}} \|\tilde{Y} - A\beta\|^2$$

This is a regression linear problem.

Let us solve this problem in the general case. Let $\tilde{Y}_i = A_i\beta^* + \epsilon'_i$, with $\epsilon'_i \sim \mathcal{N}(0, \sigma'^2)$. The solution of this problem is explicitly :

$$\hat{\beta} = (A^T A)^{-1} A^T \tilde{Y}$$

This least-square estimator is unbiased :

$$\mathbb{E}[\hat{\beta}] = \beta^*$$

$$\hat{\beta} = \left(\frac{1}{n} \sum_{i=1}^n A_i^T A_i \right)^{-1} \times \left(\frac{1}{n} \sum_{i=1}^n A_i^T \tilde{Y}_i \right)$$

Let us note :

$$\hat{D} = \frac{1}{n} \sum_{i=1}^n A_i^T A_i, \quad \text{and} \quad \hat{\delta} = \left(\frac{1}{n} \sum_{i=1}^n A_i^T \tilde{Y}_i \right)$$

We have :

$$\hat{\beta} = \hat{D}^{-1} \hat{\delta}$$

$$\hat{D} \xrightarrow{a.s} \mathbb{E}[A_i^T A_i]$$

$$\hat{\delta} \xrightarrow{a.s} \mathbb{E}[A_i^T \tilde{Y}_i]$$

$$\hat{\beta} = \hat{D}^{-1} \hat{\delta} \xrightarrow{a.s} D^{-1} \delta$$

as $\phi : A \rightarrow A^{-1}$ is continuous on $\mathcal{GL}_n(\mathbb{R})$.

Now, let us show that $\hat{\beta}$ is asymptotically normal :

$$\begin{aligned} \sqrt{n}(\hat{\beta} - \beta^*) &= \sqrt{n}(\hat{D}^{-1} \hat{\delta} - \beta^*) \\ &= \sqrt{n}(\hat{D}^{-1} \hat{\delta} - \hat{D}^{-1} \hat{D} \beta^*) \\ &= \sqrt{n} \hat{D}^{-1} (\hat{\delta} - \hat{D} \beta^*) \\ &= \sqrt{n} \hat{D}^{-1} \left(\frac{1}{n} \sum_{i=1}^n A_i^T \tilde{Y}_i - \frac{1}{n} \sum_{i=1}^n A_i^T A_i \beta^* \right) \\ &= \frac{\sqrt{n}}{n} \hat{D}^{-1} \left(\sum_{i=1}^n A_i^T (\tilde{Y}_i - A_i \beta^*) \right) \\ &= \frac{1}{\sqrt{n}} \hat{D}^{-1} \left(\sum_{i=1}^n A_i^T \epsilon_i \right) \end{aligned}$$

This line is made of two terms. Let's show that each one of them converges in law.

$$\begin{aligned} \frac{1}{\sqrt{n}} \left(\sum_{i=1}^n A_i^T \epsilon'_i \right) &= \sqrt{n} \left(\frac{1}{n} \sum_{i=1}^n A_i^T \epsilon'_i \right) \\ &= \sqrt{n} \left(\frac{1}{n} \sum_{i=1}^n A_i^T \epsilon'_i - 0 \right) \\ &\xrightarrow{a.s} \mathcal{N}(0, \text{Var}(A_i^T \epsilon_i)) \end{aligned}$$

Yet, as $\epsilon_i \perp A_i$ and $\mathbb{E}[A_i^T \epsilon'_i] = 0$, $\text{Var}(A_i^T \epsilon_i) = \mathbb{E}[A_i A_i^T \epsilon_i^2] = \mathbb{E}[A_i A_i^T] \sigma'^2$.
 Finally, $\frac{1}{\sqrt{n}} (\sum_{i=1}^n A_i^T \epsilon'_i) \xrightarrow{a.s.} \mathcal{N}(0, D \sigma'^2)$.

On the other hand, $\hat{D}^{-1} \xrightarrow{\mathcal{L}} D^{-1} = Cte$.

Finally, with Slutsky, we obtain that :

$$\begin{aligned} \sqrt{n}(\hat{\beta} - \beta^*) &\xrightarrow{\mathcal{L}} D^{-1} \mathcal{N}(0, D \sigma'^2) \\ &\xrightarrow{\mathcal{L}} \mathcal{N}(0, D^{-1} \sigma'^2) \\ &\xrightarrow{\mathcal{L}} \mathcal{N}(0, \sigma'^2 (A^T A)^{-1}) \end{aligned}$$

Let's get back to the first problem :

As $\beta^* = 0$ and $\hat{\beta} = \hat{\theta} - \theta^*$, we have :

$$\sqrt{n}(\hat{\theta} - \theta^*) \xrightarrow{\mathcal{L}} \mathcal{N}(0, \sigma^2 (A^T A)^{-1})$$

and,

$$\hat{\theta} \sim \mathcal{N}(\theta^*, \frac{\sigma^2}{n} (A^T A)^{-1})$$

As a first conclusion, we have that $\hat{\theta}$ is asymptotically normal.

Let Σ be the covariance matrix estimated from the computation of $\hat{\theta}$. In our case, $\Sigma = \frac{\sigma^2}{n} (A^T A)^{-1}$.

As $\hat{\theta}$ is asymptotically normal, we can apply the delta-method :

$$\begin{aligned} \sqrt{n}(\hat{\theta} - \theta^*) &\xrightarrow{\mathcal{L}} \mathcal{N}(0, \Sigma) \\ \sqrt{n}(h_{\hat{\theta}} - h_{\theta^*}) &\xrightarrow{\mathcal{L}} \mathcal{N}(0, \nabla_{\theta} h_{\theta}^T \Sigma \nabla_{\theta} h_{\theta}) \end{aligned}$$

And finally :

$$h_{\hat{\theta}} \rightarrow \mathcal{N}(h_{\theta^*}, \frac{1}{n} \nabla_{\theta} h_{\theta}^T \Sigma \nabla_{\theta} h_{\theta})$$

By estimating $\frac{1}{n} \Sigma$ from `curve_fit`, we can compute the confidence interval of the prediction with the quantiles of the normal distribution.