


Parameter estimation details

1	1	Clarkia life history	2
2	2	Creating the dataset	3
3	2.1	Seed bag experiment	5
4	2.2	Seedling and fruiting plant surveys in permanent plots	7
5	2.3	Surveys for fruiting plant density, fruits per plant, seeds per fruit	8
6	2.4	Additional assumptions	10
7	3	Model	11
8	3.1	Seed persistence, germination, and viability	12
9	3.2	Seedling survival to fruiting	13
10	3.3	Fecundity: fruits per plant and seeds per fruit	15
11	4	Computing vital rates	16

1 *Clarkia* life history

Clarkia xantiana is a winter annual. *C. xantiana* seeds germinate with late fall and winter rains; in the Kern Valley this is between October and late February or early March. Seedlings grow throughout the winter and spring, and surviving plants flower in late spring and early summer, April into early July. Pollinated fruits set seed in the early summer, June to July. Seeds of *C. xantiana* are produced in early summer, with fruits that dry out and gradually split open. Most seeds appear to be shed from fruits within 3-4 months after production, but can remain on the plant for up to a year. Seeds are small (< 1 mm in width) and have no structures to aid in aerial dispersal. Seeds can remain viable in the field and laboratory for at least 5 years, contrary to previous suggestions that seeds are unlikely to persist past a few years (Lewis 1962).

We represent the *Clarkia xantiana* as a life cycle graph (Figure 1A) that describes transitions from October of year t to October of year $t + 1$ in terms of underlying vital rates. The census period occurs when the entire population is seeds, and corresponds to the time at which seed bags are placed into the field (see below). Seeds are grouped into three stages: age 0 seeds, which were produced in the current year; age 1 seeds, which were produced in the previous year; age 2+ seeds, which were produced two or more years ago. Persistence of seeds in the seed bank is represented by transitions from



younger to older seeds. Production of new seeds is captured by transition to the age 0 seed state.

Transitions in the life cycle graph are the product of age-specific seed survival and germination, and aboveground seedling survival to fruiting, fruit production, and seeds per fruit. Seed-related rates are represented separately for age 0, 1, and 2+ seeds. Germination for each age class is given as g_1 , g_2 , and g_3 , respectively. Seed survival from seed production to the first October is given as s_0 , and survival from October to February is given as s_1 , s_3 , and s_5 for age 0, 1, and 2+ seeds, respectively. Survival from February to October is given as s_2 , s_4 , and s_6 for age 0, 1, and 2+ seeds, respectively. We assume that vital rates remain unchanged after age 2. We also we assume that all plants experience the same vital rates upon germination seed age at germination does not affect seedling survival to fruiting (σ), fruits per plant (F), or seeds per fruit (ϕ).

The life cycle graph (Figure 1A) corresponds to the annual projection matrix

$$\mathbf{A} = \begin{bmatrix} s_1 g_1 \sigma F \phi s_0 & s_3 g_2 \sigma F \phi s_0 & s_5 g_3 \sigma F \phi s_0 \\ s_1 (1 - g_1) s_2 & 0 & 0 \\ 0 & s_3 (1 - g_2) s_4 & s_5 (1 - g_3) s_6 \end{bmatrix} \quad (1)$$

that summarizes transitions between stages.

2 Creating the dataset

We used field surveys and experiments to assemble the demographic observations necessary to provide insight into the population dynamics of *Clarkia xantiana* (Table 1). Specifically, we sought to parameterize the structured population model for *Clarkia xantiana* populations represented by Equation 1.

Starting in 2006, Monica Geber and collaborators collected annual estimates for demo-

graphic data on the winter annual plant *Clarkia xantiana* ssp. *xantiana*. The data include annual estimates for survival of seedlings to fruiting adults, fruits per adult plant, and seeds per fruit. In addition the data on above-ground vital rates, experiments have been used to infer germination and seed survival in the seed bank. The demographic study has been used to test hypotheses about the geography of demography (Eckhart et al. 2011) and species distributions (Pironon et al. 2018).

We used an experiment to estimate transitions in the seed bank. The experiment has been previously described in Eckhart et al. 2011 and we reanalyze the data here. From 2006-2009, Geber and collaborators buried seeds in bags and periodically dug them up to count seedlings and intact, viable seeds. This data estimates transitions leading to germination or survival of seeds that are 1, 2, and 3 years old.

Starting in 2007, there are (1) estimates of fruits/plant and seeds/fruit that provide an estimate of seed input into a plot and (2) estimates of germinants the following year. For most plots, the number of seeds entering a plot in year $t - 1$ is much greater than the number of seedlings emerging in a plot in year t . However, this is not uniformly true, and there is also experimental data suggesting these seeds may survive in the seed bank for at least 10 years at some locations.

Key: experiments from 2006-2009 to estimate age-specific seed rates.

Field surveys in permanent plots to estimate seedling survival to fruiting.

Combine data from permanent plots and additional grids to estimate fruits per plant.

Collect fruits to estimate seeds per fruit.

TABLE 1: Summary of data sets used to estimate parameters.

Parameter data	Description	Data set	Time span
SEED VITAL RATES	—	—	—
Seed survival and germination	Seed bag burial	\mathbf{Y}_1	2006-2009
Seed viability	Viability trials	\mathbf{Y}_2	2006-2009
Seed survival and germination	Seed pots	\mathbf{Y}_3	2013-2019
SEEDLING SURVIVAL	—	—	—
Seedling survival to fruiting	Field surveys	\mathbf{Y}_4	2006-2019
FRUITS PER PLANT	—	—	—
Total fruit equivalents per plant	Field surveys	\mathbf{Y}_5	2006-2012
Undamaged and damaged fruits per plant	Field surveys	\mathbf{Y}_6	2013-2019
Total fruit equivalents per plant	Extra plots	\mathbf{Y}_7	2006-2012
Undamaged and damaged fruits per plant	Extra plots	\mathbf{Y}_8	2013-2019
SEEDS PER FRUIT	—	—	—
Seeds per undamaged fruit	Lab counts	\mathbf{Y}_9	2006-2019
Seeds per damaged fruit	Lab counts	\mathbf{Y}_{10}	2013-2019

2.1 Seed bag burial experiment

Note to self about how to turn this into Methods. The key pieces are Figure 2, columns 1, 3, and 6 of Table 2 and columns 1 & 3 in Table 3. Figure 2 illustrates the relationship between the seed bag burial experiment, viability trials, and survival/germination probabilities. Table 2 gives the mathematical relationship among the components of Figure 2. Table 3 translates the survival and germination components to parameters for the structured population model. I would to pare this down I would pare down Section ?? to describe the experiment and reduce the tables to the relevant parameters, likely combining them into 1 table (top section being the survival functions, bottom section being the model parameters). I would keep the description of the full model and the viability models. I would move any discussion of how the models were constructed and DAGs to a separate document. I would reduce the section on translating estimated parameters to model parameters to generally describe how this was done and move the rest to a separate document.

We used a field experiment in which we buried known numbers of seeds ($n = 100$) to estimate the probability of germination and survival for seeds of different ages (Figure 2A, gray panel).

Buried bags were unearthed two times each during their first, second, or third year. Bags that were dug up in a given year were only used to count intact seeds in that year; bags were removed from the field at the end of the year. The seed bag burial experiment was initiated in 3 consecutive years (3 rounds) at the 20 study populations. In round 1 (started in 2006), bags were dug up in year 1, 2, and 3. In round 2 (started in 2007), bags were dug up in year 1 and 2. In round 3 (started in 2008), bags were dug up in year 1. There were thus 3 sets of counts for age 1 seeds, 2 sets of counts for age 2 seeds, and 1 set of counts for age 3 seeds. In summary, during each experimental round we counted the number of seeds (y_{ijkm}) in the seed bags for up to 3 years. We obtained estimates from the i^{th} bag, from the j^{th} population, from the k^{th} year in at several time points indexed by m corresponding to the times at which bags were unearthed. These counts represent the number of seeds that remain in the soil seed bank. **UPDATE** We thus define a persistent seed as one that remains intact in the seed bags (i.e. was countable in the seed bags); seeds were lost through physical destruction, decay, or germination. In (Figure 2A), points represent the average seed numbers associated with hypothetical counts in seed bags across an experimental round. We conducted viability trials in each year we conducted seed burial experiments. At the end of each experimental year, bags were brought to the lab and intact seeds were tested in a two-stage viability trial (Figure 2B, gray panel). In the lab, we conducted germination trials and viability assays on subsets of the seeds from each bag to estimate the viability of the intact seeds. First, we placed up to 15 seeds from each bag on to moist filter paper in a disposable cup and observed the number of germinants over 10 days; we counted and removed germinants every 2 days. For each bag, we summed the number of seeds tested and germinating to obtain the number of trials ($n_{ijk}^{\text{test}_g}$) and successes (y_{ijk}^{germ}) summarizing the germination trials. After 10 days, all remaining ungerminated seeds (up to a total of 10 seeds) were sliced in half and individually placed into the wells of 96-well plates filled with a solution of tetrazolium

chloride, which stains viable tissue red. [Eckhart et al. 2011: not all ungerminated seeds were tested; most were] We covered the plates with foil. Each 96-well plate contained seed from at least one bag per population of a given seed-age class. Two or three tests of up to 15 seeds each were conducted for each bag. We checked and counted for viable seeds every 2 days for 10 days. For each bag, we summed the number of seeds tested and staining to obtain the number of trials ($n_{ijk}^{\text{test}_v}$) and successes (y_{ijk}^{viab}) summarizing the viability trials.

2.2 Seedling and fruiting plant surveys in permanent plots

To assess the survival of seedlings to fruiting plants, we use counts of seedlings and fruiting plants in 30 0.5 m² plots at 20 populations from 2006–2018 (?). Seedlings and fruiting plants were counted in February and June, respectively. We observed counts of seedlings (n_{ijk}) and fruiting plants (y_{ijk}) in the i^{th} plot, from the j^{th} population, from the k^{th} year. Seedlings and fruiting plants in each plot are counted by a single person at each visit.

Discuss measurement error due to undercounting While we assume that the data on seedlings is measured perfectly (i.e. we did not under- or over-count seedlings), there are at least two possible sources of error: (1) measurement error that arises because we failed to count seedlings that were present and (2) error that arises because seedlings germinated after we visited the population. Germination phenology varies from year to year and with the position of the population in the range. For now, I ignore data that involved undercounting by filtering out those rows in the dataset. We may want to develop a model that relates our estimate of seedlings to the true number of seedlings in a plot because we sometimes observe more fruiting plants than seedlings but I had trouble developing such a model for the time being.

We assume that the data on fruiting plants is measured perfectly (i.e. we did not under- or over-count) because plants stand out from the background vegetation in June. Our model estimates the proportion of seedlings that survive to become fruiting plants.




FIGURE 2: Summary of the seed bag burial experiments and viability trials. **Figure will be labeled as (A: seed bag trials , B: viability trials, C: germination probability, D: survival probability).** (A) The gray panel contains a graphical representation of the seed bag trials. Seeds were buried at the start of each experiment (100 seeds in month 0). Seed bags were unearthed and intact seeds (y_{\cdot}) and germinants ($y_{g,\cdot}$) counted. The graph below the panel shows a hypothetical survival function associated with persistence of seeds in the soil seed bank. (B) The gray panel contains a graphical representation of the viability trials. Seeds were tested in two rounds; germination trials were performed and then some or all of the ungerminated seeds were tested for viability. The graph below the panel shows hypothetical data from a series of viability trials and the interpolated, inferred viabilities at times when viability was unobserved. (C) Age-specific germination probability is summarized in three ways. (D) The graph shows the survival function for persistence of seeds in the soil seed bank (black line) and the estimated discrete survival probabilities for persistence and viability of seeds (orange points).

2.3 Surveys for fruiting plant density, fruits per plant, seeds per fruit

139 better address permanent vs. extra plot

140 To assess the number of fruits per plant for *C. xantiana*, we use data on counts of the number

141 of fruits per plant at 20 populations (?). At each population, we made two sets of counts.
 142 First, we counted the number of fruits per plant on all plants in the 0.5m² permanent plots.
 143 Second, we counted the number of fruits per plant on plants that we sampled haphazardly
 144 across the site using throws of a 0.5m² grid. We used these counts to estimate the number
 145 fruits produced per plant.
 146 From 2006–2012, we counted the number of undamaged fruits on a plant. We then took
 147 the damaged fruits on a plant and visually stacked them end to end to estimate how many
 148 additional undamaged fruits that was equivalent to (e.g. two half fruits corresponded to one
 149 undamaged fruit). We used this as our count (y_{ijk}^{TFE}) of total fruit equivalents on the i^{th}
 150 plant, from the j^{th} population, from the k^{th} year.
 151 From 2013–present, we counted and separately recorded the number of undamaged (y_{ijk}^{UF})
 152 and damaged (y_{ijk}^{DF}) fruits on a plant plant. We used these counts to estimate the number
 153 of fruits produced per plant.
 154 To assess the number of seeds per fruit for *C. xantiana*, we use data on counts of the seeds
 155 per fruit of fruits that were haphazardly collected in 20 populations (?). In the field, we
 156 collected fruits that were undamaged. In the lab, we broke open the fruits to count the
 157 number of seeds per fruit. For each population in each year, we attempted to obtain 20-30
 158 counts of seeds produced per undamaged fruit. The plants were outside permanent plots
 159 to avoid affecting seed input. We used these fruits to estimate the mean number of seeds
 160 produced by undamaged and damaged fruits.
 161 From 2006–present, we collected one undamaged fruit from each of 20-30 plants that were
 162 haphazardly collected across each population. We counted the number of seeds in the fruit
 163 (y_{ijk}^{US}), corresponding to the i^{th} undamaged fruit, from the j^{th} population, from the k^{th} year.
 164 From 2013–present, we additionally collected a damaged fruit from the same plant whenever
 165 available. We counted the number of seeds in the fruit (y_{ijk}^{DS}), corresponding to the i^{th}
 166 undamaged fruit, from the j^{th} population, from the k^{th} year.

2.4 Additional assumptions

167 Address the data exclusion issues under additional assumptions. This includes undercounting
 168 in the seedling survival to fruiting dataset. See figure and table for this. We assume that
 169 this is likely due to only visiting the plots once each year; we can exclude these rows from
 170 the dataset or keep them and assume high survival of 1.




FIGURE 3: Graphical summary of undercounting in the dataset on seedling survival to fruiting. Each panel summarizes the datasets on seedling survival to fruiting (Tables ?? and ??). The black lines correspond to total plots with data on seedling survival. The red lines correspond to the number of plots with fewer seedlings than fruiting plants in a plot (corresponding to undercounting).

3 Model

We use observational and experimental data from 20 populations of *Clarkia xantiana* to estimate transition probabilities across the life cycle. We fit multilevel models to obtain population-specific estimates for belowground vital rates, and model year- and population-specific estimates for aboveground vital rates. Because we were interested in describing the life histories of individual populations, we built separate models for each population. Our general approach was to include a population-level mean as the top hierarchy, and have the next level be year-specific means drawn from the population mean. Although we used different likelihoods for different observations (e.g. binomial for seed bag experiments; Poisson for counts of seed per fruit), for a single population our approach can be summarized by the following model structure

$$[\theta_j, \theta_0, \sigma_j^2, \varsigma^2 | y_{ij}] \propto [y_{ij} | \theta_j, \sigma_j^2][\theta_j | \theta_0, \varsigma^2][\theta_0][\sigma_j^2][\varsigma^2]. \quad (2)$$

The distribution of the observations y_{ij} is conditional on the year-specific parameters θ_j and σ_j^2 . In turn, the year-specific parameter θ_j is conditional on the population-specific parameters θ_0 and ς^2 . All parameters found only on the right of conditional statements are also given priors $(\theta_0, \sigma_j^2, \varsigma^2)$. In practice, we implemented this model by writing the upper levels of the model as Normal-Normal distributions; we thus drew year-specific means from a population-specific mean.

For each model, we include the expression for the posterior proportional to the joint distribution in [Appendix: Joint Posterior](#). Details about the choice of priors are give in [Appendix: Priors](#).

3.1 Seed persistence and germination

Describe full model at once; introduce model for germination then model for seed persistence
as the product of absence of germination and destruction which combines the germination
history and the deterministic survival function; then add the piece of estimating s_0 .




FIGURE 4: Directed acyclic graphs for the full model. Models for each set of data are encapsulated in a gray box; links among the datasets are shown by arrows that cross over between boxes. Solid arrows depict the relationships among random variables, and dashed arrows depict the deterministic relationships.

3.2 Viability

All data from viability trials is in the form of binomial trials: we have counts of seeds at the
start and end of an experimental window of time. All models have the same structure for
seeds in bag i in population j in experimental year k . If the number of seeds starting the trial
(trials) is n_{ijk} and the number of seeds at the end of the trial (successes) is y_{ijk} , we write a

197 model that has a population-level mean and year-level means drawn from the population-level
 198 distribution. Broadly, this is two-level hierarchical model with a population-level mean, and
 199 year-level means drawn from the population-level distribution. The probability of success
 200 for each bag is drawn from this year- and population-level distribution. The model uses a
 201 binomial likelihood.

3.3 Seedling survival to fruiting

202 Address missing years, low data years, pooling

We estimated survival across all populations taking into account both temporal and between-population variability with the following model. We write a model that has a population-level mean and year-level means drawn from the population-level distribution. The probability of success (seedling survival to fruiting) for each plot is drawn from this year- and population-level distribution. The model has a binomial likelihood, and thus thus has a similar structure as the model for data on seed survival:

$$\begin{aligned}
 [\alpha_S, \mu_S, \sigma_S, \mu_S^{\text{pop}}, \sigma_S^{\text{pop}} | \mathbf{y}^{\text{fruiting}}] &\propto \prod_{i=1}^I \prod_{j=1}^J \prod_{k=1}^K \text{binomial}(y_{ijk}^{\text{fruiting}} | n_{ijk}^{\text{seedling}}, \text{logit}^{-1}(\alpha_{S,ijk})) \\
 &\times \text{normal}(\alpha_{S,ijk} | \mu_{S,jk}, \sigma_{S,jk}) \\
 &\times \text{normal}(\mu_{S,jk} | \mu_{S,j}^{\text{pop}}, \sigma_{S,j}^{\text{pop}}) \\
 &\times \text{half-normal}(\sigma_{S,jk} | 0, 1) \\
 &\times \text{normal}(\mu_{S,j}^{\text{pop}} | 0, 1000) \text{half-normal}(\sigma_{S,j}^{\text{pop}} | 0, 1).
 \end{aligned} \tag{3}$$

203 We obtain the population- and population-by-year posterior distributions of seedling survival
 204 to fruiting (σ) by marginalization. We transform these posteriors to $[0, 1]$ by taking the
 205 inverse logit; this transforms the parameters into the probability of success.




FIGURE 5: Directed acyclic graphs for the model for seedling survival to fruiting.

3.4 Fecundity: fruits per plant and seeds per fruit

We estimated fruits per plant across all populations taking into account both temporal and between-population variability with the following model. We constructed models with population-level mean and year-level means drawn from the population-level distribution. We used a Poisson likelihood to model the count data of fruits per plant. We modeled sampling uncertainty with a lognormal, which draws the true number of fruits (the zs) from a distribution with the median (μ). Each combination of year and population is assigned its own sampling variance (σ^2). The general form of the model is the following:

$$\begin{aligned}
& [z, \mu_{\text{TF}}, \sigma_{\text{TF}}^2, (\sigma_{\text{TF}}^{\text{pop}})^2, \nu_{\text{TF}} | y^{\text{TF}}] \propto \\
& \prod_{j=1}^J \prod_{i=1}^{N_1} \prod_{k=1}^{K_1} \text{Poisson}(y_{ijk}^{\text{TF}} | z_{\text{TF},ijk}) \text{lognormal}(z_{\text{TF},ijk} | \log(\mu_{\text{TF},jk}), \sigma_{\text{TF},jk}^2) \\
& \times \text{lognormal}(\mu_{\text{TF},jk} | \log(g(\nu_{\text{TF},j}), (\sigma_{\text{TF},j}^{\text{pop}})^2)) \\
& \times \text{inverse gamma}(\sigma_{\text{TF},jk}^2 | 0.001, 0.001) \\
& \times \text{gamma}(\nu_{\text{TF},j} | 1, 1) \text{inverse gamma}((\sigma_{\text{TF},j}^{\text{pop}})^2 | 0.001, 0.001).
\end{aligned} \tag{4}$$

206 We obtain the population- and population-by-year posterior distributions of fruits per plant
 207 (F) by marginalization.

208 Estimating total fruit equivalents per plant required combining two types of data. First, I
 209 used counts of total fruit equivalents per plant from (2006-2012). I modeled these counts as
 210 observations (zs). Second, I used counts of undamaged and damaged fruits per plant from
 211 (2013-present). I separately modeled counts of each as observations to obtain estimates of
 212 undamaged and damaged fruits per plant.

We estimated seeds per fruit across all populations taking into account both temporal and between-population variability with the following model. We constructed models with population-level mean and year-level means drawn from the population-level distribution.




FIGURE 6: Directed acyclic graphs for the models for fecundity.

We used a Poisson likelihood to model the count data of seeds per fruit. We modeled sampling uncertainty with a lognormal, which draws the true number of seeds (the z s) from a distribution with the median (μ). Each combination of year and population is assigned its own sampling variance (σ^2). The general form of the model is the following:

$$\begin{aligned}
& [z, \mu_{\text{TF}}, \sigma_{\text{TF}}^2, (\sigma_{\text{TF}}^{\text{pop}})^2, \nu_{\text{TF}} | \mathbf{y}^{\text{TF}}] \propto \\
& \prod_{j=1}^J \prod_{i=1}^{N_1} \prod_{k=1}^{K_1} \text{Poisson}(y_{ijk}^{\text{TF}} | z_{\text{TF},ijk}) \text{ lognormal}(z_{\text{TF},ijk} | \log(\mu_{\text{TF},jk}), \sigma_{\text{TF},jk}^2) \\
& \times \text{ lognormal}(\mu_{\text{TF},jk} | \log(g(\nu_{\text{TF},j}), (\sigma_{\text{TF},j}^{\text{pop}})^2) \\
& \times \text{ inverse gamma}(\sigma_{\text{TF},jk}^2 | 0.001, 0.001) \\
& \times \text{ gamma}(\nu_{\text{TF},j} | 1, 1) \text{ inverse gamma}((\sigma_{\text{TF},j}^{\text{pop}})^2 | 0.001, 0.001).
\end{aligned} \tag{5}$$

213 We obtain the population- and population-by-year posterior distributions of seeds per fruit

214 (ϕ) by marginalization.

215 First, I used counts of seeds per undamaged fruit (2006-present). I modeled these counts
216 as observations (zs). Second, I used counts of seeds from damaged fruits per plant from
217 (2013-present). I separately modeled counts of each as observations to obtain estimates of
218 seeds per damaged fruit.

4 Computing vital rates

219 General summary should cover link of parameter estimates from observational and exper-
220 imental data to structured model parameters. Key issues to address are different kinds of
221 data collection in different years, as well as combining datasets to get a coherent estimate
222 for a parameter (correct support).

223 Use model checking and compare outputs to look at effect of accounting for viability in
224 belowground rates, and understand that certain years in aboveground rates had no data in
225 permanent plots and were thus more strongly pooled.

4.1 Belowground vital rates

226 Describe calculating persistence + persistence/viability. Then describe linking these to the
227 structured parameters

TABLE 2: Seed persistence and viability in the soil seed bank

	Persistence	Persistence & viability
Time (x_i)	$S(x_i)$	$S(x_i)$
Oct ₀	θ_0	$\phi_0 = \theta_0$
Jan _{1,total}	θ_1	$\phi_1 = \theta_1(\gamma_1 + (1 - \gamma_1)\nu_1^{1/3})$
Jan _{1,intact}	θ_2	$\phi_2 = \theta_2\nu_1^{1/3}$
Oct ₁	θ_3	$\phi_3 = \theta_3\nu_1$
Jan _{2,total}	θ_4	$\phi_4 = \theta_4(\gamma_2 + (1 - \gamma_2)\nu_1(\nu_2/\nu_1)^{1/3})$
Jan _{2,intact}	θ_5	$\phi_5 = \theta_5\nu_1(\nu_2/\nu_1)^{1/3}$
Oct ₂	θ_6	$\phi_6 = \theta_6\nu_2$
Jan _{3,total}	θ_7	$\phi_7 = \theta_7(\gamma_3 + (1 - \gamma_3)\nu_2(\nu_3/\nu_2)^{1/3})$
Jan _{3,intact}	θ_8	$\phi_8 = \theta_8\nu_2(\nu_3/\nu_2)^{1/3}$
Oct ₃	θ_9	$\phi_9 = \theta_9\nu_3$

The survival function for viable seeds (ϕ) is composed of estimates of persistence over time (θ), estimates of viability (ν), and estimates of germination conditional on persistence (γ). We used the survival function and germination probabilities to define the parameters in the structured population model for *Clarkia xantiana* ssp. *xantiana*. Table 3 defines the age-specific germination probabilities and survival probabilities for the Leslie matrix in Eckhart et al. 2011 in terms of the survival function (Table 2) and germination probabilities.

TABLE 3: Structured model parameters conditional on persistence and viability

Parameter	Probability
g_1	γ_1/ϕ_1
g_2	γ_2/ϕ_4
g_3	γ_3/ϕ_7
s_1	ϕ_1
s_2	ϕ_3/ϕ_2
s_3	ϕ_4/ϕ_3
s_4	ϕ_6/ϕ_5
s_5	ϕ_7/ϕ_6
s_6	ϕ_9/ϕ_8

Figure 2C illustrates the relationship among the pieces discussed here. Estimates of germination from the seed bag experiment correspond to the probability of germination conditional on persistence (e.g. γ_1). Multiplying these estimates by the probability of persistence in

239 January before germination gives the unconditional probability of germination (e.g. $\theta_1 \times \gamma_1$).
240 Finally, the probability of germination conditional on viability is estimated by incorporating
241 loss of viability into the survival function (e.g. γ_1/ϕ_1).
242 We tested the viability of seeds in October, and were thus able to estimate the proportion
243 of viable seeds (Figure 2B; filled points). We further inferred the viability of intact seeds
244 in January by assuming that seeds lost viability at a constant rate (exponential decay).
245 Further, we interpolated between estimates by assuming that viability changed at a constant
246 rate between years, and that all seeds were viable at the start of the experiment (Figure 2B;
247 open points).

4.2 Aboveground vital rates

248 Describe using the posteriors to calculate ratio of and get TFE equivalents in later years.