

Spectrum-Dependent Learning Curves in Kernel Regression and Wide Neural Networks

Blake Bordelon, Abdulkadir Canatar, Cengiz Pehlevan

Notation

Notation

- We have training data $\{\mathbf{x}_i, y_i\}_{i=1}^p \in \mathcal{X} \times \mathbb{R}$

Notation

- We have training data $\{\mathbf{x}_i, y_i\}_{i=1}^p \in \mathcal{X} \times \mathbb{R}$
- We consider some function space \mathcal{F}

Notation

- We have training data $\{\mathbf{x}_i, y_i\}_{i=1}^p \in \mathcal{X} \times \mathbb{R}$
- We consider some function space \mathcal{F}
- Assume that the data are labeled by some $f^* \in \mathcal{F}$:

$$y_i = f^*(\mathbf{x}_i)$$

Notation

- We have training data $\{\mathbf{x}_i, y_i\}_{i=1}^p \in \mathcal{X} \times \mathbb{R}$
- We consider some function space \mathcal{F}
- Assume that the data are labeled by some $f^* \in \mathcal{F}$:

$$y_i = f^*(\mathbf{x}_i)$$

- Data follows some distribution $\mathbf{x} \sim p$

Reproducing Kernel Hilbert Space

Reproducing Kernel Hilbert Space

- Choose \mathcal{F} to be a Reproducing Kernel Hilbert Space (RKHS), denote $\mathcal{H} = \mathcal{F}$

Reproducing Kernel Hilbert Space

- Choose \mathcal{F} to be a Reproducing Kernel Hilbert Space (RKHS), denote $\mathcal{H} = \mathcal{F}$
- Hilbert space \implies Banach space (also called complete) and inner product $\langle \cdot, \cdot \rangle_{\mathcal{H}}$

Every Cauchy sequence converges



Reproducing Kernel Hilbert Space

- Choose \mathcal{F} to be a Reproducing Kernel Hilbert Space (RKHS), denote $\mathcal{H} = \mathcal{F}$
- Hilbert space \implies Banach space (also called complete) and inner product $\langle \cdot, \cdot \rangle_{\mathcal{H}}$
- Being RKHS means:
 $\exists!$ $K(\cdot, \cdot)$ symmetric, positive semi-definite fulfilling the reproducing property:

Every Cauchy sequence converges



$$\forall f \in \mathcal{F} : f(\mathbf{x}) = \langle K(\mathbf{x}, \cdot), f \rangle_{\mathcal{H}}$$

Reproducing Kernel Hilbert Space

- Choose \mathcal{F} to be a Reproducing Kernel Hilbert Space (RKHS), denote $\mathcal{H} = \mathcal{F}$
- Hilbert space \implies Banach space (also called complete) and inner product $\langle \cdot, \cdot \rangle_{\mathcal{H}}$
- Being RKHS means:
 $\exists!$ $K(\cdot, \cdot)$ symmetric, positive semi-definite fulfilling the reproducing property:

$$\forall f \in \mathcal{F} : f(\mathbf{x}) = \langle K(\mathbf{x}, \cdot), f \rangle_{\mathcal{H}}$$

- Equivalent to requiring that the evaluation functional $L_{\mathbf{x}}(f) = f(\mathbf{x})$ is continuous (which in turn is equivalent to boundedness of the linear functional):

$$\forall \mathbf{x} \quad |L_{\mathbf{x}}(f)| \leq M \|f\|_{\mathcal{H}}$$

Kernel Ridge(less) Regression

Kernel Ridge(less) Regression

- $\min_{f \in \mathcal{H}} \sum_{i=1}^p (f(\mathbf{x}_i) - y_i)^2 + \lambda \|f\|_{\mathcal{H}}^2$

Kernel Ridge(less) Regression

- $\min_{f \in \mathcal{H}} \sum_{i=1}^p (f(\mathbf{x}_i) - y_i)^2 + \lambda \|f\|_{\mathcal{H}}^2$
- Define the subspace $\mathbb{L} = \left\{ \sum_{i=1}^p \alpha_i K(\mathbf{x}_i, \cdot) : \boldsymbol{\alpha} \in \mathbb{R}^p \right\}$ and decompose any $f \in \mathcal{H}$ as $f = f_{\mathbb{L}} + f_{\perp}$ with f_{\perp} orthogonal to \mathbb{L}

Kernel Ridge(less) Regression

- $\min_{f \in \mathcal{H}} \sum_{i=1}^p (f(\mathbf{x}_i) - y_i)^2 + \lambda \|f\|_{\mathcal{H}}^2$
- Define the subspace $\mathbb{L} = \left\{ \sum_{i=1}^p \alpha_i K(\mathbf{x}_i, \cdot) : \alpha \in \mathbb{R}^p \right\}$ and decompose any $f \in \mathcal{H}$ as $f = f_{\mathbb{L}} + f_{\perp}$ with f_{\perp} orthogonal to \mathbb{L}
- **RKHS Magic:** $f(\mathbf{x}_i) = \langle f_{\mathbb{L}} + f_{\perp}, K(\cdot, \mathbf{x}_i) \rangle_{\mathcal{H}} = f_{\mathbb{L}}(\mathbf{x}_i)$

For ridgeless case ($\lambda = 0$), infinite dimensional problem reduced to finding the best p -dimensional parameter α

Kernel Ridge(less) Regression

- $\min_{f \in \mathcal{H}} \sum_{i=1}^p (f(\mathbf{x}_i) - y_i)^2 + \lambda \|f\|_{\mathcal{H}}^2$
- Define the subspace $\mathbb{L} = \left\{ \sum_{i=1}^p \alpha_i K(\mathbf{x}_i, \cdot) : \alpha \in \mathbb{R}^p \right\}$ and decompose any $f \in \mathcal{H}$ as $f = f_{\mathbb{L}} + f_{\perp}$ with f_{\perp} orthogonal to \mathbb{L}
- **RKHS Magic:** $f(\mathbf{x}_i) = \langle f_{\mathbb{L}} + f_{\perp}, K(\cdot, \mathbf{x}_i) \rangle_{\mathcal{H}} = f_{\mathbb{L}}(\mathbf{x}_i)$

For ridgeless case ($\lambda = 0$), infinite dimensional problem reduced to finding the best p -dimensional parameter α

- $f(\mathbf{x}) = \mathbf{y}^T (\mathbf{K}(\mathbf{X}, \mathbf{X}) + \lambda \mathbb{1})^{-1} \mathbf{K}(\mathbf{x}, \mathbf{X})$

Kernel Ridge(less) Regression

- $\min_{f \in \mathcal{H}} \sum_{i=1}^p (f(\mathbf{x}_i) - y_i)^2 + \lambda \|f\|_{\mathcal{H}}^2$
- Define the subspace $\mathbb{L} = \left\{ \sum_{i=1}^p \alpha_i K(\mathbf{x}_i, \cdot) : \alpha \in \mathbb{R}^p \right\}$ and decompose any $f \in \mathcal{H}$ as $f = f_{\mathbb{L}} + f_{\perp}$ with f_{\perp} orthogonal to \mathbb{L}
- **RKHS Magic:** $f(\mathbf{x}_i) = \langle f_{\mathbb{L}} + f_{\perp}, K(\cdot, \mathbf{x}_i) \rangle_{\mathcal{H}} = f_{\mathbb{L}}(\mathbf{x}_i)$

For ridgeless case ($\lambda = 0$), infinite dimensional problem reduced to finding the best p -dimensional parameter α

- $f(\mathbf{x}) = \mathbf{y}^T (\mathbf{K}(\mathbf{X}, \mathbf{X}) + \lambda \mathbb{1})^{-1} \mathbf{K}(\mathbf{x}, \mathbf{X})$
- **Remember:** $f(\mathbf{x}) = \mathbf{y}^T \Theta(\mathbf{X}, \mathbf{X})^{-1} \Theta(\mathbf{x}, \mathbf{X})$

 Predictive function of NTK

Uniform Convergence vs Average Case Analysis

Uniform Convergence vs Average Case Analysis

- Entire statistical learning theory based on a **worst case** analysis:

$$\mathbb{P}_{\mathbf{X} \sim p} \left(\sup_{h \in \mathcal{H}} |L_{\text{train}}(h) - L_{\text{gen}}(h)| \right) \geq 1 - \delta$$

Uniform Convergence vs Average Case Analysis

- Entire statistical learning theory based on a **worst case** analysis:

$$\mathbb{P}_{\mathbf{x} \sim p} \left(\sup_{h \in \mathcal{H}} |L_{\text{train}}(h) - L_{\text{gen}}(h)| \right) \geq 1 - \delta$$

- The goal here is to provide an expression for the **average case** analysis of the generalization error. This is very common in statistical mechanics. We will use some of its tools here.

Uniform Convergence vs Average Case Analysis

- Entire statistical learning theory based on a **worst case** analysis:

$$\mathbb{P}_{\mathbf{X} \sim p} \left(\sup_{h \in \mathcal{H}} |L_{\text{train}}(h) - L_{\text{gen}}(h)| \right) \geq 1 - \delta$$

- The goal here is to provide an expression for the **average case** analysis of the generalization error. This is very common in statistical mechanics. We will use some of its tools here.
- Sometimes, expectations will be denoted in physics-fashion:

$$\langle f(\mathbf{X}) \rangle_{\mathbf{X}} = \mathbb{E}_{\mathbf{X} \sim p}[f(\mathbf{X})]$$

Generalization Errors

Generalization Errors

- Denote by $f_{\mathbf{K}}(\mathbf{x}; \mathbf{X}, f^*)$ the function learnt with kernel regression

Generalization Errors

- Denote by $f_K(\mathbf{x}; \mathbf{X}, f^*)$ the function learnt with kernel regression
- $E_g(\mathbf{X}, f^*) = \mathbb{E}_{\mathbf{x} \sim p} \left[(f_K(\mathbf{x}; \mathbf{X}, f^*) - f^*(\mathbf{x}))^2 \right]$

Generalization Errors

- Denote by $f_K(\mathbf{x}; \mathbf{X}, f^*)$ the function learnt with kernel regression
- $E_g(\mathbf{X}, f^*) = \mathbb{E}_{\mathbf{x} \sim p} \left[(f_K(\mathbf{x}; \mathbf{X}, f^*) - f^*(\mathbf{x}))^2 \right]$
- $E_g = \langle E_g(\mathbf{X}, f^*) \rangle_{\mathbf{X}, f^*}$ (Integrate over data and teacher)

Generalization Errors

- Denote by $f_K(\mathbf{x}; \mathbf{X}, f^*)$ the function learnt with kernel regression
- $E_g(\mathbf{X}, f^*) = \mathbb{E}_{\mathbf{x} \sim p} \left[(f_K(\mathbf{x}; \mathbf{X}, f^*) - f^*(\mathbf{x}))^2 \right]$
- $E_g = \langle E_g(\mathbf{X}, f^*) \rangle_{\mathbf{X}, f^*}$ (Integrate over data and teacher)
- **Observe:** Fixed teacher f^* is included in this analysis by setting $p_{f^*} \sim \delta_{f^*}$

Generalization Errors

- Denote by $f_{\mathbf{K}}(\mathbf{x}; \mathbf{X}, f^*)$ the function learnt with kernel regression
- $E_g(\mathbf{X}, f^*) = \mathbb{E}_{\mathbf{x} \sim p} \left[(f_{\mathbf{K}}(\mathbf{x}; \mathbf{X}, f^*) - f^*(\mathbf{x}))^2 \right]$
- $E_g = \langle E_g(\mathbf{X}, f^*) \rangle_{\mathbf{X}, f^*}$ (Integrate over data and teacher)
- **Observe:** Fixed teacher f^* is included in this analysis by setting $p_{f^*} \sim \delta_{f^*}$
- **Goal:** Calculate E_g for any kernel \mathbf{K} and any teacher distribution

Mercer Decomposition

Mercer Decomposition

- Due to Mercer, we can decompose any kernel into a (possibly) infinite eigenbasis:

$$K(\mathbf{x}, \mathbf{x}') = \sum_{i=1}^M \lambda_i \phi_i(\mathbf{x}) \phi_i(\mathbf{x}') = \sum_{i=1}^M \psi_i(\mathbf{x}) \psi_i(\mathbf{x}') = \psi(\mathbf{x})^T \psi(\mathbf{x}')$$

Handwritten notes:
An arrow points from the equation $\psi_i(\mathbf{x}) = \sqrt{\lambda_i} \phi_i(\mathbf{x})$ to the $\psi_i(\mathbf{x})$ term in the sum.
Another arrow points from the text "Feature representation of sample \mathbf{x} " to the $\psi(\mathbf{x})^T$ term.

where ϕ_i fulfills $\int_{\mathcal{X}} K(\mathbf{x}, \mathbf{y}) \phi_i(\mathbf{y}) p(\mathbf{y}) d\mathbf{y} = \lambda_i \phi_i(\mathbf{x})$

Mercer Decomposition

- Due to Mercer, we can decompose any kernel into a (possibly) infinite eigenbasis:

$$K(\mathbf{x}, \mathbf{x}') = \sum_{i=1}^M \lambda_i \phi_i(\mathbf{x}) \phi_i(\mathbf{x}') = \sum_{i=1}^M \psi_i(\mathbf{x}) \psi_i(\mathbf{x}') = \psi(\mathbf{x})^T \psi(\mathbf{x}')$$

$\psi_i(\mathbf{x}) = \sqrt{\lambda_i} \phi_i(\mathbf{x})$

Feature representation of sample \mathbf{x}

where ϕ_i fulfills $\int_{\mathcal{X}} K(\mathbf{x}, \mathbf{y}) \phi_i(\mathbf{y}) p(\mathbf{y}) d\mathbf{y} = \lambda_i \phi_i(\mathbf{x})$

- $\{\psi_i(\cdot)\}_{i=1}^M$ form a basis of \mathcal{H}

Mercer Decomposition

- Due to Mercer, we can decompose any kernel into a (possibly) infinite eigenbasis:

$$K(\mathbf{x}, \mathbf{x}') = \sum_{i=1}^M \lambda_i \phi_i(\mathbf{x}) \phi_i(\mathbf{x}') = \sum_{i=1}^M \psi_i(\mathbf{x}) \psi_i(\mathbf{x}') = \psi(\mathbf{x})^T \psi(\mathbf{x}')$$

Handwritten notes:
A red arrow points from the equation $\psi_i(\mathbf{x}) = \sqrt{\lambda_i} \phi_i(\mathbf{x})$ to the $\psi_i(\mathbf{x})$ term in the sum.
Another red arrow points from the text "Feature representation of sample \mathbf{x} " to the $\psi(\mathbf{x})^T$ term.

where ϕ_i fulfills $\int_{\mathcal{X}} K(\mathbf{x}, \mathbf{y}) \phi_i(\mathbf{y}) p(\mathbf{y}) d\mathbf{y} = \lambda_i \phi_i(\mathbf{x})$

- $\{\psi_i(\cdot)\}_{i=1}^M$ form a basis of \mathcal{H}
- Write $f^*(\mathbf{x}) = \sum_{i=1}^M \bar{w}_i \psi_i(\mathbf{x})$ and $f(\mathbf{x}) = \sum_{i=1}^M w_i \psi_i(\mathbf{x})$

Kernel Regression in Feature Space

Kernel Regression in Feature Space

- Define the feature matrix

$$\mathbf{\Psi} = \begin{bmatrix} \text{---} & \psi(\mathbf{x}_1) & \text{---} \\ & \vdots & \\ \text{---} & \psi(\mathbf{x}_p) & \text{---} \end{bmatrix} = \begin{bmatrix} \psi_1(\mathbf{x}_1) & \dots & \psi_M(\mathbf{x}_1) \\ \vdots & & \vdots \\ \psi_1(\mathbf{x}_p) & \dots & \psi_M(\mathbf{x}_p) \end{bmatrix} \in \mathbb{R}^{M \times p}$$

Kernel Regression in Feature Space

- Define the feature matrix

$$\mathbf{\Psi} = \begin{bmatrix} \text{---} & \psi(\mathbf{x}_1) & \text{---} \\ & \vdots & \\ \text{---} & \psi(\mathbf{x}_p) & \text{---} \end{bmatrix} = \begin{bmatrix} \psi_1(\mathbf{x}_1) & \dots & \psi_M(\mathbf{x}_1) \\ \vdots & & \vdots \\ \psi_1(\mathbf{x}_p) & \dots & \psi_M(\mathbf{x}_p) \end{bmatrix} \in \mathbb{R}^{M \times p}$$

- Rewrite the kernel as $K(\mathbf{X}, \mathbf{X}) = \mathbf{\Psi}\mathbf{\Psi}^T$

Kernel Regression in Feature Space

- Define the feature matrix

$$\Psi = \begin{bmatrix} \text{---} & \psi(\mathbf{x}_1) & \text{---} \\ & \vdots & \\ \text{---} & \psi(\mathbf{x}_p) & \text{---} \end{bmatrix} = \begin{bmatrix} \psi_1(\mathbf{x}_1) & \dots & \psi_M(\mathbf{x}_1) \\ \vdots & & \vdots \\ \psi_1(\mathbf{x}_p) & \dots & \psi_M(\mathbf{x}_p) \end{bmatrix} \in \mathbb{R}^{M \times p}$$

- Rewrite the kernel as $K(\mathbf{X}, \mathbf{X}) = \Psi\Psi^T$
- $\min_{\mathbf{w} \in \mathbb{R}^M} \|\Psi^T \mathbf{w} - \mathbf{y}\|_2^2 + \lambda \|\mathbf{w}\|_2^2$

Kernel Regression in Feature Space

- Define the feature matrix

$$\Psi = \begin{bmatrix} \text{---} & \psi(\mathbf{x}_1) & \text{---} \\ & \vdots & \\ \text{---} & \psi(\mathbf{x}_p) & \text{---} \end{bmatrix} = \begin{bmatrix} \psi_1(\mathbf{x}_1) & \dots & \psi_M(\mathbf{x}_1) \\ \vdots & & \vdots \\ \psi_1(\mathbf{x}_p) & \dots & \psi_M(\mathbf{x}_p) \end{bmatrix} \in \mathbb{R}^{M \times p}$$

- Rewrite the kernel as $K(\mathbf{X}, \mathbf{X}) = \Psi\Psi^T$
- $\min_{\mathbf{w} \in \mathbb{R}^M} \|\Psi^T \mathbf{w} - \mathbf{y}\|_2^2 + \lambda \|\mathbf{w}\|_2^2$
- $\mathbf{w} = (\Psi\Psi^T + \lambda \mathbf{1})^{-1} \Psi \mathbf{y} \in \mathbb{R}^M$

Decomposition of Generalization Error

Decomposition of Generalization Error

Lemma 1:

We have the following decomposition of the generalization error:

$$E_g = \sum_{i=1}^M E_i$$

where the modes are given by $E_i = \lambda_i \langle (w_i - \bar{w}_i)^2 \rangle_{\mathbf{x}, \bar{\mathbf{w}}}$

Decomposition of Generalization Error

Lemma 1:

We have the following decomposition of the generalization error:

$$E_g = \sum_{i=1}^M E_i$$

where the modes are given by $E_i = \lambda_i \langle (w_i - \bar{w}_i)^2 \rangle_{\mathbf{x}, \bar{\mathbf{w}}}$

This provides in some sense a **spectral decomposition** of the generalization error as each term is weighted by the corresponding eigenvalue of the kernel.

Derivation

Derivation

$$E_g = \langle (f(\mathbf{x}) - f^*(\mathbf{x}))^2 \rangle_{\mathbf{x}, \mathbf{X}, f^*}$$

Derivation

$$E_g = \langle (f(\mathbf{x}) - f^*(\mathbf{x}))^2 \rangle_{\mathbf{x}, \mathbf{X}, f^*} = \left\langle \left(\sum_{i=1}^M (w_i - \bar{w}_i) \psi_i(\mathbf{x}) \right)^2 \right\rangle_{\mathbf{x}, \mathbf{X}, f^*}$$

Derivation

$$\begin{aligned} E_g &= \langle (f(\mathbf{x}) - f^*(\mathbf{x}))^2 \rangle_{\mathbf{x}, \mathbf{X}, f^*} = \left\langle \left(\sum_{i=1}^M (w_i - \bar{w}_i) \psi_i(\mathbf{x}) \right)^2 \right\rangle_{\mathbf{x}, \mathbf{X}, f^*} \\ &= \left\langle \left(\sum_{i,j=1}^M (w_i - \bar{w}_i)(w_j - \bar{w}_j) \psi_i(\mathbf{x}) \psi_j(\mathbf{x}) \right)^2 \right\rangle_{\mathbf{x}, \mathbf{X}, f^*} \end{aligned}$$

Derivation

$$\begin{aligned} E_g &= \langle (f(\mathbf{x}) - f^*(\mathbf{x}))^2 \rangle_{\mathbf{x}, \mathbf{X}, f^*} = \left\langle \left(\sum_{i=1}^M (w_i - \bar{w}_i) \psi_i(\mathbf{x}) \right)^2 \right\rangle_{\mathbf{x}, \mathbf{X}, f^*} \\ &= \left\langle \left(\sum_{i,j=1}^M (w_i - \bar{w}_i)(w_j - \bar{w}_j) \psi_i(\mathbf{x}) \psi_j(\mathbf{x}) \right)^2 \right\rangle_{\mathbf{x}, \mathbf{X}, f^*} \\ &= \sum_{i,j=1}^M \left\langle (w_i - \bar{w}_i)(w_j - \bar{w}_j) \langle \psi_i(\mathbf{x}) \psi_j(\mathbf{x}) \rangle_{\mathbf{x}} \right\rangle_{\mathbf{X}, f^*} \end{aligned}$$

Derivation

$$\begin{aligned}E_g &= \langle (f(\mathbf{x}) - f^*(\mathbf{x}))^2 \rangle_{\mathbf{x}, \mathbf{X}, f^*} = \left\langle \left(\sum_{i=1}^M (w_i - \bar{w}_i) \psi_i(\mathbf{x}) \right)^2 \right\rangle_{\mathbf{x}, \mathbf{X}, f^*} \\&= \left\langle \left(\sum_{i,j=1}^M (w_i - \bar{w}_i)(w_j - \bar{w}_j) \psi_i(\mathbf{x}) \psi_j(\mathbf{x}) \right)^2 \right\rangle_{\mathbf{x}, \mathbf{X}, f^*} \\&= \sum_{i,j=1}^M \left\langle (w_i - \bar{w}_i)(w_j - \bar{w}_j) \langle \psi_i(\mathbf{x}) \psi_j(\mathbf{x}) \rangle_{\mathbf{x}} \right\rangle_{\mathbf{X}, f^*}\end{aligned}$$

Now $\langle \psi_i(\mathbf{x}) \psi_j(\mathbf{x}) \rangle_{\mathbf{x}} = \int_{\mathcal{X}} \psi_i(\mathbf{x}) \psi_j(\mathbf{x}) p(\mathbf{x}) d\mathbf{x} = \mathbb{1}_{\{i=j\}} \lambda_i$

Derivation

$$\begin{aligned}E_g &= \langle (f(\mathbf{x}) - f^*(\mathbf{x}))^2 \rangle_{\mathbf{x}, \mathbf{X}, f^*} = \left\langle \left(\sum_{i=1}^M (w_i - \bar{w}_i) \psi_i(\mathbf{x}) \right)^2 \right\rangle_{\mathbf{x}, \mathbf{X}, f^*} \\&= \left\langle \left(\sum_{i,j=1}^M (w_i - \bar{w}_i)(w_j - \bar{w}_j) \psi_i(\mathbf{x}) \psi_j(\mathbf{x}) \right)^2 \right\rangle_{\mathbf{x}, \mathbf{X}, f^*} \\&= \sum_{i,j=1}^M \left\langle (w_i - \bar{w}_i)(w_j - \bar{w}_j) \langle \psi_i(\mathbf{x}) \psi_j(\mathbf{x}) \rangle_{\mathbf{x}} \right\rangle_{\mathbf{X}, f^*}\end{aligned}$$

Now $\langle \psi_i(\mathbf{x}) \psi_j(\mathbf{x}) \rangle_{\mathbf{x}} = \int_{\mathcal{X}} \psi_i(\mathbf{x}) \psi_j(\mathbf{x}) p(\mathbf{x}) d\mathbf{x} = \mathbb{1}_{\{i=j\}} \lambda_i$

$$E_g = \sum_{i=1}^M \lambda_i \left\langle (w_i - \bar{w}_i)^2 \right\rangle_{\mathbf{X}, f^*} = \sum_{i=1}^M E_i$$



Theorem 2:

Theorem 2:

We have the following expression for the generalization error under the minimizer \mathbf{w} :

$$E_g = \text{tr}(\mathbf{D} \langle \mathbf{G}^2 \rangle_{\mathbf{x}})$$

Theorem 2:

We have the following expression for the generalization error under the minimizer \mathbf{w} :

$$E_g = \text{tr}(\mathbf{D} \langle \mathbf{G}^2 \rangle_{\mathbf{x}})$$

where we define the matrices

- $\mathbf{G} = \left(\frac{1}{\lambda} \Phi \Phi^T + \Lambda^{-1} \right)^{-1}$
- $\Phi = \Lambda^{-\frac{1}{2}} \Psi$
- $\mathbf{D} = \Lambda^{-\frac{1}{2}} \langle \bar{\mathbf{w}} \bar{\mathbf{w}}^T \rangle_{\bar{\mathbf{w}}} \Lambda^{-\frac{1}{2}}$

$\Lambda = \text{diag}(\lambda_1, \dots, \lambda_M)$



Theorem 2:

We have the following expression for the generalization error under the minimizer \mathbf{w} :

$$E_g = \text{tr}(\mathbf{D} \langle \mathbf{G}^2 \rangle_{\mathbf{x}})$$

where we define the matrices

- $\mathbf{G} = \left(\frac{1}{\lambda} \Phi \Phi^T + \Lambda^{-1} \right)^{-1}$
 - $\Phi = \Lambda^{-\frac{1}{2}} \Psi$
 - $\mathbf{D} = \Lambda^{-\frac{1}{2}} \langle \bar{\mathbf{w}} \bar{\mathbf{w}}^T \rangle_{\bar{\mathbf{w}}} \Lambda^{-\frac{1}{2}}$
- $\Lambda = \text{diag}(\lambda_1, \dots, \lambda_M)$

Observe that:

Theorem 2:

We have the following expression for the generalization error under the minimizer \mathbf{w} :

$$E_g = \text{tr}(\mathbf{D} \langle \mathbf{G}^2 \rangle_{\mathbf{x}})$$

where we define the matrices

- $\mathbf{G} = \left(\frac{1}{\lambda} \Phi \Phi^T + \Lambda^{-1} \right)^{-1}$
 - $\Phi = \Lambda^{-\frac{1}{2}} \Psi$
 - $\mathbf{D} = \Lambda^{-\frac{1}{2}} \langle \bar{\mathbf{w}} \bar{\mathbf{w}}^T \rangle_{\bar{\mathbf{w}}} \Lambda^{-\frac{1}{2}}$
- $\Lambda = \text{diag}(\lambda_1, \dots, \lambda_M)$

Observe that:

- Nice separation of teacher-dependence (\mathbf{D}) and data-dependence (\mathbf{G}).

Theorem 2:

We have the following expression for the generalization error under the minimizer \mathbf{w} :

$$E_g = \text{tr}(\mathbf{D} \langle \mathbf{G}^2 \rangle_{\mathbf{x}})$$

where we define the matrices

- $\mathbf{G} = \left(\frac{1}{\lambda} \Phi \Phi^T + \Lambda^{-1}\right)^{-1}$
 - $\Phi = \Lambda^{-\frac{1}{2}} \Psi$
 - $\mathbf{D} = \Lambda^{-\frac{1}{2}} \langle \bar{\mathbf{w}} \bar{\mathbf{w}}^T \rangle_{\bar{\mathbf{w}}} \Lambda^{-\frac{1}{2}}$
- $\Lambda = \text{diag}(\lambda_1, \dots, \lambda_M)$

Observe that:

- Nice separation of teacher-dependence (\mathbf{D}) and data-dependence (\mathbf{G}).
- Calculating $\langle \bar{\mathbf{w}} \bar{\mathbf{w}}^T \rangle_{\bar{\mathbf{w}}}$ is relatively easy (if the distribution is centered, this is just the covariance matrix)

Theorem 2:

We have the following expression for the generalization error under the minimizer \mathbf{w} :

$$E_g = \text{tr}(\mathbf{D} \langle \mathbf{G}^2 \rangle_{\mathbf{x}})$$

where we define the matrices

- $\mathbf{G} = \left(\frac{1}{\lambda} \Phi \Phi^T + \Lambda^{-1} \right)^{-1}$
 - $\Phi = \Lambda^{-\frac{1}{2}} \Psi$
 - $\mathbf{D} = \Lambda^{-\frac{1}{2}} \langle \bar{\mathbf{w}} \bar{\mathbf{w}}^T \rangle_{\bar{\mathbf{w}}} \Lambda^{-\frac{1}{2}}$
- $\Lambda = \text{diag}(\lambda_1, \dots, \lambda_M)$

Observe that:

- Nice separation of teacher-dependence (\mathbf{D}) and data-dependence (\mathbf{G}).
- Calculating $\langle \bar{\mathbf{w}} \bar{\mathbf{w}}^T \rangle_{\bar{\mathbf{w}}}$ is relatively easy (if the distribution is centered, this is just the covariance matrix)
- Calculating $\langle \mathbf{G}^2 \rangle_{\mathbf{x}}$ is difficult


Derivation

Derivation

$$E_g(\mathbf{X}, f^*) = \left\langle (f(\mathbf{x}) - f^*(\mathbf{x}))^2 \right\rangle_{\mathbf{x}} = (\mathbf{w} - \bar{\mathbf{w}})^T \Lambda (\mathbf{w} - \bar{\mathbf{w}})$$

same steps as in Lemma

Derivation

$$E_g(\mathbf{X}, f^*) = \left\langle (f(\mathbf{x}) - f^*(\mathbf{x}))^2 \right\rangle_{\mathbf{x}} = (\mathbf{w} - \bar{\mathbf{w}})^T \Lambda (\mathbf{w} - \bar{\mathbf{w}})$$


same steps as in Lemma

Let's recall the minimal solution (using $\mathbf{y} = f^*(\mathbf{x}) = \Psi^T \bar{\mathbf{w}}$)

Derivation


$$E_g(\mathbf{X}, f^*) = \left\langle (f(\mathbf{x}) - f^*(\mathbf{x}))^2 \right\rangle_{\mathbf{x}} = (\mathbf{w} - \bar{\mathbf{w}})^T \Lambda (\mathbf{w} - \bar{\mathbf{w}})$$

same steps as in Lemma

Let's recall the minimal solution (using $\mathbf{y} = f^*(\mathbf{x}) = \Psi^T \bar{\mathbf{w}}$)

$$\mathbf{w} = \left(\Psi \Psi^T + \lambda \mathbf{1} \right)^{-1} \Psi \Psi^T \bar{\mathbf{w}} = \bar{\mathbf{w}} - \lambda \left(\Psi \Psi^T + \lambda \mathbf{1} \right)^{-1} \bar{\mathbf{w}}$$

Derivation



$$E_g(\mathbf{X}, f^*) = \left\langle (f(\mathbf{x}) - f^*(\mathbf{x}))^2 \right\rangle_{\mathbf{x}} = (\mathbf{w} - \bar{\mathbf{w}})^T \Lambda (\mathbf{w} - \bar{\mathbf{w}})$$

Let's recall the minimal solution (using $\mathbf{y} = f^*(\mathbf{x}) = \Psi^T \bar{\mathbf{w}}$)

$$\mathbf{w} = \left(\Psi \Psi^T + \lambda \mathbf{1} \right)^{-1} \Psi \Psi^T \bar{\mathbf{w}} = \bar{\mathbf{w}} - \lambda \left(\Psi \Psi^T + \lambda \mathbf{1} \right)^{-1} \bar{\mathbf{w}}$$

Then we get that

Derivation

$$E_g(\mathbf{X}, f^*) = \left\langle (f(\mathbf{x}) - f^*(\mathbf{x}))^2 \right\rangle_{\mathbf{x}} = (\mathbf{w} - \bar{\mathbf{w}})^T \mathbf{\Lambda} (\mathbf{w} - \bar{\mathbf{w}})$$


same steps as in Lemma

Let's recall the minimal solution (using $\mathbf{y} = f^*(\mathbf{x}) = \Psi^T \bar{\mathbf{w}}$)

$$\mathbf{w} = \left(\Psi \Psi^T + \lambda \mathbf{1} \right)^{-1} \Psi \Psi^T \bar{\mathbf{w}} = \bar{\mathbf{w}} - \lambda \left(\Psi \Psi^T + \lambda \mathbf{1} \right)^{-1} \bar{\mathbf{w}}$$

Then we get that

$$E_g(\mathbf{X}, f^*) = \lambda^2 \bar{\mathbf{w}}^T \left(\Psi \Psi^T + \lambda \mathbf{1} \right)^{-1} \mathbf{\Lambda} \left(\Psi \Psi^T + \lambda \mathbf{1} \right)^{-1} \bar{\mathbf{w}}$$

Derivation

same steps as in Lemma

$$E_g(\mathbf{X}, f^*) = \left\langle (f(\mathbf{x}) - f^*(\mathbf{x}))^2 \right\rangle_{\mathbf{x}} = (\mathbf{w} - \bar{\mathbf{w}})^T \Lambda (\mathbf{w} - \bar{\mathbf{w}})$$


Let's recall the minimal solution (using $\mathbf{y} = f^*(\mathbf{x}) = \Psi^T \bar{\mathbf{w}}$)

$$\mathbf{w} = \left(\Psi \Psi^T + \lambda \mathbb{1} \right)^{-1} \Psi \Psi^T \bar{\mathbf{w}} = \bar{\mathbf{w}} - \lambda \left(\Psi \Psi^T + \lambda \mathbb{1} \right)^{-1} \bar{\mathbf{w}}$$

Then we get that

$$\begin{aligned} E_g(\mathbf{X}, f^*) &= \lambda^2 \bar{\mathbf{w}}^T \left(\Psi \Psi^T + \lambda \mathbb{1} \right)^{-1} \Lambda \left(\Psi \Psi^T + \lambda \mathbb{1} \right)^{-1} \bar{\mathbf{w}} \\ &= \lambda^2 \bar{\mathbf{w}}^T \Lambda^{-\frac{1}{2}} \Lambda^{\frac{1}{2}} \left(\Psi \Psi^T + \lambda \mathbb{1} \right)^{-1} \Lambda^{\frac{1}{2}} \Lambda^{\frac{1}{2}} \left(\Psi \Psi^T + \lambda \mathbb{1} \right)^{-1} \Lambda^{-\frac{1}{2}} \Lambda^{\frac{1}{2}} \bar{\mathbf{w}} \end{aligned}$$

Derivation

$$E_g(\mathbf{X}, f^*) = \left\langle (f(\mathbf{x}) - f^*(\mathbf{x}))^2 \right\rangle_{\mathbf{x}} = (\mathbf{w} - \bar{\mathbf{w}})^T \Lambda (\mathbf{w} - \bar{\mathbf{w}})$$


Let's recall the minimal solution (using $\mathbf{y} = f^*(\mathbf{x}) = \Psi^T \bar{\mathbf{w}}$)

$$\mathbf{w} = \left(\Psi \Psi^T + \lambda \mathbb{1} \right)^{-1} \Psi \Psi^T \bar{\mathbf{w}} = \bar{\mathbf{w}} - \lambda \left(\Psi \Psi^T + \lambda \mathbb{1} \right)^{-1} \bar{\mathbf{w}}$$

Then we get that

$$\begin{aligned} E_g(\mathbf{X}, f^*) &= \lambda^2 \bar{\mathbf{w}}^T \left(\Psi \Psi^T + \lambda \mathbb{1} \right)^{-1} \Lambda \left(\Psi \Psi^T + \lambda \mathbb{1} \right)^{-1} \bar{\mathbf{w}} \\ &= \lambda^2 \bar{\mathbf{w}}^T \Lambda^{-\frac{1}{2}} \Lambda^{\frac{1}{2}} \left(\Psi \Psi^T + \lambda \mathbb{1} \right)^{-1} \Lambda^{\frac{1}{2}} \Lambda^{\frac{1}{2}} \left(\Psi \Psi^T + \lambda \mathbb{1} \right)^{-1} \Lambda^{-\frac{1}{2}} \Lambda^{\frac{1}{2}} \bar{\mathbf{w}} \\ &= \bar{\mathbf{w}}^T \Lambda^{-\frac{1}{2}} \mathbf{G} \mathbf{G} \Lambda^{-\frac{1}{2}} \bar{\mathbf{w}} = \text{tr} \left(\bar{\mathbf{w}}^T \Lambda^{-\frac{1}{2}} \mathbf{G} \mathbf{G} \Lambda^{-\frac{1}{2}} \bar{\mathbf{w}} \right) \end{aligned}$$

Derivation

$$E_g(\mathbf{X}, f^*) = \left\langle (f(\mathbf{x}) - f^*(\mathbf{x}))^2 \right\rangle_{\mathbf{x}} = (\mathbf{w} - \bar{\mathbf{w}})^T \Lambda (\mathbf{w} - \bar{\mathbf{w}})$$

same steps as in Lemma

Let's recall the minimal solution (using $\mathbf{y} = f^*(\mathbf{x}) = \Psi^T \bar{\mathbf{w}}$)

$$\mathbf{w} = \left(\Psi \Psi^T + \lambda \mathbb{1} \right)^{-1} \Psi \Psi^T \bar{\mathbf{w}} = \bar{\mathbf{w}} - \lambda \left(\Psi \Psi^T + \lambda \mathbb{1} \right)^{-1} \bar{\mathbf{w}}$$

Then we get that

$$\begin{aligned} E_g(\mathbf{X}, f^*) &= \lambda^2 \bar{\mathbf{w}}^T \left(\Psi \Psi^T + \lambda \mathbb{1} \right)^{-1} \Lambda \left(\Psi \Psi^T + \lambda \mathbb{1} \right)^{-1} \bar{\mathbf{w}} \\ &= \lambda^2 \bar{\mathbf{w}}^T \Lambda^{-\frac{1}{2}} \Lambda^{\frac{1}{2}} \left(\Psi \Psi^T + \lambda \mathbb{1} \right)^{-1} \Lambda^{\frac{1}{2}} \Lambda^{\frac{1}{2}} \left(\Psi \Psi^T + \lambda \mathbb{1} \right)^{-1} \Lambda^{-\frac{1}{2}} \Lambda^{\frac{1}{2}} \bar{\mathbf{w}} \\ &= \bar{\mathbf{w}}^T \Lambda^{-\frac{1}{2}} \mathbf{G} \mathbf{G} \Lambda^{-\frac{1}{2}} \bar{\mathbf{w}} = \text{tr} \left(\bar{\mathbf{w}}^T \Lambda^{-\frac{1}{2}} \mathbf{G} \mathbf{G} \Lambda^{-\frac{1}{2}} \bar{\mathbf{w}} \right) \\ &= \text{tr} \left(\Lambda^{-\frac{1}{2}} \bar{\mathbf{w}} \bar{\mathbf{w}}^T \Lambda^{-\frac{1}{2}} \mathbf{G} \mathbf{G} \right) = \text{tr} (\mathbf{D} \mathbf{G}^2) \end{aligned}$$



How to calculate $\langle G^2 \rangle_x$?

How to calculate $\langle G^2 \rangle_x$?

Two approaches outlined in the paper:

How to calculate $\langle G^2 \rangle_x$?

Two approaches outlined in the paper:

- Turn this expectation into a **PDE** by deriving a recursion in the number of samples. Viewing the sample size p as a **continuous parameter** will lead to the PDE formulation

How to calculate $\langle G^2 \rangle_x$?

Two approaches outlined in the paper:

- Turn this expectation into a **PDE** by deriving a recursion in the number of samples. Viewing the sample size p as a **continuous parameter** will lead to the PDE formulation
- Use the (infamous) **Replica trick** to brute-force calculate/approximate the expectation

How to calculate $\langle G^2 \rangle_x$?

Two approaches outlined in the paper:

- Turn this expectation into a **PDE** by deriving a recursion in the number of samples. Viewing the sample size p as a **continuous parameter** will lead to the PDE formulation
- Use the (infamous) **Replica trick** to brute-force calculate/approximate the expectation
- Both approximations **agree!**

PDE Formulation (I)

PDE Formulation (I)

- **Trick:** Introduce $\tilde{\mathbf{G}}(p, \nu) = \left(\frac{1}{\lambda} \mathbf{\Phi} \mathbf{\Phi}^T + \mathbf{\Lambda}^{-1} + \nu \mathbf{1} \right)^{-1}$, then

PDE Formulation (I)

- **Trick:** Introduce $\tilde{\mathbf{G}}(p, \nu) = \left(\frac{1}{\lambda}\mathbf{\Phi}\mathbf{\Phi}^T + \mathbf{\Lambda}^{-1} + \nu\mathbb{1}\right)^{-1}$, then

$$\langle \mathbf{G}^2 \rangle_{\mathbf{x}} = -\frac{\partial}{\partial \nu} \left\langle \tilde{\mathbf{G}}(p, \nu) \right\rangle_{\mathbf{x}} \Big|_{\nu=0}$$

PDE Formulation (I)

- **Trick:** Introduce $\tilde{\mathbf{G}}(p, v) = \left(\frac{1}{\lambda}\mathbf{\Phi}\mathbf{\Phi}^T + \mathbf{\Lambda}^{-1} + v\mathbf{1}\right)^{-1}$, then

$$\langle \mathbf{G}^2 \rangle_{\mathbf{x}} = -\frac{\partial}{\partial v} \left\langle \tilde{\mathbf{G}}(p, v) \right\rangle_{\mathbf{x}} \Big|_{v=0}$$

Just the basis matrix from Mercer

- **Recall:** $\mathbf{\Phi} = \mathbf{\Lambda}^{-\frac{1}{2}}\mathbf{\Psi} \in \mathbb{R}^{M \times p}$. Add a sample: $\tilde{\mathbf{\Phi}} \in \mathbb{R}^{M \times (p+1)}$

PDE Formulation (I)

- **Trick:** Introduce $\tilde{\mathbf{G}}(p, v) = (\frac{1}{\lambda} \Phi \Phi^T + \Lambda^{-1} + v \mathbb{1})^{-1}$, then

$$\langle \mathbf{G}^2 \rangle_{\mathbf{x}} = - \frac{\partial}{\partial v} \langle \tilde{\mathbf{G}}(p, v) \rangle_{\mathbf{x}} \Big|_{v=0}$$

Just the basis matrix from Mercer

- **Recall:** $\Phi = \Lambda^{-\frac{1}{2}} \Psi \in \mathbb{R}^{M \times p}$. Add a sample: $\tilde{\Phi} \in \mathbb{R}^{M \times (p+1)}$

$$\tilde{\Phi} \tilde{\Phi}^T = \begin{bmatrix} | & & | & & | \\ \phi_1 & \dots & \phi_p & & \phi_{p+1} \\ | & & | & & | \end{bmatrix} \begin{bmatrix} - & \phi_1^T & - \\ & \vdots & \\ - & \phi_p^T & - \\ - & \phi_{p+1}^T & - \end{bmatrix}$$

PDE Formulation (I)

- **Trick:** Introduce $\tilde{\mathbf{G}}(p, \nu) = \left(\frac{1}{\lambda} \Phi \Phi^T + \Lambda^{-1} + \nu \mathbb{1} \right)^{-1}$, then

$$\langle \mathbf{G}^2 \rangle_{\mathbf{x}} = - \frac{\partial}{\partial \nu} \left\langle \tilde{\mathbf{G}}(p, \nu) \right\rangle_{\mathbf{x}} \Big|_{\nu=0}$$

Just the basis matrix from Mercer

- **Recall:** $\Phi = \Lambda^{-\frac{1}{2}} \Psi \in \mathbb{R}^{M \times p}$. Add a sample: $\tilde{\Phi} \in \mathbb{R}^{M \times (p+1)}$

$$\begin{aligned} \tilde{\Phi} \tilde{\Phi}^T &= \begin{bmatrix} | & & | & | \\ \phi_1 & \dots & \phi_p & \phi_{p+1} \\ | & & | & | \end{bmatrix} \begin{bmatrix} - & \phi_1^T & - \\ - & \vdots & - \\ - & \phi_p^T & - \\ - & \phi_{p+1}^T & - \end{bmatrix} \\ &= \begin{bmatrix} \Phi & | \\ \phi_{p+1} \\ | \end{bmatrix} \begin{bmatrix} - & \Phi^T & - \\ - & \phi_{p+1}^T & - \end{bmatrix} \end{aligned}$$

PDE Formulation (I)

- **Trick:** Introduce $\tilde{\mathbf{G}}(p, \nu) = \left(\frac{1}{\lambda} \Phi \Phi^T + \Lambda^{-1} + \nu \mathbb{1} \right)^{-1}$, then

$$\langle \mathbf{G}^2 \rangle_{\mathbf{x}} = - \frac{\partial}{\partial \nu} \left\langle \tilde{\mathbf{G}}(p, \nu) \right\rangle_{\mathbf{x}} \Big|_{\nu=0}$$

Just the basis matrix from Mercer

- **Recall:** $\Phi = \Lambda^{-\frac{1}{2}} \Psi \in \mathbb{R}^{M \times p}$. Add a sample: $\tilde{\Phi} \in \mathbb{R}^{M \times (p+1)}$

$$\begin{aligned} \tilde{\Phi} \tilde{\Phi}^T &= \begin{bmatrix} | & & | & | \\ \phi_1 & \dots & \phi_p & \phi_{p+1} \\ | & & | & | \end{bmatrix} \begin{bmatrix} - & \phi_1^T & - \\ & \vdots & \\ - & \phi_p^T & - \\ - & \phi_{p+1}^T & - \end{bmatrix} \\ &= \begin{bmatrix} \Phi & | \\ \phi_{p+1} \\ | \end{bmatrix} \begin{bmatrix} - & \Phi^T & - \\ & \phi_{p+1}^T & - \end{bmatrix} = \Phi \Phi^T + \phi_{p+1} \phi_{p+1}^T \end{aligned}$$

PDE Formulation (II)

PDE Formulation (II)


- We can use the Sherman-Woodbury formula:

PDE Formulation (II)

- We can use the Sherman-Woodbury formula:

$$\tilde{\mathbf{G}}(p+1, \nu) = \left(\frac{1}{\lambda} \tilde{\mathbf{\Phi}} \tilde{\mathbf{\Phi}}^T + \mathbf{\Lambda}^{-1} + \nu \mathbb{1} \right)^{-1}$$

PDE Formulation (II)


$$(A + \underline{u}\underline{v}^T)^{-1} = A^{-1} + \frac{A^{-1}\underline{u}\underline{v}^T A^{-1}}{1 + \underline{v}^T A^{-1} \underline{u}}$$

- We can use the Sherman-Woodbury formula:

$$\begin{aligned}\tilde{\mathbf{G}}(p+1, v) &= \left(\frac{1}{\lambda} \tilde{\mathbf{\Phi}} \tilde{\mathbf{\Phi}}^T + \mathbf{\Lambda}^{-1} + v \mathbf{1} \right)^{-1} \\ &= \left(\frac{1}{\lambda} \mathbf{\Phi} \mathbf{\Phi}^T + \frac{1}{\lambda} \phi_{p+1} \phi_{p+1}^T + \mathbf{\Lambda}^{-1} + v \mathbf{1} \right)^{-1}\end{aligned}$$

PDE Formulation (II)

- We can use the Sherman-Woodbury formula:

$$\begin{aligned}\tilde{\mathbf{G}}(p+1, v) &= \left(\frac{1}{\lambda} \tilde{\mathbf{\Phi}} \tilde{\mathbf{\Phi}}^T + \mathbf{\Lambda}^{-1} + v \mathbf{1} \right)^{-1} \\ &= \left(\frac{1}{\lambda} \mathbf{\Phi} \mathbf{\Phi}^T + \frac{1}{\lambda} \phi_{p+1} \phi_{p+1}^T + \mathbf{\Lambda}^{-1} + v \mathbf{1} \right)^{-1} \\ &= \left(\tilde{\mathbf{G}}(p, v)^{-1} + \frac{1}{\lambda} \phi_{p+1} \phi_{p+1}^T \right)^{-1}\end{aligned}$$

PDE Formulation (II)

- We can use the Sherman-Woodbury formula:

$$\begin{aligned}\tilde{\mathbf{G}}(p+1, v) &= \left(\frac{1}{\lambda} \tilde{\mathbf{\Phi}} \tilde{\mathbf{\Phi}}^T + \mathbf{\Lambda}^{-1} + v \mathbf{1} \right)^{-1} \\ &= \left(\frac{1}{\lambda} \mathbf{\Phi} \mathbf{\Phi}^T + \frac{1}{\lambda} \phi_{p+1} \phi_{p+1}^T + \mathbf{\Lambda}^{-1} + v \mathbf{1} \right)^{-1} \\ &= \left(\tilde{\mathbf{G}}(p, v)^{-1} + \frac{1}{\lambda} \phi_{p+1} \phi_{p+1}^T \right)^{-1} \\ &= \tilde{\mathbf{G}}(p, v) - \frac{\tilde{\mathbf{G}}(p, v) \frac{1}{\lambda} \phi_{p+1} \phi_{p+1}^T \tilde{\mathbf{G}}(p, v)}{1 + \phi_{p+1}^T \tilde{\mathbf{G}}(p, v) \phi_{p+1}}\end{aligned}$$

PDE Formulation (II)

- We can use the Sherman-Woodbury formula:

$$\begin{aligned}\tilde{\mathbf{G}}(p+1, v) &= \left(\frac{1}{\lambda} \tilde{\mathbf{\Phi}} \tilde{\mathbf{\Phi}}^T + \mathbf{\Lambda}^{-1} + v \mathbf{1} \right)^{-1} \\ &= \left(\frac{1}{\lambda} \mathbf{\Phi} \mathbf{\Phi}^T + \frac{1}{\lambda} \phi_{p+1} \phi_{p+1}^T + \mathbf{\Lambda}^{-1} + v \mathbf{1} \right)^{-1} \\ &= \left(\tilde{\mathbf{G}}(p, v)^{-1} + \frac{1}{\lambda} \phi_{p+1} \phi_{p+1}^T \right)^{-1} \\ &= \tilde{\mathbf{G}}(p, v) - \frac{\tilde{\mathbf{G}}(p, v) \frac{1}{\lambda} \phi_{p+1} \phi_{p+1}^T \tilde{\mathbf{G}}(p, v)}{1 + \phi_{p+1}^T \tilde{\mathbf{G}}(p, v) \phi_{p+1}}\end{aligned}$$

- Taking the expectation over the data \mathbf{X} :

PDE Formulation (II)

- We can use the Sherman-Woodbury formula:

$$\begin{aligned}\tilde{\mathbf{G}}(p+1, \nu) &= \left(\frac{1}{\lambda} \tilde{\mathbf{\Phi}} \tilde{\mathbf{\Phi}}^T + \mathbf{\Lambda}^{-1} + \nu \mathbf{1} \right)^{-1} \\ &= \left(\frac{1}{\lambda} \mathbf{\Phi} \mathbf{\Phi}^T + \frac{1}{\lambda} \phi_{p+1} \phi_{p+1}^T + \mathbf{\Lambda}^{-1} + \nu \mathbf{1} \right)^{-1} \\ &= \left(\tilde{\mathbf{G}}(p, \nu)^{-1} + \frac{1}{\lambda} \phi_{p+1} \phi_{p+1}^T \right)^{-1} \\ &= \tilde{\mathbf{G}}(p, \nu) - \frac{\tilde{\mathbf{G}}(p, \nu) \frac{1}{\lambda} \phi_{p+1} \phi_{p+1}^T \tilde{\mathbf{G}}(p, \nu)}{1 + \phi_{p+1}^T \tilde{\mathbf{G}}(p, \nu) \phi_{p+1}}\end{aligned}$$

- Taking the expectation over the data \mathbf{X} :

$$\langle \tilde{\mathbf{G}}(p+1, \nu) \rangle_{\tilde{\mathbf{\Phi}}} = \langle \tilde{\mathbf{G}}(p, \nu) \rangle_{\mathbf{\Phi}} - \left\langle \frac{\tilde{\mathbf{G}}(p, \nu) \phi_{p+1} \phi_{p+1}^T \tilde{\mathbf{G}}(p, \nu)}{\lambda + \phi_{p+1}^T \tilde{\mathbf{G}}(p, \nu) \phi_{p+1}} \right\rangle_{\tilde{\mathbf{\Phi}}}$$

PDE Formulation (III)

PDE Formulation (III)

- First **approximation**, take expectation of quotient separately:

PDE Formulation (III)

- First **approximation**, take expectation of quotient separately:

$$\langle \tilde{\mathbf{G}}(p+1, v) \rangle_{\tilde{\Phi}} \approx \langle \tilde{\mathbf{G}}(p, v) \rangle_{\Phi} - \frac{\langle \tilde{\mathbf{G}}(p, v) \phi_{p+1} \phi_{p+1}^T \tilde{\mathbf{G}}(p, v) \rangle_{\tilde{\Phi}}}{\langle \lambda + \phi_{p+1}^T \tilde{\mathbf{G}}(p, v) \phi_{p+1} \rangle_{\tilde{\Phi}}}$$

PDE Formulation (III)

- First **approximation**, take expectation of quotient separately:

$$\begin{aligned}
 \langle \tilde{\mathbf{G}}(p+1, v) \rangle_{\tilde{\Phi}} &\approx \langle \tilde{\mathbf{G}}(p, v) \rangle_{\Phi} - \frac{\langle \tilde{\mathbf{G}}(p, v) \phi_{p+1} \phi_{p+1}^T \tilde{\mathbf{G}}(p, v) \rangle_{\tilde{\Phi}}}{\langle \lambda + \phi_{p+1}^T \tilde{\mathbf{G}}(p, v) \phi_{p+1} \rangle_{\tilde{\Phi}}} \\
 &= \langle \tilde{\mathbf{G}}(p, v) \rangle_{\Phi} - \frac{\langle \tilde{\mathbf{G}}(p, v) \langle \phi_{p+1} \phi_{p+1}^T \rangle_{\phi_{p+1}} \tilde{\mathbf{G}}(p, v) \rangle_{\Phi}}{\lambda + \langle \text{tr} \left(\langle \phi_{p+1} \phi_{p+1}^T \rangle_{\phi_{p+1}} \tilde{\mathbf{G}}(p, v) \right) \rangle_{\Phi}}
 \end{aligned}$$

PDE Formulation (III)

- First **approximation**, take expectation of quotient separately:

$$\begin{aligned}
 \langle \tilde{\mathbf{G}}(p+1, v) \rangle_{\tilde{\Phi}} &\approx \langle \tilde{\mathbf{G}}(p, v) \rangle_{\Phi} - \frac{\langle \tilde{\mathbf{G}}(p, v) \phi_{p+1} \phi_{p+1}^T \tilde{\mathbf{G}}(p, v) \rangle_{\tilde{\Phi}}}{\langle \lambda + \phi_{p+1}^T \tilde{\mathbf{G}}(p, v) \phi_{p+1} \rangle_{\tilde{\Phi}}} \\
 &= \langle \tilde{\mathbf{G}}(p, v) \rangle_{\Phi} - \frac{\langle \tilde{\mathbf{G}}(p, v) \langle \phi_{p+1} \phi_{p+1}^T \rangle_{\phi_{p+1}} \tilde{\mathbf{G}}(p, v) \rangle_{\Phi}}{\lambda + \langle \text{tr} \left(\langle \phi_{p+1} \phi_{p+1}^T \rangle_{\phi_{p+1}} \tilde{\mathbf{G}}(p, v) \right) \rangle_{\Phi}} \\
 &= \langle \tilde{\mathbf{G}}(p, v) \rangle_{\Phi} - \frac{\langle \tilde{\mathbf{G}}(p, v)^2 \rangle_{\Phi}}{\lambda + \langle \text{tr} \left(\tilde{\mathbf{G}}(p, v) \right) \rangle_{\Phi}}
 \end{aligned}$$

$\langle \phi_{p+1} \phi_{p+1}^T \rangle_{\phi_{p+1}} = \mathbb{1}$

PDE Formulation (IV)

PDE Formulation (IV)

Equivalently:

PDE Formulation (IV)

Equivalently:

$$\langle \tilde{\mathbf{G}}(p+1, \nu) \rangle_{\tilde{\Phi}} - \langle \tilde{\mathbf{G}}(p, \nu) \rangle_{\Phi} \approx - \frac{\langle \tilde{\mathbf{G}}(p, \nu)^2 \rangle_{\Phi}}{\lambda + \langle \text{tr} \left(\tilde{\mathbf{G}}(p, \nu) \right) \rangle_{\Phi}}$$

PDE Formulation (IV)

Equivalently:

$$\langle \tilde{\mathbf{G}}(p+1, \nu) \rangle_{\tilde{\Phi}} - \langle \tilde{\mathbf{G}}(p, \nu) \rangle_{\Phi} \approx - \frac{\langle \tilde{\mathbf{G}}(p, \nu)^2 \rangle_{\Phi}}{\lambda + \langle \text{tr} \left(\tilde{\mathbf{G}}(p, \nu) \right) \rangle_{\Phi}}$$
$$\stackrel{\text{"1} \rightarrow \text{0"}}{\Longleftrightarrow} \frac{\partial}{\partial p} \langle \tilde{\mathbf{G}}(p, \nu) \rangle_{\Phi} \approx - \frac{\frac{\partial}{\partial \nu} \langle \tilde{\mathbf{G}}(p, \nu) \rangle_{\Phi}}{\lambda + \langle \text{tr} \left(\tilde{\mathbf{G}}(p, \nu) \right) \rangle_{\Phi}}$$

PDE Formulation (IV)

Equivalently:

$$\langle \tilde{\mathbf{G}}(p+1, \nu) \rangle_{\Phi} - \langle \tilde{\mathbf{G}}(p, \nu) \rangle_{\Phi} \approx - \frac{\langle \tilde{\mathbf{G}}(p, \nu)^2 \rangle_{\Phi}}{\lambda + \langle \text{tr} \left(\tilde{\mathbf{G}}(p, \nu) \right) \rangle_{\Phi}}$$
$$\stackrel{\text{"1} \rightarrow \text{0"}}{\Longleftrightarrow} \frac{\partial}{\partial p} \langle \tilde{\mathbf{G}}(p, \nu) \rangle_{\Phi} \approx - \frac{\frac{\partial}{\partial \nu} \langle \tilde{\mathbf{G}}(p, \nu) \rangle_{\Phi}}{\lambda + \langle \text{tr} \left(\tilde{\mathbf{G}}(p, \nu) \right) \rangle_{\Phi}}$$

where we used that $\frac{\partial}{\partial \nu} \langle \tilde{\mathbf{G}}(p, \nu) \rangle_{\Phi} = \langle \tilde{\mathbf{G}}(p, \nu)^2 \rangle_{\Phi}$

PDE Formulation (IV)

Equivalently:

$$\langle \tilde{\mathbf{G}}(p+1, \nu) \rangle_{\Phi} - \langle \tilde{\mathbf{G}}(p, \nu) \rangle_{\Phi} \approx - \frac{\langle \tilde{\mathbf{G}}(p, \nu)^2 \rangle_{\Phi}}{\lambda + \langle \text{tr}(\tilde{\mathbf{G}}(p, \nu)) \rangle_{\Phi}}$$
$$\stackrel{\text{"1} \rightarrow \text{0"}}{\Longleftrightarrow} \frac{\partial}{\partial p} \langle \tilde{\mathbf{G}}(p, \nu) \rangle_{\Phi} \approx - \frac{\frac{\partial}{\partial \nu} \langle \tilde{\mathbf{G}}(p, \nu) \rangle_{\Phi}}{\lambda + \langle \text{tr}(\tilde{\mathbf{G}}(p, \nu)) \rangle_{\Phi}}$$

where we used that $\frac{\partial}{\partial \nu} \langle \tilde{\mathbf{G}}(p, \nu) \rangle_{\Phi} = \langle \tilde{\mathbf{G}}(p, \nu)^2 \rangle_{\Phi}$

We hence reduced the problem to the PDE

PDE Formulation (IV)

Equivalently:

$$\langle \tilde{\mathbf{G}}(p+1, v) \rangle_{\Phi} - \langle \tilde{\mathbf{G}}(p, v) \rangle_{\Phi} \approx - \frac{\langle \tilde{\mathbf{G}}(p, v)^2 \rangle_{\Phi}}{\lambda + \langle \text{tr}(\tilde{\mathbf{G}}(p, v)) \rangle_{\Phi}}$$
$$\stackrel{"1 \rightarrow 0"}{\Longleftrightarrow} \frac{\partial}{\partial p} \langle \tilde{\mathbf{G}}(p, v) \rangle_{\Phi} \approx - \frac{\frac{\partial}{\partial v} \langle \tilde{\mathbf{G}}(p, v) \rangle_{\Phi}}{\lambda + \langle \text{tr}(\tilde{\mathbf{G}}(p, v)) \rangle_{\Phi}}$$

where we used that $\frac{\partial}{\partial v} \langle \tilde{\mathbf{G}}(p, v) \rangle_{\Phi} = \langle \tilde{\mathbf{G}}(p, v)^2 \rangle_{\Phi}$

We hence reduced the problem to the PDE

$$\frac{\partial}{\partial p} \langle \tilde{\mathbf{G}}(p, v) \rangle_{\Phi} = - \frac{\frac{\partial}{\partial v} \langle \tilde{\mathbf{G}}(p, v) \rangle_{\Phi}}{\lambda + \langle \text{tr}(\tilde{\mathbf{G}}(p, v)) \rangle_{\Phi}}$$

PDE Formulation (V)

PDE Formulation (V)

- **Observe:** This is a matrix PDE

PDE Formulation (V)

- **Observe:** This is a matrix PDE
- We have the initial condition $\tilde{\mathbf{G}}(0, \nu) = (\mathbf{\Lambda}^{-1} + \nu \mathbf{1})^{-1}$

no samples $\Rightarrow \Phi \Phi^T = \mathbf{0}$

PDE Formulation (V)

- **Observe:** This is a matrix PDE
- We have the initial condition $\tilde{\mathbf{G}}(0, \nu) = (\mathbf{\Lambda}^{-1} + \nu \mathbf{1})^{-1}$
- $\tilde{\mathbf{G}}(0, \nu)$ is a diagonal matrix for any $\nu \implies \tilde{\mathbf{G}}(0, \nu)_{ij} = 0$

PDE Formulation (V)

- **Observe:** This is a matrix PDE
- We have the initial condition $\tilde{\mathbf{G}}(0, \nu) = (\mathbf{\Lambda}^{-1} + \nu \mathbf{1})^{-1}$
- $\tilde{\mathbf{G}}(0, \nu)$ is a diagonal matrix for any $\nu \implies \tilde{\mathbf{G}}(0, \nu)_{ij} = 0$
 $\implies \langle \tilde{\mathbf{G}}(p, \nu) \rangle_{\Phi}$ remains diagonal

PDE Formulation (V)

- **Observe:** This is a matrix PDE
- We have the initial condition $\tilde{\mathbf{G}}(0, \nu) = (\mathbf{\Lambda}^{-1} + \nu \mathbf{1})^{-1}$
- $\tilde{\mathbf{G}}(0, \nu)$ is a diagonal matrix for any $\nu \implies \tilde{\mathbf{G}}(0, \nu)_{ij} = 0$
 $\implies \langle \tilde{\mathbf{G}}(p, \nu) \rangle_{\Phi}$ remains diagonal
- Hence introduce $g_i(p, \nu) = \langle \tilde{\mathbf{G}}(p, \nu_{ii}) \rangle_{\Phi}$ and
 $t(p, \nu) = \sum_{i=1}^M g_i(t, \nu)$

PDE Formulation (V)

- **Observe:** This is a matrix PDE
- We have the initial condition $\tilde{\mathbf{G}}(0, \nu) = (\mathbf{\Lambda}^{-1} + \nu \mathbf{1})^{-1}$
- $\tilde{\mathbf{G}}(0, \nu)$ is a diagonal matrix for any $\nu \implies \tilde{\mathbf{G}}(0, \nu)_{ij} = 0$
 $\implies \langle \tilde{\mathbf{G}}(p, \nu) \rangle_{\Phi}$ remains diagonal
- Hence introduce $g_i(p, \nu) = \langle \tilde{\mathbf{G}}(p, \nu)_{ii} \rangle_{\Phi}$ and
 $t(p, \nu) = \sum_{i=1}^M g_i(t, \nu)$

By summing up all the individual PDEs we get the PDE

PDE Formulation (V)

- **Observe:** This is a matrix PDE
- We have the initial condition $\tilde{\mathbf{G}}(0, v) = (\Lambda^{-1} + v\mathbf{1})^{-1}$
- $\tilde{\mathbf{G}}(0, v)$ is a diagonal matrix for any $v \implies \tilde{\mathbf{G}}(0, v)_{ij} = 0$
 $\implies \langle \tilde{\mathbf{G}}(p, v) \rangle_{\Phi}$ remains diagonal
- Hence introduce $g_i(p, v) = \langle \tilde{\mathbf{G}}(p, v)_{ii} \rangle_{\Phi}$ and
 $t(p, v) = \sum_{i=1}^M g_i(p, v)$

By summing up all the individual PDEs we get the PDE

$$\frac{\partial t(p, v)}{\partial p} = \frac{1}{\lambda + t(p, v)} \frac{\partial t(p, v)}{\partial v}$$

Method of Characteristics (I)

Method of Characteristics (I)

- **Assume** we have the general PDE

$$a(p, v) \frac{\partial t}{\partial p} + b(p, v) \frac{\partial t}{\partial v} = c(p, v)$$

Method of Characteristics (I)

- **Assume** we have the general PDE

$$a(p, v) \frac{\partial t}{\partial p} + b(p, v) \frac{\partial t}{\partial v} = c(p, v)$$

- The solution forms a **surface**

$$S = \{(t(p, v), p, v) : (p, v) \in \mathbb{R}^2\}$$

Method of Characteristics (I)

- **Assume** we have the general PDE

$$a(p, v) \frac{\partial t}{\partial p} + b(p, v) \frac{\partial t}{\partial v} = c(p, v)$$

- The solution forms a **surface**

$$S = \{(t(p, v), p, v) : (p, v) \in \mathbb{R}^2\}$$

- We have the **tangent** vectors

$$1) \quad \frac{\partial}{\partial p}(t, p, v) = \left(\frac{\partial t}{\partial p}, 1, 0\right)$$

$$2) \quad \frac{\partial}{\partial v}(t, p, v) = \left(\frac{\partial t}{\partial v}, 0, 1\right)$$

Method of Characteristics (I)

- **Assume** we have the general PDE

$$a(p, v) \frac{\partial t}{\partial p} + b(p, v) \frac{\partial t}{\partial v} = c(p, v)$$

- The solution forms a **surface**

$$S = \{(t(p, v), p, v) : (p, v) \in \mathbb{R}^2\}$$

- We have the **tangent** vectors

$$1) \frac{\partial}{\partial p}(t, p, v) = \left(\frac{\partial t}{\partial p}, 1, 0\right)$$

$$2) \frac{\partial}{\partial v}(t, p, v) = \left(\frac{\partial t}{\partial v}, 0, 1\right)$$

- We can form the **normal vector** $n(p, v) = \left(-1, \frac{\partial t}{\partial p}, \frac{\partial t}{\partial v}\right)$

Method of Characteristics (II)

Method of Characteristics (II)

- **Observe:** $(c(p, v), a(p, v), b(p, v)) \bullet \mathbf{n}(p, v) = 0$
 $\implies (c(p, v), a(p, v), b(p, v))$ is in the tangent plane

Method of Characteristics (II)

- **Observe:** $(c(p, v), a(p, v), b(p, v)) \bullet \mathbf{n}(p, v) = 0$
 $\implies (c(p, v), a(p, v), b(p, v))$ is in the tangent plane
- **Idea:** Construct S such that
 $\forall (z, p, v) \in S : (c(p, v), a(p, v), b(p, v))$ is in the tangent plane to S

Method of Characteristics (II)

- **Observe:** $(c(p, v), a(p, v), b(p, v)) \bullet n(p, v) = 0 \quad \forall (p, v)$
 $\implies (c(p, v), a(p, v), b(p, v))$ is in the tangent plane
- **Idea:** Construct S such that
 $\forall (z, p, v) \in S : (c(p, v), a(p, v), b(p, v))$ is in the tangent plane to S
- **Assume:** We have initial data $t(p, v_0) = f(p, v_0)$

Method of Characteristics (III)

Method of Characteristics (III)

- If we want to construct a curve $\Gamma(s) = (z(s), p(s), v(s))$ in S , we can start from the initial data and make sure that its tangent vector agrees with $(c(p, v), a(p, v), b(p, v))$:

$$1) \quad (z(0), p(0), v(0)) = (f(p_0, v_0), p_0, v_0)$$

$$2) \quad \left(\frac{\partial z}{\partial s}, \frac{\partial p}{\partial s}, \frac{\partial v}{\partial s} \right) = (c(p, v), a(p, v), b(p, v))$$

Method of Characteristics (III)

- If we want to construct a curve $\Gamma(s) = (z(s), p(s), v(s))$ in S , we can start from the initial data and make sure that its tangent vector agrees with $(c(p, v), a(p, v), b(p, v))$:

$$1) (z(0), p(0), v(0)) = (f(p_0, v_0), p_0, v_0)$$

$$2) \left(\frac{\partial z}{\partial s}, \frac{\partial p}{\partial s}, \frac{\partial v}{\partial s} \right) = (c(p, v), a(p, v), b(p, v))$$

- This is an ODE, so called **characteristic ODE**

Method of Characteristics (III)


- If we want to construct a curve $\Gamma(s) = (z(s), p(s), v(s))$ in S , we can start from the initial data and make sure that its tangent vector agrees with $(c(p, v), a(p, v), b(p, v))$:

$$1) (z(0), p(0), v(0)) = (f(p_0, v_0), p_0, v_0)$$

$$2) \left(\frac{\partial z}{\partial s}, \frac{\partial p}{\partial s}, \frac{\partial v}{\partial s} \right) = (c(p, v), a(p, v), b(p, v))$$

- This is an ODE, so called **characteristic ODE**
- Its solution is called **characteristic curve**, taking the unions of all such curves constructs entire S

Visualization



Method of Characteristics: Applied Here (I)

Method of Characteristics: Applied Here (I)

- $\frac{\partial t}{\partial p} = \frac{1}{\lambda+t} \frac{\partial t}{\partial v}$ and $t(0, v) = \text{tr} (\Lambda^{-1} + v\mathbb{1})^{-1} = f(v)$

Method of Characteristics: Applied Here (I)

- $\frac{\partial t}{\partial p} = \frac{1}{\lambda+t} \frac{\partial t}{\partial v}$ and $t(0, v) = \text{tr}(\Lambda^{-1} + v\mathbb{1})^{-1} = f(v)$
- The parameter vector is given by $\left(0, 1, -\frac{1}{\lambda+t}\right)$

Method of Characteristics: Applied Here (I)

- $\frac{\partial t}{\partial p} = \frac{1}{\lambda+t} \frac{\partial t}{\partial v}$ and $t(0, v) = \text{tr}(\Lambda^{-1} + v\mathbb{1})^{-1} = f(v)$
- The parameter vector is given by $\left(0, 1, -\frac{1}{\lambda+t}\right)$
- **Initial data** $(f(v), 0, v)$

Method of Characteristics: Applied Here (I)

- $\frac{\partial t}{\partial p} = \frac{1}{\lambda+t} \frac{\partial t}{\partial v}$ and $t(0, v) = \text{tr}(\Lambda^{-1} + v\mathbb{1})^{-1} = f(v)$
- The parameter vector is given by $\left(0, 1, -\frac{1}{\lambda+t}\right)$
- **Initial data** $(f(v), 0, v)$
- Let's take a characteristic curve $(z(s), p(s), v(s))$:

Method of Characteristics: Applied Here (I)

- $\frac{\partial t}{\partial p} = \frac{1}{\lambda+t} \frac{\partial t}{\partial v}$ and $t(0, v) = \text{tr}(\Lambda^{-1} + v\mathbb{1})^{-1} = f(v)$
- The parameter vector is given by $\left(0, 1, -\frac{1}{\lambda+t}\right)$
- **Initial data** $(f(v), 0, v)$
- Let's take a characteristic curve $(z(s), p(s), v(s))$:
 - 1) $(z(0), p(0), v(0)) = (f(v_0), 0, v_0)$

Method of Characteristics: Applied Here (I)

- $\frac{\partial t}{\partial p} = \frac{1}{\lambda+t} \frac{\partial t}{\partial v}$ and $t(0, v) = \text{tr}(\Lambda^{-1} + v\mathbb{1})^{-1} = f(v)$
- The parameter vector is given by $\left(0, 1, -\frac{1}{\lambda+t}\right)$
- **Initial data** $(f(v), 0, v)$
- Let's take a characteristic curve $(z(s), p(s), v(s))$:
 - 1) $(z(0), p(0), v(0)) = (f(v_0), 0, v_0)$
 - 2) $\left(\frac{\partial z}{\partial s}, \frac{\partial p}{\partial s}, \frac{\partial v}{\partial s}\right) = \left(0, 1, -\frac{1}{\lambda+t}\right)$

Method of Characteristics: Applied Here (I)

- $\frac{\partial t}{\partial p} = \frac{1}{\lambda+t} \frac{\partial t}{\partial v}$ and $t(0, v) = \text{tr}(\Lambda^{-1} + v\mathbb{1})^{-1} = f(v)$
- The parameter vector is given by $\left(0, 1, -\frac{1}{\lambda+t}\right)$
- **Initial data** $(f(v), 0, v)$
- Let's take a characteristic curve $(z(s), p(s), v(s))$:
 - 1) $(z(0), p(0), v(0)) = (f(v_0), 0, v_0)$
 - 2) $\left(\frac{\partial z}{\partial s}, \frac{\partial p}{\partial s}, \frac{\partial v}{\partial s}\right) = \left(0, 1, -\frac{1}{\lambda+t}\right)$
- $(z(s), p(s), v(s)) = \left(c_1(v_0), s + c_2(v_0), -\frac{s}{\lambda+t} + c_3(v_0)\right)$
 $= (f(v_0), s, -\frac{s}{\lambda+t} + v_0)$

Method of Characteristics: Applied Here (II)

Method of Characteristics: Applied Here (II)

- $(z(s), p(s), v(s)) = (f(v_0), s, -\frac{s}{\lambda+t} + v_0)$

Method of Characteristics: Applied Here (II)

- $(z(s), p(s), v(s)) = (f(v_0), s, -\frac{s}{\lambda+t} + v_0)$
- Inverting the equations gives:
 - 1) $v_0 = v + \frac{s}{\lambda+t}$
 - 2) $s = p$
 - 3) $z = f(v + \frac{p}{\lambda+t})$

Method of Characteristics: Applied Here (II)

- $(z(s), p(s), v(s)) = (f(v_0), s, -\frac{s}{\lambda+t} + v_0)$
- Inverting the equations gives:
 - 1) $v_0 = v + \frac{s}{\lambda+t}$
 - 2) $s = p$
 - 3) $z = f(v + \frac{p}{\lambda+t})$
- We hence get the solution finally:

Method of Characteristics: Applied Here (II)

- $(z(s), p(s), v(s)) = (f(v_0), s, -\frac{s}{\lambda+t} + v_0)$
- Inverting the equations gives:
 - 1) $v_0 = v + \frac{s}{\lambda+t}$
 - 2) $s = p$
 - 3) $z = f(v + \frac{p}{\lambda+t})$
- We hence get the solution finally:

$$t(p, v) = f\left(v + \frac{p}{\lambda + t}\right) = \text{tr} \left(\left(\Lambda^{-1} + \left(v + \frac{p}{\lambda + t(p, v)}\right) \mathbb{1} \right)^{-1} \right)$$

\Rightarrow Still an implicit equation

Generalization Error Formula

Generalization Error Formula

Plugging in the PDE approximation to obtain $\langle \mathbf{G}^2 \rangle_{\mathbf{x}}$, after some algebraic manipulations, leads to:

Generalization Error Formula

Plugging in the PDE approximation to obtain $\langle \mathbf{G}^2 \rangle_{\mathbf{x}}$, after some algebraic manipulations, leads to:

Theorem 3:

The average generalization error can be approximated like

$$E_i(p) = \frac{\langle \bar{w}_i^2 \rangle}{\lambda_i} \left(\frac{1}{\lambda_i} + \frac{p}{\lambda + t(p)} \right)^{-2} \left(1 - \frac{p\gamma(p)}{(\lambda + t(p))^2} \right)^{-1}$$

where $\gamma(p) = \sum_{i=1}^M \left(\frac{1}{\lambda_i} + \frac{p}{\lambda + t(p)} \right)^{-2}$

Algorithm

Algorithm

The implicit equation needs to be solved numerically. If one is able to do that, we have the following algorithm:

Algorithm

The implicit equation needs to be solved numerically. If one is able to do that, we have the following algorithm:

Algorithm 1 Computing Theoretical Learning Curves

Input: RKHS spectrum $\{\lambda_\rho\}$, target function weights $\{\bar{w}_\rho\}$, regularizer λ , sample sizes $\{p_i\}$, $i = 1, \dots, m$;

for $i = 1$ **to** m **do**

Solve numerically $t_i = \sum_\rho \left(\frac{1}{\lambda_\rho} + \frac{p_i}{\lambda + t_i} \right)^{-1}$

Compute $\gamma_i = \sum_\rho \left(\frac{1}{\lambda_\rho} + \frac{p_i}{\lambda + t_i} \right)^{-2}$

$E_{\rho,i} = \frac{\langle \bar{w}_\rho^2 \rangle}{\lambda_\rho} \left(\frac{1}{\lambda_\rho} + \frac{p_i}{\lambda + t_i} \right)^{-2} \left(1 - \frac{p_i \gamma_i}{(\lambda + t_i)^2} \right)^{-1}$

end for

Algorithm

The implicit equation needs to be solved numerically. If one is able to do that, we have the following algorithm:

Algorithm 1 Computing Theoretical Learning Curves

Input: RKHS spectrum $\{\lambda_\rho\}$, target function weights $\{\bar{w}_\rho\}$, regularizer λ , sample sizes $\{p_i\}$, $i = 1, \dots, m$;

for $i = 1$ **to** m **do**

Solve numerically $t_i = \sum_\rho \left(\frac{1}{\lambda_\rho} + \frac{p_i}{\lambda + t_i} \right)^{-1}$

Compute $\gamma_i = \sum_\rho \left(\frac{1}{\lambda_\rho} + \frac{p_i}{\lambda + t_i} \right)^{-2}$

$E_{\rho,i} = \frac{\langle \bar{w}_\rho^2 \rangle}{\lambda_\rho} \left(\frac{1}{\lambda_\rho} + \frac{p_i}{\lambda + t_i} \right)^{-2} \left(1 - \frac{p_i \gamma_i}{(\lambda + t_i)^2} \right)^{-1}$

end for

This allows one to plot learning curves with the sample size as the varying parameter

Statistical Mechanics

Statistical Mechanics

- Use variation of **Replica trick** to estimate $\langle \mathbf{G}^2 \rangle_{\mathbf{x}}$

Statistical Mechanics

- Use variation of **Replica trick** to estimate $\langle \mathbf{G}^2 \rangle_{\mathbf{x}}$
- Idea is to write the inverse of the partition function as

$$Z^{-1} = \lim_{n \rightarrow 0} Z^{n-1}$$

Statistical Mechanics

- Use variation of **Replica trick** to estimate $\langle \mathbf{G}^2 \rangle_{\mathbf{x}}$
- Idea is to write the inverse of the partition function as

$$Z^{-1} = \lim_{n \rightarrow 0} Z^{n-1}$$

- Then calculate the right-hand side for **integer** n and then analytically continue to found formula to take the limit

Statistical Mechanics

- Use variation of **Replica trick** to estimate $\langle \mathbf{G}^2 \rangle_{\mathbf{x}}$
- Idea is to write the inverse of the partition function as

$$Z^{-1} = \lim_{n \rightarrow 0} Z^{n-1}$$

- Then calculate the right-hand side for **integer** n and then analytically continue to found formula to take the limit
- As usual the calculations are **very bizarre**, order parameter definitions are enforced via Dirac deltas, those get replaced by their Fourier integral representation and then the whole thing simplifies somehow...

Statistical Mechanics

- Use variation of **Replica trick** to estimate $\langle \mathbf{G}^2 \rangle_{\mathbf{x}}$
- Idea is to write the inverse of the partition function as

$$Z^{-1} = \lim_{n \rightarrow 0} Z^{n-1}$$

- Then calculate the right-hand side for **integer** n and then analytically continue to found formula to take the limit
- As usual the calculations are **very bizarre**, order parameter definitions are enforced via Dirac deltas, those get replaced by their Fourier integral representation and then the whole thing simplifies somehow...
- Using a saddle point approximation they finally find the same solution as with the PDE approach

Experiments (I)

Experiments (I)

- Experiment mainly with NTK

Experiments (I)


- Experiment mainly with NTK
- Assume target is given by $f^*(\mathbf{x}) = \sum_{i=1}^{p'} \bar{\alpha}_i \Theta(\mathbf{x}, \bar{\mathbf{x}}_i)$,
 $\bar{\alpha}_i \sim \text{Bernoulli}(\frac{1}{2})$ and $\mathbf{x} \sim \mathbb{S}^{d-1}$

Experiments (I)


- Experiment mainly with NTK
- Assume target is given by $f^*(\mathbf{x}) = \sum_{i=1}^{p'} \bar{\alpha}_i \Theta(\mathbf{x}, \bar{\mathbf{x}}_i)$,
 $\bar{\alpha}_i \sim \text{Bernoulli}(\frac{1}{2})$ and $\mathbf{x} \sim \mathbb{S}^{d-1}$
- This form allows for perfect calculation of E_i leveraging the spherical harmonics $Y_{k,m}$

Experiments (I)

- Experiment mainly with NTK
- Assume target is given by $f^*(\mathbf{x}) = \sum_{i=1}^{p'} \bar{\alpha}_i \Theta(\mathbf{x}, \bar{\mathbf{x}}_i)$,
 $\bar{\alpha}_i \sim \text{Bernoulli}(\frac{1}{2})$ and $\mathbf{x} \sim \mathbb{S}^{d-1}$
- This form allows for perfect calculation of E_i leveraging the spherical harmonics $Y_{k,m}$



(b) 3-layer NTK $k = 1$ $\lambda = 1$



(a) 3-layer NTK $d = 15$ $\lambda = 0$

Experiments (II)

Experiments (II)

- Compare NTK learning curves with finite-width NNs

Experiments (II)

- Compare NTK learning curves with finite-width NNs
- $f^*(\mathbf{x}) = \sum_{i=1}^{p'} \bar{\alpha}_i Q_k(\mathbf{x}^T \bar{\mathbf{x}}_i)$ where

$$Q_k(\mathbf{x}^T \mathbf{x}') = \frac{1}{N(d, k)} \sum_{m=1}^{N(d, k)} Y_{km}(\mathbf{x}) Y_{km}(\mathbf{x}')$$

Experiments (II)

- Compare NTK learning curves with finite-width NNs
- $f^*(\mathbf{x}) = \sum_{i=1}^{p'} \bar{\alpha}_i Q_k(\mathbf{x}^T \bar{\mathbf{x}}_i)$ where

$$Q_k(\mathbf{x}^T \mathbf{x}') = \frac{1}{N(d, k)} \sum_{m=1}^{N(d, k)} Y_{km}(\mathbf{x}) Y_{km}(\mathbf{x}')$$

- Recall that the Mercer decomposition for NTK for $\mathbf{x}_i \sim \mathcal{U}(\mathbb{S}^{d-1})$ is

$$\Theta(\mathbf{x}, \mathbf{x}') = \sum_{k=0}^{\infty} \lambda_k \sum_{m=1}^{N(d, k)} Y_{km}(\mathbf{x}) Y_{km}(\mathbf{x}')$$

Experiments (II)

- Compare NTK learning curves with finite-width NNs
- $f^*(\mathbf{x}) = \sum_{i=1}^{p'} \bar{\alpha}_i Q_k(\mathbf{x}^T \bar{\mathbf{x}}_i)$ where

$$Q_k(\mathbf{x}^T \mathbf{x}') = \frac{1}{N(d, k)} \sum_{m=1}^{N(d, k)} Y_{km}(\mathbf{x}) Y_{km}(\mathbf{x}')$$

- Recall that the Mercer decomposition for NTK for $\mathbf{x}_i \sim \mathcal{U}(\mathbb{S}^{d-1})$ is

$$\Theta(\mathbf{x}, \mathbf{x}') = \sum_{k=0}^{\infty} \lambda_k \sum_{m=1}^{N(d, k)} Y_{km}(\mathbf{x}) Y_{km}(\mathbf{x}')$$

- f^* is hence composed of harmonics belonging to the same eigenvalue


Experiments (III)

Experiments (III)


- Train 2 and 4-layer NNs with widths 500, $d = 30$.

Experiments (III)

- Train 2 and 4-layer NNs with widths 500, $d = 30$.




(a) 2-layer NN $N = 10000$




(b) 4-layer NN $N = 500$

Experiments (III)

- Train 2 and 4-layer NNs with widths 500, $d = 30$.



(a) 2-layer NN $N = 10000$



(b) 4-layer NN $N = 500$

- Good agreement for two layer network, but seems to get worse with depth. No deeper depth experiments included.

Experiments (IV)

Experiments (IV)


- Consider more complicated teacher functions:

$$f^*(\mathbf{x}) = \bar{\mathbf{r}}^T \sigma(\bar{\mathbf{W}}\mathbf{x}) \text{ and } f(\mathbf{x}) = \mathbf{r}^T \sigma(\mathbf{W}\mathbf{x})$$

Experiments (IV)

- Consider more complicated teacher functions:

$$f^*(\mathbf{x}) = \bar{\mathbf{r}}^T \sigma(\bar{\mathbf{W}}\mathbf{x}) \text{ and } f(\mathbf{x}) = \mathbf{r}^T \sigma(\mathbf{W}\mathbf{x})$$



(c) 2-Layer NN Student-Teacher; $N = 8000$

Discussion

Discussion

- I really enjoyed the **mathematics/tools** in this paper

Discussion

- I really enjoyed the **mathematics/tools** in this paper
- Learning curves very general as they hold for **any kernel**

Discussion

- I really enjoyed the **mathematics/tools** in this paper
- Learning curves very general as they hold for **any kernel**
- Also a bit disappointing, would have been nicer to really **specialize** the theory to NTK

Discussion

- I really enjoyed the **mathematics/tools** in this paper
- Learning curves very general as they hold for **any kernel**
- Also a bit disappointing, would have been nicer to really **specialize** the theory to NTK
- Would also have been nicer to have an **analytical** expression (instead of these implicit equations) but probably hard to do

Discussion

- I really enjoyed the **mathematics/tools** in this paper
- Learning curves very general as they hold for **any kernel**
- Also a bit disappointing, would have been nicer to really **specialize** the theory to NTK
- Would also have been nicer to have an **analytical** expression (instead of these implicit equations) but probably hard to do
- Nice that the same solution pops out under two different approximations. But maybe they are **more similar** than one can see at first glance. In both cases, two integer parameters are **made continuous** and limits are taken