# Equivalence of Standardized Score Sum and F1 Score Optimization

Greg Jennings

December 2024

Equivalence of Standardized Score Sum and F1 Score Optimization

### 1 Setup and Definitions

Let's consider a binary classification problem with:

- $\bullet$  Total dataset size: N
- $\bullet$  Number of positive examples (setosa): P
- Number of negative examples: N P

For standardized scores:

- Let  $\mu = \frac{P}{N}$  be the mean (proportion of positive examples)
- $\bullet \ \mbox{Let} \ \sigma = \sqrt{\frac{P(N-P)}{N^2}} \ \mbox{be the standard deviation}$

Therefore:

- Positive class gets score:  $a = \frac{1-\mu}{\sigma}$
- Negative class gets score:  $-b = \frac{0-\mu}{\sigma}$

# 2 For Any Fixed Window Size

For a window containing n points:

- Let TP be true positives
- FP = n TP will be false positives

#### 2.1 Standardized Score Sum

The sum S of standardized scores in the window:

$$S = TP \cdot a + FP \cdot (-b)$$

$$= TP \cdot a + (n - TP) \cdot (-b)$$

$$= TP(a + b) - nb$$

#### 2.2 F1 Score

F1 score is defined as:

$$F1 = 2 \cdot \frac{precision \cdot recall}{precision + recall}$$

where:

$$precision = \frac{TP}{TP + FP} = \frac{TP}{n}$$
 
$$recall = \frac{TP}{P}$$

Therefore:

$$F1 = 2 \cdot \frac{\frac{TP}{n} \cdot \frac{TP}{P}}{\frac{TP}{n} + \frac{TP}{P}}$$
$$= 2 \cdot \frac{TP^2}{nP} \cdot \frac{nP}{TP(n+P)}$$
$$= 2 \cdot \frac{TP}{n+P}$$

# 3 Proof of Optimization Equivalence

- 1. For fixed window size n, the standardized score sum S is:
  - $\bullet$  Linear in TP
  - Has slope (a + b) which is positive (since a > 0 and b < 0)
  - Has constant term -nb
- 2. Similarly, F1 score for fixed n is:
  - $\bullet$  Linear in TP
  - Has slope  $\frac{2}{n+P}$  (positive)
  - Has zero constant term
- 3. Therefore, for any fixed window size n:

- ullet Both metrics are linear functions of TP
- Both have positive slopes
- The constant terms don't affect optimization
- 4. As we slide our window:
  - $\bullet$  For each window size n
  - $\bullet$  Both metrics will achieve their maximum at the same value of TP
  - Therefore they will select the same optimal interval

### 4 Why This Matters

The equivalence shows that:

- 1. Maximizing the sum of standardized scores
- 2. Maximizing F1 score

Are functionally equivalent optimization problems when finding optimal intervals. This means we can use the computationally efficient dynamic programming approach to find optimal F1 score intervals.

#### 5 Additional Note

The key insight is that standardization creates a scoring system where:

- Positive examples contribute positive scores
- Negative examples contribute negative scores
- The ratio of these scores is optimally balanced to align with F1 score optimization