

## Abstract Algebra Notes

**Definition.** A **map**  $f : A \rightarrow B$  is a subset  $f \subset A \times B$  such that for all  $a \in A$ , there exists a  $b \in B$  such that  $b$  is unique with  $(a, b) \in f$ .

**Definition.** We write  $f(a) = b$  if  $(a, b) \in f$ .  $A$  is the **domain** of  $f$  and  $B$  is the **codomain**.

**Definition.** A **binary operation** on  $A$  is a map  $\star : A \times A \rightarrow A$  such that  $\star(a_1, a_2) = a_1 \star a_2$  for  $a_1, a_2 \in A$ .

**Definition.** A binary operation  $\star$  is **associative** on  $A$  if for all  $a, b, c \in A$ ,  $a \star (b \star c) = (a \star b) \star c$ .

**Definition.** An element  $e \in A$  is an **identity** element of  $\star$  if for each  $a \in A$ ,  $e \star a = a \star e = a$ .

**Definition.** An element  $a \in A$  has an **inverse** under  $\star$  if there exists a  $b \in A$  such that  $a \star b = b \star a = e$ .

**Definition.** A set  $A$  with an associative binary operation  $\star$  is a **group** if  $A$  has an identity element under  $\star$  and every  $a \in A$  has an inverse.

### Definition

A group is a pair  $(G, \star)$  where  $G$  is a set and  $\star$  is a binary operation on  $G$  such that

1. For all  $a, b, c \in A$ ,  $a \star (b \star c) = (a \star b) \star c$ .
2. There exists an  $e \in G$  such that  $a \star e = e \star a = a$  for all  $a \in G$ .
3. For all  $a \in G$ , there exists a  $b \in G$  such that  $a \star b = b \star a = e$ .

**Definition.** A group  $(G, \star)$  is **abelian** or commutative if for all  $g, h \in G$ ,  $g \star h = h \star g$ .

**Theorem.** Let  $(G, \star)$  be a group.

1.  $e$  is unique.
2.  $g^{-1}$  is unique.
3.  $\forall g \in G, (g^{-1})^{-1} = g$ .
4.  $\forall g, h \in G, (g \star h)^{-1} = h^{-1} \star g^{-1}$ .

**Proof.** We may prove each part separately.

1. Suppose  $e, e'$  are identity elements. Then for all  $a \in G$ ,

$$a \star e = e \star a = a \quad (\text{i})$$

$$a \star e' = e' \star a = a \quad (\text{ii})$$

By (i),  $e' = e \star e'$  and by (ii),  $e = e \star e'$ . Therefore,  $e = e'$ .

2. Supposed  $a \star b = b \star a = e$ , then

$$\begin{aligned} b &= b \star e \\ &= b \star (a \star a^{-1}) \\ &= (b \star a) \star a^{-1} \\ &= e \star a^{-1} \\ &= a^{-1} \end{aligned}$$

Thus,  $b = a^{-1}$ .

3.  $g^{-1} \star (g^{-1})^{-1} = e = g^{-1} \star g$ . By (ii),  $g = (g^{-1})^{-1}$ .

4. Consider  $(a \star b) \star (b^{-1} \star a^{-1})$ .

$$\begin{aligned} (a \star b) \star (b^{-1} \star a^{-1}) &= a \star (b \star b^{-1}) \star a^{-1} \\ &= a \star e \star a^{-1} \\ &= a \star a^{-1} \\ &= e \end{aligned}$$

Thus,  $(b^{-1} \star a^{-1}) = (a \star b)^{-1}$ .

**Definition.** Let  $[n] = \{1, 2, \dots, n\}$ . The **symmetric group** denoted  $S_n$  of degree  $n$  is the set of all bijections on  $[n]$  under the operation of composition.

$$S_n = \{\sigma : [n] \rightarrow [n] \mid \sigma \text{ is a bijection}\}$$

**Definition.** The **order** of  $(G, \star)$  is the number of elements in  $G$  denoted  $|G|$ .

**Definition.** Let  $n \geq 2$ . The **dihedral group** of index  $n$  is the group of all symmetries of a regular polygon  $P_n$  with  $n$  vertices in the Euclidean plane.

Symmetries of  $P_n$  consist of rotations and reflections.

Choose a vertex  $v$ . Let  $L_0$  be the line from the center of  $P_n$  through  $v$ . Let  $L_k$  be  $L_0$  rotated by  $\frac{\pi k}{n}$  for  $1 \leq k \leq n$ . Let  $\sigma_k$  be a reflection about  $L_k$ . Let  $\rho_k$  be a rotation about  $\frac{2\pi k}{n}$ ,  $1 \leq k \leq n$ .

**Definition.** A subset  $S \subseteq G$  of a group  $(G, \star)$  is a set of **generators**, denoted  $\langle S \rangle = G$ , if and only if every element of  $G$  can be written as a product of elements of  $S$  and their inverses.

**Definition.** Any equation satisfied by generators is called a relation.

**Definition.** A presentation of  $G$ , denoted  $\langle S \mid R \rangle$ , is a set of generators of  $G$  and relations such that any other relation can be derived by those given.

**Example.**

$$D_{2n} = \langle r, s \mid r^n = s^2 = 1, rs = sr^{-1} \rangle$$

**Definition.** The cycles  $\sigma = (\sigma_1 \sigma_2 \dots \sigma_n)$  and  $\tau = (\tau_1 \tau_2 \dots \tau_m)$  are disjoint if  $\sigma_i \neq \tau_j$  for  $1 \leq i \leq n$  and  $1 \leq j \leq m$ .

**Definition.** A cycle of length 2 is called a transposition.

**Definition.** An expression of the form  $(a_1 a_2 \dots a_m)$  is called a cycle of length m or an m-cycle.

**Proposition.** Let  $\alpha = (a_1 a_2 \dots a_m)$  and  $\beta = (b_1 b_2 \dots b_n)$ . If  $a_i \neq b_j$  for any  $i, j$ , then  $\alpha\beta = \beta\alpha$ .

**Proposition.** Every permutation can be written as a product of disjoint cycles.

**Proposition.** A cycle of length  $n$  has order  $n$ .

**Proposition.** Let  $\alpha_1, \alpha_2, \dots, \alpha_n$  be disjoint cycles. Then,

$$|\alpha_1 \alpha_2 \dots \alpha_n| = \text{lcm}(|\alpha_1|, |\alpha_2|, \dots, |\alpha_n|)$$

**Proposition.** Every permutation in  $S_n$  is a product of 2-cycles (which are not necessarily disjoint).

**Proposition.** If  $\alpha = \beta_1 \beta_2 \dots \beta_r = \gamma_1 \gamma_2 \dots \gamma_s$  where  $\beta_i, \gamma_j$  are transpositions, then  $r$  and  $s$  have the same parity.

**Definition.** If  $r$  and  $s$  are both odd,  $\alpha$  is called an odd permutation. If  $r$  and  $s$  are both even,  $\alpha$  is called an even permutation.

**Definition.** The set of even permutations in  $S_n$  form a group called the alternating group, denoted  $A_n$ .

**Note.**  $|A_n| = \frac{n!}{2}$  for  $n > 1$ .

### Definition

Let  $(G, \star)$  and  $(G', *)$  be groups. A map of sets  $\varphi : G \rightarrow G'$  is a group homomorphism if for all  $a, b \in G$ ,

$$\varphi(a \star b) = \varphi(a) * \varphi(b)$$

**Example.** The following are two very simple examples of homomorphisms.

Trivial Homomorphism

$$\varphi : G \rightarrow G', \varphi(g) = e, \forall g \in G$$

Identity Homomorphism

$$\varphi : G \rightarrow G', \varphi(g) = g, \forall g \in G$$

**Definition.** If  $\varphi : G \rightarrow G'$  is a homomorphism, the **domain** of  $\varphi$  is  $\text{Dom}(\varphi) = G$ , the **codomain** of  $\varphi$  is  $\text{Codom}(\varphi) = G'$ , the **range** or **image** of  $\varphi$  is  $\varphi(G) = \{\varphi(g) : g \in G\} \subseteq G'$  denoted  $\text{Range}(\varphi)$  or  $\text{Im}(\varphi)$ .

### Definition

A homomorphism which is bijective is called an **isomorphism**.

$\varphi : G \rightarrow G'$  is an isomorphism if and only if there exists  $\psi : G' \rightarrow G$  such that  $\psi$  is a homomorphism and  $\varphi \circ \psi = 1_{G'}$ ,  $\psi \circ \varphi = 1_G$ , i.e.  $\psi$  is an inverse homomorphism to  $\varphi$ . We say  $G$  is isomorphic to  $G'$  by  $G \cong G'$  or  $\varphi : G \xrightarrow{\sim} G'$ .

**Definition.** Let  $(G, \star)$  be a group. A subset  $H \subseteq G$  is a **subgroup** if  $(H, \star)$  is also a group.

If  $H \neq \emptyset$  and  $H \subseteq G$ ,  $H \leq G$  or  $H$  is a subgroup of  $G$  if and only if

1.  $H$  is closed under  $\star$  ( $\forall h_1, h_2 \in H, h_1 \star h_2 \in H$ ).
2.  $H$  is closed under inverses ( $h \in H \Rightarrow h^{-1} \in H$ ).

**Note.** The following is notation for arbitrary and abelian groups.

$$\begin{aligned} x \star y &\rightarrow xy \text{ for arbitrary } G, x + y \text{ for abelian } G \\ e &\rightarrow 1 \text{ for arbitrary } G, 0 \text{ for abelian } G \end{aligned}$$

For an arbitrary subset  $A \subseteq G$ , and  $g \in G$ ,

$$gA = \{ga : a \in A\} \quad Ag = \{ag : a \in A\} \quad gAg^{-1} = \{gag^{-1} : a \in A\}$$

### Theorem

Let  $\emptyset \neq H \subseteq G$ ,  $H \leq G$  if and only if  $\forall x, y \in H, xy^{-1} \in H$ .