

# STATISTICS FUNDAMENTALS

*Insert Instructor Name*

*Title, Company*

---

# STATISTICS FUNDAMENTALS

---

## LEARNING OBJECTIVES

- Use NumPy and Pandas libraries to analyze datasets using basic summary statistics: mean, median, mode, max, min, quartile, inter-quartile range, variance, standard deviation, and correlation
- Create data visualizations - including: line graphs, box plots, and histograms- to discern characteristics and trends in a dataset
- Identify a normal distribution within a dataset using summary statistics and visualization
- ID variable types and complete dummy coding by hand

---

**COURSE**

---

**PRE-WORK**

---

# **PRE-WORK REVIEW**

---

• Create and open an iPython Notebook

• Complete the Python pre-work

---

**OPENING**


---

# STATISTICS FUNDAMENTALS

# LET'S REVIEW THE DATA SCIENCE WORKFLOW

The steps:

1. Identify the problem
2. Acquire the data
3. Parse the data
4. Mine the data
5. Refine the data
6. Build a data model
7. Present the results



---

# TODAY

---

• We're going to begin to talk about step 3: Parsing the Data

• We'll begin to talk about the fundamentals of Statistics

---

## **INTRODUCTION**

---

# **LAYING THE GROUND WORK**



---

# WE'RE GOING TO COVER SEVERAL TOPICS

---

• Mean

• Median

• Mode

• Max

• Min

• Quartile

• Interquartile Range

• Variance


• Standard Deviation

• Correlation

# MEAN

The mean of a set of values is the sum of the values divided by the number of values. It is also called the average.

$$\overline{X} = \frac{\sum X}{N}$$



---

## **MEAN EXAMPLE**

---

Find the mean of 19, 13, 15, 25, and 18.

---

## MEAN EXAMPLE

---

Find the mean of 19, 13, 15, 25, and 18.

$$\frac{19 + 13 + 15 + 25 + 18}{5} = \frac{90}{5} = 18$$

# MEDIAN


• The median refers to the midpoint in a series of numbers.

• To find the median

• Arrange the numbers in order smallest to largest.

• If there is an odd number of values, the middle value is the median.

• If there is an even number of values, the average of the middle two values is the median.



---

## **MEDIAN EXAMPLE**

---

Find the median of 19, 29, 36, 15, and 20.

---

## **MEDIAN EXAMPLE**

---

Find the median of 19, 29, 36, 15, and 20.

Ordered Values:

15, 19, 20, 29, 36

20 is the median

---

## **MEDIAN EXAMPLE**

---

Find the median of 67, 28, 92, 37, 81, 75.



---

## MEDIAN EXAMPLE

---

Find the median of 67, 28, 92, 37, 81, 75.

Ordered Values:

28, 37, 67, 75, 81, 92

67 and 75 are the middle values.


$$\frac{67 + 75}{2} = \frac{142}{2} = 71$$

71 is the median.

# MODE

The mode of a set of values is the value that occurs most often.

A set of values may have more than one mode or no mode.



---

## **MODE EXAMPLE**

---


Find the mode of 15, 21, 26, 25, 21, 23, 28, and 21.

---

## MODE EXAMPLE

---

Find the mode of 15, 21, 26, 25, 21, 23, 28, and 21.



21 is the mode because it occurs most frequently

---

## **MODE EXAMPLE**

---


Find the mode of 12, 15, 18, 26, 15, 9, 12, and 27.

---

## MODE EXAMPLE

---

Find the mode of 12, 15, 18, 26, 15, 9, 12, and 27.



12 and 15 are the modes since the both occur twice.

---

## **MODE EXAMPLE**

---


Find the mode of 4, 8, 15, 21, and 23.

---

## MODE EXAMPLE

---

Find the mode of 4, 8, 15, 21, and 23.



There is no mode since all values occur the same number of times.



---

# ACTIVITY: KNOWLEDGE CHECK

---

## ANSWER THE FOLLOWING QUESTIONS (5 minutes)



### EXERCISE

1. For the following groups of numbers, calculate the mean, median and mode by hand. Also determine the min and max.
  - a. 18, 24, 17, 21, 24, 16, 29, 18
  - b. 75, 87, 49, 68, 75, 84, 98, 92
  - c. 55, 47, 38, 66, 56, 64, 44, 39

### DELIVERABLE

Answers to the above questions

---

**CODEALONG**

---

# SUMMARY STATISTICS IN PANDAS

---

## **CODEALONG: SUMMARY STATISTICS IN PANDAS**

---

Open the starter-code notebook located in lessons/lesson-03/code/starter-code of the class repo.

---

# CODEALONG PART 1: BASIC STATS

---

We can use Pandas to calculate the mean, median, mode, min, and max.

Methods available include:

`.min()` - Compute minimum value

`.max()` - Compute maximum value

`.mean()` - Compute mean value

`.median()` - Compute median value

`.mode()` - Compute mode value

`.count()` - Count the number of observations

---


# QUARTILES AND INTERQUARTILE RANGE

---

Quartiles divide a rank-ordered data set into four equal parts.

The values that divide each part are called first, second, and third quartiles, denoted  $Q_1$ ,  $Q_2$ , and  $Q_3$ , respectively.

The interquartile range (IQR) is  $Q_3 - Q_1$ , a measure of variability.




---

## CODEALONG PART 2: BOX PLOT

---

RBox plots give a nice visual of min, max, mean, median, and the quartile and interquartile range.





# BIAS VS. VARIANCE

---

Error due to **bias** is calculated as the difference between the *expected prediction* of our model and the *correct value* we are trying to predict.

Imagine creating multiple models on various datasets. **Bias** measures *how far off in general* models' predictions are from the correct value.




---

# BIAS VS. VARIANCE

---


Error due to **variance** is taken as the variability of a model prediction for a given point.

Imagine creating multiple models on various datasets. The **variance** is *how much the predictions for a given point vary* between different realizations of the model.





# BIAS VS. VARIANCE



---

# STANDARD DEVIATION

---

Standard deviation (SD,  $\sigma$  for population,  $s$  for sample) is a measure that is used to quantify the amount of variation or dispersion of a set of data values.

Standard deviation is the square root of variance.

$$s = \sqrt{\frac{\sum (x - \bar{x})^2}{n - 1}}$$

---

# STANDARD ERROR

---

- The standard error of the mean (SEM) quantifies the precision of the mean.
- It is a measure of how far your sample mean is likely to be from the true population mean.
- It generally increases with the size of an estimate, meaning a large standard error may not indicate the estimate of the mean is unreliable.
- It's often better to compare the error in relation to the size of the estimate.

---

## STANDARD ERROR

---

$$SE_{\bar{x}} = \frac{s}{\sqrt{n}}$$

---

## **CODEALONG PART 3: STANDARD DEVIATION & VARIANCE**

---

You can calculate variance and standard deviation easily in Pandas.

Methods include:

`.std()` - Compute Standard Deviation


`.var()` - Compute variance

`.describe()` - short cut that prints out count, mean, std, min, quartiles, max

# CORRELATION

The correlation measures the extent of interdependence of variable quantities.

Example correlation values



---

# CONTEXT

---

• For most projects, descriptive stats will come first. These help you get to know your dataset better.

• Sometimes, descriptive stats may be all you need to answer your question.

---

## INTRODUCTION

---

**IS THIS  
NORMAL?**



---


# THE NORMAL DISTRIBUTION

---

A normal distribution is often a key assumption to many models.

The normal distribution depends upon the *mean* and the *standard deviation*.

The *mean* determines the center of the distribution. The *standard deviation* determines the height and width of the distribution.



---


# THE NORMAL DISTRIBUTION

---

Normal distributions are symmetric, bell-shaped curves.

When the standard deviation is large, the curve is short and wide.


When the standard deviation is small, the curve is tall and narrow.



# SKEWNESS

Skewness is a measure of the asymmetry of the distribution of a random variable about its mean.

Skewness can be positive or negative, or even undefined.




---

# KURTOSIS

---

Kurtosis is a measure of whether the data are peaked or flat relative to a normal distribution.

Datasets with high kurtosis tend to have a distinct peak near the mean, decline rather rapidly, and have heavy tails.



---

**DEMO**

---

# **DETERMINING THE DISTRIBUTION OF YOUR DATA**

---

# **DETERMINING THE DISTRIBUTION OF YOUR DATA**

---

Follow along as we walk through this in an iPython Notebook.

---

## **GUIDED PRACTICE**

---

**IS THIS  
SKEWED?**

---

# ACTIVITY: IS THIS SKEWED?

---

## DIRECTIONS (10 minutes)



### EXERCISE

1. We're going to walk through several images of datasets.
2. For each image, vote on whether the image is:
  - a. Normal
  - b. Positively, negatively, or not skewed
  - c. Has positive, negative, or zero kurtosis
3. Determine how you would correct the issue with each dataset to return it to the normal distribution.



---

## INTRODUCTION

---

# VARIABLE TYPES

---

# **VARIABLE TYPES**

---

• Numeric variables can take on a large range of non-predetermined, quantitative values. These are things such as height, income, etc.

• Categorical variables can take on a specific set of variables. These are things such as race, gender, paint colors, movie titles, etc.

---

**DEMO**

---

**CLASSES**

---

## **CLASS/DUMMY VARIABLES**

---

Let's say we have the categorical variable `area`, which takes on one of the following values: `rural`, `suburban`, and `urban`.

We need to represent these numerically for a model. So how do we code them?

---

# CLASS/DUMMY VARIABLES

---

How about 0=rural, 1=suburban, and 2=urban?

---

## **CLASS/DUMMY VARIABLES**

---

But this implies an ordered relationship - is urban twice suburban?  
That doesn't make sense.

However, we can represent this information by converting the one area variable into two new variables:

area\_urban and area\_suburban.

---

## **CLASS/DUMMY VARIABLES**

---

• We'll draw out how categorical variables can be represented without implying order.

• First, let's choose a reference category. This will be our “base” category.

• It's often good to choose the category with the largest sample size and a criteria that will help model interpretation. If we are testing for a disease, the reference category would be people without the disease.

---

## **CLASS/DUMMY VARIABLES**

---

- Step 1: Select a reference category. We'll choose rural as our reference category.
- Step 2: Convert the values urban, suburban, and urban into a numeric representation that does not imply order.
- Step 3: Create two new variables: area\_urban and area\_suburban.



---

## CLASS/DUMMY VARIABLES

---

Why do we need only two dummy variables?

rural	urban	suburban
-------	-------	----------

We can derive all of the possible values from these two. If an area isn't urban or suburban, we know it must be rural.

In general, if you have a categorical feature with  $k$  categories, you need to create  $k-1$  dummy variable to represent all of the information.

---

# CLASS/DUMMY VARIABLES

---

Let's see our dummy variables.

	area_urban	area_suburban
rural	0	0
suburban	0	1
urban	1	0

As mentioned before, if we know  $\text{area\_urban}=0$  and  $\text{area\_suburban}=0$ , then the area must be rural.

---

## **CLASS/DUMMY VARIABLES**

---

• We can do this for a gender variable with two categories: male and female.

• How many dummy variables need to be created?

---

# **CLASS/DUMMY VARIABLES**

---

$\text{R\# of categories} - 1 = 2 - 1 = 1$

---

## CLASS/DUMMY VARIABLES

---

We will make female our reference category. Thus, female=0 and male=1.

	gender_male
female	0
male	1

This can be done in Pandas with the `get_dummies` method.

---

**INDEPENDENT PRACTICE**

---

# DUMMY COLORS

---

# ACTIVITY: DUMMY COLORS

---



## EXERCISE

### DIRECTIONS (15 minutes)

It's important to understand the concept before we use the Pandas function `get_dummies` to create dummy variables. So today, we'll create our dummy variables by hand.

1. Draw a table like the one on the white board.
2. Create dummy variables for the variable “colors” that has 6 categories: blue, red, green, purple, grey, and brown. Use grey as the reference.

### DELIVERABLE

Dummy variables table for colors

---

**CONCLUSION**

---

# TOPIC REVIEW



---

# REVIEW

---

- Let's go through the process for creating dummy variables for “colors”.
- We talked about several different types of summary statistics, what are they?
- We covered several different types of visualizations; which ones?
- We talked about the normal distribution; how do we determine your data's distribution?
- Any other questions?

---

**COURSE**

---

**BEFORE NEXT  
CLASS**

---

**BEFORE NEXT CLASS**

---

**DUE DATE**

Project: Unit Project 2

---

**LESSON**

---

**CREDITS**

---

**THANKS FOR THE FOLLOWING**

---

# **CITATIONS**

RTitle, Author: link

RTitle, Author: link

RTitle, Author: link

---

# LESSON

---

Q & A

---

**LESSON**

---

# EXIT TICKET

**DON'T FORGET TO FILL OUT YOUR  
EXIT TICKET**

---

**THANKS!**

---

**NAME**

Optional Information:

Email?

Website?

Twitter?