

Drought frequency predicts life history strategies in African *Heliophila*

J. Grey Monroe^{1,2}, Brian Gill³, Kathryn G. Turner⁴, & John K. McKay²

³ ¹ Graduate Degree Program in Ecology, Colorado State University, Fort Collins, CO 80521,
⁴ USA

⁵ ² College of Agriculture, Colorado State University, Fort Collins, CO 80521, USA

⁶ ³ Institute for Environment and Society, Brown University, Providence, RI 02912, USA

⁷ ⁴ Biology Department, Pennsylvania State University, State College, PA 16802, USA

⁷ ⁴ Biology Department, Pennsylvania State University, State College, PA 16802, USA

Author Note

Correspondence concerning this article should be addressed to J. Grey Monroe, 307
University Ave, Fort Collins, CO 80521. E-mail: monroejg@colostate.edu

11

Abstract

12 Explaining variation in life history strategies is a long-standing goal of evolutionary biology.
13 For plants, annual and perennial life histories are thought to reflect adaptation to
14 environments that differ in the frequency of environmental stress such as drought. Here we
15 test this hypothesis in *Heliophila* (Brassicaceae), a diverse genus of flowering plants native to
16 Africa, by integrating 34 years of satellite-based drought measurements with 2192 herbaria
17 occurrence records. Consistent with predictions from classic life history theory, we find that
18 perennial *Heliophila* species occur in environments where droughts are significantly less
19 frequent compared to annuals. These associations are predictive while controlling for
20 phylogeny, lending support to the hypothesis that drought related natural selection has
21 influenced the distributions of these strategies. Additionally, the difference in drought
22 frequency between annual and perennial species distributions is greatest during the summer
23 and fall, which also appears to be when annuals are in the seed phase of their life cycle based
24 on collection dates of annual species. Together, these finding provide empirical support for
25 classic hypotheses about the drivers of life history strategy in plants - that perennials out
26 compete annuals in environments with less frequent drought and that annuals are adapted to
27 environments with more frequent drought by escaping drought prone seasons as seeds.

28

Keywords: drought adaptation, life history evolution, remote sensing, phylogeography,
29 herbaria records

30 Drought frequency predicts life history strategies in African *Heliophila*

31 **Introduction**

32 Understanding the causes and consequences of life history variation is a longstanding
33 goal of ecology and evolutionary biology (Cole, 1954). In plants, life histories are especially
34 diverse, with herbaceous species that complete their life cycle in a number of weeks to trees
35 that live for thousands of years (Brown, 1996). Along this continuum an important division
36 exists, distinguishing annuals which complete their seed to seed life cycle within a single
37 calendar year from perennials which can persist over multiple years. Annual plants flower
38 once, set seed, senesce, and then die, spending at least some portion of the year as a seed,
39 where they are relatively protected from environmental stress. In contrast, perennial plants
40 can continue vegetative growth after reproduction and must survive conditions experienced
41 during all seasons. These represent fundamentally different life history strategies, but the
42 ecological factors that explain their evolution and distributions remain empirically unresolved
43 (Friedman and Rubin, 2015).

44 Classical theory predict shorter life spans in environments where adult mortality is
45 high (Charnov and Schaffer, 1973; Franco and Silvertown, 1996; Stearns, 1992). In plants,
46 this has been extended to the hypothesis that annuality is adaptive when it allows plants to
47 escape drought (Schaffer and Gadgil, 1975). Lack of water is perhaps the greatest threat to
48 survival during vegetative or reproductive growth and annuals can remain dormant (and
49 protected as a seed) during drought. Thus, environments with greater seasonal drought
50 frequency may select for annual life histories that complete reproduction prior to drought
51 prone seasons. Conversely, environments with less frequent drought may select for perennial
52 species, which may benefit from multiple bouts of reproduction and competitive advantage
53 by preventing recruitment of annual species (Corbin and D'Antonio, 2004). These
54 predictions have been supported by the observation of annuals in arid environments in *Oryza*
55 *perennis* (Morishima et al., 1984) and *Oenothera* (Evans et al., 2005). Additionally, annual

56 and perennial species of *Nemesia* were qualitatively associated with winter rather and
57 summer rainfall environments respectively (Datson et al., 2008) and annual species of
58 *Scorzonerooides* were associated with environments classified as unpredictable (Cruz-Mazo et
59 al., 2009). However, whether the history frequency of drought events indeed predicts the
60 distributions annual or perennial life history strategies has yet to be tested in a phylogenetic
61 context with suitable sample sizes of distribution and climate data.

62 Here we combine a long-term global dataset of satellite detected drought events with
63 metadata from natural history collections to test these classic hypotheses about the
64 evolution of life history strategies within the African endemic mustard genus, *Heliophila* L.
65 (Brassicaceae). If annuality is an adaptive strategy allowing plants to escape drought prone
66 seasons, then drought frequency should predict the distribution of life history strategies
67 across landscapes, and annual species should be more commonly associated with drought
68 prone regions than perennial species. Furthermore, if annual species have adapted to escape
69 drought prone seasons, observations of growing annual species (i.e. occurring in forms other
70 than seed) should be rare during drought prone seasons. Phylogenetic relatedness can have
71 significant non-random effects on species distributions and life history traits (Barrett et al.,
72 1996), and therefore we assessed the relationship between life history distribution and
73 drought frequency in a phylogenetically controlled background.

74

Materials and Methods

75 **Data**

76 **Availability.** All analyses were performed using R. All data and the source code to
77 produce this manuscript are available at <https://github.com/greymonroe/heliophila>.
78 Software used is listed in the supplement.

79 **Satellite-detected drought data.** Remotely sensed data is a powerful tool for
 80 characterizing seasonal patterns in drought because it is less limited in spatial and temporal
 81 scope and resolution than weather stations or field observations (AghaKouchak et al., 2015).
 82 To quantify the frequency of drought during different seasons across landscapes, we used the
 83 remotely sensed Vegetative Health Index (VHI), which measures landscape scale reductions
 84 in plant cover and temperature conditions characteristic of drought (Kogan, 2001).
 85 Generated from data collected by NOAA AVHRR satellites since 1981, the VHI combines
 86 Normalized Difference Vegetation Index (NDVI) derived measures of vegetative stress
 87 (Vegetative Condition Index - VCI) with temperature stress indicated by anomalies in
 88 thermal spectra (Temperature Condition Index - TCI). The VHI of year y during week w of
 89 [1, 52] at pixel i is derived from the following equations, where n is the number of years
 90 observed.

$$VCI_{y,w,i} = 100 \frac{NDVI_{y,w,i} - NDVI_{min}}{NDVI_{max} - NDVI_{min}}$$

$$TCI_{y,w,i} = 100 \frac{T_{y,w,i} - T_{min}}{T_{max} - T_{min}}$$

$$VHI_{y,w,i} = 0.5(VCI_{y,w,i}) + 0.5(TCI_{y,w,i})$$

91 where $NDVI_{min} = min(NDVI_{1981,w,i} \dots NDVI_{1981+n,w,i})$ and
 92 $NDVI_{max} = max(NDVI_{1981,w,i} \dots NDVI_{1981+n,w,i})$ and $T_{min} = min(T_{1981,w,i} \dots T_{1981+n,w,i})$
 93 and $T_{max} = max(T_{1981,w,i} \dots T_{1981+n,w,i})$

94 Thus, VHI measurements are standardized according to conditions historically
 95 observed at each locations. These measurements have been validated and generally used for
 96 evaluating drought risk and predicting crop yields in agriculture (e.g., Rojas et al., 2011;
 97 Kogan et al., 2016). But they also present a new tool to study seasonal patterns in the

frequency of drought across environments and to test hypotheses about the effect of drought on ecological and evolutionary processes (Kerr and Ostrovsky, 2003). As such, the VHI has been applied recently to study drought related ecology of natural species and proven useful for predicting intraspecific variation in drought tolerance traits and genes (Dittberner et al., 2018; Mojica et al., 2016; John G Monroe et al., 2018). Here, we accessed VHI data at 16km² resolution from 1981 to 2015 (https://www.star.nesdis.noaa.gov/smcd/emb/vci/VH/vh_ftp.php) to characterize the seasonal drought frequencies experienced by annual and perennial *Heliophila* species.

Life history data for *Heliophila*. *Heliophila* is a genus of flowering plants endemic to the southern portion of Africa including the Cape Floristic and Succulent Karoo Regions. These are among the most botanically diverse environments on Earth and the estimated ~50 *Heliophila* species occurring there are considered to make up the most diverse genus of the family Brassicaceae (Mandáková et al., 2012; Mummenhoff et al., 2005). This genus includes both perennial and annual species and this change in life history strategy has likely arisen multiple independent times (Appel and Al-Shehbaz, 1997; Mummenhoff et al., 2005). Furthermore, the fine scale climatic heterogeneity of Southern Africa is ideal for studying the distribution of traits in relation to environmental parameters (Sayre et al., 2013). We used life histories reported by Mummenhoff et al. (2005), grouping species with annual or perennial life histories. Perenniality was defined based any form of perennial life history (e.g., herbs, shrubs, mixed, etc). Because the nature of species reported with mixed traits were unknown (i.e. plasticity vs. genetic variation), we classified these species here as perennial since they can maintain vegetative growth after reproduction at least to some capacity.

***Heliophila* occurrence records.** Botanists have collected and maintained over 350 million botanical specimens worldwide over the past 300 years. Herbarium specimens and their associated metadata have been used since the 1960s to study species' geographical distributions (reviewed by Willis et al. (2017) and Lang et al. (2018)). And as they become

124 digitized (Soltis, 2017), these collections have been used to study relationships between trait
125 distributions, geography, and climate (Davis et al., 2015; Stropp et al., 2016; Václavík et al.,
126 2017; Wolf et al., 2016). To characterize the distributions of annual and perennial *Heliophila*
127 species, all records for the genus *Heliophila* were downloaded from the Global Biodiversity
128 Information Facility (gbif.org) on July 21, 2018 (GBIF, 2018).

129 **Sequence data for phylogeny.** Aligned *Heliophila* ITS sequences were obtained
130 from previous work by Mandáková et al. (2012). *Aethionema*, *Alliaria*, *Cardamine*, *Chamira*,
131 and *Rorippa* ITS records from were downloaded from Genbank.

132 Analyses

133 **Drought frequency calculations.** To characterize drought regimens across the
134 distributions of annual and perennial species of *Heliophila*, we calculated drought during
135 different seasons at the location of observations for *Heliophila* records using the VHI.
136 Specifically, we created global maps of the frequencies of observing drought conditions
137 (VHI<40, NOAA) during the winter (quarter surrounding winter solstice), spring (quarter
138 surrounding spring equinox), summer (quarter surrounding summer solstice) and fall (quarter
139 surrounding fall equinox) from 1981 to 2015. From these maps, the drought frequency during
140 the winter, spring, summer, and fall were extracted for the locations of all GBIF records.

141 **Filtering of occurrence records.** To avoid instances with spurious location data,
142 we filtered raw GBIF by restricting our analyses to include only:

- 143 • records for species with reported life history
- 144 • records with geospatial data
- 145 • records without known geospatial coordinate issues (i.e., coordinates reported are those
146 of herbarium)
- 147 • records from collection sites classified as land pixels

- 148 • records from Africa (to exclude locations of cultivation)
149 • records without duplicates (i.e., identical species, location, collection date)

150 **Phylogeny construction.** Out group species were aligned together with *Helophilus*
151 ITS sequences using MAFFT. Model selection for construction of phylogeny was performed
152 in jModeltest2 with CIPRES. Based on this analysis, *GTR + L* were selected. Ultrametric
153 phylogeny was estimated with branch lengths as relative time.

154 **Comparison of drought frequency between annual and perennial species.**

155 To evaluate the hypothesis that annual and perennial life history strategies reflect
156 adaptations to alternative drought regimes, we tested the corresponding prediction that the
157 observed distributions of annual and perennial *Helophilus* species would be significantly
158 associated with historic drought frequency. First, we compared the frequency of drought
159 during the winter, spring, summer, and fall between raw occurrence records of annual and
160 perennial species by t-tests. To account for variation in the number of occurrence records per
161 species, we next calculated the mean drought frequency during the winter, spring, summer
162 and fall for each species. The relationships between species mean values of drought frequency
163 during each season and life habitat (annual or perennial) were tested using Firth's
164 penalized-likelihood logistic regressions. Because demographic histories caused by ancestry
165 can confound trait - environment associations, we then tested for the relationships between
166 drought frequency and life history while controlling for relationships between species using
167 phylogenetic logistic regressions.

168 **Collection dates.** To test the hypothesis that annual species have adapted to
169 escape drought prone seasons as seeds, collection dates for herbarium specimens were
170 compared between annual and perennial species. Comparisons of distributions were made by
171 Two-sample Kolmogorov-Smirnov test, t-test, and Barlett variance test.

Results




Figure 1. Species and examples of herbaria specimens of *Heliophila* (a) Phylogeny and life history strategies of species studied. Orange circles at branch tips mark annual species and blue circles mark perennial species. Example herbaria specimens accessed via GBIF of (a) *H. minima*, (b) *H. deserticola*, (c) *H. coronopifolia* and (d) *H. ephemera*. Images (a,c,d) courtesy of The Bavarian Natural History Collections (CC BY-SA 4.0) and (b) The London Natural History Museum (CC BY 4.0). Links to images are found in the supplement.




Figure 2. Locations of occurrence records of (a) annual and (b) perennial *Heliophila*. Drought frequency during the (c) winter, (d) spring, (e) summer and (f) fall measured using the VHI. (g) Drought frequencies during each season at the observation locations of annual and perennial *Heliophila* (t tests, ** = $p < 0.01$).

173 Out of 8670 *Heliophila* GBIF records, 6634 were for species with reported life history

174 (Mummenhoff et al., 2005), 2856 had geospatial data, 2833 did not have geospatial issues,

175 2684 were located on pixels classified as land having drought measurements, 2543 were

176 located in Africa, 2192 were not duplicated. Thus, after all filtering steps, 2192 records for

177 42 species (Figure 1, Table S1) passed for further analyses. The number of samples varied

178 between species, with a mean of 52.19 samples per species. *Heliophila rigidiuscula* had the

179 most records, 201, and *Heliophila cornellsbergia* the fewest, 2 (Table S1).

180 There were clear visual differences between the distributions of the 960 annual and the

181 1232 perennial *Heliophila* observation records (see Figure S1 for maps of individual species).

182 While annual species were generally found in the western regions of South Africa and

183 Namibia, primarily in the Cape Floristic Region and Succulent Karoo (Figure 2a), the

184 occurrence of perennials extended to the east coast of South Africa (Figure 2b).

185 The frequency of drought varied considerably across the ranges of *Heliophila* species

186 (Figure 2c-f). This heterogeneity is expected, given that this is one of the most climatically

187 diverse regions of the Earth (Sayre et al., 2013). It is worth noting the east to west cline in

188 drought frequency observed during the summer, which distinguishes the high drought

189 frequency of the Kalahari Sands and Namid Desert phytogeographic regions from the low

190 drought frequency of the Drakensberg Mountains and Coastal Zambesian phytogeographic

191 regions. In the Cape phytogeographic region there was finer scale heterogeneity in drought

192 frequency during the summer.

193 Theory predicts that annuality should be adaptive in places where stresses such as

194 drought are more common. Conversely, perenniability should be adaptive in places where such

195 stresses are less frequent. We found that the frequency of drought was significantly higher at

196 the locations of occurrence records for annual species. In terms of raw observation records

197 (Figure 2g), the frequency of drought was significantly higher at the location of annuals

198 during the winter ($t = 10.65$, $p = 0.00$), spring ($t = 10.73$, $p = 0.00$), summer ($t = 12.67$, p

= 0.00), and fall ($t = 15.26$, $p = 0.00$). Because raw occurrence records do not account for variation in the number of records per species (Table S1), we also tested whether mean drought frequency values of each species were significantly different between annuals and perennials by a Firths-penalized logistic regression. We found that the mean drought frequencies were significantly higher ($\alpha = 0.05$) in annual species during the spring, summer, and fall (Table 1, Figure 3a). We further tested whether annual species are found in places where droughts occur more frequently while controlling for the phylogenetic relatedness of *Heliophila* species. This is important, because environmental differences in species distributions can be confounded with demographic history caused by ancestry. We found that while controlling for phylogeny, the mean drought frequencies were significantly higher ($\alpha = 0.05$) in annual species during the spring, summer, and fall (Table 1, Figure 3a). These findings indicate that common ancestry alone does not explain differences in the drought frequencies experienced in the environments of annual and perennial *Heliophila*.

The preceding results indicated that annual species are found in environments where droughts are significantly more frequent, especially in the summer and fall. Classic life history theory hypothesizes that annuality reflects adaptation to such environments because it allows species to escape stressful conditions. If this is the case, we would expect that annuals spend the drought prone seasons of summer and fall as seeds. To test this hypothesis, we compared the dates of occurrence records between annual and perennial *Heliophila* species. The distributions reveal a considerable difference in the timing of observation of these two life histories. In comparison to perennials, which appear to be collected throughout the year, annuals are almost exclusively observed during the winter and spring (Figure 3b). The differences between the distribution of collection dates were significant by all tests ($ks.test D = 0.25$, $p = 0$; $bartlett.test K2 = 503.18$, $p = 0.00$) This is consistent with a model of life history in which annual species flower in the spring, set seed, senesce, and die before the summer. Thus, these annual species are likely to remain dormant during the summer and fall, when drought is the strongest predictor of the distributions of

Table 1

*Logistic regressions between life history, and the mean drought frequency observed at herbaria collection sites of *Heliophila* species the winter, spring, summer, and fall.*

Predictor	Estimate‘	P‘	Estimate*	P*
Intercept	2.2575	0.1739	0.7231	0.6636
Winter drought freq.	-6.7484	0.1661	-1.5452	0.7274
Intercept	4.5594	0.0443	5.0107	0.0534
Spring drought freq.	-11.7895	0.0423	-12.9014	0.0464
Intercept	7.1742	0.0011	7.7093	0.0054
Summer drought freq.	-18.2999	0.0010	-19.9056	0.0042
Intercept	6.4226	0.0029	7.0162	0.0082
Fall drought freq.	-19.0512	0.0026	-20.8174	0.0067

Note. ‘ = Firth’s penalized logistic regression. * = Phylogenetically constrained logistic regression. Annual species were scored as 0 and perennial species as 1.

226 annual and perennial life histories (Figure 3a).

227

Discussion

228 To test the hypothesis that annual and perennial plants reflect adaptation to
 229 alternative drought environments we examined the landscape distribution of life history
 230 strategies in the large and diverse mustard genus, *Heliophila*. Using metadata of 2192
 231 occurrence records and a 34 year dataset of satellite-detected droughts, we tested the
 232 prediction that annual species are more often observed in drought-prone locations than




Figure 3. (a) Comparison (mean + SE) of drought frequency across seasons measured at the GBIF records of annual and perennial species of *Heliophila*. (phylogenetic logistic regression, * = $p < 0.05$, ** = $p < 0.01$) (b) Collection dates of GBIF records of annual and perennial species of *Heliophila*.

perennial species, when controlling for phylogenetic relatedness. We found that drought frequency is significantly different between the distributions of annual and perennial species, with annuals being found in environments with more frequent drought, and that this signal is strongest during the seasons when annuals are likely escaping via seed dormancy. These results remain significant while controlling for the phylogenetic relationships of *Heliophila* species, yielding support for the role that natural selection has played in driving contemporary distributions of these alternative strategies in relation to drought regimens.

We cannot eliminate the possibility that confounding traits or environmental variables are the causative factors explaining variation in the distributions of annual and perennial species. Nevertheless, these results reveal quantitative support for the classic prediction that annual species are found in environments that experience more frequent drought than perennial species. These findings complement previous reports of qualitative associations between annuality with environments characterized as having increased aridity (Evans et al., 2005), alternative precipitation defined habitats (Datson et al., 2008; Morishima et al., 1984), or greater unpredictability (Cruz-Mazo et al., 2009). However, to our knowledge this is the first study to demonstrate a significant association between life history and drought in a phylogenetic context informed by large scale species distribution data and long term drought measures.

Unfortunately, herbarium collections and their associated data do not represent systematic or random sampling of a species distribution. Significant biases in collecting exist, which we have not necessarily controlled for here, and may have some effect on our findings, such as a bias toward collecting near roads or near the locations of natural history collections (Daru et al., 2018). Future research will benefit from systematic sampling efforts to avoid these noted biases. However, the ecosystems of southern Africa include several biodiversity hotspots and are among the most botanically well sampled regions on Earth (Daru et al., 2018), suggesting that this may currently be the optimal region for our analyses of life history distribution. Indeed, we were able to use 2192 occurrence records to study 42 species, which represents a significant advance over relying on personal observations to characterize species distributions.

These findings support classical theoretical predictions about the adaptive value of annual and perennial life history strategies. Taken together, they suggest that in *Helophilus*, annual species are adapted to environments with increased summer droughts by avoiding these seasons in a dormant seed phase of their life cycle. They also suggest that perenniality

266 is adaptive in environments where droughts are less frequent. While most previous work has
267 focused on describing the evolutionary origins of annuality (Andreasen and Baldwin, 2001;
268 Barrett et al., 1996; Conti et al., 1999; Friedman and Rubin, 2015; Verboom et al., 2004)
269 there are at least a few other cases where perenniability appears to have arisen from an annual
270 ancestor (Bena et al., 1998; Tank and Olmstead, 2008). And while early theory predicted
271 selection for annuality when adult mortality is high (Stearns, 1992), we also find evidence
272 that the transition to perenniability could be explained by the historical frequency of drought,
273 a likely cause of mortality in plants. The phylogeny reveals several transitions from annual
274 to perennial life history (Figure 1a) and that the distributions of perennial *Heliophila* extend
275 into regions where drought frequency is low (Figure 2b, Figure S1). Perennials may be able to
276 out compete annual relatives in environments where the infrequency of drought favors
277 strategies that allow plants to benefit from growth over many seasons. It may also indicate
278 that annuals rely on drought as a source of disturbance for seedling recruitment when
279 competing with perennials (Corbin and D'Antonio, 2004). Indeed, no annual species were
280 observed in the low drought regions of eastern South Africa 2a, Figure S1).

281 These findings suggest that species with locally adaptive life history strategies could be
282 threatened by rapidly changing drought regimens (Dai, 2011). This could have impacts on
283 ecosystem functioning and processes such as carbon cycling if the composition of annual and
284 perennial species changes as a response (Garnier et al., 1997; J Grey Monroe et al., 2018;
285 Roumet et al., 2006). Furthermore, the frequency of drought may be an important factor
286 when considering the use of perennial cropping systems (Lelièvre and Volaire, 2009; Parry et
287 al., 2005).

288 In conclusion, we find strong support for classic life history theory which predicts that
289 annuality is adaptive in environments where droughts occur more frequently. Additionally,
290 we report evidence consistent with a life history model in annuals in which they escape
291 drought prone seasons during the seed phase of their life cycle. Finally, we find evidence that

292 the distributions of perennial lineages may indicate a competitive advantage in areas where
293 droughts are infrequent. More broadly, this work highlights the irreplaceable value of natural
294 history collections and demonstrates the power of combining such information with large
295 scale remote sensing data to address outstanding classic hypotheses in ecology and evolution.

296 **Acknowledgments**

297 We thank Jesse Lasky and members of the Sloan lab for generous feedback that
298 improved the quality of this work. This research was supported by NSF Award 1701918 and
299 USDA-NIFA Award 2014-38420-21801 to JGM.

300 **Author contributions**

301 JGM, BG, KGT and JKM contributed to the design of the research, interpretation,
302 and writing the manuscript. JGM, BG, and KGT contributed to the performance of the
303 research and data analysis.

304 **References**

- 305 AghaKouchak A, Farahmand A, Melton F, Teixeira J, Anderson M, Wardlow BD, Hain C.
306 2015. Remote sensing of drought: Progress, challenges and opportunities. *Reviews of*
307 *Geophysics* **53**:452–480.
- 308 Alfaro M, Santini F, Brock C, Alamillo H, Dornburg A, Rabosky D, Carnevale G, Harmon L.
309 2009. Nine exceptional radiations plus high turnover explain species diversity in
310 jawed vertebrates. *Proceedings of the National Academy of Sciences of the United*
311 *States of America* **106**:13410–13414.
- 312 Andreasen K, Baldwin BG. 2001. Unequal evolutionary rates between annual and perennial
313 lineages of checker mallows (sidalcea, malvaceae): Evidence from 18S–26S rDNA
314 internal and external transcribed spacers. *Molecular Biology and Evolution*
315 **18**:936–944.
- 316 Appel O, Al-Shehbaz IA. 1997. Generic limits and taxonomy of hornungia, pritzelago, and
317 hymenolobus (brassicaceae). *Novon* 338–340.
- 318 Aust F, Barth M. 2018. papaja: Create APA manuscripts with R Markdown.
- 319 Barrett SCH, Harder LD, Worley AC. 1996. The comparative biology of pollination and
320 mating in flowering plants. *Phil Trans R Soc Lond B* **351**:1271–1280.
- 321 Bates D, Maechler M. 2018. Matrix: Sparse and dense matrix classes and methods.
- 322 Bena G, Lejeune B, Prosperi J-M, Olivieri I. 1998. Molecular phylogenetic approach for
323 studying life-history evolution: The ambiguous example of the genus medicago l.
324 *Proceedings of the Royal Society of London B: Biological Sciences* **265**:1141–1151.
- 325 Brown PM. 1996. OLDLIST: A database of maximum tree ages. *Tree rings, environment,*
326 *and humanity Radiocarbon* **1996**:727–731.

- 327 Charnov EL, Schaffer WM. 1973. Life-history consequences of natural selection: Cole's result
328 revisited. *The American Naturalist* **107**:791–793.
- 329 Cole LC. 1954. The population consequences of life history phenomena. *The Quarterly
330 Review of Biology* **29**:103–137.
- 331 Conti E, Soltis DE, Hardig TM, Schneider J. 1999. Phylogenetic relationships of the silver
332 saxifrages (saxifraga, sect. Ligulatae haworth): Implications for the evolution of
333 substrate specificity, life histories, and biogeography. *Molecular Phylogenetics and
334 Evolution* **13**:536–555.
- 335 Corbin JD, D'Antonio CM. 2004. Competition between native perennial and exotic annual
336 grasses: Implications for an historical invasion. *Ecology* **85**:1273–1283.
- 337 Cruz-Mazo G, Buidé M, Samuel R, Narbona E. 2009. Molecular phylogeny of scorzoneroideae
338 (asteraceae): Evolution of heterocarpy and annual habit in unpredictable
339 environments. *Molecular phylogenetics and evolution* **53**:835–847.
- 340 Dai A. 2011. Drought under global warming: A review. *Wiley Interdisciplinary Reviews:
341 Climate Change* **2**:45–65.
- 342 Daru BH, Park DS, Primack RB, Willis CG, Barrington DS, Whitfeld TJ, Seidler TG,
343 Sweeney PW, Foster DR, Ellison AM, others. 2018. Widespread sampling biases in
344 herbaria revealed from large-scale digitization. *New Phytologist* **217**:939–955.
- 345 Datson P, Murray B, Steiner K. 2008. Climate and the evolution of annual/perennial
346 life-histories in nemesia (scrophulariaceae). *Plant Systematics and Evolution*
347 **270**:39–57.
- 348 Davis CC, Willis CG, Connolly B, Kelly C, Ellison AM. 2015. Herbarium records are
349 reliable sources of phenological change driven by climate and provide novel insights

- 350 into species' phenological cueing mechanisms. *American Journal of Botany*
351 **102**:1599–1609.
- 352 Dittberner H, Korte A, Mettler-Altmann T, Weber A, Monroe G, Meaux J de. 2018.
353 Natural variation in stomata size contributes to the local adaptation of water-use
354 efficiency in *arabidopsis thaliana*. *bioRxiv* 253021.
- 355 Eastman J, Alfaro M, Joyce P, Hipp A, Harmon L. 2011. A novel comparative method for
356 identifying shifts in the rate of character evolution on trees. *Evolution* **65**:3578–3589.
- 357 Evans ME, Hearn DJ, Hahn WJ, Spangle JM, Venable DL. 2005. Climate and life-history
358 evolution in evening primroses (*oenothera*, *onagraceae*): A phylogenetic comparative
359 analysis. *Evolution* **59**:1914–1927.
- 360 Franco M, Silvertown J. 1996. Life history variation in plants: An exploration of the
361 fast-slow continuum hypothesis. *Phil Trans R Soc Lond B* **351**:1341–1348.
- 362 Friedman J, Rubin MJ. 2015. All in good time: Understanding annual and perennial
363 strategies in plants. *American journal of botany* **102**:497–499.
- 364 Garnier E, Cordonnier P, Guillerm J-L, Sonié L. 1997. Specific leaf area and leaf nitrogen
365 concentration in annual and perennial grass species growing in mediterranean
366 old-fields. *Oecologia* **111**:490–498.
- 367 GBIF. 2018. GBIF occurrence download.
- 368 Genz A, Bretz F. 2009. Computation of multivariate normal and t probabilities, Lecture
369 notes in statistics. Heidelberg: Springer-Verlag.
- 370 Hadfield JD. 2010. MCMC methods for multi-response generalized linear mixed models: The
371 MCMCglmm R package. *Journal of Statistical Software* **33**:1–22.

- 372 Harmon L, Weir J, Brock C, Glor R, Challenger W. 2008. GEIGER: Investigating
373 evolutionary radiations. *Bioinformatics* **24**:129–131.
- 374 Heinze G, Ploner M. 2018. Logistf: Firth's bias-reduced logistic regression.
- 375 Henry L, Wickham H. 2018. Purrr: Functional programming tools.
- 376 Hijmans RJ. 2018. Raster: Geographic data analysis and modeling.
- 377 Ho LST, Ane C. 2014. A linear-time algorithm for gaussian and non-gaussian trait evolution
378 models. *Systematic Biology* **63**:397–408.
- 379 Kerr JT, Ostrovsky M. 2003. From space to species: Ecological applications for remote
380 sensing. *Trends in ecology & evolution* **18**:299–305.
- 381 Kogan F, Guo W, Strashnaia A, Kleshenko A, Chub O, Virchenko O. 2016. Modelling and
382 prediction of crop losses from noaa polar-orbiting operational satellites. *Geomatics,*
383 *Natural Hazards and Risk* **7**:886–900.
- 384 Kogan FN. 2001. Operational space technology for global vegetation assessment. *Bulletin of*
385 *the American Meteorological Society* **82**:1949–1964.
- 386 Lang PL, Willems FM, Scheepens J, Burbano HA, Bosdorf O. 2018. Using herbaria to
387 study global environmental change. PeerJ Preprints.
- 388 Lelièvre F, Volaire F. 2009. Current and potential development of perennial grasses in
389 rainfed mediterranean farming systems. *Crop Science* **49**:2371–2378.
- 390 Mandáková T, Mummenhoff K, Al-Shehbaz IA, Mucina L, Mühlhausen A, Lysak MA. 2012.
391 Whole-genome triplication and species radiation in the southern african tribe
392 heliophileae (brassicaceae). *Taxon* **61**:989–1000.

- 393 Mojica JP, Mullen J, Lovell JT, Monroe JG, Paul JR, Oakley CG, McKay JK. 2016.
394 Genetics of water use physiology in locally adapted *Arabidopsis thaliana*. *Plant*
395 *Science*. doi:10.1016/j.plantsci.2016.03.015
- 396 Monroe JG, Markman DW, Beck WS, Felton AJ, Vahsen ML, Pressler Y. 2018.
397 Ecoevolutionary dynamics of carbon cycling in the anthropocene. *Trends in ecology &*
398 *evolution* **33**:213–225.
- 399 Monroe JG, Powell T, Price N, Mullen J, Howard A, Evans K, Lovell J, McKay J. 2018.
400 Drought adaptation in nature by extensive genetic loss-of-function. *bioRxiv* 372854.
- 401 Morishima H, Sano Y, Oka H. 1984. Differentiation of perennial and annual types due to
402 habitat conditions in the wild rice *Oryza perennis*. *Plant Systematics and Evolution*
403 **144**:119–135.
- 404 Mummenhoff K, Al-Shehbaz IA, Bakker FT, Linder HP, Mühlhausen A. 2005. Phylogeny,
405 morphological evolution, and speciation of endemic brassicaceae genera in the cape
406 flora of southern africa. *Annals of the Missouri Botanical Garden* 400–424.
- 407 Müller K. 2018. Bindrcpp: An 'rcpp' interface to active bindings.
- 408 Müller K, Wickham H. 2018. Tibble: Simple data frames.
- 409 Orme D, Freckleton R, Thomas G, Petzoldt T, Fritz S, Isaac N, Pearse W. 2018. Caper:
410 Comparative analyses of phylogenetics and evolution in r.
- 411 Paradis E, Schliep K. 2018. Ape 5.0: An environment for modern phylogenetics and
412 evolutionary analyses in R. *Bioinformatics* **xx**:xxx–xxx.
- 413 Parry M, Flexas J, Medrano H. 2005. Prospects for crop production under drought:
414 Research priorities and future directions. *Annals of Applied Biology* **147**:211–226.

- 415 Pebesma EJ, Bivand RS. 2005. Classes and methods for spatial data in R. *R News* **5**:9–13.
- 416 Plummer M, Best N, Cowles K, Vines K. 2006. CODA: Convergence diagnosis and output
417 analysis for mcmc. *R News* **6**:7–11.
- 418 R by Thomas Lumley VJCP to, author. BRN that maintainers are not available to give
419 advice on using a package they did not. 2015. Gee: Generalized estimation equation
420 solver.
- 421 R Core Team. 2018. R: A language and environment for statistical computing. Vienna,
422 Austria: R Foundation for Statistical Computing.
- 423 Revell LJ. 2012. Phytools: An r package for phylogenetic comparative biology (and other
424 things). *Methods in Ecology and Evolution* **3**:217–223.
- 425 Richard A. Becker OS code by, Ray Brownrigg. Enhancements by Thomas P Minka ARWR
426 version by, Deckmyn. A. 2018. Maps: Draw geographical maps.
- 427 Rojas O, Vrieling A, Rembold F. 2011. Assessing drought probability for agricultural areas
428 in africa with coarse resolution remote sensing imagery. *Remote sensing of
429 Environment* **115**:343–352.
- 430 Roumet C, Urcelay C, Di'az S. 2006. Suites of root traits differ between annual and
431 perennial species growing in the field. *New phytologist* **170**:357–368.
- 432 Sayre RG, Comer P, Hak J, Josse C, Bow J, Warner H, Larwanou M, Kelbessa E, Bekele T,
433 Kehl H, others. 2013. A new map of standardized terrestrial ecosystems of africa.
434 *African Geographical Review*.
- 435 Schaffer W, Gadgil M. 1975. Selection for optimal life histories in plants. *Ecology and
436 evolution of communities* 142–157.

- 437 Slater G, Harmon L, Wegmann D, Joyce P, Revell L, Alfaro M. 2012. Fitting models of
438 continuous trait evolution to incompletely sampled comparative data using
439 approximate bayesian computation. *Evolution* **66**:752–762.
- 440 Soetaert K. 2018. Shape: Functions for plotting graphical shapes, colors.
- 441 Soetaert K. 2017. Diagram: Functions for visualising simple graphs (networks), plotting flow
442 diagrams.
- 443 Soltis PS. 2017. Digitization of herbaria enables novel research. *American journal of botany*
444 **104**:1281–1284.
- 445 Stearns SC. 1992. The evolution of life histories.
- 446 Stropp J, Ladle RJ, M. Malhado AC, Hortal J, Gaffuri J, H. Temperley W, Olav Skøien J,
447 Mayaux P. 2016. Mapping ignorance: 300 years of collecting flowering plants in africa.
448 *Global Ecology and Biogeography* **25**:1085–1096.
- 449 Tank DC, Olmstead RG. 2008. From annuals to perennials: Phylogeny of subtribe
450 castillejinae (orobanchaceae). *American Journal of Botany* **95**:608–625.
- 451 Václavík T, Beckmann M, Cord AF, Bindewald AM. 2017. Effects of uv-b radiation on leaf
452 hair traits of invasive plants—combining historical herbarium records with novel
453 remote sensing data. *PloS one* **12**:e0175671.
- 454 Venables WN, Ripley BD. 2002. Modern applied statistics with s, Fourth. ed. New York:
455 Springer.
- 456 Verboom GA, Linder HP, Stock WD. 2004. Testing the adaptive nature of radiation:
457 Growth form and life history divergence in the african grass genus ehrharta (poaceae:
458 Ehrhartoideae). *American Journal of Botany* **91**:1364–1370.

- ⁴⁵⁹ Wickham H. 2018a. Forcats: Tools for working with categorical variables (factors).
- ⁴⁶⁰ Wickham H. 2018b. Stringr: Simple, consistent wrappers for common string operations.
- ⁴⁶¹ Wickham H. 2017. Tidyverse: Easily install and load the 'tidyverse'.
- ⁴⁶² Wickham H. 2016. Ggplot2: Elegant graphics for data analysis. Springer-Verlag New York.
- ⁴⁶³ Wickham H, François R, Henry L, Müller K. 2018. Dplyr: A grammar of data manipulation.
- ⁴⁶⁴ Wickham H, Henry L. 2018. Tidyr: Easily tidy data with 'spread()' and 'gather()' functions.
- ⁴⁶⁵ Wickham H, Hester J, Francois R. 2017. Readr: Read rectangular text data.
- ⁴⁶⁶ Willis CG, Ellwood ER, Primack RB, Davis CC, Pearson KD, Gallinat AS, Yost JM, Nelson
⁴⁶⁷ G, Mazer SJ, Rossington NL, others. 2017. Old plants, new tricks: Phenological
⁴⁶⁸ research using herbarium specimens. *Trends in ecology & evolution* **32**:531–546.
- ⁴⁶⁹ Wolf A, Zimmerman NB, Anderegg WR, Busby PE, Christensen J. 2016. Altitudinal shifts
⁴⁷⁰ of the native and introduced flora of California in the context of 20th-century
⁴⁷¹ warming. *Global ecology and biogeography* **25**:418–429.

472

Supplement

473

Images used. <https://www.gbif.org/occurrence/1099023487>

474

<https://www.gbif.org/occurrence/1057389408> <https://www.gbif.org/occurrence/1099023562>

475

<https://www.gbif.org/occurrence/1099023490>

476

Software used. We used R (Version 3.5.1; R Core Team, 2018) and the R-packages

477 *ape* (Version 5.2; Paradis and Schliep, 2018; Orme et al., 2018; Soetaert, 2018), *bindrcpp*

478 (Version 0.2.2; Müller, 2018), *caper* (Version 1.0.1; Orme et al., 2018), *coda* (Version 0.19.2;

479 Plummer et al., 2006), *diagram* (Version 1.6.4; Soetaert, 2017), *dplyr* (Version 0.7.8;

480 Wickham et al., 2018), *forcats* (Version 0.3.0; Wickham, 2018a), *gee* (Version 4.13.19; R by

481 Thomas Lumley and author., 2015), *geiger* (Version 2.0.6; Alfaro et al., 2009; Eastman et al.,

482 2011; Harmon et al., 2008; Slater et al., 2012), *ggplot2* (Version 3.1.0; Wickham, 2016),

483 *logistf* (Version 1.23; Heinze and Ploner, 2018), *maps* (Version 3.3.0; Richard A. Becker et al.,

484 2018), *MASS* (Version 7.3.51.1; Venables and Ripley, 2002), *Matrix* (Version 1.2.15; Bates

485 and Maechler, 2018), *MCMCglmm* (Version 2.26; Hadfield, 2010), *mvtnorm* (Version 1.0.8;

486 Genz and Bretz, 2009), *papaja* (Version 0.1.0.9842; Aust and Barth, 2018), *phylobolm* (Version

487 2.6; Ho and Ane, 2014), *phytools* (Version 0.6.60; Revell, 2012), *purrr* (Version 0.2.5; Henry

488 and Wickham, 2018), *raster* (Version 2.8.4; Hijmans, 2018), *readr* (Version 1.2.1; Wickham

489 et al., 2017), *shape* (Version 1.4.4; Soetaert, 2018), *sp* (Version 1.3.1; Pebesma and Bivand,

490 2005), *stringr* (Version 1.3.1; Wickham, 2018b), *tibble* (Version 1.4.2; Müller and Wickham,

491 2018), *tidyverse* (Version 1.2.1; Wickham, 2017), and *tidyverse* (Version 1.2.1;

492 Wickham, 2017) for all our analyses.

493

Supplementary tables and figures.

Table S1

Heliophila species records and the mean drought frequencies during different seasons at the location of records

Species	LH	n	Winter	Spring	Summer	Fall
<i>Heliophila acuminata</i>	a	28	0.32	0.38	0.41	0.36
<i>Heliophila africana</i>	a	91	0.33	0.35	0.34	0.34
<i>Heliophila amplexicaulis</i>	a	60	0.32	0.36	0.39	0.33
<i>Heliophila arenaria</i>	a	65	0.34	0.37	0.38	0.34
<i>Heliophila carnosa</i>	p	129	0.33	0.37	0.39	0.31
<i>Heliophila cedarbergensis</i>	p	3	0.40	0.43	0.32	0.27
<i>Heliophila collina</i>	a	16	0.35	0.47	0.48	0.45
<i>Heliophila cornellsbergia</i>	a	2	0.33	0.42	0.35	0.21
<i>Heliophila cornuta</i>	p	101	0.35	0.40	0.40	0.34
<i>Heliophila coronopifolia</i>	a	40	0.37	0.42	0.40	0.37
<i>Heliophila crithmifolia</i>	a	97	0.35	0.42	0.45	0.38
<i>Heliophila descurva</i>	a	12	0.36	0.38	0.38	0.29
<i>Heliophila deserticola</i>	a	133	0.48	0.48	0.46	0.45
<i>Heliophila digitata</i>	a	30	0.33	0.38	0.44	0.38
<i>Heliophila dregeana</i>	p	17	0.33	0.37	0.33	0.32
<i>Heliophila elongata</i>	p	82	0.26	0.32	0.30	0.25
<i>Heliophila ephemera</i>	a	3	0.14	0.27	0.31	0.26
<i>Heliophila esterhuyseniae</i>	p	3	0.21	0.30	0.37	0.27
<i>Heliophila eximia</i>	p	12	0.42	0.41	0.32	0.34
<i>Heliophila gariepina</i>	a	12	0.50	0.53	0.48	0.41
<i>Heliophila glauca</i>	p	35	0.29	0.35	0.34	0.33
<i>Heliophila juncea</i>	p	150	0.32	0.37	0.39	0.35
<i>Heliophila linearis</i>	p	94	0.32	0.33	0.28	0.30
<i>Heliophila macowaniana</i>	a	31	0.33	0.38	0.44	0.39
<i>Heliophila macra</i>	p	22	0.30	0.30	0.32	0.29
<i>Heliophila macrosperma</i>	p	5	0.28	0.36	0.35	0.25
<i>Heliophila minima</i>	p	35	0.36	0.45	0.51	0.39

<i>Heliophila namaquana</i>	a	16	0.39	0.46	0.48	0.39
<i>Heliophila nubigena</i>	p	19	0.31	0.36	0.43	0.38
<i>Heliophila pectinata</i>	a	16	0.27	0.34	0.50	0.34
<i>Heliophila polygaloides</i>	p	12	0.40	0.48	0.42	0.34
<i>Heliophila pubescens</i>	a	9	0.31	0.40	0.48	0.39
<i>Heliophila pusilla</i>	a	45	0.32	0.38	0.38	0.34
<i>Heliophila rigidiuscula</i>	p	201	0.30	0.33	0.28	0.24
<i>Heliophila scoparia</i>	p	106	0.31	0.37	0.36	0.31
<i>Heliophila seselifolia</i>	a	80	0.36	0.42	0.45	0.40
<i>Heliophila suavissima</i>	p	92	0.30	0.39	0.42	0.31
<i>Heliophila subulata</i>	p	103	0.29	0.33	0.31	0.29
<i>Heliophila tricuspidata</i>	p	8	0.28	0.33	0.38	0.30
<i>Heliophila trifurca</i>	a	77	0.45	0.48	0.48	0.43
<i>Heliophila tulbaghensis</i>	p	3	0.36	0.41	0.36	0.35
<i>Heliophila variabilis</i>	a	97	0.35	0.41	0.40	0.37

Note. LH = Life history (a = annual, p = perennial). n=sample size of GBIF records. Seasons are mean drought frequencies observed at locations of records.



Figure S1. Maps of occurrence records for individual species. Orange points indicate annual species. Blue points indicate perennial species.