Foreman notebook, §15

The centre ray read off the list

22 October 2026

This entry follows §14.17 of the running notebook, which was not available in this session; it is written to be appended there. Handbook citations are by printed number and page of the handbook PDF of 20 October 2026; "the note" is the tree-coaction note of 22 October 2026. Every object is a member of the serif family or a construction of the handbook, written with its slots. The receipts are in the script centre_ray.py.

15 The centre ray read off the list

The neighbours of the five-mark centre are the values of the centre ray of the list (Def. 35.9, p. 309), the period $P_{\mathcal{B}}(\sigma\mathbf{1};\sigma,\sigma)$. The interest of the list is one Gramma (Thm. 8.3, p. 54), the period is its residue on the balance wall, and the residue is the Gramma of the letter columns (Thm. 2.9, p. 15). Its share side (Prop. 4.2, p. 25) is the working form of this entry: a fibre integral of Gammas, one per basis for each letter, pinned by equal incidence. The pin is what the list contributes. Closing the fibre at the negative strengths (Thm. 4.5, p. 26) lands on the pairs of k-tuples of bases with equal incidence; the residue at zero shares is the corner of Thm. 38.5(i); the native shares (Thm. 4.3, p. 25) are uniform, the share side of the table's gauge; and the banana is the list whose pin identifies the two letters mark by mark, which turns its share side into the Mellin read of a power of one Gramma.

15.1 The centre period on its share side

Proposition 15.1 (The share side of the centre)**.** [derived]*; receipts* [computed] *Let* $(\mathcal{E},\mathcal{B})$ *be a list,* $n=|\mathcal{E}|\geq 2$*,* $e_0\in\mathcal{E}$*, and* $\operatorname{Re}\sigma>0$*. Then*

$$\Gamma(\sigma)^2\,P_{\mathcal{B}}(\sigma\,\mathbf{1};\sigma,\sigma)=\int_{\substack{|\mathbf{a}|=\sigma,\ |\mathbf{b}|=\sigma\\ \sum_T a_T\mathbf{1}_T=\sum_T b_T\mathbf{1}_T}}\ \prod_{T\in\mathcal{B}}\Gamma(a_T)\,\Gamma(b_T),$$

the fibre integral of Def. 4.1, p. 25, with one share a_T *for the letter* K *and one share* b_T *for the letter* Ψ *at every basis, pinned by the two letter totals and by the incidence at the marks* $e\neq e_0$*; it has dimension* $2|\mathcal{B}|-n-1$*, and the uniform shares* $a_T=b_T=\sigma/|\mathcal{B}|$ *are an admissible real point.*

Proof. By Thm. 8.3, p. 54, $\Gamma(s)\Gamma(t)\,J_{\mathcal{B}}(\mathbf{m};s,t)=\mathsf{G}_{(\boldsymbol{\mu};s;t);t_{\mathcal{B}}}(\mathbf{1})$ with the unit columns $(\mathbf{e}_e;0;0)$ and the letter columns $(\chi_T/2;1;0)$, $(-\chi_T/2;0;1)$. The linear form $\ell(x;v;w)=\sum_e x_e-\frac{2r-n}{2}(v-w)$ is 1 on the unit columns and 0 on the letter columns, and at the index it is $\kappa_{\mathcal{E}}$ (Thm. 8.2(iii), p. 54): the balance wall is the first wall of this Gramma along ℓ, with the letter columns as facet terms. By Thm. 8.10, p. 56, the period is the residue of $J_{\mathcal{B}}$ there, and by Thm. 2.9, p. 15, at $p=0$ that residue is the Gramma of the facet terms. Its share side (Prop. 4.2) has one share per letter column; the pins of the two letter scales are $|\mathbf{a}|=s$, $|\mathbf{b}|=t$, and those of the marks at $\boldsymbol{\mu}=0$ are $\sum_T(a_T-b_T)\chi_T/2\in\mathbb{C}\mathbf{1}$,

which with $|\mathbf{a}| = |\mathbf{b}|$ is $\sum_T (a_T - b_T)\mathbf{1}_T = 0$. At the centre $s = t = \sigma$. The uniform shares satisfy the pins and have positive real parts. □

Receipts. For $\binom{[2]}{1}$ the fibre has dimension one and the integral is four Gammas on a line (Cor. 4.7, p. 26),

$$\frac{1}{2\pi i}\int \Gamma(a)^2\,\Gamma(\sigma-a)^2\,da = \Gamma_{\,1}^{\,4}\begin{bmatrix}\sigma,\ \sigma,\ \sigma,\ \sigma\\ 2\sigma\end{bmatrix}, \qquad \text{so} \qquad P_{\mathcal{B}}(\sigma\,\mathbf{1};\sigma,\sigma) = \Gamma_{\,1}^{\,2}\begin{bmatrix}\sigma,\ \sigma\\ 2\sigma\end{bmatrix} \quad \text{[derived]}.$$

For $\binom{[3]}{1}$ at $\sigma = 1$ the two-dimensional fibre integral is $2.343907238689463\ldots$, the value of the receipts table below, to fourteen digits [computed].

Proposition 15.2 (The banana's pin)**.** [derived]*; receipt* [computed] *For* $\binom{\mathcal{E}}{1}$ *the pin is* $a_e = b_e$ *at every mark, so each mark carries* $\Gamma(a_e)^2$*, which is the share kernel of the Gramma* $\mathsf{G}_{0;(1,-1)}(1,y) = \int_0^\infty x^{-1}e^{-x-y/x}dx$ *(Prop. 4.2), and*

$$\Gamma(\sigma)^2\,P_{\mathcal{B}}(\sigma\,\mathbf{1};\sigma,\sigma) = \int_{|\mathbf{a}|=\sigma}\prod_{e\in\mathcal{E}}\Gamma(a_e)^2 = \int_0^\infty y^{\sigma-1}\,\mathsf{G}_{0;(1,-1)}(1,y)^n\,dy.$$

At $n = 5$*,* $\sigma = 1$ *the right side is* $39.945588011996245699\ldots$ [computed].

Proof. With $T = \{e\}$ the pins are $a_e - b_e = 0$ for $e \neq e_0$ and the two totals, so $a = b$ and the fibre is the share integral $|\mathbf{a}| = \sigma$ (Def. 4.1). The share side of $\mathsf{G}_{0;(1,-1)}(1,y)$ is $\frac{1}{2\pi i}\int\Gamma(c)^2y^{-c}dc$, and the read $\int_0^\infty y^{\sigma-1}(\cdot)\,dy$ of a product of n such fibre integrals pins the sum of their shares at σ. □

So the "Bessel moments" are the banana's share side: the pin of a single mark turns the two letters into one Gramma per mark (alias $\mathsf{G}_{0;(1,-1)}(1,y) = 2K_0(2\sqrt{y})$). A general list keeps a share per basis and per letter and couples them only through the incidence; that coupling is the broader arena.

Proposition 15.3 (The corner is the residue at zero shares)**.** [derived] *With* $U_k = \{\pi(k),\ldots,\pi(n)\}$ *the tails of an order* π *and* c *the width (Def. 7.10, p. 49),*

$$\lim_{\sigma\downarrow 0}\sigma^{n-1}P_{\mathcal{B}}(\sigma\,\mathbf{1};\sigma,\sigma) = \sum_\pi\prod_{k=2}^{n}c(U_k)^{-1},$$

and $\sigma^{n-1}P_{\mathcal{B}}(\sigma\,\mathbf{1};\sigma,\sigma)$ *has no term of order* σ*. This is Thm. 38.5(i), p. 330, at the centre question, whose walls are* $\sigma c(S)$ *(Prop. 8.21(ii), p. 59); the term of order* σ *there is a sum of* $\log\theta$ *at greedy bases, zero for a list. On the share side,* $\sigma^{n+1}\Gamma(\sigma)^2P_{\mathcal{B}}(\sigma\,\mathbf{1};\sigma,\sigma)$ *tends to the same number: it is the residue of the fibre integral of Proposition 15.1 at the origin, where all* $2|\mathcal{B}|$ *Gammas have their first pole. For* $\binom{\mathcal{E}}{1}$ *the corner is* $n!$*, the* $5!$ *of §14.17; for* $\binom{[5]}{2}$ *the tails of every order have widths* $1,2,2,1$ *and it is* 30*.*

Proposition 15.4 (Closing the centre at the negative strengths)**.** [derived] *For* $k\in\mathbb{N}$*, at* $\sigma = -k$ *the share side closes (Thm. 4.5, p. 26) onto the lattice of pairs* $(\boldsymbol{\alpha},\boldsymbol{\beta})\in\mathbb{N}^{\mathcal{B}}\times\mathbb{N}^{\mathcal{B}}$ *with* $|\boldsymbol{\alpha}| = |\boldsymbol{\beta}| = k$ *and* $\sum_T\alpha_T\mathbf{1}_T = \sum_T\beta_T\mathbf{1}_T$*, where every Gamma is at a pole,* $a_T = -\alpha_T$*,* $b_T = -\beta_T$*, with the residues* $\prod_T(-1)^{\alpha_T+\beta_T}/(\alpha_T!\,\beta_T!)$ *(Prop. 1.3(iv), p. 11). The continued period has a pole of order at most* $n-1$ *at* $\sigma = -k$*, and*

$$\begin{aligned}\lim_{\sigma\to -k}(\sigma+k)^{n-1}\,P_{\mathcal{B}}(\sigma\,\mathbf{1};\sigma,\sigma) &= \sum_\pi\prod_{k=2}^{n}c(U_k)^{-1}\cdot\sum_{(\boldsymbol{\alpha},\boldsymbol{\beta})}\frac{(k!)^2}{\prod_T\alpha_T!\,\beta_T!}\\ &= \sum_\pi\prod_{k=2}^{n}c(U_k)^{-1}\cdot[\alpha^0]\big(K(\alpha)\Psi(\alpha)/\alpha^{\mathcal{E}}\big)^k,\end{aligned}$$

the sum over that lattice: it counts the pairs of k*-tuples of bases with equal incidence.*

Proof. On the balance wall the period is $\sum_\pi \int_{[0,1]^{n-1}} \prod_{k'=2}^{n} q_{k'}^{\kappa_{U_{k'}}-1} h_\pi(q)\,dq$ in the sector coordinates $\alpha_{\pi(k')} = \rho\, q_2 \cdots q_{k'}$ (Thm. 8.7(ii), p. 55, and Thm. 8.10). At the centre $\kappa_{U_{k'}} = \sigma c(U_{k'})$ and $h_\pi = (k_\pi \psi_\pi)^{-\sigma}$, and $k_\pi\psi_\pi = \prod_{k'} q_{k'}^{c(U_{k'})} K(\alpha)\Psi(\alpha)/\alpha^{\mathcal{E}}$ is a polynomial in q with nonnegative coefficients and constant term one (Lemma 8.5, p. 55, weights one). By Lemma 8.6, p. 55, each cube integral continues with at most a simple pole in each $\kappa_{U_{k'}}$ at $-\mathbb{N}$, and at $\sigma = -k$ the $n-1$ poles meet only through the term q^{kc}, $c = (c(U_{k'}))_{k'}$, of $(k_\pi\psi_\pi)^k (k_\pi\psi_\pi)^{-(\sigma+k)}$, with coefficient $\prod_{k'} c(U_{k'})^{-1} [q^{kc}](k_\pi\psi_\pi)^k$. The map $\alpha^v \mapsto q^{(v(U_{k'}))_{k'}}$ is injective on exponents of degree zero, so $[q^{kc}](k_\pi\psi_\pi)^k = [\alpha^0]\big(K(\alpha)\Psi(\alpha)/\alpha^{\mathcal{E}}\big)^k$. Expanding each letter's k-th power (the share kernel of Cor. 4.4, p. 26, closed at the exponent $-k$) gives the lattice sum. On the share side the factor $(k!)^{-2}$ is the residue of $\Gamma(\sigma)^2$ at $-k$, and the common factor of the lattice points is the residue at zero shares, the corner (Proposition 15.3). □

Corollary 15.5 (One law for the period and its closing)**.** [derived] *If polynomials $c_0, \dots, c_d$ satisfy*

$$\sum_{i=0}^{d} c_i(\sigma)\, P_{\mathcal{B}}((\sigma+i)\,\mathbf{1}; (\sigma+i), (\sigma+i)) = 0$$

identically, then for every $k \in \mathbb{N}$

$$\sum_{i=0}^{\min(d,k)} c_i(-k)\, [\alpha^0]\big(K(\alpha)\Psi(\alpha)/\alpha^{\mathcal{E}}\big)^{k-i} = 0.$$

So the least law of the period in its strength is at least as long as the least recurrence of its closings, and the closings obey the reflected law from $k = 0$, with nothing below.

Proof. Multiply by $(\sigma+k)^{n-1}$ and let $\sigma \to -k$: the terms with $i \le k$ tend to $c_i(-k)$ times the corner times the closing at $k-i$ by Proposition 15.4, and the others are regular there. □

The laws come from the family's own relations. By the letter presentation (Prop. 3.6, p. 20) the centre period is a Breta with one bunch for the shape and one singleton bunch per letter; its letter splits and scale exchanges (Thm. 3.5, p. 19) are the letter and mark relations of the interest (Thm. 8.3(i)–(iii)), and along the centre ray they are the relations of Thm. 35.3, p. 308, whose eliminations are the laws of Cor. 35.6. The converse of Corollary 15.5, that the least recurrence of the closings, read at $k = -\sigma$, is a law of the period, is not derived here; for the banana through seven marks it is not needed, the laws being derived by adding marks (Theorem 15.25). For every list below the minimal solution it gives agrees with an independent quadrature to fourteen digits, and the closings obey it from $k = 0$ with nothing below [computed].

Procedure 15.6 (The centre ray from the list)**.** (i) Close the share side at $\sigma = -k$: $x_k = [\alpha^0]\big(K(\alpha)\Psi(\alpha)/\alpha^{\mathcal{E}}\big)^k$, $k \le k_{\max}$, the letter x_k abbreviating them in this procedure. (ii) Find their least recurrence $\sum_{j=0}^{d} p_j(k) x_{k+j} = 0$ by exact linear algebra and check it on the surplus equations. (iii) Read it at $k = -\sigma$: $c_i(\sigma) = p_{d-i}(-\sigma - d)$, $\sum_i c_i(\sigma) P_{\mathcal{B}}((\sigma+i)\,\mathbf{1}; (\sigma+i), (\sigma+i)) = 0$. (iv) The period is its minimal solution: $P_{\mathcal{B}}((\sigma+1)\,\mathbf{1}; (\sigma+1), (\sigma+1))/P_{\mathcal{B}}(\sigma\,\mathbf{1}; \sigma, \sigma) \to K(\mathbf{1})^{-2}$ (Prop. 35.7, p. 309), the native value of least modulus. (v) Scale it by the residue at zero shares: c_0 has a zero of order $n-1$ at 0, and $[\sigma^{n-1}] c_0 \cdot \sum_\pi \prod_{k=2}^{n} c(U_k)^{-1} + \sum_{i \ge 1} c_i(0) P_{\mathcal{B}}(i\,\mathbf{1}; i, i) = 0$.

Receipts [computed], no quadrature in the procedure:

list	$P_{\mathcal{B}}(\mathbf{1};1,1)$	reading
$\binom{[3]}{1}$	$2.34390723868945889060156 2\ldots$	$3L(\chi_{-3},2)$ (alias)
$\binom{[4]}{1}$	$8.41439832211715999779816 7\ldots$	$(\mathsf{T}_{3;(1)}[x]\,\mathbb{I}_{\frac12+\mathbb{Z}})((0,\infty)) = 7\,(\mathsf{T}_{3;(1)}[x]\,\mathbb{I}_{\mathbb{Z}})((0,\infty))$
$\binom{[4]}{2}$	$2.34390723868945889060156 2\ldots$	as $\binom{[3]}{1}$: Cor. 19.17 (p. 143) and duality
$\binom{[5]}{1}$	$39.94558801199624569894402\ldots$	$M_5(1)$ of §14.17

The four-mark values are those of the note (Ex. 6.3 and 6.2, p. 39). There they are closings of the share side at positive strength: the Gammas of a basis meet in double poles, whose Laurent data are Trailmoments of the counting input and of lattice inputs (Prop. 5.14, p. 33; Lemma 35.20, p. 314).

15.2 The masters are the signs

A native gauge of the centre ray is a point where its direction $(\mathbf{1};1,1)$ is native (Def. 35.4, p. 308): the defect $1-R_e(\alpha)-R_e^*(\alpha)$ vanishes at every mark, and since $R_e^*(\alpha)=1-R_e(\alpha^{-1})$ (Thm. 7.14, p. 50) this reads

$$R_e(\alpha)=R_e(\alpha^{-1})\qquad(e\in\mathcal{E}):$$

inverting the gauge leaves the occupation unchanged. Its native value is $\big(K(\alpha)K(\alpha^{-1})\big)^{-1}$.

Proposition 15.7 (The banana has only signs)**.** [derived] *The native gauges of the centre ray of* $\binom{\mathcal{E}}{1}$ *are exactly the signs* $\eta\in\{\pm1\}^{\mathcal{E}}$, $\eta_{e_0}=1$, *with* $K(\eta)=\sum_e\eta_e\neq0$*; their native values are* $K(\eta)^{-2}$ *(Thm. 35.12, p. 310).*

Proof. For $K=\sum_e\alpha_e$ the occupation is $R_e(\alpha)=\alpha_e/K(\alpha)$, so $R_e(\alpha)=R_e(\alpha^{-1})$ for every e says $\alpha_e^2=K(\alpha)/K(\alpha^{-1})$ for every e. □

So along the banana sequence the native values are $(n-2j)^{-2}$, j the number of minus signs, $\lceil n/2\rceil$ of them, all simple. At $n=5$ they are $\frac{1}{25},\frac19,1$, the symbol of the ray law vanishes there (Cor. 35.6, p. 308), the order is three, and the corner leaves two masters. Every list has the signs with $K(\eta)\neq0$ among the native gauges of its centre ray; what is special about $\binom{\mathcal{E}}{1}$ is that it has nothing else.

Remark 15.8 (The lowered settings of §14.17). With $g=\mathsf{G}_{0;(1,-1)}(1,\cdot)$ and $h(y)=\sum_m y^m/\Gamma(1+m)^2$, the settings $x^{(j)}(s)=\int_0^\infty y^{s-1}g^{5-j}h^j\,dy$, $j=0,1,2$, satisfy the ray law in the normalization $M(s)=\Gamma(s)^2P_{\mathcal{B}}(s\,\mathbf{1};s,s)$, and $x^{(0)}=M$ is Proposition 15.2. Their corners are $120,0,0$ [computed], and $x^{(j)}(s+1)/\big(s^2x^{(j)}(s)\big)\to(5-2j)^{-2}$: they are the sign classes with j minus signs. "The g's outnumber the h's" is $K(\eta)\neq0$; the tie at four marks is $K(\eta)=0$, which is not a native gauge.

Proposition 15.9 (The lowered-setting block is the frame)**.** [derived]*; receipt* [computed] *Let* $\Phi(s)=\det\big[x^{(i)}(s+l)\big]_{i,l=0}^{2}$. *By Prop. 35.8, p. 309, applied to the ray law of* M *(end coefficients* s^6 *and* -225*),* $\Phi(s+1)=\frac{s^6}{225}\Phi(s)$ *and* $\Phi(s)=\Phi(1)\,\Gamma(s)^6\,225^{1-s}$. *The amplitudes at the three sign classes,* $x^{(j)}(s)\sim2\pi^{(5-2j)/2}2^{-j}\rho_j^{5/2}\,\Gamma(2s-\frac52)\,\rho_j^{-2s}$ *with* $\rho_j=2(5-2j)$*, give*

$$\Phi(1)=\frac{128\sqrt{15}}{16875}\,\pi^3=0.91088056479820303289039718797 5\ldots,$$

which the settings computed directly (quadrature, as a check) reproduce to thirty digits. The corners reduce the third column, so

$$\Phi(1)=\frac{120}{225}\Big(x^{(1)}(1)\,x^{(2)}(2)-x^{(1)}(2)\,x^{(2)}(1)\Big),\qquad x^{(1)}(1)x^{(2)}(2)-x^{(1)}(2)x^{(2)}(1)=\frac{16\sqrt{15}}{1125}\,\Gamma(\tfrac12)^6:$$

the block of §14.17 is the frame of the ray over its corner. The masters do not enter Φ*; their coefficients are minors that the zero corners of the lowered settings kill. So the frame does not shorten the last minor of §14.17.*

15.3 A parallel class closes through the banana

Proposition 15.10. [derived] *Let $\mathcal{B}$ be obtained from a list $\mathcal{H}$ on the marks $1,\dots,m$ by replacing mark i with a parallel class P_i of s_i marks, so that $K_{\mathcal{B}}(\alpha) = K_{\mathcal{H}}\big((\alpha(P_i))_i\big)$ with $\alpha(P_i) = \sum_{e\in P_i}\alpha_e$. Then $[\alpha^v]K_{\mathcal{B}}^k = \big([A^w]K_{\mathcal{H}}^k\big)\prod_i \big(w_i!/\prod_{e\in P_i} v_e!\big)$ with $w_i = v(P_i)$, and the closings of $\mathcal{B}$ are*

$$\sum_{w}\big([A^w]K_{\mathcal{H}}(A)^k\big)^2\prod_i[\alpha^0]\Big(K_{P_i}(\alpha)\Psi_{P_i}(\alpha)/\alpha^{P_i}\Big)^{w_i},$$

with $K_{P_i} = \sum_{e\in P_i}\alpha_e$ and Ψ_{P_i} the letters of the banana $\binom{P_i}{1}$: its closings at $-w_i$.

The banana enters the closing only through its own closings, as the parallel node enters the network through its full leans (Cor. 25.9, p. 209).

15.4 The five-mark centres

All connected lists on five marks, up to duality, which leaves the centre unchanged (Prop. 8.14, p. 57); "doubled" and "tripled" are parallel classes. [computed]

list	$\lvert\mathcal{B}\rvert$	$K(\eta)^2$ at the signs (mult.)	roots of the symbol	order	corner	masters
$\binom{[5]}{1}$	5	25 (1), 9 (5), 1 (10)	$25, 9, 1$	3	120	2
$\binom{[4]}{2}$, a mark doubled	9	81 (1), 9 (6), 1 (9)	$81, 9, 1$	3	36	2
$\binom{[5]}{2}$	10	100 (1), 4 (15)	$100, 4, 1, -8$	4	30	3
$\binom{[3]}{2}$, two marks doubled	8	64 (1), 16 (2), 4 (8), 0 (5)	$64, 16, 4, 4, 1$	5	44	$3^\dagger$
$\binom{[3]}{2}$, a mark tripled	7	49 (1), 25 (1), 9 (3), 1 (11)	$49, 25, 9, 1, 1, 1$	6	60	$4^\dagger$

list	$P_{\mathcal{B}}(\mathbf{1};1,1)$
$\binom{[5]}{1}$	39.945588011996245698944028498423995891849782924 77…
$\binom{[4]}{2}$, a mark doubled	5.493901143580844012444257607820691195853638640 6801…
$\binom{[5]}{2}$	4.029714170711049607320695734428406933139867563 6396…
$\binom{[3]}{2}$, two marks doubled	7.865837368364021592421974697446168210982382809 4581…
$\binom{[3]}{2}$, a mark tripled	12.848722141915555957312339931253534636619123934 709…

The roots of the symbol are those of the recurrence of the closings, the values $K(\alpha)K(\alpha^{-1})$ at the native gauges; the ray law of the period has their reciprocals. "Masters" is $m-1$ for the least m with a rational relation among $1, P_{\mathcal{B}}(\mathbf{1};1,1),\dots,P_{\mathcal{B}}(m\,\mathbf{1};m,m)$ (PSLQ at 120 digits, coefficients below 10^{15}, values good to 150 digits): over $\mathbb{Q}$, for the symmetric insertion, on the centre ray. In the first three rows the one relation is the corner.

† In the last two rows c_0 and c_1 both carry $(\sigma+1)^2$, so the ray law at $\sigma=-1$ reaches the period at neither -1 nor 0, and it is a second relation, a *second corner*: the boundary term, the constant term at $\sigma=-1$ (Thm. 1.6, p. 12),

$$\operatorname*{CT}_{\sigma=-1}\Big(c_0(\sigma)P_{\mathcal{B}}(\sigma\,\mathbf{1};\sigma,\sigma)+c_1(\sigma)P_{\mathcal{B}}((\sigma{+}1)\,\mathbf{1};(\sigma{+}1),(\sigma{+}1))\Big) = -\sum_{i\ge 2}c_i(-1)\,P_{\mathcal{B}}((i{-}1)\,\mathbf{1};(i{-}1),(i{-}1)),$$

is rational [computed]. In lowest terms the two relations read

$$2503 - 566\,P_{\mathcal{B}}(\mathbf{1};1,1) + 88248\,P_{\mathcal{B}}(2\,\mathbf{1};2,2) - 908872\,P_{\mathcal{B}}(3\,\mathbf{1};3,3) + 1425600\,P_{\mathcal{B}}(4\,\mathbf{1};4,4) = 0,$$
$$13800002 - 1848045\,P_{\mathcal{B}}(\mathbf{1};1,1) + 227950815\,P_{\mathcal{B}}(2\,\mathbf{1};2,2) - 3601110627\,P_{\mathcal{B}}(3\,\mathbf{1};3,3)$$
$$+ 11177654265\,P_{\mathcal{B}}(4\,\mathbf{1};4,4) - 8565102000\,P_{\mathcal{B}}(5\,\mathbf{1};5,5) = 0,$$

so the masters are $5-2=3$ and $6-2=4$. Write $c_0=(\sigma+1)^2A$ and $c_1=(\sigma+1)^2B$. Proposition 15.4 at $\sigma=-1$ requires $A(-1)\,|\mathcal{B}|+B(-1)=0$, the closing at 1 being $|\mathcal{B}|$; it holds, $B(-1)/A(-1)=-8$ and -7 [computed]. There the recursion of Corollary 15.5 leaves the corner at -1 free, so this is a check of Proposition 15.4 that the recursion does not supply.

$\binom{[5]}{2}$ has native gauges off the signs: on the pattern $\alpha\propto(a,a,b,b,1)$ the native equations have solutions with $K(\alpha)K(\alpha^{-1})=1$ and -8. There is no rational relation among 1, the two masters of $\binom{[5]}{1}$ and the three of $\binom{[5]}{2}$ (coefficients below 10^8, 150 digits), nor with π^2 adjoined.

15.5 The native shares, the roots, and an exact pairing

Every root of the symbol of the closings' recurrence is the reciprocal native value $K(\alpha_c)K(\alpha_c^{-1})$ of a native gauge α_c of the centre ray (Cor. 35.6, p. 308).

Proposition 15.11 (Both reads sit at the native shares)**.** [derived]*; receipts* [computed] *Let the list be connected, and let* $H=2\big(\kappa_{\mathbf{1}}(X_e,X_{e'})\big)_{e,e'\neq e_0}$*, twice the covariances of the indicators under the weight law (Def. 9.2, p. 61), the Hessian of Thm. 35.11, p. 310.*

(i) On the real fibre of Proposition 15.1 the native shares of Thm. 4.3(ii), p. 25, are the uniform shares $a_T=b_T=\sigma/|\mathcal{B}|$*: every basis takes the same share from each letter. This is the table's gauge* $\mathbf{1}$ *read on the share side.*

(ii) The period reads the fibre along its imaginary directions. By Thm. 4.3(iii), which at the period is Thm. 38.5(ii), p. 330,

$$P_{\mathcal{B}}(k\,\mathbf{1};k,k)\sim K(\mathbf{1})^{-2k}\Big(\frac{2\pi}{k}\Big)^{\frac{n-1}{2}}(\det H)^{-\frac12}.$$

(iii) The closing reads the same fibre on its integer points. Dropping the pin, the multinomial theorem gives $\sum_{|\boldsymbol{\alpha}|=|\boldsymbol{\beta}|=k}(k!)^2/\prod_T\alpha_T!\beta_T!=K(\mathbf{1})^k\Psi(\mathbf{1})^k=|\mathcal{B}|^{2k}$*; the pin keeps the Gaussian of the fibre at the native shares scaled by* k*, and*

$$[\alpha^0]\big(K(\alpha)\Psi(\alpha)/\alpha^{\mathcal{E}}\big)^k\sim K(\mathbf{1})^{2k}(2\pi k)^{-\frac{n-1}{2}}(\det H)^{-\frac12}.$$

(iv) So $k^{n-1}\,P_{\mathcal{B}}(k\,\mathbf{1};k,k)\,[\alpha^0]\big(K(\alpha)\Psi(\alpha)/\alpha^{\mathcal{E}}\big)^k\to(\det H)^{-1}$*: one Gaussian, read once along a contour and once on a lattice, so that the powers of* 2π *cancel and the determinant appears twice.*

Proof. (i) The uniform point satisfies the pins, and $\nabla S=-\log\mathbf{w}$ is constant there, a combination of the two rows of the letter totals; S is strictly concave on the real fibre (Thm. 4.3(ii)). (ii) Thm. 38.5(ii) at the centre: $X(0)=-2\log K(\mathbf{1})$ and $-\nabla_z^2X(0)=8\kappa_{\mathbf{1}}=4H$ on the chart, $y=2z$; Thm. 35.11 is the same edge read on the generating function of the ray. (iii) The terms $(k!)^2/\prod\alpha_T!\beta_T!$ on the pinned lattice peak at the uniform shares scaled by k; their Gaussian there is the quadratic form of $\sum\log\Gamma$ at the native shares restricted to the fibre, the same form as in (ii), now summed over the integer points of the fibre; $\nabla_z^2E(0)=8\kappa_{\mathbf{1}}$ (Thm. 7.14, p. 50) identifies it with H. □

Receipts [computed]: $(\det H)^{-1}=\frac{3125}{16},\frac{729}{16},\frac{3125}{81},\frac{512}{9},\frac{16807}{192}$ in the order of the table of §15.4. At $k=70$ the product agrees with these to about 10^{-4}, the error falling like k^{-2}. For $\binom{[2]}{1}$ the product is exact at every k: $k\,\Gamma\begin{bmatrix}k,\ k\\ 2k\end{bmatrix}\binom{2k}{k}=2=(\det H)^{-1}$ [derived].

Remark 15.12 (The signs are the pin read modulo two). Averaging over the signs detects the incidence modulo two: for every k,

$$2^{-n}\sum_{\eta\in\{\pm1\}^{\mathcal{E}}}K(\eta)^{2k}=\sum_{\substack{|\boldsymbol{\alpha}|=|\boldsymbol{\beta}|=k\\ \sum_T\alpha_T\mathbf{1}_T\equiv\sum_T\beta_T\mathbf{1}_T\ (2)}}\frac{(k!)^2}{\prod_T\alpha_T!\,\beta_T!}\ \ \geq\ [\alpha^0]\big(K(\alpha)\Psi(\alpha)/\alpha^{\mathcal{E}}\big)^k\quad[\text{derived}],$$

since $K(\eta)\Psi(\eta)/\eta^{\mathcal{E}}=K(\eta)^2$. So the sign values $K(\eta)^2$ are the growth rates of the pin's reduction modulo two, with $K(\mathbf{1})^2$ the largest. The full pin keeps them as roots (Thm. 35.12) and adds the native gauges off the signs, below.

Proposition 15.13 (An exact pairing of the period with its closing for the five-mark banana)**.** [derived] *from the ray law; receipt* [computed]. *For* $\binom{[5]}{1}$ *the ray law* $\sum_{i=0}^{3}c_i(\sigma)P_{\mathcal{B}}((\sigma+i)\,\mathbf{1};(\sigma+i),(\sigma+i))=0$ *has*

$$c_0=-\sigma^4,\qquad c_1=35\sigma^4+70\sigma^3+63\sigma^2+28\sigma+5,$$
$$c_2=-(\sigma+1)^2(259\sigma^2+518\sigma+285),\qquad c_3=225(\sigma+1)^2(\sigma+2)^2,$$

and it is self-adjoint under the reflection of the strength: $c_i(-\sigma-i)=c_i(\sigma)$ *for every* i. *So the closings, read as a function of* σ, *solve the adjoint law, and for every* $\sigma\in\{1,2,3,\dots\}$

$$\sum_{i=1}^{3}\sum_{j=1}^{i}c_i(\sigma-j)\ [\alpha^0]\big(K(\alpha)\Psi(\alpha)/\alpha^{\mathcal{E}}\big)^{\sigma-j}\ P_{\mathcal{B}}((\sigma-j+i)\,\mathbf{1};(\sigma-j+i),(\sigma-j+i))=120,$$

the residue at zero shares, with the closings at negative indices read as 0.

Proof. The left side, $B(\sigma)$ in this proof, is the bilinear concomitant of the law and its adjoint: $B(\sigma+1)-B(\sigma)$ is the law applied to the period times the closing at σ, less the period at σ times the adjoint law applied to the closings. The first vanishes by the law. The adjoint law at σ is $\sum_i c_i(\sigma-i)x_{\sigma-i}$, which by $c_i(\sigma-i)=c_i(-\sigma)$ is the recurrence of Corollary 15.5 at $k=\sigma$, zero from $k=0$ on. At $\sigma=1$ only $j=1$ survives, and $B(1)=\sum_i c_i(0)P_{\mathcal{B}}(i\,\mathbf{1};i,i)=120$ is the corner relation of Procedure 15.6(v). The self-adjointness is an identity of the four polynomials; $B(\sigma)=120$ holds to seventy digits for $\sigma\leq55$ [computed]. □

At large σ the closings and the period move by the factors 25 and $\frac{1}{25}$ per step, so $B(\sigma)\to\lambda\chi'(\lambda)\cdot\lim_k k^4\,P_{\mathcal{B}}(k\,\mathbf{1};k,k)\,[\alpha^0]\big(K(\alpha)\Psi(\alpha)/\alpha^{\mathcal{E}}\big)^k$, with $\chi(\lambda)=\sum_i[\sigma^4]c_i\,\lambda^i=-1+35\lambda-259\lambda^2+225\lambda^3$ the symbol and $\lambda=\frac{1}{25}$. With Proposition 15.11:

$$120\cdot\det H=\lambda\chi'(\lambda)\big|_{\lambda=1/25}=\frac{384}{625},\qquad\det H=\frac{16}{3125}\quad[\text{derived}].$$

The residue at zero shares (the low end of strength, Thm. 38.5(i)) and the Gaussian at the native shares (the high end, Thm. 38.5(ii)) are the two readings of one exact identity between the fibre and its lattice. For the other four five-mark lists the closings solve no adjoint of the ray law in the forms tried: no combination $\sum_{l<d}r_l(\sigma)[\alpha^0]\big(K(\alpha)\Psi(\alpha)/\alpha^{\mathcal{E}}\big)^{\sigma+l}$ with polynomial r_l of degree at most 20, nor with the apparent factors of c_0 as denominators up to degree 14, solves it [computed]. In that framework the pairing of the period with its closing is not a single product there; it must carry other solution classes.

Proposition 15.14 (Signs through parallel classes)**.** [derived] *In the setting of Proposition 15.10,* $K_{\mathcal{B}}(\eta)=K_{\mathcal{H}}\big(s_1(\eta),\dots,s_m(\eta)\big)$ *with* $s_i(\eta)=\sum_{e\in P_i}\eta_e\in\{s_i,s_i-2,\dots,-s_i\}$*: the sign values of the bananas* $\binom{P_i}{1}$ *feed the letter of* $\mathcal{H}$.

For the triangle $\mathcal{H} = \binom{[3]}{2}$, $K_{\mathcal{H}} = e_2$, and with one mark tripled, $s = s_1(\eta) \in \{3, 1, -1, -3\}$:

$$e_2(s, 1, 1) = 2s + 1 \in \{7, 3, -1, -5\}, \qquad e_2(s, 1, -1) = -1 \quad \text{for every } s,$$

which is the census 49 (1), 25 (1), 9 (3), 1 (11) of §15.4. The second form does not see s.

Proposition 15.15 (Letters that forget a class)**.** [derived] *Write* $K_{\mathcal{B}} = A_i\, K_{\mathcal{H}/i}(A) + K_{\mathcal{H}\setminus i}(A)$ *with* $A_i = \alpha(P_i)$ *(Def. 7.3, p. 48), and* $A'_i = \sum_{e \in P_i} \alpha_e^{-1}$.

(i) Tripled triangle. On $\alpha_3 = -\alpha_2$ *the contraction letter* $K_{\mathcal{H}/1} = A_2 + A_3$ *vanishes, so every copy has occupation zero at* α *and at* α^{-1}*, and* $K(\alpha) = -\alpha_2^2$*,* $K(\alpha^{-1}) = -\alpha_2^{-2}$ *whatever the copies are. The native gauges there are*

$$\big\{\alpha_3 = -\alpha_2,\ A_1 = \alpha_2^2 A'_1\big\},$$

a family of dimension two modulo scale, all of native value 1*, containing the eight sign classes with* $\eta_3 = -\eta_2$.

(ii) Two marks doubled. On $P_2 = (t, -t)$ *the class letter vanishes,* $A_2 = A'_2 = 0$*, and* $K = A_1\alpha_3$ *is the letter of* $\mathcal{H} \setminus 2$*, in which both remaining classes are coloops. The native gauges there are*

$$\Big\{P_1 = (c, c),\ P_2 = (t, -t),\ t^2 = \frac{c\,\alpha_3\,(2\alpha_3 + c)}{2c + \alpha_3}\Big\},$$

a curve of native value $\frac{1}{4}$ *which on the torus* $|\alpha_e| = 1$ *is a circle through the signs with* $(s_1, s_2) = (\pm 2, 0)$.

Proof. Nativity is $R_e(\alpha) = R_e(\alpha^{-1})$ at every mark. (i) At a copy $x \in P_1$, $R_x(\alpha) = \alpha_x(A_2 + A_3)/K(\alpha)$ and $R_x(\alpha^{-1}) = \alpha_x^{-1}(A'_2 + A'_3)/K(\alpha^{-1})$, both zero on $\alpha_3 = -\alpha_2$; at α_2 and α_3 the two sides differ by $\mp(A_1/\alpha_2 - \alpha_2 A'_1)$. (ii) At α_3 both occupations are 1 on $A_2 = A'_2 = 0$; at P_1 the equation holds when $\alpha_{1a}^2 = \alpha_{1b}^2 = A_1/A'_1$, and $\alpha_{1b} = -\alpha_{1a}$ gives $K = 0$; at P_2 it holds exactly at the stated t^2. When $|c| = |\alpha_3| = 1$, $|2\alpha_3 + c| = |2c + \alpha_3|$, so $|t| = 1$. □

Remark 15.16 (Local exponents). Put $x_k \approx \lambda^k k^\beta$ into the recurrence of the closings [computed]. At every simple root of every row, $\beta = -2 = -\frac{n-1}{2}$, the exponent of a nondegenerate native gauge (Thm. 35.5(i), p. 308). At the repeated roots the indicial polynomials are

$$(\beta+1)(\beta+2)(\beta+3) \quad (\text{tripled triangle},\ \lambda = 1), \qquad (2\beta+3)(2\beta+5) \quad (\text{two marks doubled},\ \lambda = 4).$$

A family of real dimension d on the torus $|\alpha_e| = 1$, transversally nondegenerate, gives $\beta = -\frac{n-1-d}{2}$. The Hessian of $\log\big(K(\alpha)K(\alpha^{-1})\big)$ there has rank two along the family of (i) and rank three along the circle of (ii) [computed], so these are -1 and $-\frac{3}{2}$. The isolated sign orbit at $\lambda = 1$ (one copy plus, two minus, $\eta_2 = \eta_3$) has full rank and gives -2. The exponents -3 and $-\frac{5}{2}$: [not yet computed]. These families are integral directions in the exceptional set of Thm. 35.19(i), p. 313: the order of the ray law counts native values with these multiplicities, and the families raise the order, and with it the masters, past the number of native values.

Remark 15.17 (Native gauges at complex points off the signs [computed]). $\binom{[5]}{2}$: $K(\alpha)K(\alpha^{-1}) = 1$ at $(\omega^2, \omega^2, \omega, \omega, 1)$, $\omega = e^{2\pi \mathrm{i}/3}$, where $K = K(\alpha^{-1}) = 1$; and -8 at $(1, 1, b, b, 1)$, $b = -(2 + \sqrt{3})$, and at its inverse, where $K = -2 - 2\sqrt{3}$ and $K(\alpha^{-1}) = -2 + 2\sqrt{3}$. Two marks doubled: 1 at $(\omega, \omega^2, \omega, \omega^2, 1)$. The signs are the points of order two of the torus; points of order three are the next layer that occurs here.

15.6 Closing the banana at strength one

Proposition 15.18 (Four marks close on the half-lattice)**.** [derived] *For* $\binom{[4]}{1}$*, merging the marks in pairs by four Gammas on a line (Cor. 4.7, p. 26),* $\int_{a_1+a_2=c}\Gamma(a_1)^2\Gamma(a_2)^2=\Gamma^{\,4}_{1}\!\left[\begin{smallmatrix}c,\ c,\ c,\ c\\ 2c\end{smallmatrix}\right]$*, leaves a fibre of dimension one, and*

$$P_{\mathcal{B}}(\mathbf{1};1,1)=\frac{1}{2\pi i}\int_{(1/2)}\Gamma^{\,4}_{1}\!\left[\begin{smallmatrix}c,\ c,\ c,\ c\\ 2c\end{smallmatrix}\right]\Gamma^{\,4}_{1}\!\left[\begin{smallmatrix}1-c,\ 1-c,\ 1-c,\ 1-c\\ 2-2c\end{smallmatrix}\right]dc$$
$$=(\mathsf{T}_{3;(1)}[x]\,\mathbb{I}_{\frac12+\mathbb{Z}})((0,\infty))=7\,(\mathsf{T}_{3;(1)}[x]\,\mathbb{I}_{\mathbb{Z}})((0,\infty)).$$

Proof. By Lemma 5.16, p. 34, and the shift and reflection of the stacked Gamma (Prop. 1.3(iii),(v), p. 11) the integrand is

$$\Gamma^{\,4}_{0}\big[c,\ c,\ 1-c,\ 1-c\big]^2\ \frac{\sin 2\pi c}{\pi(1-2c)}=\frac{2\pi^3\cos\pi c}{(1-2c)\sin^3\pi c}.$$

On the lines $\operatorname{Re}c=N+\frac12$ it is $O\big(N^{-1}e^{-2\pi|\operatorname{Im}c|}\big)$, so it closes to the right (Thm. 4.5, p. 26). At $c=m\ge1$, $2\pi^3\cos\pi c/\sin^3\pi c=2(c-m)^{-3}+O(c-m)$, with no parts of orders one and two, as in Lemma 5.16; the half-integers are zeros. So the residue is $2\,[\varepsilon^2](1-2m-2\varepsilon)^{-1}=-8/(2m-1)^3$, and the closing is $\sum_{m\ge1}8(2m-1)^{-3}=\sum_{x\in\frac12+\mathbb{Z},\,x>0}x^{-3}$. The last equality is the coset relation at the step 2 (Thm. 5.12(i), p. 32). The half-lattice is the doubled share $2c$ of a merged pair, the lower entry of its stacked Gamma. □

This is the note's value (Ex. 6.2, p. 39), now read as a closing of the share side.

Proposition 15.19 (A mark added is a critical-line kernel)**.** [derived]*; receipts* [computed] *For every* $n\ge1$*,*

$$P_{\binom{[n+1]}{1}}(\mathbf{1};1,1)=-\sum_{m\ge1}\partial_\sigma P_{\binom{[n]}{1}}(\sigma\mathbf{1};\sigma,\sigma)\big|_{\sigma=m}.$$

Proof. In the share side of $\binom{[n+1]}{1}$ (Proposition 15.2) merge the first n marks: their fibre at the total d is $\Gamma(d)^2P_{\binom{[n]}{1}}(d\mathbf{1};d,d)$, by the same proposition at n marks. The last mark carries $\Gamma(1-d)^2$, and $\Gamma(d)^2\Gamma(1-d)^2=\Gamma^{\,4}_{0}\big[d,\ d,\ 1-d,\ 1-d\big]$, whose right closing reads the derivative of the rest at the positive integers (Lemma 5.16(i)). The rest is holomorphic on $\operatorname{Re}d>0$, bounded on each vertical line by its value at the real part, and decays as $\operatorname{Re}d\to\infty$, so the closing applies (Thm. 4.5). □

Receipts [computed]. At $n=2$, with $\partial_\sigma\Gamma^{\,2}_{1}\!\left[\begin{smallmatrix}\sigma,\ \sigma\\ 2\sigma\end{smallmatrix}\right]=2\Gamma^{\,2}_{1}\!\left[\begin{smallmatrix}\sigma,\ \sigma\\ 2\sigma\end{smallmatrix}\right]\big((\log\Gamma)'(\sigma)-(\log\Gamma)'(2\sigma)\big)$ and $(\log\Gamma)'(j+1)=-(\mathsf{T}_{1;(1)}[x]\,\mathbb{I}_{\mathbb{Z}})((j,\infty))$ (Prop. 5.14, p. 33),

$$P_{\binom{[3]}{1}}(\mathbf{1};1,1)=2\sum_{m\ge1}\Gamma^{\,2}_{1}\!\left[\begin{smallmatrix}m,\ m\\ 2m\end{smallmatrix}\right]\,(\mathsf{T}_{1;(1)}[x]\,\mathbb{I}_{\mathbb{Z}})((m-1,2m-1])=2.343907238689458890601562\ldots,$$

the period of two marks times a head of the counting input, to 25 digits. At $n=3$ the sum is $7(\mathsf{T}_{3;(1)}[x]\,\mathbb{I}_{\mathbb{Z}})((0,\infty))$ and at $n=4$ it is $39.945588011996245699\ldots$, both to 20 digits.

Remark 15.20 (The five-mark centre is a Trailmoment of a live input). By Proposition 15.19,

$$P_{\binom{[5]}{1}}(\mathbf{1};1,1)=\big(\mathsf{T}_{0;(1)}[x]\,f\,\mathbb{I}_{\mathbb{Z}}\big)((0,\infty)),\qquad f(m)=-\partial_\sigma P_{\binom{[4]}{1}}(\sigma\mathbf{1};\sigma,\sigma)\big|_{\sigma=m},$$

a live input (Chapter 6): its density is read from the four-mark centre period, and the four-mark ray law differentiated in σ gives it a recurrence of order two whose inhomogeneity is the values. At $\sigma = 0$ the differentiated law is a relation among the first two slopes, the corner and the values, the term of order σ of Proposition 15.3 being zero. So every $f(m)$ lies in $\mathbb{Q} + \mathbb{Q}(\mathsf{T}_{3;(1)}[x]\,\mathbb{I}_{\mathbb{Z}})((0,\infty)) + \mathbb{Q}f(1)$ [derived]. For example $P_{\binom{[4]}{1}}(2\,\mathbf{1}; 2, 2) = (7\zeta(3) - 6)/16$ and $f(2) = -\big(36 + 21\zeta(3) - 2f(1)\big)/32$ (alias $\zeta(3) = (\mathsf{T}_{3;(1)}[x]\,\mathbb{I}_{\mathbb{Z}})((0,\infty))$) [computed]. The first slope is

$$-f(1) = 48\,\mathrm{Li}_4(\tfrac{1}{2}) + 2\log^4 2 - 2\pi^2\log^2 2 - \tfrac{17}{30}\pi^4 = -39.381582123122165624939809345 08\ldots$$

by PSLQ at 60 digits [computed]. In the family these aliases are Trailmoments of index at most four of the counting input at the frequencies 0 and $\frac{1}{2}$ and of the lattice input of density 2^{-x} (Def. 5.2, p. 28). So the four-mark slope sits on the half-lattice of Proposition 15.18, at index four. The masters of the five-mark centre are values of this live Trailmoment, and this is where the closing stops landing on plain lattice inputs.

15.7 Closing the live input of five marks

Throughout this subsection $p(\sigma) = P_{\binom{[4]}{1}}(\sigma\mathbf{1}; \sigma, \sigma)$ for real $\sigma > 0$ and $f = -p'$; $p\,\mathbb{I}_{\mathbb{Z}}$ and $f\,\mathbb{I}_{\mathbb{Z}}$ are the *values input* and the *slope input* (Def. 5.4, p. 29). The four-mark ray law is

$$\sum_{i=0}^{2} c_i(\sigma)\,p(\sigma + i) = 0, \qquad c_0 = \sigma^3, \quad c_1 = -2(2\sigma+1)(5\sigma^2+5\sigma+2), \quad c_2 = 64(\sigma+1)^3,$$

read off the closings by Procedure 15.6 and derived in Theorem 15.25 below. For a function R on a coset of $\mathbb{Z}$ put $(L^*R)(x) = \sum_i c_i(x-i)\,R(x-i)$, and let L'^* be the same with the derivatives c_i' in place of the c_i.

Proposition 15.21 (The live recurrence)**.** [derived] *The slope input has a recurrence of order two whose inhomogeneity is the values input:*

$$\sum_{i=0}^{2} c_i(\sigma)\,f(\sigma + i) = \sum_{i=0}^{2} c_i'(\sigma)\,p(\sigma + i), \qquad c_0' = 3\sigma^2, \quad c_1' = -6(10\sigma^2+10\sigma+3), \quad c_2' = 192(\sigma+1)^2.$$

Proof. The law holds identically in σ (Theorem 15.25); differentiate it. □

Proposition 15.22 (A mark added, at every strength)**.** [derived]*; receipts* [computed] *For real $s > 0$ let $W_s(d) = \Gamma_2^2\begin{bmatrix} d,\ d \\ 1+d-s,\ 1+d-s \end{bmatrix}$. Then*

$$\Gamma(s)^2\,P_{\binom{[5]}{1}}(s\mathbf{1}; s, s) = -\big(\mathsf{T}_{0;(1)}[x]\,(W_s p)'\,\mathbb{I}_{s+\mathbb{Z}}\big)((s-1,\infty)),$$
$$x^{(1)}(s) = \quad \big(\mathsf{T}_{0;(1)}[x]\,W_s p\,\mathbb{I}_{s+\mathbb{Z}}\big)((s-1,\infty)),$$

with $x^{(1)}$ the lowered setting with one h of Remark 15.8. The period reads the slopes of the word $W_s p$ on the coset of its strength, and the lowered setting reads its values. At $s \in \{1, 2, \ldots\}$ the word $W_s(x) = \prod_{j=1}^{s-1}(x-j)^2$ is a polynomial with double zeros at $1, \ldots, s-1$, the cut moves to 0, and

$$\Gamma(s)^2 P_{\binom{[5]}{1}}(s\mathbf{1}; s, s) = (\mathsf{T}_{0;(1)}[x]\,(W_s f - W_s' p)\,\mathbb{I}_{\mathbb{Z}})((0,\infty)), \qquad x^{(1)}(s) = (\mathsf{T}_{0;(1)}[x]\,W_s p\,\mathbb{I}_{\mathbb{Z}})((0,\infty)).$$

Proof. Merge the first four marks in the share side of five (Proposition 15.2): $\Gamma(s)^2 P_{\binom{[5]}{1}}(s\mathbf{1}; s, s) = \frac{1}{2\pi i}\int_{(c)} \Gamma(d)^2 p(d)\,\Gamma(s-d)^2\,dd$ with $0 < c < s$, as in the proof of Proposition 15.19. By the reflection (Prop. 1.3(v), p. 11), $\Gamma(s-d)\Gamma(1-s+d) = \pi/\sin\pi(s-d)$, so

$$\Gamma(d)^2\,\Gamma(s-d)^2 = W_s(d)\,\Gamma_{\binom{4}{0}}\big[s-d,\ s-d,\ 1-s+d,\ 1-s+d\big]\,,$$

the stacked Gamma of Lemma 5.16, p. 34, at $a = s-d$: double poles at $d \in s+\mathbb{Z}$, with no simple parts. The double zeros of W_s at $d = s-1, s-2, \dots$ remove the poles between the contour and s. The rest is as in Proposition 15.19: $W_s p$ is holomorphic on $\operatorname{Re} d > 0$, of polynomial growth along vertical lines, where the kernel decays like $e^{-2\pi|\operatorname{Im} d|}$, and it decays like $16^{-\operatorname{Re} d}$ to the right; closing to the right (Thm. 4.5, p. 26) reads $-(W_s p)'$ at $d = s, s+1, \dots$. For $x^{(1)}$, expand $h = \sum_k y^k/\Gamma(1+k)^2$ against g^4 (Proposition 15.2 at four marks): $x^{(1)}(s) = \sum_{k\ge 0}\Gamma(s+k)^2 p(s+k)/\Gamma(1+k)^2$, and $\Gamma(s+k)^2/\Gamma(1+k)^2 = W_s(s+k)$. □

Receipts [computed]: at $s = \frac{3}{2}$ both sides of the first identity are 2.27912905076377, and at $s = \frac{9}{4}$ both are 0.129204079141924 (the left side by one-dimensional quadrature of the share side; fifteen digits).

Proposition 15.23 (Summation by parts at the cut: what lands on lattice inputs)**.** [derived] *For every polynomial R,*

$$(\mathsf{T}_{0;(1)}[x]\,(L^*R)\,p\,\mathbb{I}_{\mathbb{Z}})((0,\infty)) = -24\,R(0),$$
$$(\mathsf{T}_{0;(1)}[x]\,(L^*R)\,f\,\mathbb{I}_{\mathbb{Z}})((0,\infty)) = (\mathsf{T}_{0;(1)}[x]\,(L'^*R)\,p\,\mathbb{I}_{\mathbb{Z}})((0,\infty)).$$

Proof. By the shifted inputs (Thm. 6.2, p. 42) a shift of the input is a shift of the word, and the cut relation (Prop. 5.3, p. 29) returns the points that cross the cut: for a density φ,

$$\sum_{x\ge 1} R(x)\sum_i c_i(x)\,\varphi(x+i) = (\mathsf{T}_{0;(1)}[x]\,(L^*R)\,\varphi\,\mathbb{I}_{\mathbb{Z}})((0,\infty)) - R(0)\sum_{i\ge 1} c_i(0)\,\varphi(i) - R(-1)\,c_2(-1)\,\varphi(1),$$

and $c_2(-1) = c_2'(-1) = 0$. For $\varphi = p$ the left side vanishes by the law. For $\varphi = f$ it is $\sum_x R(x)\sum_i c_i'(x)\,p(x+i)$ by Proposition 15.21, to which the same move applies. So the boundaries are

$$\sum_{i\ge 1} c_i(0)\,p(i) = -4p(1)+64p(2), \qquad \sum_{i\ge 1}\big(c_i(0)f(i) - c_i'(0)p(i)\big) = -4f(1)+64f(2)+18p(1)-192p(2).$$

The first is $-[\sigma^3]c_0$ times the corner 24 (Procedure 15.6(v)); the second is the term of order σ of $\sigma^3 p(\sigma)$, which vanishes (Proposition 15.3). With the values of Remark 15.20 they are -24 and 0: the lattice Trailmoments in them, $p(1) = 7(\mathsf{T}_{3;(1)}[x]\,\mathbb{I}_{\mathbb{Z}})((0,\infty))$, $f(1)$ and $f(2)$, cancel identically. □

So what the closing of the live Trailmoment puts on lattice inputs is the residue at zero shares, and nothing else: neither $(\mathsf{T}_{3;(1)}[x]\,\mathbb{I}_{\mathbb{Z}})((0,\infty))$ nor the four-mark slope $f(1)$ survives.

Proposition 15.24 (The fixed block)**.** [derived]*; receipts* [computed] *L^* raises the degree of a polynomial by three, with leading factor $1-20+64 = 45$, the symbol of the law at $\lambda = 1$, which is not a native value of four marks; so the polynomials are $L^*(\mathbb{Q}[x]) \oplus \operatorname{span}\{1, x-1, (x-1)^2\}$. By Proposition 15.23, every Trailmoment at index zero of a polynomial word against the values input or the slope input, cut at 0, is a rational number plus a rational combination of the six*

$$(\mathsf{T}_{0;(1)}[x]\,(x-1)^j\,p\,\mathbb{I}_{\mathbb{Z}})((0,\infty)), \qquad (\mathsf{T}_{0;(1)}[x]\,(x-1)^j\,f\,\mathbb{I}_{\mathbb{Z}})((0,\infty)) \qquad (j = 0,1,2).$$

The share side adds one relation:

$$5\,(\mathsf{T}_{0;(1)}[x]\,p\,\mathbb{I}_{\mathbb{Z}})((0,\infty)) - 5\,(\mathsf{T}_{0;(1)}[x]\,(x-1)\,f\,\mathbb{I}_{\mathbb{Z}})((0,\infty)) = (\mathsf{T}_{0;(1)}[x]\,f\,\mathbb{I}_{\mathbb{Z}})((0,\infty)).$$

Proof. The splitting: $\deg L^*R = \deg R + 3$, so L^* is injective on polynomials and its image misses exactly three dimensions in each degree. The relation says that each mark takes on average a fifth of the strength. On the fibre of Proposition 15.2 the shares sum to σ, and the integrand and the pins are symmetric in the five marks; so the fibre integral with one share a_e inserted is the same for every e, and the five sum to $\sigma\,\Gamma(\sigma)^2 P_{\binom{[5]}{1}}(\sigma\mathbf{1};\sigma,\sigma)$. Merging the first four marks, the fifth share is $\sigma - d$, and at $\sigma = 1$ the closing of Proposition 15.22 reads its insertion as $-\sum_{x\geq 1}\partial_d\big((1-d)p(d)\big)\big|_{d=x} = (\mathsf{T}_{0;(1)}[x]\,p\,\mathbb{I}_{\mathbb{Z}})((0,\infty)) - (\mathsf{T}_{0;(1)}[x]\,(x-1)f\,\mathbb{I}_{\mathbb{Z}})((0,\infty))$; the right side is $\frac{1}{5}P_{\binom{[5]}{1}}(\mathbf{1};1,1) = \frac{1}{5}(\mathsf{T}_{0;(1)}[x]\,f\,\mathbb{I}_{\mathbb{Z}})((0,\infty))$. □

Receipts [computed]. At the strengths $2,\dots,5$ the same insertion reduces to rational multiples of this relation. There is no rational relation among 1, $(\mathsf{T}_{3;(1)}[x]\,\mathbb{I}_{\mathbb{Z}})((0,\infty))$, $f(1)$ and the five Trailmoments other than $(\mathsf{T}_{0;(1)}[x]\,(x-1)f\,\mathbb{I}_{\mathbb{Z}})((0,\infty))$ with coefficients below 10^{30} (PSLQ, 360 digits): in that framework, over $\mathbb{Q}$, the block is five-dimensional. Its six values:

j	$(\mathsf{T}_{0;(1)}[x]\,(x-1)^j\,p\,\mathbb{I}_{\mathbb{Z}})((0,\infty))$	$(\mathsf{T}_{0;(1)}[x]\,(x-1)^j\,f\,\mathbb{I}_{\mathbb{Z}})((0,\infty))$
0	8.57028044337446126814969582424653728 6…	39.945588011996245698944028498423995 89…
1	0.16106970227667485885131066624851419 8…	0.58116284097521212836089012456173810 7…
2	0.17187458138353690484768349157529391 6…	0.61683551197162484909292454115302812 6…

Theorem 15.25 (A mark added carries the ray law)**.** [derived]*; receipts* [computed] *Normalize the ray law of* $\binom{[n]}{1}$, $\sum_{i=0}^{d_n} c_i^{(n)}(\sigma)\,P_{\binom{[n]}{1}}((\sigma+i)\mathbf{1};\sigma+i,\sigma+i) = 0$, *so that* $c_0^{(n)} = (-1)^n\sigma^{n-1}$, *as for four marks above and for five marks in Proposition 15.13, and let* L_n^* *be its adjoint. For* $n = 1,\dots,6$,

$$\sum_{i=0}^{d_{n+1}} c_i^{(n+1)}(s)\binom{x-1}{s-1+i}^2 = \big(L_n^* A_s\big)(x), \qquad A_s(x) = \big((n+1)x-(n+2)s\big)\binom{x}{s}^2,$$

an identity of rational functions of x *and* s *once both sides are divided by* $\binom{x}{s}^2$. *Read one way it produces the law: the words* $\binom{x-1}{s-1+i}^2$ *have distinct degrees, so the coefficients of* $L_n^*A_s$ *in them are the* $c_i^{(n+1)}(s)$. *Read the other way it proves it: for* $2 \leq n \leq 6$, *if the ray law of* $\binom{[n]}{1}$ *holds identically in* σ, *so does that of* $\binom{[n+1]}{1}$, *and the lowered setting* $x^{(1)}$ *of* $n+1$ *marks, divided by* $\Gamma(s)^2$, *obeys it too. The two-mark period* $P_{\binom{[2]}{1}}(\sigma\mathbf{1};\sigma,\sigma) = \Gamma_{\begin{smallmatrix}2\\1\end{smallmatrix}}\begin{bmatrix}\sigma,\ \sigma\\ 2\sigma\end{bmatrix}$ *obeys* $\sigma P(\sigma) - 2(2\sigma+1)P(\sigma+1) = 0$ *by the shift of the stacked Gamma (Prop. 1.3(iii), p. 11). So the ray laws of* $\binom{[n]}{1}$, $n \leq 7$, *are derived from the stacked Gamma, one mark at a time.*

Proof. Write p_n for the centre period of $\binom{[n]}{1}$, and let $s > 0$ be non-integral. Proposition 15.22, with n marks merged in place of four, gives $P_{\binom{[n+1]}{1}}((s+i)\mathbf{1};s+i,s+i) = -\sum_{d\in s+\mathbb{N}}(w_{s,i}\,p_n)'(d)$ with $w_{s,i} = W_{s+i}/\Gamma(s+i)^2$, the terms $d = s,\dots,s+i-1$ being zero by the double zeros of W_{s+i}. On the coset $w_{s,i}(d)$ is $\binom{d-1}{s-1+i}^2$ written with Gammas, so the displayed identity holds there. On $s+\mathbb{N}$ the move of Proposition 15.23 has no boundary for the words A_s and A_s', which vanish at $d = s-1, s-2,\dots$: $\sum_d (L_n^*A)\,\varphi = \sum_d A\,(L_n\varphi)$, and likewise with L_n'. With $(L_n^*A)' = L_n'^*A + L_n^*(A')$ and $L_np_n' = -L_n'p_n$ (the law differentiated),

$$\sum_i c_i^{(n+1)}(s)\,P_{\binom{[n+1]}{1}}((s+i)\mathbf{1};s+i,s+i) = -\sum_{d\in s+\mathbb{N}}\Big((L_n'^*A_s)\,p_n + A_s'\,L_np_n + A_s\,L_np_n'\Big)(d)$$
$$= -\sum_d (L_n'^*A_s)\,p_n + \sum_d A_s\,L_n'p_n = 0,$$

and $\sum_i c_i^{(n+1)}(s)\,\Gamma(s+i)^{-2}x^{(1)}(s+i) = \sum_d (L_n^* A_s)\,p_n = \sum_d A_s\,L_n p_n = 0$. Both sides are holomorphic in s, so the laws hold identically. The displayed identity is a finite computation with rational functions of (x, s). □

Receipts [computed]. The certificate A_s was found by exact linear algebra at each $n \le 6$ and $s \le 5$, the same at every n up to the factor $(n+1)x - (n+2)s$, and the identity was then checked with s indeterminate. At $n = 1$ ($p_1 = 1$, $c^{(1)} = (-1, 1)$) it holds as an identity, but the sums of the proof diverge. The five-mark law at $s = \frac{1}{2}$ and $s = \frac{3}{10}$, from the share side by quadrature: residuals $4.6 \cdot 10^{-14}$ and $2.3 \cdot 10^{-14}$ against terms of size 10^2. At $s = 0$, with the cut at 0, the same reduction returns the corner relation of $n+1$ marks as $A_0'(0) = n+1$ times the boundary of Proposition 15.23: the corners $(n+1)! = (n+1) \cdot n!$ of Proposition 15.3, here $120 = 5 \cdot 24$.

Corollary 15.26 (Two masters over the corner, two lowered numbers)**.** [derived] *For every $s \in \{1, 2, \dots\}$ there are rationals a_s, b_s, q_s with*

$$\Gamma(s)^2 P_{\binom{[5]}{1}}(s\mathbf{1}; s, s) = q_s + a_s\,P_{\binom{[5]}{1}}(\mathbf{1}; 1, 1) + b_s\,P_{\binom{[5]}{1}}(2\mathbf{1}; 2, 2),$$
$$x^{(1)}(s) = a_s\,x^{(1)}(1) + b_s\,x^{(1)}(2),$$

and in the block

$$P_{\binom{[5]}{1}}(\mathbf{1}; 1, 1) = (\mathsf{T}_{0;(1)}[x]\,f\,\mathbb{I}_{\mathbb{Z}})((0,\infty)),$$
$$P_{\binom{[5]}{1}}(2\mathbf{1}; 2, 2) = (\mathsf{T}_{0;(1)}[x]\,(x-1)^2 f\,\mathbb{I}_{\mathbb{Z}})((0,\infty)) - 2\,(\mathsf{T}_{0;(1)}[x]\,(x-1)\,p\,\mathbb{I}_{\mathbb{Z}})((0,\infty)),$$
$$x^{(1)}(1) = (\mathsf{T}_{0;(1)}[x]\,p\,\mathbb{I}_{\mathbb{Z}})((0,\infty)), \qquad x^{(1)}(2) = (\mathsf{T}_{0;(1)}[x]\,(x-1)^2 p\,\mathbb{I}_{\mathbb{Z}})((0,\infty)).$$

Modulo $L^*(x\,\mathbb{Q}[x])$ every word W_s lies in $\operatorname{span}\{1, (x-1)^2\}$: the ray reaches the direction $x-1$ of the block only through the value term $-2(\mathsf{T}_{0;(1)}[x]\,(x-1)p\,\mathbb{I}_{\mathbb{Z}})((0,\infty))$ of its second master. For example $(a_3, b_3, q_3) = \big(-\frac{1}{45}, \frac{19}{15}, \frac{8}{15}\big)$ and $(a_4, b_4, q_4) = \big(-\frac{113}{1125}, \frac{1907}{375}, \frac{944}{375}\big)$.

Proof. The four block identities are Proposition 15.22 at $s = 1, 2$, with $W_1 = 1$, $W_2 = (x-1)^2$, $W_2' = 2(x-1)$. Write $W_s/\Gamma(s)^2 = L^* B_s + r_s$ with $\deg r_s \le 2$. Since the display of Theorem 15.25 is an identity of polynomials and the splitting of Proposition 15.24 is direct, the r_s obey the five-mark recurrence for $s \ge 1$, and at $s = 0$, where $A_0 = 5x$, they obey $\sum_{i \ge 1} c_i^{(5)}(0)\,r_i = 0$. As $r_1 = 1$ and $r_2 = (x-1)^2$, every r_s lies in their span, and the same recursion, from $B_1 = B_2 = 0$, gives $B_s(0) = 0$. Then $x^{(1)}(s) = \Gamma(s)^2(\mathsf{T}_{0;(1)}[x]\,r_s\,p\,\mathbb{I}_{\mathbb{Z}})((0,\infty))$, and by Proposition 15.23 $P_{\binom{[5]}{1}}(s\mathbf{1}; s, s) = (\mathsf{T}_{0;(1)}[x]\,(r_s f - r_s' p)\,\mathbb{I}_{\mathbb{Z}})((0,\infty)) + 24 B_s'(0)$, where $\sum_{i \ge 1} c_i^{(5)}(0) B_i'(0) = A_0'(0) = 5$ carries the corner. □

Proposition 15.27 (The lowered half is a stacked Gamma)**.** [computed]

$$x^{(1)}(1) = (\mathsf{T}_{0;(1)}[x]\,p\,\mathbb{I}_{\mathbb{Z}})((0,\infty)) = \frac{1}{120}\Gamma^{\,8}_{\,4}\left[\begin{matrix}\frac{1}{15}, & \frac{2}{15}, & \frac{4}{15}, & \frac{8}{15}, & \frac{1}{5}, & \frac{2}{5}, & \frac{3}{5}, & \frac{4}{5} \\ & & \frac{1}{2}, & \frac{1}{2}, & \frac{1}{2}, & \frac{1}{2} & & \end{matrix}\right]$$

to 364 *digits, and*

$$x^{(1)}(2) = \frac{65\,x^{(1)}(1) - 32\,\Gamma(\frac{1}{2})^8/x^{(1)}(1)}{1125} \quad (360\ \textit{digits}),$$
$$x^{(2)}(1) = \frac{\sqrt{15}}{4\,\Gamma(\frac{1}{2})^2}\,x^{(1)}(1), \qquad x^{(2)}(2) = \frac{\sqrt{15}}{4\,\Gamma(\frac{1}{2})^2} \cdot \frac{65\,x^{(1)}(1) + 32\,\Gamma(\frac{1}{2})^8/x^{(1)}(1)}{1125} \quad (100\ \textit{digits}).$$

So the block of Proposition 15.9 is $x^{(1)}(1)x^{(2)}(2) - x^{(1)}(2)x^{(2)}(1) = \frac{\sqrt{15}}{4\Gamma(1/2)^2} \cdot \frac{64\,\Gamma(1/2)^8}{1125} = \frac{16\sqrt{15}}{1125}\Gamma(\frac{1}{2})^6$: the stacked Gamma cancels from the frame, which is why the frame saw only $\Gamma(\frac{1}{2})$.

The values half that the ray reaches is two stacked-Gamma numbers, $x^{(1)}(1)$ and $\Gamma(\frac{1}{2})^8/x^{(1)}(1)$. The direction $x-1$ is not among them: $(\mathsf{T}_{0;(1)}[x]\,(x-1)\,p\,\mathbb{I}_{\mathbb{Z}})((0,\infty))$ has no rational relation with 1, $(\mathsf{T}_{3;(1)}[x]\,\mathbb{I}_{\mathbb{Z}})((0,\infty))$, $\Gamma(\frac{1}{2})^4$, $x^{(1)}(1)$ and $\Gamma(\frac{1}{2})^8/x^{(1)}(1)$, nor with ten products of $x^{(1)}(1)^{\pm1}$, powers of $\Gamma(\frac{1}{2})^2$ and $\sqrt{15}$, with coefficients below 10^{12} [computed].

Proposition 15.28 (The block on the four-mark chart and its pencil)**.** [derived]*; receipts* [computed] *On the chart $y_{e_0}=0$ of four marks let $X=K(\alpha)K(\alpha^{-1})$, the reciprocal of the stride of the centre ray (Def. 35.9, p. 309), so that $p(\sigma)=\int X^{-\sigma}dy$, the value along the ray from the real origin $(\sigma\mathbf{1};\sigma,\sigma)$ (Def. 35.1, p. 307), with $X\geq 16$ on the chart (Thm. 35.10(i), p. 309). For a polynomial R,*

$$(\mathsf{T}_{0;(1)}[x]\,R\,p\,\mathbb{I}_{\mathbb{Z}})((0,\infty))=\int \rho_R(X)\,dy,\qquad (\mathsf{T}_{0;(1)}[x]\,R\,f\,\mathbb{I}_{\mathbb{Z}})((0,\infty))=\int \log X\,\rho_R(X)\,dy,$$
$$\rho_R(X)=\sum_{m\geq 1}R(m)X^{-m},$$

a rational function of X whose only pole is $X=1$. In particular

$$P_{\binom{[5]}{1}}(\mathbf{1};1,1)=\int\frac{\log X}{X-1}\,dy,\qquad P_{\binom{[5]}{1}}(\mathbf{21};2,2)=\int\frac{(X+1)\log X-2(X-1)}{(X-1)^3}\,dy,$$
$$x^{(1)}(1)=\int\frac{dy}{X-1},\qquad x^{(1)}(2)=\int\frac{X+1}{(X-1)^3}\,dy,\qquad (\mathsf{T}_{0;(1)}[x]\,(x-1)\,p\,\mathbb{I}_{\mathbb{Z}})((0,\infty))=\int\frac{dy}{(X-1)^2}.$$

The first is the left closing of Lemma 5.16(ii), p. 34, read on the chart. With the generating function of the four-mark ray from the origin $(\varepsilon\mathbf{1};\varepsilon,\varepsilon)$, $G_\varepsilon(q)=\sum_{k\geq1}p(k+\varepsilon)\,q^{k-1}=\int X^{-\varepsilon}\,dy/(X-q)$ for $|q|<16$, carried by the pencil $K\Psi-q\alpha^{\mathbf{1}}$ (Thm. 35.10(ii)),

$$(\mathsf{T}_{0;(1)}[x]\,(x-1)^j\,p\,\mathbb{I}_{\mathbb{Z}})((0,\infty))=(q\partial_q)^jG_0(q)\big|_{q=1},$$
$$(\mathsf{T}_{0;(1)}[x]\,(x-1)^j\,f\,\mathbb{I}_{\mathbb{Z}})((0,\infty))=-\partial_\varepsilon(q\partial_q)^jG_\varepsilon(q)\big|_{q=1,\ \varepsilon=0}:$$

the block is the jet at $q=1$ of the four-mark generating function and of its slope in the origin.

Proof. Def. 35.1 along the ray from $(\sigma\mathbf{1};\sigma,\sigma)$, the positivity of every term, and $\sum_{m\geq1}R(m)X^{-m}$ summed for $X>1$: $\rho_1=1/(X-1)$, $\rho_{x-1}=1/(X-1)^2$, $\rho_{(x-1)^2}=(X+1)/(X-1)^3$. For the masters use Corollary 15.26; for the pencil, Thm. 35.10(ii) with $k_0=1$ and the origin form $X^{-\varepsilon}dy$. □

Receipts [computed], by three-dimensional quadrature on the chart, to twelve digits:

$$\int\frac{dy}{X}=8.414398322115,\qquad \int\frac{\log X}{X-1}\,dy=39.945588011991,\qquad \int\frac{dy}{X-1}=8.570280443372,$$

against $7(\mathsf{T}_{3;(1)}[x]\,\mathbb{I}_{\mathbb{Z}})((0,\infty))$, the five-mark centre at strength one, and $x^{(1)}(1)$.

Remark 15.29 (What the closing found). (i) The live Trailmoment lands on lattice inputs only through the residue at zero shares (Proposition 15.23).

(ii) The words of the five-mark centre ray reach a plane of the five-dimensional block on each side: on the values side the lowered setting, two stacked-Gamma numbers (Proposition 15.27); on the slopes side the period, whose two masters are

$$P_{\binom{[5]}{1}}(\mathbf{1};1,1)=(\mathsf{T}_{0;(1)}[x]\,f\,\mathbb{I}_{\mathbb{Z}})((0,\infty)),$$
$$P_{\binom{[5]}{1}}(\mathbf{21};2,2)=(\mathsf{T}_{0;(1)}[x]\,(x-1)^2f\,\mathbb{I}_{\mathbb{Z}})((0,\infty))-2\,(\mathsf{T}_{0;(1)}[x]\,(x-1)\,p\,\mathbb{I}_{\mathbb{Z}})((0,\infty)):$$

the slope in the origin of the four-mark generating function at $q = 1$, and its second jet less twice the first jet of the generating function itself. They have no rational relation with 1, $(\mathsf{T}_{3;(1)}[x]\,\mathbb{I}_{\mathbb{Z}})((0,\infty))$, $f(1)$, $\Gamma(\frac{1}{2})^4$, $\Gamma(\frac{1}{2})^8$, $(\mathsf{T}_{5;(1)}[x]\,\mathbb{I}_{\mathbb{Z}})((0,\infty))$, $\Gamma(\frac{1}{2})^4(\mathsf{T}_{3;(1)}[x]\,\mathbb{I}_{\mathbb{Z}})((0,\infty))$, the two stacked-Gamma numbers and the direction $x-1$ of the block, with coefficients below 10^{20} (PSLQ, 360 digits) [computed]: in that framework, over $\mathbb{Q}$, they are new numbers.

(iii) The last minor of §14.17 is now written in the masters over the stacked Gamma (Proposition 15.27):

$$P_{\binom{[5]}{1}}(\mathbf{1};1,1)\,x^{(1)}(2) - P_{\binom{[5]}{1}}(2\mathbf{1};2,2)\,x^{(1)}(1)$$
$$= \frac{65\,x^{(1)}(1) - 32\,\Gamma(\frac{1}{2})^8/x^{(1)}(1)}{1125}\,P_{\binom{[5]}{1}}(\mathbf{1};1,1) - x^{(1)}(1)\,P_{\binom{[5]}{1}}(2\mathbf{1};2,2)$$
$$= 4.34000293153566203407\ldots,$$

and it is not a rational combination of 1, $\Gamma(\frac{1}{2})^4$, $\Gamma(\frac{1}{2})^8$, $(\mathsf{T}_{3;(1)}[x]\,\mathbb{I}_{\mathbb{Z}})((0,\infty))$ and seven products of $x^{(1)}(1)^{\pm1}$ or $x^{(1)}(1)^{\pm2}$ with powers of $\Gamma(\frac{1}{2})^4$, with coefficients below 10^{15} [computed]. Its scale is the masters' scale; nothing smaller is missing.

(iv) The rational words and the pole columns of Chapter 6 did not enter: every word of the centre ray is a polynomial, read at index zero. What closed the live Trailmoment is the shifted inputs (Thm. 6.2), the cut relation (Prop. 5.3), summation by parts in its adjoint form (Thm. 6.5, p. 43), and a fixed block at nonpositive index, the counterpart of Thm. 6.9, p. 45.

(v) Aliases, last: $x^{(1)}(1) = 8\int_0^\infty tI_0(t)K_0(t)^4\,dt$, whose evaluation by the Gammas at fifteenths is the classical one through the singular modulus of $\mathbb{Q}(\sqrt{-15})$ [recalled]; Theorem 15.25 gives the recurrences of the moments $\int_0^\infty t^{2s-1}K_0(t)^n\,dt$, $n \le 7$, by adding marks.

15.8 Open

(i) The certificate for every n. Theorem 15.25 for all n would derive every banana ray law from the stacked Gamma, the law of each list produced by the one before. At each $n \le 6$ the same A_s appeared; an operator proof for general n: [not yet computed]. From five to six marks the symbol of the five-mark law vanishes at $\lambda = 1$ (a native value of five marks), L_5^* raises degrees by less than four, and the block of six marks is built differently: [not yet computed]. The analogue of the certificate for a mark added in parallel to a mark of another list (Proposition 15.10), which would carry the second row of the table of §15.4 from $\binom{[4]}{2}$: [not yet computed]

(ii) The direction $x-1$ of the block, $(\mathsf{T}_{0;(1)}[x]\,(x-1)\,p\,\mathbb{I}_{\mathbb{Z}})((0,\infty)) = (q\partial_q)G_0(q)|_{q=1} = \int dy/(X-1)^2 = 0.16106970227667485885\ldots$, which the ray reaches only through the value term of its second master. Expanding h as in Proposition 15.22, it is the lowered setting with h replaced by $y\,\partial_y h$: $\int_0^\infty g^4\,y\,h'(y)\,dy$ [derived] (alias $4\int_0^\infty t^2 I_1(t)K_0(t)^4 dt$, to 25 digits [computed]). Which member of the family it is, and whether a lean of the five-mark list reads it as a period: [not yet computed]

(iii) Three marks: reduce the live Trailmoment of the receipts of Proposition 15.19 at $n = 2$ to the Trailmoment of the odd periodic density modulo 3 at index 2 (alias $3L(\chi_{-3},2)$) inside the family; Thm. 5.17 does not reach it, the parity being mismatched. [not yet computed]

(iv) The three masters of $\binom{[5]}{2}$: no identification yet. [not yet computed]

(v) The second corners, from the list. The boundary term at $\sigma = -1$ is a constant term of the continued period (Thm. 1.6), assembled from its Laurent coefficients of orders $-(n-1)$, $-(n-2)$, $-(n-3)$ at $\sigma = 0$ and $\sigma = -1$. The leading ones are the corner and the corner times

$|\mathcal{B}|$ (Proposition 15.4); the next vanishes at 0 (Proposition 15.3) and the ray law fixes it at -1. On the share side the third ones are Laurent data of the Gammas at the lattice points of the closing, which are heads of the counting input and Trailmoments of the counting input cut at the data (Prop. 5.13, p. 32; Prop. 5.14, p. 33); the rationality of the two boundary terms says that the Trailmoments cancel and the heads remain. Carrying this out: [not yet computed]. Also: which lists put $(\sigma+1)^2$ into both c_0 and c_1. Here they are the two triangles $\binom{[3]}{2}$ with parallel classes, and not $\binom{[4]}{2}$ with a mark doubled. [not yet computed]

(vi) The last minor of §14.17, $M_5(1)x^{(1)}(2) - M_5(2)x^{(1)}(1) = 4.34000293153566\ldots$, is the minor of the solutions of native values $\frac{1}{25}$ and $\frac{1}{9}$, so it is the minimal solution of the exterior square of the ray law, of native value $\frac{1}{225}$ [derived]. It is now the masters over the stacked Gamma (Remark 15.29(iii)), so a relation of degree two fixing it would be a relation between the masters and $x^{(1)}(1)$. The exact pairing of Proposition 15.13 is a relation of degree one between solutions, fixed by the residue at zero shares. Whether pairings of the sign classes with closings of the share side twisted by the signs fix relations of degree two such as this minor: [not yet computed]

(vii) The pairing for the other lists: the partner of the period in the adjoint law, when the closings are not. [not yet computed]

(viii) Lean rays. At a lean with uniform masses only $\eta = \mathbf{1}$ stays native for $\binom{\mathcal{E}}{1}$: at a sign the native equation reads $(m-t)K(\eta) = (s-t)\eta_e$ for every e [derived]. The gauges that move off the signs and their values. [not yet computed]

(ix) Families of native gauges. In both cases of Proposition 15.15 the letter collapses at the gauge to the letter of a minor that leaves directions free: a class whose contraction letter vanishes is free, and a class whose own letter vanishes leaves a deletion with coloops, whose relative scales are free. A rule for every list, and the exponents -3 and $-\frac{5}{2}$ of the remark on local exponents. [not yet computed]

(x) A third flavour: $\mathrm{M}(K_4)$ on six marks, an atom, whose full cobasis lean $(0,2)$ at unit masses is $6\,(\mathsf{T}_{3;(1)}[x]\,\mathbb{I}_{\mathbb{Z}})((0,\infty))$ (alias $6\zeta(3)$) [recalled]; its centre ray. [not yet computed]