

# Molten Salt Reactor Transients

## How Power of The Future Performs In Accidents

Gavin Ridley  
Advanced Reactors and Fuel Cycles Group

University of Illinois at Urbana-Champaign

07.20.2017



I L L I N O I S



# Outline

## 1 Motivation

Molten Salt Reactors  
What We Simulated  
Multiphysics Coupling

## 2 Methods

MOOSE  
Sustainable, open software  
Group Constant Generation

## 3 Results & Conclusion



# Energy for the future

*Cheap and abundant nuclear energy is no longer a luxury; it will eventually be a necessity for the maintenance of the human condition. – Alvin Weinberg*



Molten salt reactors offer a convincing solution to the problem of fossil fuels.

- Potentially much cheaper than normal nuclear
- Make meltdowns impossible
- Better natural resource utilization



# MSR Comparison

## Topaz Solar Farm

- 9.5 sq. mi of California desert
- Max output of 550 MW(e), on average makes 132 MW(e)
- \$2.5B construction




Figure 1: Topaz solar farm in California, credit GigaOM media

## Terrestrial Energy IMSR concept

- About 300 MW(e) output, more than double Topaz farm
- Initial cost estimates rank IMSR cheaper than coal




Figure 2: IMSR and some other small nuclear designs, credit Terrestrial Energy



## Benchmark Case: MSRE

- Constructed at Oak Ridge National Lab, ran reliably 1965-1969 at 7.4 MW(th)
- Various tests proved theory and tech-readiness for full-scale power plants
- Stopped operation since all needed experiments were done





Figure 3: MSRE reactor core diagram from ORNL technical reports





# MSRs: An Intrinsically Coupled System I

**Neutrons'** changing concentrations can be described approximately with coupled diffusion equations:

$$\frac{1}{v_g} \frac{\partial \phi_g}{\partial t} = \nabla \cdot D_g \nabla \phi_g + \sum_{g' \neq g}^G \Sigma_{g' \rightarrow g}^s \phi_{g'} + \chi_g^p \sum_{g'=1}^G (1-\beta) \nu \Sigma_{g'}^f \phi_{g'} + \chi_g^d \sum_i^I \lambda_i C_i - \Sigma_g^r \phi_g \quad (1)$$

where  $v_g$  = speed of neutrons in group  $g$

$\phi_g$  = flux of neutrons in group  $g$

$t$  = time

$D_g$  = Diffusion coefficient for neutrons in group  $g$

$\Sigma_g^r$  = macroscopic cross-section for removal of neutrons from group  $g$

$\Sigma_{g' \rightarrow g}^s$  = macroscopic cross-section of scattering from  $g'$  to  $g$

$\chi_g^p$  = prompt fission spectrum, neutrons in group  $g$

$G$  = number of discrete groups,  $g$

$\nu$  = number of neutrons produced per fission

$\Sigma_g^f$  = macroscopic cross section for fission due to neutrons in group  $g$

$\chi_g^d$  = delayed fission spectrum, neutrons in group  $g$

$I$  = number of delayed neutron precursor groups

$\beta$  = delayed neutron fraction

$\lambda_i$  = average decay constant of delayed neutron precursors in precursor group  $i$



## MSRs: An Intrinsically Coupled System II

**Delayed neutron precursors** are products of freshly split uranium that emit a new neutron after a delay. *critical* to reactor control; they shift power change timescales from picoseconds to seconds despite only being a few tenths of a percent of emitted neutrons.

$$\frac{\partial C_i}{\partial t} = \sum_{g'=1}^G \beta_i \nu \Sigma_{g'}^f \phi_{g'} - \lambda_i C_i - \frac{\partial}{\partial z} u C_i \quad (2)$$

**Heat and temperature** affect the coefficients in Equation 1 significantly. Energy conservation must be solved:

$$\rho_f c_{p,f} \frac{\partial T_f}{\partial t} + \nabla \cdot (\rho_f c_{p,f} \vec{u} \cdot T_f - k_f \nabla T_f) = Q_f \quad (3)$$

- $\rho_f$  = density of fuel salt
- $c_{p,f}$  = specific heat capacity of fuel salt
- $T_f$  = temperature of fuel salt
- $\vec{u}$  = velocity of fuel salt
- $k_f$  = thermal conductivity of fuel salt
- $Q_f$  = source term





# Outline

## 1 Motivation

Molten Salt Reactors  
What We Simulated  
Multiphysics Coupling

## 2 Methods

MOOSE  
Sustainable, open software  
Group Constant Generation


## 3 Results & Conclusion

$$\vec{\Omega} \cdot \vec{\nabla} \Psi + \sigma_t(\vec{r}) \Psi(\vec{r}, \vec{\Omega}) = \frac{1}{4\pi} (\sigma_s(\vec{r}) \Phi(\vec{r}) + S(\vec{r}))$$

$$\nabla \cdot k \nabla T = 0$$

$$\frac{\partial c}{\partial t} - \nabla \cdot (\vec{v}c) = 0$$

 MOOSE





# MOOSE Physics Representation

- Highly object-oriented code solves weak form of PDE using finite element method
- PETSc solves resulting system of nonlinear equations using generalized minimal residual method (GMRES)
- Some of the world's most cutting-edge numerical algorithms and scalable parallel computing are made painlessly accessible to the everyday user



## MOOSE Example

In MOOSE, the term  $D\nabla^2 u$  is easily represented by:

```
Real
GroupDiffusion::computeQpResidual()
{
    return _D[_qp][_group] * _grad_test[_i][_qp] *
           computeConcentrationGradient(_u, _grad_u, _qp);
}
```


A vacuum boundary condition in neutronics calculations can easily be represented by:

```
Real
VacuumConcBC::computeQpResidual()
{
    return _test[_i][_qp] * computeConcentration(_u, _qp) / 2.;
}
```




# Moltres

- [github.com/arfc/moltres](https://github.com/arfc/moltres)
- Publicly developed on github
- Continuous integration by Civet
- Includes detailed guide for contributing





# Group constant generation





# Outline

## ① Motivation

- Molten Salt Reactors
- What We Simulated
- Multiphysics Coupling

## ② Methods

- MOOSE
- Sustainable, open software
- Group Constant Generation

## ③ Results & Conclusion



# Results

asfd





# Conclusion

We showed many things.

- Cats are peculiar
- Blue and Orange are fierce colors
- Math can be rendered nicely
- Cite your sources



# Acknowledgement

Acknowledgements should include both people who helped and funding streams.  
If you are funded by an NEUP grant, that number usually goes here. .



**NUCLEAR**  
ENERGY IS COOL, KIDS