

# Marginalia to a Theorem of Asperó and Schindler

Obrad Kasum<sup>1</sup>

Institut de Mathématiques de Jussieu-Paris Rive Gauche  
Université Paris Cité

Simon's Semester, December 18, 2023

---

<sup>1</sup>Received funding from the European Union's Horizon 2020 research and innovation program under the Marie Skłodowska-Curie grant agreement No. 945322 



- $\phi$  is a consistent infinitary propositional formula,

- $\phi$  is a consistent infinitary propositional formula,
- $\kappa < \text{rank}(\phi)$  is an inaccessible cardinal,

- $\phi$  is a consistent infinitary propositional formula,
- $\kappa < \text{rank}(\phi)$  is an inaccessible cardinal,
- $\mathbb{H}$  is the poset consisting of consistent finite sets of subformulas of  $\phi$ , ordered by  $\supseteq$ .

- $\phi$  is a consistent infinitary propositional formula,
- $\kappa < \text{rank}(\phi)$  is an inaccessible cardinal,
- $\mathbb{H}$  is the poset consisting of consistent finite sets of subformulas of  $\phi$ , ordered by  $\supseteq$ .

## Proposition

In  $V^{\mathbb{H}}$ , there exists a model for  $\phi$ .

- $\phi$  is a consistent infinitary propositional formula,
- $\kappa < \text{rank}(\phi)$  is an inaccessible cardinal,
- $\mathbb{H}$  is the poset consisting of consistent finite sets of subformulas of  $\phi$ , ordered by  $\supseteq$ .

## Proposition

In  $V^{\mathbb{H}}$ , there exists a model for  $\phi$ .

- However, in general,  $\mathbb{H}$  collapses  $\omega_1$ .

- $\phi$  is a consistent infinitary propositional formula,
- $\kappa < \text{rank}(\phi)$  is an inaccessible cardinal,
- $\mathbb{H}$  is the poset consisting of consistent finite sets of subformulas of  $\phi$ , ordered by  $\supseteq$ .

## Proposition

In  $V^{\mathbb{H}}$ , there exists a model for  $\phi$ .

- However, in general,  $\mathbb{H}$  collapses  $\omega_1$ .
- We want to define a poset  $\mathbb{P}_\kappa$  which also adds a model for  $\phi$ , but which is in addition stationary set preserving.

$$\blacksquare \mathcal{E} := \{\lambda < \kappa : \phi \in V_\lambda \prec V_\kappa\}$$



- $\mathcal{E} := \{\lambda < \kappa : \phi \in V_\lambda \prec V_\kappa\}$
- For all  $\lambda \in \mathcal{E}$ ,  $\mathcal{C}_\lambda$  denotes the set of all countable  $M$  satisfying

- $\mathcal{E} := \{\lambda < \kappa : \phi \in V_\lambda \prec V_\kappa\}$
- For all  $\lambda \in \mathcal{E}$ ,  $\mathcal{C}_\lambda$  denotes the set of all countable  $M$  satisfying
  - $M \prec \hat{M} \models \text{ZFC}^-$ ,

- $\mathcal{E} := \{\lambda < \kappa : \phi \in V_\lambda \prec V_\kappa\}$
- For all  $\lambda \in \mathcal{E}$ ,  $\mathcal{C}_\lambda$  denotes the set of all countable  $M$  satisfying
  - $M \prec \hat{M} \models \text{ZFC}^-$ ,
  - $\phi, V_\lambda \in M$ ,

- $\mathcal{E} := \{\lambda < \kappa : \phi \in V_\lambda \prec V_\kappa\}$
- For all  $\lambda \in \mathcal{E}$ ,  $\mathcal{C}_\lambda$  denotes the set of all countable  $M$  satisfying
  - $M \prec \hat{M} \models \text{ZFC}^-$ ,
  - $\phi, V_\lambda \in M$ ,
  - $\hat{M} = \text{Hull}(M, V_\lambda)$ .

- $\mathcal{E} := \{\lambda < \kappa : \phi \in V_\lambda \prec V_\kappa\}$
- For all  $\lambda \in \mathcal{E}$ ,  $\mathcal{C}_\lambda$  denotes the set of all countable  $M$  satisfying
  - $M \prec \hat{M} \models \text{ZFC}^-$ ,
  - $\phi, V_\lambda \in M$ ,
  - $\hat{M} = \text{Hull}(M, V_\lambda)$ .
- For all  $M \prec \hat{M} \models \text{ZFC}^-$  and for all  $\lambda \in \text{Ord}$ , we denote by  $M \downarrow \lambda$  the image of  $M$  under the transitive collapse of  $\text{Hull}(M, V_\lambda)$ .

- $\mathcal{E} := \{\lambda < \kappa : \phi \in V_\lambda \prec V_\kappa\}$
- For all  $\lambda \in \mathcal{E}$ ,  $\mathcal{C}_\lambda$  denotes the set of all countable  $M$  satisfying
  - $M \prec \hat{M} \models \text{ZFC}^-$ ,
  - $\phi, V_\lambda \in M$ ,
  - $\hat{M} = \text{Hull}(M, V_\lambda)$ .
- For all  $M \prec \hat{M} \models \text{ZFC}^-$  and for all  $\lambda \in \text{Ord}$ , we denote by  $M \downarrow \lambda$  the image of  $M$  under the transitive collapse of  $\text{Hull}(M, V_\lambda)$ .
  - If  $M \in \mathcal{C}_\lambda$  and  $\xi \in \mathcal{E} \cap \lambda$ , do we have  $M \downarrow \xi \in \mathcal{C}_\xi$ ?

- $\mathcal{E} := \{\lambda < \kappa : \phi \in V_\lambda \prec V_\kappa\}$
- For all  $\lambda \in \mathcal{E}$ ,  $\mathcal{C}_\lambda$  denotes the set of all countable  $M$  satisfying
  - $M \prec \hat{M} \models \text{ZFC}^-$ ,
  - $\phi, V_\lambda \in M$ ,
  - $\hat{M} = \text{Hull}(M, V_\lambda)$ .
- For all  $M \prec \hat{M} \models \text{ZFC}^-$  and for all  $\lambda \in \text{Ord}$ , we denote by  $M \downarrow \lambda$  the image of  $M$  under the transitive collapse of  $\text{Hull}(M, V_\lambda)$ .
  - If  $M \in \mathcal{C}_\lambda$  and  $\xi \in \mathcal{E} \cap \lambda$ , do we have  $M \downarrow \xi \in \mathcal{C}_\xi$ ?
- For all  $S \subseteq \text{Ord}$ , we denote  $\mathcal{C}_S := \bigcup_{\lambda \in S \cap \mathcal{E}} \mathcal{C}_\lambda$ .

- $\mathcal{E} := \{\lambda < \kappa : \phi \in V_\lambda \prec V_\kappa\}$
- For all  $\lambda \in \mathcal{E}$ ,  $\mathcal{C}_\lambda$  denotes the set of all countable  $M$  satisfying
  - $M \prec \hat{M} \models \text{ZFC}^-$ ,
  - $\phi, V_\lambda \in M$ ,
  - $\hat{M} = \text{Hull}(M, V_\lambda)$ .
- For all  $M \prec \hat{M} \models \text{ZFC}^-$  and for all  $\lambda \in \text{Ord}$ , we denote by  $M \downarrow \lambda$  the image of  $M$  under the transitive collapse of  $\text{Hull}(M, V_\lambda)$ .
  - If  $M \in \mathcal{C}_\lambda$  and  $\xi \in \mathcal{E} \cap \lambda$ , do we have  $M \downarrow \xi \in \mathcal{C}_\xi$ ?
- For all  $S \subseteq \text{Ord}$ , we denote  $\mathcal{C}_S := \bigcup_{\lambda \in S \cap \mathcal{E}} \mathcal{C}_\lambda$ .
- For all  $M \in \mathcal{C}_{[0, \kappa]}$ , we denote by  $\lambda_M$  the unique  $\lambda$  such that  $M \in \mathcal{C}_\lambda$ .



Suppose that  $\lambda \in \mathcal{E} \cup \{\kappa\}$ .

Suppose that  $\lambda \in \mathcal{E} \cup \{\kappa\}$ .

- $p \in \mathbb{P}_\lambda^*$  iff  $p = (w_p, \mathcal{M}_p)$  where

Suppose that  $\lambda \in \mathcal{E} \cup \{\kappa\}$ .

- $p \in \mathbb{P}_\lambda^*$  iff  $p = (w_p, \mathcal{M}_p)$  where
  - $w_p \in \mathbb{H}$ ,

Suppose that  $\lambda \in \mathcal{E} \cup \{\kappa\}$ .

- $p \in \mathbb{P}_\lambda^*$  iff  $p = (w_p, \mathcal{M}_p)$  where
  - $w_p \in \mathbb{H}$ ,
  - $\mathcal{M}_p$  is a finite subset of  $\mathcal{C}_{<\lambda}$ ,

Suppose that  $\lambda \in \mathcal{E} \cup \{\kappa\}$ .

- $p \in \mathbb{P}_\lambda^*$  iff  $p = (w_p, \mathcal{M}_p)$  where
  - $w_p \in \mathbb{H}$ ,
  - $\mathcal{M}_p$  is a finite subset of  $\mathcal{C}_{<\lambda}$ ,
  - for all  $M, N \in \mathcal{M}_p$ ,

Suppose that  $\lambda \in \mathcal{E} \cup \{\kappa\}$ .

- $p \in \mathbb{P}_\lambda^*$  iff  $p = (w_p, \mathcal{M}_p)$  where
  - $w_p \in \mathbb{H}$ ,
  - $\mathcal{M}_p$  is a finite subset of  $\mathcal{C}_{<\lambda}$ ,
  - for all  $M, N \in \mathcal{M}_p$ ,
    - $\delta_M = \delta_N \implies M = N$ ,

Suppose that  $\lambda \in \mathcal{E} \cup \{\kappa\}$ .

- $p \in \mathbb{P}_\lambda^*$  iff  $p = (w_p, \mathcal{M}_p)$  where
  - $w_p \in \mathbb{H}$ ,
  - $\mathcal{M}_p$  is a finite subset of  $\mathcal{C}_{<\lambda}$ ,
  - for all  $M, N \in \mathcal{M}_p$ ,
    - $\delta_M = \delta_N \implies M = N$ ,
    - $\delta_M < \delta_N \implies M \in N \wedge \lambda_M < \lambda_N$ .

Suppose that  $\lambda \in \mathcal{E} \cup \{\kappa\}$ .

- $p \in \mathbb{P}_\lambda^*$  iff  $p = (w_p, \mathcal{M}_p)$  where
  - $w_p \in \mathbb{H}$ ,
  - $\mathcal{M}_p$  is a finite subset of  $\mathcal{C}_{<\lambda}$ ,
  - for all  $M, N \in \mathcal{M}_p$ ,
    - $\delta_M = \delta_N \implies M = N$ ,
    - $\delta_M < \delta_N \implies M \in N \wedge \lambda_M < \lambda_N$ .
- $p \leq q$  iff  $w_p \supseteq w_q$  and for all  $N \in \mathcal{M}_q$ , there exists  $M \in \mathcal{M}_p$  such that  $\delta_M = \delta_N$ ,  $\lambda_M = \lambda_N$ , and  $M \supseteq N$ .



Suppose that  $\lambda \in \mathcal{E} \cup \{\kappa\}$ .

- $p \in \mathbb{P}_\lambda^*$  iff  $p = (w_p, \mathcal{M}_p)$  where
  - $w_p \in \mathbb{H}$ ,
  - $\mathcal{M}_p$  is a finite subset of  $\mathcal{C}_{<\lambda}$ ,
  - for all  $M, N \in \mathcal{M}_p$ ,
    - $\delta_M = \delta_N \implies M = N$ ,
    - $\delta_M < \delta_N \implies M \in N \wedge \lambda_M < \lambda_N$ .
- $p \leq q$  iff  $w_p \supseteq w_q$  and for all  $N \in \mathcal{M}_q$ , there exists  $M \in \mathcal{M}_p$  such that  $\delta_M = \delta_N$ ,  $\lambda_M = \lambda_N$ , and  $M \supseteq N$ .
- For  $p \in \mathbb{P}_\lambda^*$  and for  $\bar{\lambda} \in \lambda \cap \mathcal{E}$ , we define

$$p \restriction \lambda := (w_p, \{M \in \mathcal{M}_p : \lambda_M < \lambda\}).$$

Suppose that  $p \in \mathbb{P}_\lambda^*$ . Then  $\mathcal{G}_\lambda(p)$  is the game of length  $\omega$  played as follows.

Suppose that  $p \in \mathbb{P}_\lambda^*$ . Then  $\mathcal{G}_\lambda(p)$  is the game of length  $\omega$  played as follows.

- Set first  $p_{-1} := p$ .

Suppose that  $p \in \mathbb{P}_\lambda^*$ . Then  $\mathcal{G}_\lambda(p)$  is the game of length  $\omega$  played as follows.

- Set first  $p_{-1} := p$ .
- In the round  $n < \omega$ , Player I plays  $Q_n$  and Player II answers by  $p_n \in \mathbb{P}_\lambda^*$  satisfying  $p_n \leq p_{n-1}$ .

Suppose that  $p \in \mathbb{P}_\lambda^*$ . Then  $\mathcal{G}_\lambda(p)$  is the game of length  $\omega$  played as follows.

- Set first  $p_{-1} := p$ .
- In the round  $n < \omega$ , Player I plays  $Q_n$  and Player II answers by  $p_n \in \mathbb{P}_\lambda^*$  satisfying  $p_n \leq p_{n-1}$ .
- If  $Q_n = \phi \in w_{p_{n-1}}$  where  $\phi \equiv \bigvee_{i \in I} \psi_i$ , then for some  $i \in I$ ,  $\psi_i \in w_{p_n}$ .

Suppose that  $p \in \mathbb{P}_\lambda^*$ . Then  $\mathcal{G}_\lambda(p)$  is the game of length  $\omega$  played as follows.

- Set first  $p_{-1} := p$ .
- In the round  $n < \omega$ , Player I plays  $Q_n$  and Player II answers by  $p_n \in \mathbb{P}_\lambda^*$  satisfying  $p_n \leq p_{n-1}$ .
- If  $Q_n = \phi \in w_{p_{n-1}}$  where  $\phi \equiv \bigvee_{i \in I} \psi_i$ , then for some  $i \in I$ ,  $\psi_i \in w_{p_n}$ .
- If  $Q_n = (\phi, i) \in w_{p_{n-1}} \times I$ , where  $\phi \equiv \bigwedge_{i \in I} \psi_i$ , then  $\psi_i \in w_{p_n}$ .

Suppose that  $p \in \mathbb{P}_\lambda^*$ . Then  $\mathcal{G}_\lambda(p)$  is the game of length  $\omega$  played as follows.

- Set first  $p_{-1} := p$ .
- In the round  $n < \omega$ , Player I plays  $Q_n$  and Player II answers by  $p_n \in \mathbb{P}_\lambda^*$  satisfying  $p_n \leq p_{n-1}$ .
- If  $Q_n = \phi \in w_{p_{n-1}}$  where  $\phi \equiv \bigvee_{i \in I} \psi_i$ , then for some  $i \in I$ ,  $\psi_i \in w_{p_n}$ .
- If  $Q_n = (\phi, i) \in w_{p_{n-1}} \times I$ , where  $\phi \equiv \bigwedge_{i \in I} \psi_i$ , then  $\psi_i \in w_{p_n}$ .
- If  $Q_n = (M, D)$  where  $M \in \mathcal{M}_{p_{n-1}}$  and  $D \in M$  is dense in  $\mathbb{P}_{\lambda_M}$ , then there exists  $q \in D$  such that  $\delta(\text{Hull}(M, q)) = \delta(M)$  and  $p_n \leq p_{n-1}, q$ .

Suppose that  $p \in \mathbb{P}_\lambda^*$ . Then  $\mathcal{G}_\lambda(p)$  is the game of length  $\omega$  played as follows.

- Set first  $p_{-1} := p$ .
- In the round  $n < \omega$ , Player I plays  $Q_n$  and Player II answers by  $p_n \in \mathbb{P}_\lambda^*$  satisfying  $p_n \leq p_{n-1}$ .
- If  $Q_n = \phi \in w_{p_{n-1}}$  where  $\phi \equiv \bigvee_{i \in I} \psi_i$ , then for some  $i \in I$ ,  $\psi_i \in w_{p_n}$ .
- If  $Q_n = (\phi, i) \in w_{p_{n-1}} \times I$ , where  $\phi \equiv \bigwedge_{i \in I} \psi_i$ , then  $\psi_i \in w_{p_n}$ .
- If  $Q_n = (M, D)$  where  $M \in \mathcal{M}_{p_{n-1}}$  and  $D \in M$  is dense in  $\mathbb{P}_{\lambda_M}$ , then there exists  $q \in D$  such that  $\delta(\text{Hull}(M, q)) = \delta(M)$  and  $p_n \leq p_{n-1}, q$ .
- Player II wins infinite plays such that the set  $\bigcup_{n < \omega} w_{p_n}$  does not contain both an atomic formula and its negation.



## Definition

$\mathbb{P}_\lambda$  is the suborder of  $(\mathbb{P}_\lambda^*, \leq)$  consisting of those  $p \in \mathbb{P}_\lambda^*$  for which Player II wins  $\mathcal{G}_\lambda(p)$ .

## Definition

$\mathbb{P}_\lambda$  is the suborder of  $(\mathbb{P}_\lambda^*, \leq)$  consisting of those  $p \in \mathbb{P}_\lambda^*$  for which Player II wins  $\mathcal{G}_\lambda(p)$ .

## Definition (De Bondt, Veličković)

Suppose that  $\theta \gg \kappa$  is regular and  $M \prec (H_\theta, \in, \kappa, \phi)$  is countable. Then  $M$  is *good* iff for all  $p \in \mathbb{P}_\kappa \cap M$ , there exist  $q \in \mathbb{P}_\kappa$  and  $\lambda \in \mathcal{E}$  such that

## Definition

$\mathbb{P}_\lambda$  is the suborder of  $(\mathbb{P}_\lambda^*, \leq)$  consisting of those  $p \in \mathbb{P}_\lambda^*$  for which Player II wins  $\mathcal{G}_\lambda(p)$ .

## Definition (De Bondt, Veličković)

Suppose that  $\theta \gg \kappa$  is regular and  $M \prec (H_\theta, \in, \kappa, \phi)$  is countable. Then  $M$  is *good* iff for all  $p \in \mathbb{P}_\kappa \cap M$ , there exist  $q \in \mathbb{P}_\kappa$  and  $\lambda \in \mathcal{E}$  such that

$$\blacksquare \quad \kappa \cap M \subseteq \lambda,$$

## Definition

$\mathbb{P}_\lambda$  is the suborder of  $(\mathbb{P}_\lambda^*, \leq)$  consisting of those  $p \in \mathbb{P}_\lambda^*$  for which Player II wins  $\mathcal{G}_\lambda(p)$ .

## Definition (De Bondt, Veličković)

Suppose that  $\theta \gg \kappa$  is regular and  $M \prec (H_\theta, \in, \kappa, \phi)$  is countable. Then  $M$  is *good* iff for all  $p \in \mathbb{P}_\kappa \cap M$ , there exist  $q \in \mathbb{P}_\kappa$  and  $\lambda \in \mathcal{E}$  such that

- $\kappa \cap M \subseteq \lambda$ ,
- $q \leq p$ ,

## Definition

$\mathbb{P}_\lambda$  is the suborder of  $(\mathbb{P}_\lambda^*, \leq)$  consisting of those  $p \in \mathbb{P}_\lambda^*$  for which Player II wins  $\mathcal{G}_\lambda(p)$ .

## Definition (De Bondt, Veličković)

Suppose that  $\theta \gg \kappa$  is regular and  $M \prec (H_\theta, \in, \kappa, \phi)$  is countable. Then  $M$  is *good* iff for all  $p \in \mathbb{P}_\kappa \cap M$ , there exist  $q \in \mathbb{P}_\kappa$  and  $\lambda \in \mathcal{E}$  such that

- $\kappa \cap M \subseteq \lambda$ ,
- $q \leq p$ ,
- $M \downarrow \lambda \in \mathcal{M}_q$ .

## Proposition (De Bondt, Veličković)

Suppose that  $\theta \gg \kappa$  is regular and  $M \prec H_\theta$  is good. Then  $\mathbb{P}_\kappa$  is semiproper for  $M$ .

## Proposition (De Bondt, Veličković)

Suppose that  $\theta \gg \kappa$  is regular and  $M \prec H_\theta$  is good. Then  $\mathbb{P}_\kappa$  is semiproper for  $M$ .

## Corollary

If there exists a local club of good models  $M \prec H_\theta$ , then  $\mathbb{P}_\kappa$  is stationary set preserving.

## Proposition (De Bondt, Veličković)

Suppose that  $\theta \gg \kappa$  is regular and  $M \prec H_\theta$  is good. Then  $\mathbb{P}_\kappa$  is semiproper for  $M$ .

## Corollary

If there exists a local club of good models  $M \prec H_\theta$ , then  $\mathbb{P}_\kappa$  is stationary set preserving.

## Local club

$C \subseteq [X]^\omega$  is a *local club* iff for weak-club many  $\bar{X} \in [X]^{\omega_1}$ , the set  $C \cap [\bar{X}]^\omega$  contains a club.



## Definition

$\phi$  is *AS good* at  $\kappa$  iff

## Definition

$\phi$  is *AS good* at  $\kappa$  iff

$\forall S$  which are stationary subsets of  $\omega_1$ ,

## Definition

$\phi$  is *AS good* at  $\kappa$  iff

$\forall S$  which are stationary subsets of  $\omega_1$ ,  
 $V^{\text{Col}(\omega, < \kappa)}$  satisfies that:

## Definition

$\phi$  is *AS good* at  $\kappa$  iff

$\forall S$  which are stationary subsets of  $\omega_1$ ,  
 $V^{\text{Col}(\omega, < \kappa)}$  satisfies that:

$\forall \mu$  which are models for  $\phi$ ,

## Definition

$\phi$  is *AS good* at  $\kappa$  iff

$\forall S$  which are stationary subsets of  $\omega_1$ ,  
 $V^{\text{Col}(\omega, < \kappa)}$  satisfies that:

$\forall \mu$  which are models for  $\phi$ ,

$\exists \tau$  which is an elementary embedding from  $V$  to some  $W$  with  
critical point  $\omega_1^V$  and which satisfies that  $\omega_1^V \in \tau(S)$ , and

## Definition

$\phi$  is *AS good* at  $\kappa$  iff

$\forall S$  which are stationary subsets of  $\omega_1$ ,  
 $V^{\text{Col}(\omega, < \kappa)}$  satisfies that:

$\forall \mu$  which are models for  $\phi$ ,

$\exists \tau$  which is an elementary embedding from  $V$  to some  $W$  with  
critical point  $\omega_1^V$  and which satisfies that  $\omega_1^V \in \tau(S)$ , and

$\exists \hat{\mu}$  which is a model for  $\tau(\phi)$  satisfying that

## Definition

$\phi$  is *AS good* at  $\kappa$  iff

$\forall S$  which are stationary subsets of  $\omega_1$ ,  
 $V^{\text{Col}(\omega, < \kappa)}$  satisfies that:

$\forall \mu$  which are models for  $\phi$ ,

$\exists \tau$  which is an elementary embedding from  $V$  to some  $W$  with  
critical point  $\omega_1^V$  and which satisfies that  $\omega_1^V \in \tau(S)$ , and

$\exists \hat{\mu}$  which is a model for  $\tau(\phi)$  satisfying that  
for all  $\psi \in \phi \downarrow$ ,  $\hat{\mu}(\tau(\psi)) = \mu(\psi)$ .

## Theorem

If  $\phi$  is AS-good at  $\kappa$ , then  $\mathbb{P}_\kappa$  is stationary set preserving.



## Theorem

If  $\phi$  is AS-good at  $\kappa$ , then  $\mathbb{P}_\kappa$  is stationary set preserving.

*Proof.*

## Theorem

If  $\phi$  is AS-good at  $\kappa$ , then  $\mathbb{P}_\kappa$  is stationary set preserving.

*Proof.*

- 1 Let  $\theta$  be large enough regular. It suffices to show that there are local-club many good models  $M \prec H_\theta$ .

## Theorem

If  $\phi$  is AS-good at  $\kappa$ , then  $\mathbb{P}_\kappa$  is stationary set preserving.

*Proof.*

- 1 Let  $\theta$  be large enough regular. It suffices to show that there are local-club many good models  $M \prec H_\theta$ .
- 2 Let  $R$  be a wellordering of  $H_\theta$  and let  $\mathcal{H} := (H_\theta, \in, R, \kappa, \phi)$ . We want to show that for all  $X \prec \mathcal{H}$  satisfying  $\omega_1 \subseteq X$  and  $|X| = \omega_1$ , we have that club many countable  $M \prec X$  are good.

## Theorem

If  $\phi$  is AS-good at  $\kappa$ , then  $\mathbb{P}_\kappa$  is stationary set preserving.

*Proof.*

- 1 Let  $\theta$  be large enough regular. It suffices to show that there are local-club many good models  $M \prec H_\theta$ .
- 2 Let  $R$  be a wellordering of  $H_\theta$  and let  $\mathcal{H} := (H_\theta, \in, R, \kappa, \phi)$ . We want to show that for all  $X \prec \mathcal{H}$  satisfying  $\omega_1 \subseteq X$  and  $|X| = \omega_1$ , we have that club many countable  $M \prec X$  are good.
- 3 Let  $X \prec \mathcal{H}$  satisfying  $\omega_1 \subseteq X$  and  $|X| = \omega_1$  be arbitrary. We denote by  $\mathcal{X}$  the structure on  $X$  inherited from  $\mathcal{H}$ .

## Theorem

If  $\phi$  is AS-good at  $\kappa$ , then  $\mathbb{P}_\kappa$  is stationary set preserving.

*Proof.*

- 1 Let  $\theta$  be large enough regular. It suffices to show that there are local-club many good models  $M \prec H_\theta$ .
- 2 Let  $R$  be a wellordering of  $H_\theta$  and let  $\mathcal{H} := (H_\theta, \in, R, \kappa, \phi)$ . We want to show that for all  $X \prec \mathcal{H}$  satisfying  $\omega_1 \subseteq X$  and  $|X| = \omega_1$ , we have that club many countable  $M \prec X$  are good.
- 3 Let  $X \prec \mathcal{H}$  satisfying  $\omega_1 \subseteq X$  and  $|X| = \omega_1$  be arbitrary. We denote by  $\mathcal{X}$  the structure on  $X$  inherited from  $\mathcal{H}$ .
- 4 Let us assume towards contradiction that there exists a stationary  $S \subseteq [X]^\omega$  such that for all  $M \in S$ , it holds that  $M \prec \mathcal{X}$  and that there exists  $p_M \in \mathbb{P}_\kappa \cap M$  witnessing that  $M$  is not good.

- 5 By pressing down applied to  $S \ni M \mapsto p_M$ , there exist  $p \in \mathbb{P}_\kappa \cap X$  and stationary  $S' \subseteq S$  such that for all  $M \in S'$ ,  $p$  witnesses that  $M$  is not good.

- 5 By pressing down applied to  $S \ni M \mapsto p_M$ , there exist  $p \in \mathbb{P}_\kappa \cap X$  and stationary  $S' \subseteq S$  such that for all  $M \in S'$ ,  $p$  witnesses that  $M$  is not good.
- 6 Let  $e : \omega_1 \rightarrow X$  be a bijection and let

$$T := \{\alpha < \omega_1 : e[\alpha] \in S', \delta(e[\alpha]) = \alpha\}.$$

Then  $T$  is stationary in  $\omega_1$ .

- 5 By pressing down applied to  $S \ni M \mapsto p_M$ , there exist  $p \in \mathbb{P}_\kappa \cap X$  and stationary  $S' \subseteq S$  such that for all  $M \in S'$ ,  $p$  witnesses that  $M$  is not good.
- 6 Let  $e : \omega_1 \rightarrow X$  be a bijection and let

$$T := \{\alpha < \omega_1 : e[\alpha] \in S', \delta(e[\alpha]) = \alpha\}.$$

Then  $T$  is stationary in  $\omega_1$ .

- 7 Let
- $\lambda \in \mathcal{E}$  be such that for all for all  $\eta < \lambda$ ,  $\kappa \cap \text{Hull}^{\mathcal{H}}(V_\eta) \subseteq \lambda$ ,



- 5 By pressing down applied to  $S \ni M \mapsto p_M$ , there exist  $p \in \mathbb{P}_\kappa \cap X$  and stationary  $S' \subseteq S$  such that for all  $M \in S'$ ,  $p$  witnesses that  $M$  is not good.
- 6 Let  $e : \omega_1 \rightarrow X$  be a bijection and let

$$T := \{\alpha < \omega_1 : e[\alpha] \in S', \delta(e[\alpha]) = \alpha\}.$$

Then  $T$  is stationary in  $\omega_1$ .

- 7 Let
- $\lambda \in \mathcal{E}$  be such that for all for all  $\eta < \lambda$ ,  $\kappa \cap \text{Hull}^{\mathcal{H}}(V_\eta) \subseteq \lambda$ ,
  - $h$  be  $V$ -generic for  $\text{Col}(\omega, < \kappa)$ ,

- 5 By pressing down applied to  $S \ni M \mapsto p_M$ , there exist  $p \in \mathbb{P}_\kappa \cap X$  and stationary  $S' \subseteq S$  such that for all  $M \in S'$ ,  $p$  witnesses that  $M$  is not good.
- 6 Let  $e : \omega_1 \rightarrow X$  be a bijection and let

$$T := \{\alpha < \omega_1 : e[\alpha] \in S', \delta(e[\alpha]) = \alpha\}.$$

Then  $T$  is stationary in  $\omega_1$ .

- 7 Let
- $\lambda \in \mathcal{E}$  be such that for all  $\eta < \lambda$ ,  $\kappa \cap \text{Hull}^{\mathcal{H}}(V_\eta) \subseteq \lambda$ ,
  - $h$  be  $V$ -generic for  $\text{Col}(\omega, < \kappa)$ ,
  - $g \in V[h]$  be  $V$ -generic for  $\mathbb{P}_\lambda$  containing  $p$ ,

- 5 By pressing down applied to  $S \ni M \mapsto p_M$ , there exist  $p \in \mathbb{P}_\kappa \cap X$  and stationary  $S' \subseteq S$  such that for all  $M \in S'$ ,  $p$  witnesses that  $M$  is not good.
- 6 Let  $e : \omega_1 \rightarrow X$  be a bijection and let

$$T := \{\alpha < \omega_1 : e[\alpha] \in S', \delta(e[\alpha]) = \alpha\}.$$

Then  $T$  is stationary in  $\omega_1$ .

- 7 Let
- $\lambda \in \mathcal{E}$  be such that for all  $\eta < \lambda$ ,  $\kappa \cap \text{Hull}^{\mathcal{H}}(V_\eta) \subseteq \lambda$ ,
  - $h$  be  $V$ -generic for  $\text{Col}(\omega, < \kappa)$ ,
  - $g \in V[h]$  be  $V$ -generic for  $\mathbb{P}_\lambda$  containing  $p$ ,
  - $\mu := \mu_g \models \phi$ .

- 5 By pressing down applied to  $S \ni M \mapsto p_M$ , there exist  $p \in \mathbb{P}_\kappa \cap X$  and stationary  $S' \subseteq S$  such that for all  $M \in S'$ ,  $p$  witnesses that  $M$  is not good.
- 6 Let  $e : \omega_1 \rightarrow X$  be a bijection and let

$$T := \{\alpha < \omega_1 : e[\alpha] \in S', \delta(e[\alpha]) = \alpha\}.$$

Then  $T$  is stationary in  $\omega_1$ .

- 7 Let
- $\lambda \in \mathcal{E}$  be such that for all  $\eta < \lambda$ ,  $\kappa \cap \text{Hull}^{\mathcal{H}}(V_\eta) \subseteq \lambda$ ,
  - $h$  be  $V$ -generic for  $\text{Col}(\omega, < \kappa)$ ,
  - $g \in V[h]$  be  $V$ -generic for  $\mathbb{P}_\lambda$  containing  $p$ ,
  - $\mu := \mu_g \models \phi$ .

We work in  $V[h]$ .

- 8 Since  $\phi$  is AS good, there exist elementary  $\tau : V \rightarrow W$  and  $\hat{\mu} \models \tau(\phi)$  such that  $\text{crit}(\tau) = \omega_1^V \in \tau(T)$  and for all  $\psi \in \phi \downarrow$ ,  $\hat{\mu}(\tau(\psi)) = \mu(\psi)$ .

- 8 Since  $\phi$  is AS good, there exist elementary  $\tau : V \rightarrow W$  and  $\hat{\mu} \models \tau(\phi)$  such that  $\text{crit}(\tau) = \omega_1^V \in \tau(T)$  and for all  $\psi \in \phi \downarrow$ ,  $\hat{\mu}(\tau(\psi)) = \mu(\psi)$ .
- 9 Let  $N := \text{Hull}^{\tau(\mathcal{H})}(\omega_1^V \cup \{\tau(p)\}) \prec \tau(\mathcal{H})$ . It holds that

- 8 Since  $\phi$  is AS good, there exist elementary  $\tau : V \rightarrow W$  and  $\hat{\mu} \models \tau(\phi)$  such that  $\text{crit}(\tau) = \omega_1^V \in \tau(T)$  and for all  $\psi \in \phi \downarrow$ ,  $\hat{\mu}(\tau(\psi)) = \mu(\psi)$ .
- 9 Let  $N := \text{Hull}^{\tau(\mathcal{H})}(\omega_1^V \cup \{\tau(p)\}) \prec \tau(\mathcal{H})$ . It holds that
- $N \in W$ ,

- 8 Since  $\phi$  is AS good, there exist elementary  $\tau : V \rightarrow W$  and  $\hat{\mu} \models \tau(\phi)$  such that  $\text{crit}(\tau) = \omega_1^V \in \tau(T)$  and for all  $\psi \in \phi \downarrow$ ,  $\hat{\mu}(\tau(\psi)) = \mu(\psi)$ .
- 9 Let  $N := \text{Hull}^{\tau(\mathcal{H})}(\omega_1^V \cup \{\tau(p)\}) \prec \tau(\mathcal{H})$ . It holds that
- $N \in W$ ,
  - $|N|^W = \omega$ ,



- 8 Since  $\phi$  is AS good, there exist elementary  $\tau : V \rightarrow W$  and  $\hat{\mu} \models \tau(\phi)$  such that  $\text{crit}(\tau) = \omega_1^V \in \tau(T)$  and for all  $\psi \in \phi \downarrow$ ,  $\hat{\mu}(\tau(\psi)) = \mu(\psi)$ .
- 9 Let  $N := \text{Hull}^{\tau(\mathcal{H})}(\omega_1^V \cup \{\tau(p)\}) \prec \tau(\mathcal{H})$ . It holds that
- $N \in W$ ,
  - $|N|^W = \omega$ ,
  - $N \prec \tau[X] \prec \tau(\mathcal{X})$ ,

- 8 Since  $\phi$  is AS good, there exist elementary  $\tau : V \rightarrow W$  and  $\hat{\mu} \models \tau(\phi)$  such that  $\text{crit}(\tau) = \omega_1^V \in \tau(T)$  and for all  $\psi \in \phi \downarrow$ ,  $\hat{\mu}(\tau(\psi)) = \mu(\psi)$ .
- 9 Let  $N := \text{Hull}^{\tau(\mathcal{H})}(\omega_1^V \cup \{\tau(p)\}) \prec \tau(\mathcal{H})$ . It holds that
- $N \in W$ ,
  - $|N|^W = \omega$ ,
  - $N \prec \tau[X] \prec \tau(\mathcal{X})$ ,
  - $\delta_N = \omega_1^V \in \tau(T)$ .

- 8 Since  $\phi$  is AS good, there exist elementary  $\tau : V \rightarrow W$  and  $\hat{\mu} \models \tau(\phi)$  such that  $\text{crit}(\tau) = \omega_1^V \in \tau(T)$  and for all  $\psi \in \phi \downarrow$ ,  $\hat{\mu}(\tau(\psi)) = \mu(\psi)$ .
- 9 Let  $N := \text{Hull}^{\tau(\mathcal{H})}(\omega_1^V \cup \{\tau(p)\}) \prec \tau(\mathcal{H})$ . It holds that
- $N \in W$ ,
  - $|N|^W = \omega$ ,
  - $N \prec \tau[X] \prec \tau(\mathcal{X})$ ,
  - $\delta_N = \omega_1^V \in \tau(T)$ .
- 10 Let  $q := (\tau(w_p), \tau(\mathcal{M}_p) \cup \{N \downarrow \tau(\lambda)\})$ . We will show that  $q \in \tau(\mathbb{P}_\kappa)$  and that  $q \leq \tau(p)$ .

- 8 Since  $\phi$  is AS good, there exist elementary  $\tau : V \rightarrow W$  and  $\hat{\mu} \models \tau(\phi)$  such that  $\text{crit}(\tau) = \omega_1^V \in \tau(T)$  and for all  $\psi \in \phi \downarrow$ ,  $\hat{\mu}(\tau(\psi)) = \mu(\psi)$ .
- 9 Let  $N := \text{Hull}^{\tau(\mathcal{H})}(\omega_1^V \cup \{\tau(p)\}) \prec \tau(\mathcal{H})$ . It holds that
- $N \in W$ ,
  - $|N|^W = \omega$ ,
  - $N \prec \tau[X] \prec \tau(\mathcal{X})$ ,
  - $\delta_N = \omega_1^V \in \tau(T)$ .
- 10 Let  $q := (\tau(w_p), \tau(\mathcal{M}_p) \cup \{N \downarrow \tau(\lambda)\})$ . We will show that  $q \in \tau(\mathbb{P}_\kappa)$  and that  $q \leq \tau(p)$ .
- 11 It follows that in  $W$ ,  $\tau(p)$  does not witness that  $N$  is not good.

- 8 Since  $\phi$  is AS good, there exist elementary  $\tau : V \rightarrow W$  and  $\hat{\mu} \models \tau(\phi)$  such that  $\text{crit}(\tau) = \omega_1^V \in \tau(T)$  and for all  $\psi \in \phi \downarrow$ ,  $\hat{\mu}(\tau(\psi)) = \mu(\psi)$ .
- 9 Let  $N := \text{Hull}^{\tau(\mathcal{H})}(\omega_1^V \cup \{\tau(p)\}) \prec \tau(\mathcal{H})$ . It holds that
- $N \in W$ ,
  - $|N|^W = \omega$ ,
  - $N \prec \tau[X] \prec \tau(\mathcal{X})$ ,
  - $\delta_N = \omega_1^V \in \tau(T)$ .
- 10 Let  $q := (\tau(w_p), \tau(\mathcal{M}_p) \cup \{N \downarrow \tau(\lambda)\})$ . We will show that  $q \in \tau(\mathbb{P}_\kappa)$  and that  $q \leq \tau(p)$ .
- 11 It follows that in  $W$ ,  $\tau(p)$  does not witness that  $N$  is not good.
- 12 Since  $\delta_N = \omega_1^V \in \tau(T)$ , we have that  $N \in \tau(S')$ .

- 8 Since  $\phi$  is AS good, there exist elementary  $\tau : V \rightarrow W$  and  $\hat{\mu} \models \tau(\phi)$  such that  $\text{crit}(\tau) = \omega_1^V \in \tau(T)$  and for all  $\psi \in \phi \downarrow$ ,  $\hat{\mu}(\tau(\psi)) = \mu(\psi)$ .
- 9 Let  $N := \text{Hull}^{\tau(\mathcal{H})}(\omega_1^V \cup \{\tau(p)\}) \prec \tau(\mathcal{H})$ . It holds that
- $N \in W$ ,
  - $|N|^W = \omega$ ,
  - $N \prec \tau[X] \prec \tau(\mathcal{X})$ ,
  - $\delta_N = \omega_1^V \in \tau(T)$ .
- 10 Let  $q := (\tau(w_p), \tau(\mathcal{M}_p) \cup \{N \downarrow \tau(\lambda)\})$ . We will show that  $q \in \tau(\mathbb{P}_\kappa)$  and that  $q \leq \tau(p)$ .
- 11 It follows that in  $W$ ,  $\tau(p)$  does not witness that  $N$  is not good.
- 12 Since  $\delta_N = \omega_1^V \in \tau(T)$ , we have that  $N \in \tau(S')$ .
- 13 By definition of  $S'$  and elementarity if  $\tau$ , this means that in  $W$ ,  $\tau(p)$  witnesses that  $N$  is not good. This is a contradiction.

Hence, QED modulo Claim 10.

Hence, QED modulo Claim 10.

### Claim 10

Let  $q := (\tau(w_p), \tau(\mathcal{M}_p) \cup \{N \downarrow \tau(\lambda)\})$ . Then  $q \in \tau(\mathbb{P}_\kappa)$  and  $q \leq \tau(p)$ .



Hence, QED modulo Claim 10.

### Claim 10

Let  $q := (\tau(w_p), \tau(\mathcal{M}_p) \cup \{N \downarrow \tau(\lambda)\})$ . Then  $q \in \tau(\mathbb{P}_\kappa)$  and  $q \leq \tau(p)$ .

*Proof.*

Hence, QED modulo Claim 10.

### Claim 10

Let  $q := (\tau(w_p), \tau(\mathcal{M}_p) \cup \{N \downarrow \tau(\lambda)\})$ . Then  $q \in \tau(\mathbb{P}_\kappa)$  and  $q \leq \tau(p)$ .

*Proof.*

- a The only non-trivial part is that  $q \in \tau(\mathbb{P}_\kappa)$ . Let us assume otherwise.

Hence, QED modulo Claim 10.

### Claim 10

Let  $q := (\tau(w_p), \tau(\mathcal{M}_p) \cup \{N \downarrow \tau(\lambda)\})$ . Then  $q \in \tau(\mathbb{P}_\kappa)$  and  $q \leq \tau(p)$ .

*Proof.*

- a The only non-trivial part is that  $q \in \tau(\mathbb{P}_\kappa)$ . Let us assume otherwise.
- b Then there exists a winning strategy  $\sigma$  for Player I in  $\mathcal{G}_{\tau(\kappa)}^W(q)$ . We will defeat this strategy in  $V[g]$ , reaching a contradiction. (We use in this step that the game  $\mathcal{G}_{\tau(\kappa)}^W(q)$  is closed for Player II.)

- Let Player I play according to  $\sigma$  and let us show that Player II can play by maintaining that for all  $n \in [-1, \omega)$ , there exist  $r, w, P$  such that

- Let Player I play according to  $\sigma$  and let us show that Player II can play by maintaining that for all  $n \in [-1, \omega)$ , there exist  $r, w, P$  such that
- $p_n = (\tau(w_r) \cup w, \tau(\mathcal{M}_r) \cup \{P \downarrow \tau(\lambda)\}),$

- Let Player I play according to  $\sigma$  and let us show that Player II can play by maintaining that for all  $n \in [-1, \omega)$ , there exist  $r, w, P$  such that
- $p_n = (\tau(w_r) \cup w, \tau(\mathcal{M}_r) \cup \{P \downarrow \tau(\lambda)\})$ ,
  - $r \in g \subseteq \mathbb{P}_\lambda$ ,

- Let Player I play according to  $\sigma$  and let us show that Player II can play by maintaining that for all  $n \in [-1, \omega)$ , there exist  $r, w, P$  such that
- $p_n = (\tau(w_r) \cup w, \tau(\mathcal{M}_r) \cup \{P \downarrow \tau(\lambda)\})$ ,
  - $r \in g \subseteq \mathbb{P}_\lambda$ ,
  - there exists finite  $F \subseteq \tau[V_\lambda]$  such that  $P = \text{Hull}^{\tau(\mathcal{H})}(\omega_1^V \cup F)$ ,

- Let Player I play according to  $\sigma$  and let us show that Player II can play by maintaining that for all  $n \in [-1, \omega)$ , there exist  $r, w, P$  such that
- $p_n = (\tau(w_r) \cup w, \tau(\mathcal{M}_r) \cup \{P \downarrow \tau(\lambda)\})$ ,
  - $r \in g \subseteq \mathbb{P}_\lambda$ ,
  - there exists finite  $F \subseteq \tau[V_\lambda]$  such that  $P = \text{Hull}^{\tau(\mathcal{H})}(\omega_1^V \cup F)$ ,
  - $w \subseteq \tau(\phi) \downarrow$  is finite,  $\hat{\mu} \models w$ , and  $w \cap \tau[V] = \emptyset$ .



- c** Let Player I play according to  $\sigma$  and let us show that Player II can play by maintaining that for all  $n \in [-1, \omega)$ , there exist  $r, w, P$  such that
- $p_n = (\tau(w_r) \cup w, \tau(\mathcal{M}_r) \cup \{P \downarrow \tau(\lambda)\})$ ,
  - $r \in g \subseteq \mathbb{P}_\lambda$ ,
  - there exists finite  $F \subseteq \tau[V_\lambda]$  such that  $P = \text{Hull}^{\tau(\mathcal{H})}(\omega_1^V \cup F)$ ,
  - $w \subseteq \tau(\phi) \downarrow$  is finite,  $\hat{\mu} \models w$ , and  $w \cap \tau[V] = \emptyset$ .
- d** The above conditions are satisfied for  $n = -1$  and  $p_{-1} = q$ .  
Let us consider  $n \geq 0$  and the move  $Q_n$  of Player I.

- c** Let Player I play according to  $\sigma$  and let us show that Player II can play by maintaining that for all  $n \in [-1, \omega)$ , there exist  $r, w, P$  such that
- $p_n = (\tau(w_r) \cup w, \tau(\mathcal{M}_r) \cup \{P \downarrow \tau(\lambda)\})$ ,
  - $r \in g \subseteq \mathbb{P}_\lambda$ ,
  - there exists finite  $F \subseteq \tau[V_\lambda]$  such that  $P = \text{Hull}^{\tau(\mathcal{H})}(\omega_1^V \cup F)$ ,
  - $w \subseteq \tau(\phi) \downarrow$  is finite,  $\hat{\mu} \models w$ , and  $w \cap \tau[V] = \emptyset$ .
- d** The above conditions are satisfied for  $n = -1$  and  $p_{-1} = q$ .  
Let us consider  $n \geq 0$  and the move  $Q_n$  of Player I.
- e** Let  $r, w, P$  be as in **c**, w.r.t.  $p_{n-1}$ .

- c** Let Player I play according to  $\sigma$  and let us show that Player II can play by maintaining that for all  $n \in [-1, \omega)$ , there exist  $r, w, P$  such that
- $p_n = (\tau(w_r) \cup w, \tau(\mathcal{M}_r) \cup \{P \downarrow \tau(\lambda)\})$ ,
  - $r \in g \subseteq \mathbb{P}_\lambda$ ,
  - there exists finite  $F \subseteq \tau[V_\lambda]$  such that  $P = \text{Hull}^{\tau(\mathcal{H})}(\omega_1^V \cup F)$ ,
  - $w \subseteq \tau(\phi) \downarrow$  is finite,  $\hat{\mu} \models w$ , and  $w \cap \tau[V] = \emptyset$ .
- d** The above conditions are satisfied for  $n = -1$  and  $p_{-1} = q$ .  
Let us consider  $n \geq 0$  and the move  $Q_n$  of Player I.
- e** Let  $r, w, P$  be as in **c**, w.r.t.  $p_{n-1}$ .
- f** We will distinguish four possibilities for  $Q_n$ . In each, we show that Player II can make the next move and preserve conditions from **c**.

■ Case I.  $Q_n = \psi \in w_{p_{n-1}}$  where  $\psi \equiv \bigvee_{i \in I} \psi_i$ .

**g** **Case I.**  $Q_n = \psi \in w_{p_{n-1}}$  where  $\psi \equiv \bigvee_{i \in I} \psi_i$ .

*Proof.* We have that  $\hat{\mu} \models \psi$ , so there exists  $i \in I$  such that  $\hat{\mu} \models \psi_i$ . We set  $p_n := (w_{p_{n-1}} \cup \{\psi_i\}, \mathcal{M}_{p_{n-1}})$ . □

**g** **Case I.**  $Q_n = \psi \in w_{p_{n-1}}$  where  $\psi \equiv \bigvee_{i \in I} \psi_i$ .

*Proof.* We have that  $\hat{\mu} \models \psi$ , so there exists  $i \in I$  such that  $\hat{\mu} \models \psi_i$ . We set  $p_n := (w_{p_{n-1}} \cup \{\psi_i\}, \mathcal{M}_{p_{n-1}})$ . □

**h** **Case II.**  $Q_n = (\psi, i) \in w_{p_{n-1}} \times I$  where  $\psi \equiv \bigwedge_{i \in I} \psi_i$ .

**g** **Case I.**  $Q_n = \psi \in w_{p_{n-1}}$  where  $\psi \equiv \bigvee_{i \in I} \psi_i$ .

*Proof.* We have that  $\hat{\mu} \models \psi$ , so there exists  $i \in I$  such that  $\hat{\mu} \models \psi_i$ . We set  $p_n := (w_{p_{n-1}} \cup \{\psi_i\}, \mathcal{M}_{p_{n-1}})$ . □

**h** **Case II.**  $Q_n = (\psi, i) \in w_{p_{n-1}} \times I$  where  $\psi \equiv \bigwedge_{i \in I} \psi_i$ .

*Proof.* We can take  $p_n := (w_{p_{n-1}} \cup \{\psi_i\}, \mathcal{M}_{p_{n-1}})$ . □

**i Case III.**  $Q_n = (P \downarrow_{\tau(\lambda)}, D)$  where  $D \in P \downarrow_{\tau(\lambda)}$  is dense  $\mathbb{P}_{\tau(\lambda)}$ .



**i** **Case III.**  $Q_n = (P \downarrow \tau(\lambda), D)$  where  $D \in P \downarrow \tau(\lambda)$  is dense  $\mathbb{P}_{\tau(\lambda)}$ .

*Proof.*

**i** **Case III.**  $Q_n = (P \downarrow \tau(\lambda), D)$  where  $D \in P \downarrow \tau(\lambda)$  is dense in  $\mathbb{P}_{\tau(\lambda)}$ .

*Proof.*

**i** Let  $\rho : \text{Hull}(P, \tau(V_\lambda)) \rightarrow \widehat{P \downarrow \tau(\lambda)}$  be the transitive collapse and let  $D^+ := \rho^{-1}(D) \in P$ . Then  $D^+$  is dense in  $\tau(\mathbb{P}_\kappa)$ .

**i** **Case III.**  $Q_n = (P \downarrow \tau(\lambda), D)$  where  $D \in P \downarrow \tau(\lambda)$  is dense in  $\mathbb{P}_{\tau(\lambda)}$ .

*Proof.*

- i** Let  $\rho : \text{Hull}(P, \tau(V_\lambda)) \rightarrow \widehat{P \downarrow \tau(\lambda)}$  be the transitive collapse and let  $D^+ := \rho^{-1}(D) \in P$ . Then  $D^+$  is dense in  $\tau(\mathbb{P}_\kappa)$ .
- ii** Since  $D^+ \in P$ , there exists  $\bar{D}^+ \in V$  such that  $\tau(\bar{D}^+) = D^+$ .

**i** **Case III.**  $Q_n = (P \downarrow \tau(\lambda), D)$  where  $D \in P \downarrow \tau(\lambda)$  is dense in  $\mathbb{P}_{\tau(\lambda)}$ .

*Proof.*

- i** Let  $\rho : \text{Hull}(P, \tau(V_\lambda)) \rightarrow \widehat{P \downarrow \tau(\lambda)}$  be the transitive collapse and let  $D^+ := \rho^{-1}(D) \in P$ . Then  $D^+$  is dense in  $\tau(\mathbb{P}_\kappa)$ .
- ii** Since  $D^+ \in P$ , there exists  $\bar{D}^+ \in V$  such that  $\tau(\bar{D}^+) = D^+$ .
- iii** Let  $\bar{D} := \bar{D}^+ \cap V_\lambda$ . By elementarity, we have that  $\tau(\bar{D}) = D$  and  $\bar{D}$  is dense in  $\mathbb{P}_\lambda$ .

**i** **Case III.**  $Q_n = (P \downarrow \tau(\lambda), D)$  where  $D \in P \downarrow \tau(\lambda)$  is dense in  $\mathbb{P}_{\tau(\lambda)}$ .

*Proof.*

- i** Let  $\rho : \text{Hull}(P, \tau(V_\lambda)) \rightarrow \widehat{P \downarrow \tau(\lambda)}$  be the transitive collapse and let  $D^+ := \rho^{-1}(D) \in P$ . Then  $D^+$  is dense in  $\tau(\mathbb{P}_\kappa)$ .
- ii** Since  $D^+ \in P$ , there exists  $\bar{D}^+ \in V$  such that  $\tau(\bar{D}^+) = D^+$ .
- iii** Let  $\bar{D} := \bar{D}^+ \cap V_\lambda$ . By elementarity, we have that  $\tau(\bar{D}) = D$  and  $\bar{D}$  is dense in  $\mathbb{P}_\lambda$ .
- iv** There exists  $s \in \bar{D} \cap g$  such that  $s \leq r$ .

**i** **Case III.**  $Q_n = (P \downarrow \tau(\lambda), D)$  where  $D \in P \downarrow \tau(\lambda)$  is dense in  $\mathbb{P}_{\tau(\lambda)}$ .

*Proof.*

- i** Let  $\rho : \text{Hull}(P, \tau(V_\lambda)) \rightarrow \widehat{P \downarrow \tau(\lambda)}$  be the transitive collapse and let  $D^+ := \rho^{-1}(D) \in P$ . Then  $D^+$  is dense in  $\tau(\mathbb{P}_\kappa)$ .
- ii** Since  $D^+ \in P$ , there exists  $\bar{D}^+ \in V$  such that  $\tau(\bar{D}^+) = D^+$ .
- iii** Let  $\bar{D} := \bar{D}^+ \cap V_\lambda$ . By elementarity, we have that  $\tau(\bar{D}) = D$  and  $\bar{D}$  is dense in  $\mathbb{P}_\lambda$ .
- iv** There exists  $s \in \bar{D} \cap g$  such that  $s \leq r$ .
- v** We now have that

**i** **Case III.**  $Q_n = (P \downarrow \tau(\lambda), D)$  where  $D \in P \downarrow \tau(\lambda)$  is dense in  $\mathbb{P}_{\tau(\lambda)}$ .

*Proof.*

- i** Let  $\rho : \text{Hull}(P, \tau(V_\lambda)) \rightarrow \widehat{P \downarrow \tau(\lambda)}$  be the transitive collapse and let  $D^+ := \rho^{-1}(D) \in P$ . Then  $D^+$  is dense in  $\tau(\mathbb{P}_\kappa)$ .
- ii** Since  $D^+ \in P$ , there exists  $\bar{D}^+ \in V$  such that  $\tau(\bar{D}^+) = D^+$ .
- iii** Let  $\bar{D} := \bar{D}^+ \cap V_\lambda$ . By elementarity, we have that  $\tau(\bar{D}) = D$  and  $\bar{D}$  is dense in  $\mathbb{P}_\lambda$ .
- iv** There exists  $s \in \bar{D} \cap g$  such that  $s \leq r$ .
- v** We now have that
  - $\tau(s) \in \tau(\bar{D}) = D,$

- i** **Case III.**  $Q_n = (P \downarrow \tau(\lambda), D)$  where  $D \in P \downarrow \tau(\lambda)$  is dense in  $\mathbb{P}_{\tau(\lambda)}$ .

*Proof.*

- i** Let  $\rho : \text{Hull}(P, \tau(V_\lambda)) \rightarrow \widehat{P \downarrow \tau(\lambda)}$  be the transitive collapse and let  $D^+ := \rho^{-1}(D) \in P$ . Then  $D^+$  is dense in  $\tau(\mathbb{P}_\kappa)$ .
- ii** Since  $D^+ \in P$ , there exists  $\bar{D}^+ \in V$  such that  $\tau(\bar{D}^+) = D^+$ .
- iii** Let  $\bar{D} := \bar{D}^+ \cap V_\lambda$ . By elementarity, we have that  $\tau(\bar{D}) = D$  and  $\bar{D}$  is dense in  $\mathbb{P}_\lambda$ .
- iv** There exists  $s \in \bar{D} \cap g$  such that  $s \leq r$ .
- v** We now have that
  - $\tau(s) \in \tau(\bar{D}) = D$ ,
  - $P' := \text{Hull}^{\tau(\mathcal{H})}(\omega_1^V \cup F \cup \{\tau(s)\})$  satisfies  $\delta(P') = \delta(P)$ ,







**i** **Case IV.**  $Q_n = (M, D)$  where  $M \in \tau(\mathcal{M}_r)$  and  $D \in M$  is dense  $\mathbb{P}_{\lambda_M}$ .

**i** **Case IV.**  $Q_n = (M, D)$  where  $M \in \tau(\mathcal{M}_r)$  and  $D \in M$  is dense  $\mathbb{P}_{\lambda_M}$ .

*Proof.*

**i** **Case IV.**  $Q_n = (M, D)$  where  $M \in \tau(\mathcal{M}_r)$  and  $D \in M$  is dense  $\mathbb{P}_{\lambda_M}$ .

*Proof.*

**i** We have that  $D \in P$ , so there exists  $\bar{D} \in V$  such that  $\tau(\bar{D}) = D$ .

**i** **Case IV.**  $Q_n = (M, D)$  where  $M \in \tau(\mathcal{M}_r)$  and  $D \in M$  is dense  $\mathbb{P}_{\lambda_M}$ .

*Proof.*

- i** We have that  $D \in P$ , so there exists  $\bar{D} \in V$  such that  $\tau(\bar{D}) = D$ .
- ii** Let  $\bar{M} := \tau^{-1}(M) \in \mathcal{M}_r$ . By elementarity of  $\tau$ , we have that  $\bar{D} \in \bar{M} \in \mathcal{M}_r$  and that  $\bar{D}$  is dense in  $\mathbb{P}_{\lambda_{\bar{M}}}$ .

**i** **Case IV.**  $Q_n = (M, D)$  where  $M \in \tau(\mathcal{M}_r)$  and  $D \in M$  is dense  $\mathbb{P}_{\lambda_M}$ .

*Proof.*

- i** We have that  $D \in P$ , so there exists  $\bar{D} \in V$  such that  $\tau(\bar{D}) = D$ .
- ii** Let  $\bar{M} := \tau^{-1}(M) \in \mathcal{M}_r$ . By elementarity of  $\tau$ , we have that  $\bar{D} \in \bar{M} \in \mathcal{M}_r$  and that  $\bar{D}$  is dense in  $\mathbb{P}_{\lambda_{\bar{M}}}$ .
- iii** Let  $\tilde{D}$  consist of all  $s \in \mathbb{P}_\lambda$  satisfying that  $s \leq r$  and that there exists  $t \in \bar{D}$  such that  $s \leq t$  and  $\delta(\text{Hull}(\bar{M}, t)) = \delta(\bar{M})$ . We want to show that  $\tilde{D}$  is dense below  $r$ .

**i** **Case IV.**  $Q_n = (M, D)$  where  $M \in \tau(\mathcal{M}_r)$  and  $D \in M$  is dense  $\mathbb{P}_{\lambda_M}$ .

*Proof.*

- i** We have that  $D \in P$ , so there exists  $\bar{D} \in V$  such that  $\tau(\bar{D}) = D$ .
- ii** Let  $\bar{M} := \tau^{-1}(M) \in \mathcal{M}_r$ . By elementarity of  $\tau$ , we have that  $\bar{D} \in \bar{M} \in \mathcal{M}_r$  and that  $\bar{D}$  is dense in  $\mathbb{P}_{\lambda_{\bar{M}}}$ .
- iii** Let  $\tilde{D}$  consist of all  $s \in \mathbb{P}_\lambda$  satisfying that  $s \leq r$  and that there exists  $t \in \bar{D}$  such that  $s \leq t$  and  $\delta(\text{Hull}(\bar{M}, t)) = \delta(\bar{M})$ . We want to show that  $\tilde{D}$  is dense below  $r$ .
- iv** Let  $r' \in \mathbb{P}_\lambda$  be such that  $r' \leq r$ . Then there exists  $\bar{M}' \in \mathcal{M}_{r'}$  such that  $\delta(\bar{M}') = \delta(\bar{M})$ ,  $\lambda_{\bar{M}'} = \lambda_{\bar{M}}$ , and  $\bar{M} \subseteq \bar{M}'$ .



**i** **Case IV.**  $Q_n = (M, D)$  where  $M \in \tau(\mathcal{M}_r)$  and  $D \in M$  is dense  $\mathbb{P}_{\lambda_M}$ .

*Proof.*

- i** We have that  $D \in P$ , so there exists  $\bar{D} \in V$  such that  $\tau(\bar{D}) = D$ .
- ii** Let  $\bar{M} := \tau^{-1}(M) \in \mathcal{M}_r$ . By elementarity of  $\tau$ , we have that  $\bar{D} \in \bar{M} \in \mathcal{M}_r$  and that  $\bar{D}$  is dense in  $\mathbb{P}_{\lambda_{\bar{M}}}$ .
- iii** Let  $\tilde{D}$  consist of all  $s \in \mathbb{P}_\lambda$  satisfying that  $s \leq r$  and that there exists  $t \in \bar{D}$  such that  $s \leq t$  and  $\delta(\text{Hull}(\bar{M}, t)) = \delta(\bar{M})$ . We want to show that  $\tilde{D}$  is dense below  $r$ .
- iv** Let  $r' \in \mathbb{P}_\lambda$  be such that  $r' \leq r$ . Then there exists  $\bar{M}' \in \mathcal{M}_{r'}$  such that  $\delta(\bar{M}') = \delta(\bar{M})$ ,  $\lambda_{\bar{M}'} = \lambda_{\bar{M}}$ , and  $\bar{M} \subseteq \bar{M}'$ .
- v** By the existence of the appropriate strategy for  $\mathcal{G}_\lambda(r')$ , there exists  $s \in \mathbb{P}_\lambda$  such that  $s \leq r'$  and such that there exists  $t \in \bar{D}$  satisfying  $s \leq t$  and  $\delta(\text{Hull}(\bar{M}', t)) = \delta(\bar{M}')$ .

**i** **Case IV.**  $Q_n = (M, D)$  where  $M \in \tau(\mathcal{M}_r)$  and  $D \in M$  is dense  $\mathbb{P}_{\lambda_M}$ .

*Proof.*

- i** We have that  $D \in P$ , so there exists  $\bar{D} \in V$  such that  $\tau(\bar{D}) = D$ .
- ii** Let  $\bar{M} := \tau^{-1}(M) \in \mathcal{M}_r$ . By elementarity of  $\tau$ , we have that  $\bar{D} \in \bar{M} \in \mathcal{M}_r$  and that  $\bar{D}$  is dense in  $\mathbb{P}_{\lambda_{\bar{M}}}$ .
- iii** Let  $\tilde{D}$  consist of all  $s \in \mathbb{P}_\lambda$  satisfying that  $s \leq r$  and that there exists  $t \in \bar{D}$  such that  $s \leq t$  and  $\delta(\text{Hull}(\bar{M}, t)) = \delta(\bar{M})$ . We want to show that  $\tilde{D}$  is dense below  $r$ .
- iv** Let  $r' \in \mathbb{P}_\lambda$  be such that  $r' \leq r$ . Then there exists  $\bar{M}' \in \mathcal{M}_{r'}$  such that  $\delta(\bar{M}') = \delta(\bar{M})$ ,  $\lambda_{\bar{M}'} = \lambda_{\bar{M}}$ , and  $\bar{M} \subseteq \bar{M}'$ .
- v** By the existence of the appropriate strategy for  $\mathcal{G}_\lambda(r')$ , there exists  $s \in \mathbb{P}_\lambda$  such that  $s \leq r'$  and such that there exists  $t \in \bar{D}$  satisfying  $s \leq t$  and  $\delta(\text{Hull}(\bar{M}', t)) = \delta(\bar{M}')$ .

vi We have that

$$\delta(\bar{M}) \leq \delta(\text{Hull}(\bar{M}, t)) \leq \delta(\text{Hull}(\bar{M}', t)) \leq \delta(\bar{M}') = \delta(\bar{M}),$$

which shows that all of the above values are equal and consequently,  $s \leq r'$  belongs to  $\tilde{D}$ .

vi We have that

$$\delta(\bar{M}) \leq \delta(\text{Hull}(\bar{M}, t)) \leq \delta(\text{Hull}(\bar{M}', t)) \leq \delta(\bar{M}') = \delta(\bar{M}),$$

which shows that all of the above values are equal and consequently,  $s \leq r'$  belongs to  $\tilde{D}$ .

vii Since we have shown that  $\tilde{D}$  is dense, we get that  $g \cap \tilde{D} \neq \emptyset$ . In other words, there exist  $s \in g$  and  $t \in \tilde{D}$  such that  $s \leq r, t$  and  $\delta(\text{Hull}(\bar{M}, t)) = \delta(\bar{M})$ .

vi We have that

$$\delta(\bar{M}) \leq \delta(\text{Hull}(\bar{M}, t)) \leq \delta(\text{Hull}(\bar{M}', t)) \leq \delta(\bar{M}') = \delta(\bar{M}),$$

which shows that all of the above values are equal and consequently,  $s \leq r'$  belongs to  $\tilde{D}$ .

vii Since we have shown that  $\tilde{D}$  is dense, we get that  $g \cap \tilde{D} \neq \emptyset$ . In other words, there exist  $s \in g$  and  $t \in \tilde{D}$  such that  $s \leq r, t$  and  $\delta(\text{Hull}(\bar{M}, t)) = \delta(\bar{M})$ .

viii Let  $P' := \text{Hull}^{\tau(\mathcal{H})}(\omega_1^V \cup F \cup \{\tau(s)\})$  and let

$$p_n := (\tau(w_s) \cup w, \tau(\mathcal{M}_s) \cup \{P' \downarrow \tau(\lambda)\}) \in \tau(\mathbb{P}_\lambda^*).$$

Note that  $\delta_{P'} = \omega_1^V$ .

vi We have that

$$\delta(\bar{M}) \leq \delta(\text{Hull}(\bar{M}, t)) \leq \delta(\text{Hull}(\bar{M}', t)) \leq \delta(\bar{M}') = \delta(\bar{M}),$$

which shows that all of the above values are equal and consequently,  $s \leq r'$  belongs to  $\tilde{D}$ .

vii Since we have shown that  $\tilde{D}$  is dense, we get that  $g \cap \tilde{D} \neq \emptyset$ . In other words, there exist  $s \in g$  and  $t \in \tilde{D}$  such that  $s \leq r, t$  and  $\delta(\text{Hull}(\bar{M}, t)) = \delta(\bar{M})$ .

viii Let  $P' := \text{Hull}^{\tau(\mathcal{H})}(\omega_1^V \cup F \cup \{\tau(s)\})$  and let

$$p_n := (\tau(w_s) \cup w, \tau(\mathcal{M}_s) \cup \{P' \downarrow \tau(\lambda)\}) \in \tau(\mathbb{P}_\lambda^*).$$

Note that  $\delta_{P'} = \omega_1^V$ .

ix We have  $\tau(t) \in D$  and  $p_n \leq \tau(t)$ , while

$$\begin{aligned} \delta(\text{Hull}(M, \tau(t))) &= \delta(\tau(\text{Hull}(\bar{M}, t))) = \delta(\text{Hull}(\bar{M}, t)) = \\ &= \delta(\bar{M}) = \delta(M). \end{aligned}$$

vi We have that

$$\delta(\bar{M}) \leq \delta(\text{Hull}(\bar{M}, t)) \leq \delta(\text{Hull}(\bar{M}', t)) \leq \delta(\bar{M}') = \delta(\bar{M}),$$

which shows that all of the above values are equal and consequently,  $s \leq r'$  belongs to  $\tilde{D}$ .

vii Since we have shown that  $\tilde{D}$  is dense, we get that  $g \cap \tilde{D} \neq \emptyset$ . In other words, there exist  $s \in g$  and  $t \in \tilde{D}$  such that  $s \leq r, t$  and  $\delta(\text{Hull}(\bar{M}, t)) = \delta(\bar{M})$ .

viii Let  $P' := \text{Hull}^{\tau(\mathcal{H})}(\omega_1^V \cup F \cup \{\tau(s)\})$  and let

$$p_n := (\tau(w_s) \cup w, \tau(\mathcal{M}_s) \cup \{P' \downarrow \tau(\lambda)\}) \in \tau(\mathbb{P}_\lambda^*).$$

Note that  $\delta_{P'} = \omega_1^V$ .

ix We have  $\tau(t) \in D$  and  $p_n \leq \tau(t)$ , while

$$\begin{aligned} \delta(\text{Hull}(M, \tau(t))) &= \delta(\tau(\text{Hull}(\bar{M}, t))) = \delta(\text{Hull}(\bar{M}, t)) = \\ &= \delta(\bar{M}) = \delta(M). \end{aligned}$$

QED (Case IV)

- k We have shown that Player II can survive  $\omega$  moves against strategy  $\sigma$ , which is a contradiction.



- k We have shown that Player II can survive  $\omega$  moves against strategy  $\sigma$ , which is a contradiction.
- l This concludes the proof of Claim 10. □

- k** We have shown that Player II can survive  $\omega$  moves against strategy  $\sigma$ , which is a contradiction.
- l** This concludes the proof of Claim 10. □

Since we have shown Claim 10, we conclude the proof of the main theorem.

THANK YOU FOR YOUR ATTENTION!