


Pacific Herring preliminary data summary for West Coast of Vancouver Island 2018

Jaclyn Cleary*

Matthew Grinnell†

July 31, 2018



Pacific Herring (*Clupea pallasii*). Image credit: [Fisheries and Oceans Canada](#).

Disclaimer This report contains preliminary data collected for Pacific Herring in 2018 in the West Coast of Vancouver Island major stock assessment area (SAR). These data may differ from data used and presented in the final stock assessment.

1 Context

Pacific Herring (*Clupea pallasii*) in British Columbia are assessed as 5 major and 2 minor stock assessment regions (SARs), and data are collected and summarized on this scale (Table 1, Figure 1). The Pacific Herring data collection program includes fishery-dependent and -independent data from 1951 to 2018. This includes annual time series of commercial catch data, biological samples (providing information on proportion-at-age and weight-at-age), and spawn index data conducted using a combination of surface and SCUBA surveys. In some areas, industry- and/or First Nations-operated in-season soundings programs are also conducted, and this information is used by resource managers, First Nations, and stakeholders to locate fish and identify areas of high and low Pacific Herring biomass to plan harvesting activities. In-season acoustic soundings are not used by stock assessment to inform the estimation of spawning biomass.

*DFO Science, Pacific Biological Station (email: Jaclyn.Cleary@dfo-mpo.gc.ca).

†DFO Science, Pacific Biological Station (email: Matthew.Grinnell@dfo-mpo.gc.ca).

The following is a description of data collected for Pacific Herring in 2018 in the West Coast of Vancouver Island major SAR (Figure 2). Data collected outside the SAR boundary are not included in this summary, and are not used for the purposes of stock assessment. Although we summarise data at the scale of the SAR for stock assessments, we summarise data at finer spatial scales in this report: locations are nested within sections, sections are nested within statistical areas, and statistical areas are nested within SARs (Table 2). Note that we refer to ‘year’ instead of ‘herring season’ in this report; therefore 2018 refers to the 2017/2018 Pacific Herring season.

2 Data collection programs

Biological samples were collected by the seine charter vessel “Proud Canadian” for 21 days from February 20th to March 12th. The primary purpose of the test charter vessel was to collect biological samples from main bodies of herring from Areas 23, 24, and 25. Nearshore herring samples were collected by the Nuuchahnulth staff in Areas 23, 24, and 25 as part of a pilot sampling program and on-going collaboration between WCVI First Nations and DFO. These nearshore biological samples were collected from spawning aggregations using cast nets. WCVI First Nations spawn reconnaissance charters reported spawn activity in all 3 Areas, and conducted spawn surveys using surface survey methods in Hesquiat Harbour (Area 24).

Herring spawn locations were primarily identified with fixed-wing overflights conducted by DFO Resource Management Area staff. Ten flights were conducted this season from February to April. Three dive charter vessels operated on the WCVI:

- The “Double Decker” surveyed 18 days from March 6th to March 23rd, mainly in Areas 24 and 25,
- The “Seaveyor” surveyed 11 days from March 2nd to March 27th, and
- The “Viking Spirit” surveyed 4 days from March 20th to March 24th.

The seine test charter vessel and the dive survey vessels were funded by DFO, through a contract to the Herring Conservation Research Society.

3 Catch and biological samples

In the 1950s and 1960s, the reduction fishery dominated Pacific Herring catch; starting in the 1970s, catch has been predominantly from roe seine and gillnet fisheries. The reduction fishery is different from current fisheries in several ways. First, the reduction fishery caught Pacific Herring of all ages, whereas current fisheries target spawning (i.e., mature) fish. Thus, reduction fisheries included age-1 fish which are not typically caught in current fisheries. Second, the reduction fishery has some uncertainty regarding the quantity and location of catch; in some cases this may affect our ability to allocate catch to a specific SAR. For the roe gillnet fishery, all Pacific Herring catch has been validated by a dockside monitoring program since 1998; the catch validation program started in

1999 for the roe seine fishery. Finally, the reduction fishery operated during the winter months, whereas roe fisheries typically target spawning fish between February and April.

Landed commercial catch of Pacific Herring by year and fishery is shown in Table 3 and Figure 3. Total harvested spawn on kelp (SOK) in 2018 in the West Coast of Vancouver Island major SAR is shown in Table 4; we also calculate the estimated spawning biomass associated with SOK harvest. See the [draft spawn index technical report](#) for calculations to convert SOK harvest to spawning biomass.

In 2018, 39 Pacific Herring biological samples were collected and processed for the West Coast of Vancouver Island major SAR (Table 5, Table 6), and a total of 1,369 Pacific Herring were aged in 2018. Differences between biological data collected from two sampling protocols regarding the number-, proportion-, weight-, and length-at-age for Pacific Herring in 2018 in the West Coast of Vancouver Island major SAR are shown in Table 7, Table 8, Table 9, and Table 10, respectively. The nearshore sampling program is a multi-year pilot study (using cast nets), therefore only biological data from the seine samples were used for the purposes of stock assessment. The locations in which the biological samples were collected are presented in Figure 4. Included herein are biological summaries of observed proportion-, number-, weight-, and length-at-age (Figure 5, Table 11, and Figure 6, respectively). Biological summaries only include samples collected using seine nets (commercial and test) due to size-selectivity of other gear types such as gillnet. Only representative biological samples are included, where ‘representative’ indicates whether the Pacific Herring sample in the set accurately reflects the larger Pacific Herring school.

4 Spawn survey data

Pacific Herring spawn surveys were conducted at 37 individual locations in 2018 in the West Coast of Vancouver Island major SAR (Table 12, and Figure 7). A summary of spawn from the last decade (2008 to 2017) is shown in Figure 8. Figure 9 shows spawn start date by decade and Statistical Area. Spawn surveys are conducted to estimate the spawn length, width, number of egg layers, and substrate type, and these data are used to estimate the index of spawning biomass (i.e., the spawn index; Figure 10, Figure 11, and Table 13). We describe the calculations used to estimate the spawn index in the [draft spawn index technical report](#). The ‘spawn index’ represents the raw survey data only, and is not scaled by the spawn survey scaling parameter, q . Therefore, these data do not represent model estimates of spawning biomass, and are considered the minimum observed spawning biomass derived from egg counts. The spawn index has two distinct periods defined by the dominant survey method: surface surveys (1951 to 1987), and dive surveys (1988 to 2018).

Some Pacific Herring Sections contribute more than others to the total spawn index, and the percentage contributed by Section varies yearly (Figure 11b, Figure 12). For example, in 2018, Section 232 contributed the most to the spawn index (33%). As with Sections, some Statistical Areas contribute more than others to the total spawn index (Figure 11c, Figure 13). An animation shows the spawn index by spawn survey location from 1951 to 2018 (Figure 14).

5 General observations

General observations provide context to the data summary report. The following observations were reported by area DFO Resource Management staff, and DFO Science staff:

- Compared to last year, the spatial extent of reported spawn increased.
- In addition, there were multiple spawning events at many locations.
- In Area 23, spawning activity was reported in the usual areas from Salmon Beach to Toquart Bay, and also around the Islands on the West side of Barkley Sound.
- In Area 24, multiple spawns occurred from January through to March with the majority occurring in Hesquiaht Harbour.
- In Area 25, there was a noticeable increase in spawning activity in Nootka Sound, where there was no corresponding activity last year.
- Large schools of fish were sounded in Sydney Inlet early in the season, but the nearest significant spawns occurred in Hesquiaht Harbour.
- Significant numbers of gray whales were observed feeding in Hesquiaht Harbour post-spawn, and also quite a few seen in northeast Barkley Sound.

6 Tables

Table 1. Pacific Herring stock assessment regions (SARs) in British Columbia.

Name	Code	Type
Haida Gwaii	HG	Major
Prince Rupert District	PRD	Major
Central Coast	CC	Major
Strait of Georgia	SoG	Major
West Coast of Vancouver Island	WCVI	Major
Area 27	A27	Minor
Area 2 West	A2W	Minor

Table 2. Statistical Areas and Sections for Pacific Herring in the West Coast of Vancouver Island major stock assessment region (SAR).

Region	Statistical Area	Section
West Coast of Vancouver Island	23	230
West Coast of Vancouver Island	23	231
West Coast of Vancouver Island	23	232
West Coast of Vancouver Island	23	233
West Coast of Vancouver Island	23	239
West Coast of Vancouver Island	24	240
West Coast of Vancouver Island	24	241
West Coast of Vancouver Island	24	242
West Coast of Vancouver Island	24	243
West Coast of Vancouver Island	24	244
West Coast of Vancouver Island	24	245
West Coast of Vancouver Island	24	249
West Coast of Vancouver Island	25	250
West Coast of Vancouver Island	25	251
West Coast of Vancouver Island	25	252
West Coast of Vancouver Island	25	253
West Coast of Vancouver Island	25	259

Table 3. Total landed commercial catch of Pacific Herring in metric tonnes (t) by gear type in 2018 in the West Coast of Vancouver Island major stock assessment region (SAR). Legend: ‘Other’ represents the reduction, the food and bait, as well as the special use fishery; ‘RoeSN’ represents the roe seine fishery; and ‘RoeGN’ represents the roe gillnet fishery. Data from the spawn-on-kelp (SOK) fishery is not included. Note: ‘WP’ indicates that data are withheld due to privacy concerns.

Gear	Catch (t)
Other	0
RoeSN	0
RoeGN	0

Table 4. Total harvested Pacific Herring spawn on kelp (SOK) in pounds (lb), and the associated estimate of spawning biomass in metric tonnes (t) from 2008 to 2018 in the West Coast of Vancouver Island major stock assessment region (SAR). See the [draft spawn index technical report](#) for calculations to convert SOK harvest to spawning biomass. Note: ‘WP’ indicates that data are withheld due to privacy concerns.

Year	Harvest (lb)	Spawning biomass (t)
2008	0	0
2009	0	0
2010	0	0
2011	0	0
2012	0	0
2013	0	0
2014	0	0
2015	0	0
2016	0	0
2017	0	0
2018	0	0

Table 5. Number of Pacific Herring biological samples processed from 2008 to 2018 in the West Coast of Vancouver Island major stock assessment region (SAR). Each sample is approximately 100 fish.

Year	Number of samples		
	Commercial	Test	Total
2008	0	22	22
2009	0	29	29
2010	0	27	27
2011	0	28	28
2012	0	10	10
2013	0	5	5
2014	0	4	4
2015	0	20	20
2016	0	25	25
2017	0	19	19
2018	0	39	39

Table 6. Number and type of Pacific Herring biological samples processed in 2018 in the West Coast of Vancouver Island major stock assessment region (SAR). Each sample is approximately 100 fish.

Type	Gear	Use	Number of samples
Test	Other	Nearshore	19
Test	Seine	Test fishery	20

Table 7. Observed number-at-age of Pacific Herring by sample type in 2018 in the West Coast of Vancouver Island major stock assessment region (SAR). The age-10 class is a ‘plus group’ which includes fish ages 10 and older. Legend: ‘Nearshore’ refers to samples collected using cast nets as part of a pilot study with WCVI First Nations.

Sample type	Number-at-age								
	2	3	4	5	6	7	8	9	10
Nearshore	145	163	111	1028	162	92	18	0	1
Seine test	73	182	149	775	132	48	7	3	0
Total	218	345	260	1803	294	140	25	3	1

Table 8. Observed proportion-at-age of Pacific Herring by sample type in 2018 in the West Coast of Vancouver Island major stock assessment region (SAR). The age-10 class is a ‘plus group’ which includes fish ages 10 and older. Legend: ‘Nearshore’ refers to samples collected using cast nets as part of a pilot study with WCVI First Nations.

Sample type	Proportion-at-age								
	2	3	4	5	6	7	8	9	10
Nearshore	0.084	0.095	0.065	0.598	0.094	0.053	0.010	0.000	0.001
Seine test	0.053	0.133	0.109	0.566	0.096	0.035	0.005	0.002	0.000
Total	0.071	0.112	0.084	0.584	0.095	0.045	0.008	0.001	0.000

Table 9. Observed mean weight-at-age in grams (g) of Pacific Herring by sample type in 2018 in the West Coast of Vancouver Island major stock assessment region (SAR). The age-10 class is a ‘plus group’ which includes fish ages 10 and older. Legend: ‘Nearshore’ refers to samples collected using cast nets as part of a pilot study with WCVI First Nations.

Sample type	Mean weight-at-age (g)								
	2	3	4	5	6	7	8	9	10
Nearshore	54	78	85	96	104	109	112	NA	114
Seine test	63	87	103	116	120	129	134	144	NA
Total	57	82	95	105	111	116	118	144	114

Table 10. Observed mean length-at-age in millimetres (mm) of Pacific Herring by sample type in 2018 in the West Coast of Vancouver Island major stock assessment region (SAR). The age-10 class is a ‘plus group’ which includes fish ages 10 and older. Legend: ‘Nearshore’ refers to samples collected using cast nets as part of a pilot study with WCVI First Nations.

Sample type	Mean length-at-age (mm)								
	2	3	4	5	6	7	8	9	10
Nearshore	160	180	189	197	202	203	211	NA	220
Seine test	165	184	193	200	202	208	208	213	NA
Total	162	182	191	198	202	205	210	213	220

Table 11. Observed proportion-at-age for Pacific Herring from 2008 to 2018 in the West Coast of Vancouver Island major stock assessment region (SAR). The age-10 class is a ‘plus group’ which includes fish ages 10 and older.

Year	Proportion-at-age								
	2	3	4	5	6	7	8	9	10
2008	0.052	0.246	0.540	0.115	0.031	0.008	0.006	0.001	0.001
2009	0.024	0.664	0.157	0.127	0.023	0.006	0.000	0.000	0.000
2010	0.176	0.295	0.424	0.052	0.045	0.007	0.001	0.000	0.000
2011	0.048	0.630	0.211	0.096	0.009	0.005	0.000	0.000	0.000
2012	0.021	0.186	0.621	0.103	0.059	0.007	0.003	0.000	0.000
2013	0.034	0.241	0.150	0.475	0.050	0.045	0.005	0.000	0.000
2014	0.029	0.752	0.147	0.029	0.032	0.007	0.000	0.004	0.000
2015	0.140	0.238	0.505	0.068	0.017	0.028	0.004	0.000	0.001
2016	0.040	0.648	0.168	0.124	0.014	0.003	0.004	0.001	0.000
2017	0.025	0.076	0.664	0.132	0.082	0.016	0.002	0.002	0.000
2018	0.053	0.133	0.109	0.566	0.096	0.035	0.005	0.002	0.000

Table 12. Pacific Herring spawn survey locations, start date, and spawn index in metric tonnes (t) in 2018 in the West Coast of Vancouver Island major stock assessment region (SAR). The ‘spawn index’ represents the raw survey data only, and is not scaled by the spawn survey scaling parameter, q . Missing spawn index values (NAs) indicate incomplete spawn surveys.

Statistical Area	Section	Location name	Start date	Spawn index (t)
23	232	David Is	February 26	453
23	232	Forbes Is	February 26	235
23	232	Larkin Is	March 16	16
23	232	Macoah Pass	February 26	2,135
23	232	Maggie Rvr	March 16	1,056
23	232	Newcombe Chnl	March 16	69
23	232	Ottaway Islet	March 16	289
23	232	Spilling Islet	March 16	506
23	232	Stopper Is	March 16	23
23	232	Toquart Bay	March 16	728
23	232	Two Rivers +	February 26	3,792
24	242	Antons Spit	January 10	4,871
24	242	Hesquiat Hrbr	March 16	250
24	242	Hesquiat Hrbr Hd	March 10	NA
24	242	Hesquiat Pen	March 16	271
24	242	Hesquiat Pt	March 20	NA
24	242	Leclaire Pt	March 10	NA
24	242	Matlahaw Pt	March 19	NA
24	242	Rondeault Pt	March 10	NA
24	244	McNeil Pen	March 19	53
25	252	Boca del Infierno Bay	March 16	61
25	252	Cook Chnl	March 16	448
25	252	Friendly Cv	March 16	1,365
25	252	McKay Psg	March 12	720
25	252	Saavedra Is	March 16	1,507
25	252	Santa Gertrudis Cv	March 16	665
25	252	Strange Is	March 05	726
25	253	Garden Pt	March 15	101
25	253	Inner Nuchatlitz	March 05	1,339
25	253	Louie Bay	March 05	1
25	253	Nuchatlitz Inlt	March 12	272
25	253	Outer Nuchatlitz	March 05	2,666
25	253	Owossitsa Cr	March 15	332
25	253	Port Langford	March 05	2,106
25	253	Rosa Hrbr	March 16	327
25	253	Rosa Is	March 16	175
25	253	Unknown Sec 253	March 05	550

Table 13. Summary of Pacific Herring spawn survey data from 2008 to 2018 in the West Coast of Vancouver Island major stock assessment region (SAR). The spawn index has two distinct periods defined by the dominant survey method: surface surveys (1951 to 1987), and dive surveys (1988 to 2018). The ‘spawn index’ represents the raw survey data only, and is not scaled by the spawn survey scaling parameter, q . Units: metres (m), and metric tonnes (t).

Year	Total length (m)	Mean width (m)	Mean number of egg layers	Spawn index (t)
2008	21,300	95	0.8	2,739
2009	32,340	152	1.0	10,607
2010	17,055	65	2.1	2,464
2011	54,735	65	2.3	9,663
2012	14,953	221	2.3	5,407
2013	33,775	103	1.1	12,258
2014	35,825	144	1.1	13,937
2015	20,450	276	1.1	11,323
2016	60,575	122	1.4	20,528
2017	44,200	181	1.3	15,734
2018	61,825	137	1.5	28,107

7 Figures




Figure 1. Boundaries for the Pacific Herring stock assessment regions (SARs) in British Columbia: there are 5 major SARs (HG, PRD, CC, SoG, and WCVI), and 2 minor SARs (A27 and A2W; Table 1). Units: kilometres (km).




Figure 2. Boundaries for the West Coast of Vancouver Island major stock assessment region (SAR; thick dashed lines), associated Statistical Areas (SA; thin solid lines), and associated Sections (thin dotted lines). Units: kilometres (km).




Figure 3. Time series of total landed catch in thousands of metric tonnes ($t \times 10^3$) of Pacific Herring by gear type from 1951 to 2018 in the West Coast of Vancouver Island major stock assessment region (SAR). Legend: ‘Other’ represents the reduction, the food and bait, as well as the special use fishery; ‘RoeSN’ represents the roe seine fishery; and ‘RoeGN’ represents the roe gillnet fishery. Data from the spawn-on-kelp (SOK) fishery is not included.




Figure 4. Location and type of Pacific Herring biological samples collected in 2018 in the West Coast of Vancouver Island major stock assessment region (SAR; thick dashed lines), and associated Sections (Sec; thin solid lines). Units: kilometres (km).




Figure 5. Time series of observed proportion-at-age (a, b) and number aged in thousands (c) of Pacific Herring from 1951 to 2018 in the West Coast of Vancouver Island major stock assessment region (SAR). The black line is the mean age, and the shaded area is the approximate 90% distribution. Biological summaries only include samples collected using seine nets (commercial and test) due to size-selectivity of other gear types such as gillnet. The age-10 class is a ‘plus group’ which includes fish ages 10 and older.




Figure 6. Time series of weight-at-age in grams (g; panel a) and length-at-age in millimetres (mm; panel b) for age-3 (circles) and 5-year running mean weight- and length-at-age (lines) for Pacific Herring from 1951 to 2018 in the West Coast of Vancouver Island major stock assessment region (SAR). Missing weight- and length-at-age values (i.e., years with no biological samples) are imputed using one of two methods: missing values at the beginning of the time series are imputed by extending the first non-missing value backwards; other missing values are imputed as the mean of the previous 5 years. Biological summaries only include samples collected using seine nets (commercial and test) due to size-selectivity of other gear types such as gillnet. The age-10 class is a 'plus group' which includes fish ages 10 and older.




Figure 7. Pacific Herring spawn survey locations, and spawn index in metric tonnes (t) in 2018 in the West Coast of Vancouver Island major stock assessment region (SAR; thick dashed lines), and associated Sections (Sec; thin solid lines). The ‘spawn index’ represents the raw survey data only, and is not scaled by the spawn survey scaling parameter, q . Missing spawn index values (grey circles) indicate incomplete spawn surveys. Units: kilometres (km).




Figure 8. Pacific Herring spawn survey locations, mean spawn index in metric tonnes (t), and spawn frequency from 2008 to 2017 in the West Coast of Vancouver Island major stock assessment region (SAR; thick dashed lines), and associated Sections (Sec; thin solid lines). The ‘spawn index’ represents the raw survey data only, and is not scaled by the spawn survey scaling parameter, q . Missing spawn index values (grey circles) indicate incomplete spawn surveys. Units: kilometres (km).




Figure 9. Pacific Herring spawn start date by decade and Statistical Area. Grey shaded regions indicate March 1st to 31st. Note that spawn size and intensity varies; therefore the number of spawns is not directly proportional to spawn extent or biomass.




Figure 10. Time series of total spawn length in thousands of metres ($m \times 10^3$; panel a), mean spawn width in metres (b), and mean number of egg layers (c) for Pacific Herring from 1983 to 2018 in the West Coast of Vancouver Island major stock assessment region (SAR). The spawn index has two distinct periods defined by the dominant survey method: surface surveys (1951 to 1987), and dive surveys (1988 to 2018).




Figure 11. Time series of spawn index in thousands of metric tonnes ($t \times 10^3$) for Pacific Herring from 1983 to 2018 in the West Coast of Vancouver Island major stock assessment region (SAR; panel a), as well as percent contributed by Statistical Area (SA), and Section (b, & c, respectively). The spawn index has two distinct periods defined by the dominant survey method: surface surveys (1951 to 1987), and dive surveys (1988 to 2018). The ‘spawn index’ represents the raw survey data only, and is not scaled by the spawn survey scaling parameter, q .




Figure 12. Time series of percent of spawn index by Section for Pacific Herring from 1983 to 2018 in the West Coast of Vancouver Island major stock assessment region (SAR). The year 2018 has a darker bar to facilitate interpretation. The spawn index has two distinct periods defined by the dominant survey method: surface surveys (1951 to 1987), and dive surveys (1988 to 2018). The ‘spawn index’ represents the raw survey data only, and is not scaled by the spawn survey scaling parameter, q .




Figure 13. Time series of percent of spawn index by Statistical Area (SA) and Section for Pacific Herring from 1983 to 2018 in the West Coast of Vancouver Island major stock assessment region (SAR). The spawn index has two distinct periods defined by the dominant survey method: surface surveys (1951 to 1987), and dive surveys (1988 to 2018). The 'spawn index' represents the raw survey data only, and is not scaled by the spawn survey scaling parameter, q .

Figure 14. Animation of Pacific Herring spawn survey locations and spawn index in metric tonnes (t) from 1951 to 2018 in the West Coast of Vancouver Island major stock assessment region (SAR; thick dashed lines), and associated Sections (Sec; thin solid lines). The spawn index has two distinct periods defined by the dominant survey method: surface surveys (1951 to 1987), and dive surveys (1988 to 2018). The ‘spawn index’ represents the raw survey data only, and is not scaled by the spawn survey scaling parameter, q . Missing spawn index values (grey circles) indicate incomplete spawn surveys. The inset shows the total spawn index by year. Units: kilometres (km).