


Pacific Herring preliminary data summary for Prince Rupert District 2026

Jaclyn Cleary*

Matthew Grinnell†

July 20, 2026



Pacific Herring (*Clupea pallasii*). Image credit: [Fisheries and Oceans Canada](#).

Disclaimer This report contains preliminary data collected for Pacific Herring in 2026 in the Prince Rupert District major stock assessment region (SAR). These data may differ from data used and presented in the final stock assessment.

1 Context

Pacific Herring (*Clupea pallasii*) in British Columbia are assessed as 5 major and 2 minor stock assessment regions (SARs), and data are collected and summarized on this scale (Table 1, Figure 1). Note that formal stock assessments are only done for major SARs. The Pacific Herring data collection program includes fishery-dependent and -independent data from 1951 to 2026. This includes annual time series of commercial catch data, biological samples (providing information on proportion-at-age and weight-at-age), and spawn index data conducted using a combination of surface and SCUBA surveys. In some areas, industry- and/or First Nations-operated in-season soundings programs are also conducted, and this information is used by resource managers, First Nations, and

*DFO Science, Pacific Biological Station (email: Jaclyn.Cleary@dfo-mpo.gc.ca).

†DFO Science, Pacific Biological Station (email: Matthew.Grinnell@dfo-mpo.gc.ca).

stakeholders to locate fish and identify areas of high and low Pacific Herring biomass to plan harvesting activities. In-season acoustic soundings are not used by stock assessment to inform the estimation of spawning biomass.

The following is a description of data collected for Pacific Herring in 2026 in the Prince Rupert District major SAR (Figure 2). Data collected outside the SAR boundary are not included in this summary, and are not used for the purposes of stock assessment. Although we summarise data at the scale of the SAR for stock assessments, we summarise data at finer spatial scales in this report: Locations are nested within Sections, Sections are nested within Statistical Areas, and Statistical Areas are nested within SARs (Table 2). Note that we refer to ‘year’ instead of ‘herring season’ in this report; therefore 2026 refers to the 2025/2026 Pacific Herring season.

2 Data collection programs

Biological samples were collected by two seine test vessels:

- The *Nita Maria* collected samples in Big Bay for 13 days from March 13th to March 25th, and
- The *Viking Spirit* collected samples in Kitkatla for 13 days from March 15th to March 27th.

The primary purpose of the test charter vessels was to collect biological samples from main bodies of herring from Big Bay and Kitkatla, identified from soundings.

- The dive charter vessel *Royal Pride* surveyed spawn for 19 days from March 27th to April 14th.

Herring spawn locations were primarily identified with fixed-wing overflights and on-grounds observations from test and dive vessels. Three spawn flights were conducted this season on March 20th, March 31st, and April 7th. Additional on-grounds spawn observations were provided by Lax Kw’alaams Band, Metlakatla, and Gixaala First Nation.

3 Catch and biological samples

In the 1950s and 1960s, the reduction fishery dominated Pacific Herring catch; starting in the 1970s, catch has been predominantly from roe seine and gillnet fisheries. The reduction fishery is different from current fisheries in several ways. First, the reduction fishery caught Pacific Herring of all ages, whereas current fisheries target spawning (i.e., mature) fish. Thus, reduction fisheries included an unknown number of age-1 fish which are not typically caught in current fisheries. Second, the reduction fishery has some uncertainty regarding the quantity and location of catch; however catches have been resolved to SAR and Statistical Area using fish slips as best as possible. For the roe gillnet fishery, all Pacific Herring catch has been validated by a dockside monitoring

program since 1998; the catch validation program started in 1999 for the roe seine fishery. Finally, the reduction fishery operated during the summer and winter months, whereas roe fisheries typically target spawning fish between February and April.

Landed commercial catch of Pacific Herring by year and fishery is shown in Table 3 and Figure 3. In addition to annual catch variability, catch varies among Statistical Areas (Figure 4). Total harvested spawn-on-kelp (SOK) in 2026 in the Prince Rupert District major SAR is shown in Table 4; we also calculate the estimated spawning biomass associated with SOK harvest. See the [spawn index technical report](#) for calculations to convert SOK harvest to spawning biomass.

In 2026, 15 Pacific Herring biological samples were collected and processed for the Prince Rupert District major SAR (Table 5, Table 6). The locations in which the biological samples were collected are presented in Figure 5. Included herein are biological summaries of observed proportion-, number-, weight-, and length-at-age (Figure 6, Table 7, and Figure 7, respectively). We also show the percent change in weight and length for age-3 and age-6 fish (Figure 8 & Figure 9, respectively). Biological summaries only include samples collected using seine nets (commercial and test) due to size-selectivity of other gear types such as gillnet. Only representative biological samples are included, where ‘representative’ indicates whether the Pacific Herring sample in the set accurately reflects the larger Pacific Herring school.

4 Spawn survey data

Pacific Herring spawn surveys were conducted at 29 individual locations in 2026 in the Prince Rupert District major SAR (Table 8, and Figure 10). A summary of spawn from the last decade (2016 to 2025) is shown in Figure 11. Figure 12 shows spawn start date by decade and Statistical Area. Spawn surveys are conducted to estimate the spawn length, width, number of egg layers, and substrate type, and these data are used to estimate the index of spawning biomass (i.e., the spawn index; Figure 13, Figure 14, Figure 15, Figure 17, Table 9, and Figure 16). See the [spawn index technical report](#) for calculations to convert SOK harvest to spawning biomass. The ‘spawn index’ is not scaled by the spawn survey scaling parameter, q . Therefore, these data do not represent model estimates of spawning biomass, and are considered the minimum observed spawning biomass derived from egg counts. The spawn index has two distinct periods defined by the dominant coastwide survey method: surface surveys (1951 to 1987), and dive surveys (1988 to 2026).

Some Pacific Herring Sections contribute more than others to the total spawn index, and the percentage contributed by Section varies yearly (Figure 17b, Figure 18). For example, in 2026, Section 042 contributed the most to the spawn index (56%). As with Sections, some Statistical Areas contribute more than others to the total spawn index (Figure 17c, Figure 19). An animation shows the spawn index by spawn survey location from 1951 to 2026 (Figure 20).

5 First Nations observations

The following observations were contributed by representatives of First Nations communities. These observations provide context and perspectives to this data report. In some cases we make edits for clarity and brevity, but we do not change the intent or substance of responses. Note: if your observations are missing, submit them [here](#).

- No observations were available at the time of distribution.

6 General observations

The following observations were reported by DFO Resource Management staff and DFO Science staff. These observations provide additional context to this data report.

6.1 Areas 3 and 4 (Big Bay)

- Nothing unusual this year in terms of spawning; the season followed a typical pattern. Juvenile component holding and spawning in the North.
- Light layers recorded in the southern and northern portions of the assessment area.
- Average of 4 layers observed around the Tree Bluff area leading into Big Bay along the southern shore.
- Average of 6 layers observed in the North throughout Stumaun Bay and Rush Brook Passage.

6.2 Area 5 (Kitkatla)

- Kitkatla had strong spawn abundance and spawn started later than in previous years.

7 Tables

Table 1. Pacific Herring stock assessment regions (SARs) in British Columbia.

Name	Code	Type
Haida Gwaii	HG	Major
Prince Rupert District	PRD	Major
Central Coast	CC	Major
Strait of Georgia	SoG	Major
West Coast of Vancouver Island	WCVI	Major
Area 27	A27	Minor
Area 2 West	A2W	Minor

Table 2. Statistical Areas and Sections for Pacific Herring in the Prince Rupert District major stock assessment region (SAR).

Region	Statistical Area	Section
Prince Rupert District	03	031
Prince Rupert District	03	032
Prince Rupert District	03	033
Prince Rupert District	04	040
Prince Rupert District	04	041
Prince Rupert District	04	042
Prince Rupert District	04	043
Prince Rupert District	04	049
Prince Rupert District	05	050
Prince Rupert District	05	051
Prince Rupert District	05	052
Prince Rupert District	05	053
Prince Rupert District	05	059

Table 3. Total landed commercial catch of Pacific Herring in metric tonnes (t) by gear type in 2026 in the Prince Rupert District major stock assessment region (SAR). Legend: ‘Other’ represents the reduction (1951 to 1970 only), the food and bait, as well as the special use fishery; ‘RoeSN’ represents the roe seine fishery; and ‘RoeGN’ represents the roe gillnet fishery. Data from the spawn-on-kelp (SOK) fishery are not included. Note: data may be withheld due to privacy concerns (WP).

Gear	Catch (t)
Other	0
RoeSN	0
RoeGN	0

Table 4. Total harvested Pacific Herring spawn-on-kelp (SOK) in pounds (lb), and the associated estimate of spawning biomass in metric tonnes (t) from 2016 to 2026 in the Prince Rupert District major stock assessment region (SAR). See the [spawn index technical report](#) for calculations to convert SOK harvest to spawning biomass. Note: data may be withheld due to privacy concerns (WP).

Year	Harvest (lb)	Spawning biomass (t)
2016	WP	WP
2017	82,597	122
2018	20,832	31
2019	WP	WP
2020	0	0
2021	0	0
2022	0	0
2023	0	0
2024	0	0
2025	0	0
2026	0	0

Table 5. Number of Pacific Herring biological samples processed from 2016 to 2026 in the Prince Rupert District major stock assessment region (SAR). Each sample is approximately 100 fish. Note: Nearshore samples are not used in stock assessments.

Year	Number of samples			
	Commercial	Test	Nearshore	Total
2016	33	11	0	44
2017	40	11	0	51
2018	17	14	0	31
2019	2	13	0	15
2020	0	20	0	20
2021	0	13	0	13
2022	0	10	0	10
2023	9	16	0	25
2024	0	10	0	10
2025	0	19	0	19
2026	0	15	0	15

Table 6. Number and type of Pacific Herring biological samples processed in 2026 in the Prince Rupert District major stock assessment region (SAR). Each sample is approximately 100 fish.

Type	Gear	Use	Number of samples
Test	Other	Test fishery	1
Test	Seine	Test fishery	14

Table 7. Observed proportion-at-age for Pacific Herring from 2016 to 2026 in the Prince Rupert District major stock assessment region (SAR). The age-10 class is a ‘plus group’ which includes fish ages 10 and older.

Year	Proportion-at-age								
	2	3	4	5	6	7	8	9	10
2016	0.017	0.162	0.485	0.060	0.165	0.050	0.039	0.016	0.006
2017	0.014	0.151	0.179	0.438	0.063	0.099	0.029	0.020	0.007
2018	0.048	0.413	0.132	0.158	0.189	0.029	0.017	0.009	0.006
2019	0.014	0.527	0.249	0.069	0.081	0.048	0.008	0.005	0.000
2020	0.004	0.023	0.762	0.154	0.031	0.017	0.007	0.002	0.001
2021	0.005	0.148	0.028	0.660	0.108	0.028	0.015	0.007	0.000
2022	0.007	0.121	0.200	0.052	0.521	0.076	0.015	0.005	0.002
2023	0.003	0.550	0.116	0.103	0.033	0.162	0.026	0.005	0.002
2024	0.021	0.280	0.498	0.066	0.027	0.026	0.069	0.012	0.001
2025	0.004	0.428	0.167	0.290	0.053	0.023	0.011	0.021	0.003
2026	0.005	0.256	0.373	0.125	0.166	0.032	0.017	0.016	0.011

Table 8. Pacific Herring spawn survey locations, start date, and spawn index in metric tonnes (t) in 2026 in the Prince Rupert District major stock assessment region (SAR). The ‘spawn index’ is not scaled by the spawn survey scaling parameter, q . Missing spawn index values indicate incomplete surveys (NAs).

Statistical Area	Section	Location name	Start date	Spawn index (t)
03	033	Birnie Is	March 28	85
03	033	Finlayson Is E	March 28	377
03	033	Flewin Pt	March 28	84
03	033	Grassy Pt	March 28	313
04	042	Belletti Pt	March 28	1,712
04	042	Big Bay	March 28	4,240
04	042	Burnt Cliff Is	March 28	2,734
04	042	Duncan Bay	March 28	2,130
04	042	Finlayson Is W	March 28	183
04	042	Pearl Hrbr	March 28	117
04	042	Reeks Pt	March 28	915
04	042	Swamp Is	March 28	7,896
04	042	Tree Bluff	March 28	12,965
04	043	Fan Is	April 01	8,056
04	043	Hearndon Pt	March 31	174
04	043	Henry Isl	March 31	1,639
04	043	Oval Pt	March 31	7,380
04	043	Welcome Hrb	March 31	20
05	051	Goschen Pt	March 31	846
05	051	Joachim Pt	March 31	680
05	052	Absalom Is	March 20	15
05	052	Cape George	March 31	1,659
05	052	Freeman Pass	March 31	783
05	052	Goschen Is	March 31	1,059
05	052	Joachim Spit	March 31	716
05	052	Nubble Pt	March 31	1,407
05	052	Shakes Is	March 31	61
05	052	Viscount Pt	March 31	189
05	052	Willis Bay	March 31	690

Table 9. Summary of Pacific Herring spawn survey data from 2016 to 2026 in the Prince Rupert District major stock assessment region (SAR). The spawn index has two distinct periods defined by the dominant coastwide survey method: surface surveys (1951 to 1987), and dive surveys (1988 to 2026). The ‘spawn index’ is not scaled by the spawn survey scaling parameter, q . Units: metres (m), and metric tonnes (t).

Year	Total length (m)	Mean width (m)	Mean number of egg layers	Spawn index (t)
2016	48,525	112	0.6	18,984
2017	45,675	174	0.7	19,235
2018	51,000	106	0.5	14,154
2019	87,125	110	0.6	27,189
2020	59,960	205	1.7	25,845
2021	72,250	117	1.3	33,061
2022	64,925	92	1.4	35,219
2023	75,875	160	1.5	42,201
2024	65,400	106	2.1	50,053
2025	83,450	93	1.8	45,416
2026	79,100	105	1.8	59,136

8 Figures




Figure 1. Boundaries for the Pacific Herring stock assessment regions (SARs) in British Columbia. There are 5 major SARs: Haida Gwaii (HG), Prince Rupert District (PRD), Central Coast (CC), Strait of Georgia (SoG), and West Coast of Vancouver Island (WCVI). There are 2 minor SARs: Area 27 (A27) and Area 2 West (A2W). Units: kilometres (km).




Figure 2. Boundaries for the Prince Rupert District major stock assessment region (SAR; thick dashed lines), associated Statistical Areas (SA; thin solid lines), and associated Sections (thin dotted lines). Units: kilometres (km).




Figure 3. Time series of total landed catch in thousands of metric tonnes ($t \times 10^3$) of Pacific Herring by gear type from 1951 to 2026 in the Prince Rupert District major stock assessment region (SAR). Legend: 'Other' represents the reduction (1951 to 1970 only), the food and bait, as well as the special use fishery; 'RoeSN' represents the roe seine fishery; and 'RoeGN' represents the roe gillnet fishery. Data from the spawn-on-kelp (SOK) fishery are not included. Bottom panel shows catch since 1991 in more detail. Note: symbols indicate years in which catch by gear type (i.e., Other, RoeSN, RoeGN) is withheld due to privacy concerns.




Figure 4. Time series of total landed catch in thousands of metric tonnes ($t \times 10^3$) of Pacific Herring by Statistical Area (SA) from 1951 to 2026 in the Prince Rupert District major stock assessment region (SAR). The year 2026 has a darker bar to facilitate interpretation. Note: symbols indicate years in which catch is withheld due to privacy concerns.




Figure 5. Location and type of Pacific Herring biological samples collected in 2026 in the Prince Rupert District major stock assessment region (SAR; thick dashed lines), and associated Sections (Sec; thin solid lines). Units: kilometres (km).




Figure 6. Time series of observed proportion-at-age (a) and number aged in thousands (b) of Pacific Herring from 1951 to 2026 in the Prince Rupert District major stock assessment region (SAR). The black line is the mean age, and the shaded area is the approximate 90% distribution. Biological summaries only include samples collected using seine nets (commercial and test) due to size-selectivity of other gear types such as gillnet. The age-10 class is a ‘plus group’ which includes fish ages 10 and older.




Figure 7. Time series of weight-at-age in grams (g) and length-at-age in millimetres (mm) for age-3 (circles) and 5-year running mean weight- and length-at-age (lines) for Pacific Herring from 1951 to 2026 in the Prince Rupert District major stock assessment region (SAR). Missing weight- and length-at-age values (i.e., years with no biological samples) are imputed using one of two methods: missing values at the beginning of the time series are imputed by extending the first non-missing value backwards; other missing values are imputed as the mean of the previous 5 years. Biological summaries only include samples collected using seine nets (commercial and test) due to size-selectivity of other gear types such as gillnet. The age-10 class is a 'plus group' which includes fish ages 10 and older.




Figure 8. Time series of percent change (%) in weight and length for age-3 fish for Pacific Herring from 1951 to 2026 in the Prince Rupert District major stock assessment region (SAR). Percent change is $\delta_t = \frac{\alpha_t - \alpha_{t-1}}{\alpha_{t-1}}$ where α_t is the weight and length of age-3 fish, respectively, in year t . Biological summaries only include samples collected using seine nets (commercial and test) due to size-selectivity of other gear types such as gillnet.




Figure 9. Time series of percent change (%) in weight and length for age-6 fish for Pacific Herring from 1951 to 2026 in the Prince Rupert District major stock assessment region (SAR). Percent change is $\delta_t = \frac{\alpha_t - \alpha_{t-1}}{\alpha_{t-1}}$ where α_t is the weight and length of age-6 fish, respectively, in year t . Biological summaries only include samples collected using seine nets (commercial and test) due to size-selectivity of other gear types such as gillnet.




Figure 10. Pacific Herring spawn survey locations, and spawn index in metric tonnes (t) in 2026 in the Prince Rupert District major stock assessment region (SAR; thick dashed lines), and associated Sections (Sec; thin solid lines). The ‘spawn index’ is not scaled by the spawn survey scaling parameter, q . Missing spawn index values indicate incomplete surveys (grey circles). Inset tracks the total spawn index. Units: kilometres (km).




Figure 11. Pacific Herring spawn survey locations, mean spawn index in metric tonnes (t), and spawn frequency from 2016 to 2025 in the Prince Rupert District major stock assessment region (SAR; thick dashed lines), and associated Sections (Sec; thin solid lines). The ‘spawn index’ is not scaled by the spawn survey scaling parameter, q . Missing spawn index values indicate incomplete surveys (grey circles). Units: kilometres (km).




Figure 12. Pacific Herring spawn start date by decade and Statistical Area. Grey shaded regions indicate March 1st to 31st. Note that spawn size and intensity varies; therefore the number of spawns is not directly proportional to spawn extent or biomass.




Figure 13. Time series of spawn index in thousands of metric tonnes ($t \times 10^3$) by type for Pacific Herring from 1951 to 2026 in the Prince Rupert District major stock assessment region (SAR). There are three types of spawn survey observations: observations of spawn taken from the surface usually at low tide, underwater observations of spawn on giant kelp, *Macrocyctis* (*Macrocyctis* spp.), and underwater observations of spawn on other types of algae and the substrate, which we refer to as ‘understory.’ The spawn index has two distinct periods defined by the dominant coastwide survey method: surface surveys (1951 to 1987), and dive surveys (1988 to 2026). See Figure 14 for the spawn survey method.




Figure 14. Time series of proportion of spawn index by surface and dive surveys for Pacific Herring from 1951 to 2026 in the Prince Rupert District major stock assessment region (SAR). The spawn index has two distinct periods defined by the dominant coastwide survey method: surface surveys (1951 to 1987), and dive surveys (1988 to 2026).




Figure 15. Time series of total spawn length in thousands of metres ($m \times 10^3$; panel a), mean spawn width in metres (b), and mean number of egg layers (c) for Pacific Herring from 1951 to 2026 in the Prince Rupert District major stock assessment region (SAR). The spawn index has two distinct periods defined by the dominant coastwide survey method: surface surveys (1951 to 1987), and dive surveys (1988 to 2026).




Figure 16. Time series of spawn index in thousands of metric tonnes ($t \times 10^3$) for Pacific Herring from 1951 to 2026 in the Prince Rupert District major stock assessment region (SAR; panel a), and percent change (b). Percent change is $\delta_t = \frac{\alpha_t - \alpha_{t-1}}{\alpha_{t-1}}$ where α_t is the spawn index in year t . The spawn index has two distinct periods defined by the dominant coastwide survey method: surface surveys (1951 to 1987), and dive surveys (1988 to 2026). The ‘spawn index’ is not scaled by the spawn survey scaling parameter, q . Note that spawn surveys in the dive survey period (1988 to 2026) are a combination of surface and dive surveys (Figures 13 and 14).




Figure 17. Time series of spawn index in thousands of metric tonnes ($t \times 10^3$) for Pacific Herring from 1951 to 2026 in the Prince Rupert District major stock assessment region (SAR; panel a), as well as percent contributed by Statistical Area (SA), and Section (b, & c, respectively). See Figure 16 for information.




Figure 18. Time series of percent of spawn index by Section for Pacific Herring from 1951 to 2026 in the Prince Rupert District major stock assessment region (SAR). The year 2026 has a darker bar to facilitate interpretation. The spawn index has two distinct periods defined by the dominant coastwide survey method: surface surveys (1951 to 1987), and dive surveys (1988 to 2026). The ‘spawn index’ is not scaled by the spawn survey scaling parameter, q .




Figure 19. Time series of percent of spawn index by Statistical Area (SA) and Section for Pacific Herring from 1951 to 2026 in the Prince Rupert District major stock assessment region (SAR). The spawn index has two distinct periods defined by the dominant coastwide survey method: surface surveys (1951 to 1987), and dive surveys (1988 to 2026). The ‘spawn index’ is not scaled by the spawn survey scaling parameter, q .




Figure 20. Animation of Pacific Herring spawn index in metric tonnes (t) by Location from 1951 to 2026 in the Prince Rupert District major stock assessment region (SAR; thick dashed lines), and associated Sections (Sec; thin solid lines). The spawn index has two distinct periods defined by the dominant coastwide survey method: surface surveys (1951 to 1987), and dive surveys (1988 to 2026). The ‘spawn index’ is not scaled by the spawn survey scaling parameter, q . Missing spawn index values indicate incomplete surveys (grey circles). Inset tracks the total spawn index. Units: kilometres (km). View the animation: [download the report](#), open with Adobe, enable Java, and click “play”.